

This is a repository copy of *Impossible measurements require impossible apparatus*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/167929/>

Version: Accepted Version

Article:

Bostelmann, Henning orcid.org/0000-0002-0233-2928, Fewster, Chris orcid.org/0000-0001-8915-5321 and Ruep, Maximilian (2021) Impossible measurements require impossible apparatus. Physical Review D. 025017. ISSN 2470-0029

<https://doi.org/10.1103/PhysRevD.103.025017>




Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Impossible measurements require impossible apparatus

Henning Bostelmann ^{*}, Christopher J. Fewster [†] and Maximilian H. Ruep [‡]
Department of Mathematics, University of York, Heslington, York YO10 5DD, United Kingdom
(Dated: October 27, 2020)

A well-recognised open conceptual problem in relativistic quantum field theory concerns the relation between measurement and causality. Naive generalisations of quantum measurement rules can allow for superluminal signalling (‘impossible measurements’). This raises the problem of delineating physically allowed quantum measurements and operations. We analyse this issue in a recently proposed framework in which local measurements (in possibly curved spacetime) are described physically by coupling the system to a probe. We show that the state-update rule in this setting is consistent with causality provided that the coupling between the system and probe is local. Thus, by establishing a well-defined framework for successive measurements, we also provide a class of physically allowed operations. Conversely, impossible measurements can only be performed using impossible (non-local) apparatus.

I. INTRODUCTION

It is a central tenet of special and general relativity that there is a maximal speed of causal influence, the *speed of light*: There can be no superluminal signalling. This should apply, in particular, to relativistic quantum field theory (QFT) and relativistic quantum information (RQI). However, as is very well known, the standard notion of *measurement* challenges this tenet: It has been argued that ‘ideal measurements’ in QFT can yield superluminal signalling [1, 2] and that ‘nondemolition’ measurements of Wilson loops in non-Abelian gauge theory can transfer charge over spacelike distances [3]. Even operations not directly associated to an ideal measurement, such as unitary transformations, can enable superluminal communication [2]. Those measurements that are in conflict with causality are called *impossible measurements* [1] and their existence naturally raises the question of delineating (a) “physically allowed quantum operations” [3], as well as (b) “observables [that] can be measured consistently with causality” [4]. These questions are not just of general conceptual importance [4] but also directly affect applications in RQI [2, 5] due to the lack of a clear-cut criterion for allowed operations that also allows an explicit construction of the latter.

One way to address the difficulties just mentioned is to adopt an operational approach to measurement, in which the *system* of interest is temporarily coupled to a measurement device (*probe*); following a measurement of a probe observable the probe is discarded (traced out). This constitutes a *measurement scheme* [6] for an induced observable of the system and, importantly, yields an associated state-update rule. Although well established in quantum mechanics, this idea was only recently adapted to QFT in possibly curved spacetimes,

thus implementing the concept of a measurement scheme in a local and covariant way [7] (see [8] for a summary). We call this the FV-framework and its elements FV-measurement schemes.

In this paper we show that, due to the locality of the coupling between *system* and *probe*, measurements in the FV-framework are *not* plagued by superluminal signalling in the sense of [1], i.e., an impossible measurement requires an impossible, non-local apparatus. As a result, FV-measurement schemes are consistent with causality and the associated state-updates provide a large and explicitly calculable class of “physically allowed quantum operations”; a significant improvement to the “case-by-case analysis” [4] which has been the approach of all previous literature on this topic to the best of our knowledge. This reinforces the usefulness of the FV-framework for treating measurements in QFT, pointing to its use as a general way to understand which operations are physically allowable, and hence forming useful underpinning for applications in RQI.

The structure of our paper is as follows: In Sec. II we recall the essence of the causality problem posed by Sorkin [1] in the form of a tripartite signalling protocol applicable in flat as well as curved spacetime; this is followed by a non-technical motivation and discussion of FV-measurement schemes and a statement of our main result. Section III comprises the precise presentation of the FV-framework in the language of algebraic quantum field theory and in Sec. IV we show that in the FV-framework, Sorkin’s protocol does not result in any acausal effects. Section V consists of a discussion of causal factorisation and forms the basis of our discussion of multiple observers covered in Sec. VI. In particular, we demonstrate how the state updates associated to selective and non-selective measurements (postulated in Sec. II and IV) can be derived from the principle of causal factorisation. The consequences on causality of this analysis are displayed in Sec. VII, where we explicitly show that the FV-framework consistently describes any finite number of causally orderable measurements without any superluminal signalling issues. As a last

^{*} henning.bostelmann@york.ac.uk

[†] chris.fewster@york.ac.uk

[‡] maximilian.ruep@york.ac.uk

point we conclude and provide an outlook in Sec. VIII.

II. HEURISTIC OVERVIEW




FIG. 1. Schematic spacetime diagram of the relative causal position of the regions O_1 , O_2 and O_3 .

a. Superluminal signalling à la Sorkin Sorkin has argued [1] that the notion of an ideal measurement conflicts with locality and causality when extended from quantum mechanics to QFT. In particular, he presented the following protocol: Let Alice, Bob and Charlie be three experimenters in three laboratories performing actions in the spacetime ‘regions of control’ O_1, O_2, O_3 such that parts of O_1 are in the past of O_2 and parts of O_2 are in the past of O_3 but such that O_1 is spacelike separated from O_3 as shown in Fig 1. Let A be a local observable of O_1 , e.g., an algebraic combination of quantum fields smeared against test functions vanishing outside O_1 . Define B, C similarly and let ρ be the initial state of the quantum field. Sorkin considers the following tripartite procedure. In step one, Alice performs a local measurement of A in her laboratory. In the absence of any post-selection in the experimental data analysis, the resulting updated state is a probabilistic mixture, i.e., a convex combination of states, each selected on a different possible outcome of the measurement, weighted by the respective probabilities. This updated state is denoted ρ_A . In step two, Bob measures B , producing a further (similar) update $\rho_A \mapsto \rho_{AB}$. In step three, Charlie measures observable C in state ρ_{AB} . Since Charlie’s laboratory is spacelike separated from Alice’s, $\text{Tr}(\rho_{AB} C)$ should (in the absence of superluminal communication) give the same result as $\text{Tr}(\rho_B C)$ - the situation where Alice does not measure at all. This condition, Sorkin argues, puts non-trivial constraints on feasible (ideal) measurements, to the extent that “it becomes a priori unclear, for quantum field theory, which observables can be measured consistently with causality and which can’t. This would seem to deprive [QFT] of any definite measurement theory, leaving the issue of what can actually be measured to (at best) a case-by-case analysis” [4]. By contrast, we will show that the FV-framework furnishes QFT with a definite measurement theory.

b. The idea behind the FV-framework A measurement scheme in quantum measurement theory is the theoretical description of a measurement on a *system*, prepared in state ρ_S , by the operational procedure of bringing it into contact with a *probe*, itself to be regarded as a quantum system, and initially prepared in state ρ_P . The ‘contact’ between system and probe is modelled by coupling them together via interactions. In quantum mechanics, this is achieved by an interacting unitary time-evolution which operates for a short period of time and is then removed. A subsequent measurement made on the probe is interpreted as a measurement of the system, and indeed it is possible to establish a correspondence between observables of the probe and induced observables of the system. One says that the combination of the probe, interacting dynamics, and probe observable, form a *measurement scheme* for the induced system observable (see [6] for a comprehensive account).

The FV-framework translates the above idea to QFT in possibly curved spacetime; equally, it can incorporate QFT under the influence of external fields. It is phrased in terms of the algebraic approach to QFT [9] (see [10] for an introduction), but for the purposes of the following discussion we use familiar terminology of QFT; the more formal algebraic version will be set out in Sec. III and used in our proof.

We consider two local relativistic QFTs, modelling the system and the probe. Taking a tensor product, they may be combined as a single theory with no coupling between them. If the two theories are obtained from Lagrangian densities \mathcal{L}_S and \mathcal{L}_P , the uncoupled combination is defined by the sum $\mathcal{L}_S + \mathcal{L}_P$. The contact between system and probe is modelled by another QFT, in which the two are coupled so that the coupling is only effective within a compact set K of spacetime, the *coupling zone*. Crucially, it is assumed that this coupled QFT is itself a local relativistic theory. For Lagrangian theories, the coupled theory would be described by a *local* coupling term such as $\mathcal{L}_I := -\lambda\alpha(x)\phi(x)\psi(x)$, where ϕ and ψ are system and probe Hermitian scalar fields respectively, and the real-valued smooth function α , perhaps representing an external field, vanishes outside K . However, we emphasise that our results are not tied to this particular coupling \mathcal{L}_I , nor is it even required that the theories involved are described by Lagrangians.

These assumptions allow for a direct identification between the free theory and the interacting theory *before* as well as *after* the coupling – or more precisely, outside K ’s causal future and past respectively. The comparison between these identifications is encoded in a unitary scattering matrix S , which takes the place of the interacting time evolution in the quantum mechanical setting. To be specific, the adjoint action $A \mapsto SAS^{-1}$ of S is obtained by mapping from the uncoupled to coupled theory using the late-time identification, followed by mapping back to the uncoupled system using the early-time identification. This corresponds to the usual

composition of Møller maps. (Assuming the coupled and uncoupled theories both have the *time-slice property* – see below – any observable can be expressed in terms of either late-time or early-time observables.) In the example above the scattering map is given by $S := \overline{T} \exp\{\frac{i\lambda}{\hbar} \int_K \alpha(x)\phi(x)\psi(x)dx\}$ to all orders in formal perturbation theory, where $\overline{T} \exp$ is the *anti*-time-ordered exponential and α functions as a smooth spacetime cutoff.

The locality of the theories under consideration is reflected in localisation properties of S , which are discussed in more detail in Lemma 1 below. In consequence, the idea of a measurement scheme can be implemented in QFT as a *local* concept. In particular, it was shown in [7, 8] how the correspondence between probe observables and induced system observables may be made, and how rules for state update appropriate to selective and non-selective measurements may be described. A non-technical outline of these results now follows.

Suppose that Z is a local observable of the probe theory, corresponding to a local observable $\mathbb{1} \otimes Z$ of the uncoupled combination of the system and probe theories. Likewise, $\rho_S \otimes \rho_P$ is an uncorrelated state of the same theory. We consider an experiment in the coupled theory, in which an observable corresponding to $\mathbb{1} \otimes Z$ at late times is measured in a state that corresponds to $\rho_S \otimes \rho_P$ at early times. The expectation value of this measurement, $E_{\rho_P}(Z; \rho_S)$, is [7]

$$E_{\rho_P}(Z; \rho_S) = \text{Tr}((\rho_S \otimes \rho_P)(S(\mathbb{1} \otimes Z)S^\dagger)). \quad (1)$$

We remark that we use the term *expectation value* in an operational way following the frequentist interpretation. For an analysis of the interpretation in terms of the first moment of an underlying probability measure we refer the reader to [11]. The induced system observable \hat{Z}_{ρ_P} corresponding to probe observable Z is, by definition, the observable whose expectation in state ρ_S matches that of the actual experiment:

$$\text{Tr}(\rho_S \hat{Z}_{\rho_P}) = E_{\rho_P}(Z; \rho_S) \quad (2)$$

It turns out that the induced observable belongs to the algebra of system observables corresponding to the coupling region (more precisely, to the causal hull of any connected neighbourhood thereof).

Turning to the issue of state updates, let us, for the sake of presentation, consider the case where there is a measurement of a probe observable Z and a (non further specified) measurement of a system observable A in the ‘out’ region (i.e., not to the past) of the Z -measurement. (A more formal discussion solely in terms of probe observables yields the same expressions for the updated states and is given in Sec. VI.) For simplicity let us assume that Z is a yes-no observable (i.e., an effect). Analysing the A and Z measurement results together over an ensemble of identical runs, we may restrict to the

subensemble in which the Z -measurement was successful (‘yes’). The expectation value of A , conditioned on success of Z , is

$$\frac{\text{Tr}((\rho_S \otimes \rho_P)(S(A \otimes Z)S^\dagger))}{\text{Prob}(Z|\rho_S)} =: \text{Tr}(\rho'_{S|Z} A), \quad (3)$$

where the system state $\rho'_{S|Z}$ defined in this way may be regarded as the updated state consequent upon successful measurement of Z , which occurs in the full ensemble with probability $\text{Prob}_{\rho_P}(Z|\rho_S) = E_{\rho_P}(Z; \rho_S)$. In a similar way, the updated state $\rho'_{S|\neg Z}$ conditioned on an unsuccessful (‘no’) Z -measurement may be obtained from the above on replacing Z by $\mathbb{1} - Z$. If no selection is made, then the updated state ρ'_S is an appropriately weighted statistical mixture of $\rho'_{S|Z}$ and $\rho'_{S|\neg Z}$, giving

$$\text{Tr}(\rho'_S C) = \text{Tr}((\rho_S \otimes \rho_P)(S(C \otimes \mathbb{1})S^\dagger)). \quad (4)$$

Notice that this expression is independent of the particular observable Z ; in tracing out the probe degrees of freedom, it is assumed that no further measurements of the probe are made.

Multiple measurements, each conducted by a different probe, may be accommodated provided that their coupling regions lie in a causal order, with each separated by a Cauchy surface from its predecessor. A crucial consistency relation established in [7] implies that the rules for state updates are independent of the choice of order when more than one is possible; this was shown explicitly in [7] for pairs of measurements and will be extended to the general case in Sec. VI below. The consistency result relies on a natural assumption called *causal factorisation*.

c. Sorkin’s protocol in the FV-framework In the FV-framework, Sorkin’s protocol is modelled as follows: Alice, Bob and Charlie are each described by *probes* which are coupled to the *system* of interest in the compact coupling zones K_1, K_2, K_3 each contained in the connected *regions* O_1, O_2, O_3 respectively, in which the experimenters perform actions. This guarantees that there is a natural causal order of K_1, K_2, K_3 , i.e., the one inherited from O_1, O_2, O_3 , see Fig 1. In particular, there are Cauchy surfaces having K_1 to their past and K_2 to their future; K_2 and K_3 are also separated by Cauchy surfaces in the same way.

The measurements of Alice and Bob in step one and two of the protocol produce an update of the system state $\rho_S \mapsto \rho_{AB}$ according to

$$\begin{aligned} & \text{Tr}(\rho_{AB} C) \\ &= \text{Tr}((\rho_S \otimes \rho_{P_1} \otimes \rho_{P_2})(S_1 S_2 (C \otimes \mathbb{1} \otimes \mathbb{1}) S_2^\dagger S_1^\dagger)), \end{aligned} \quad (5)$$

which is a straightforward generalisation of Eq. (4) using the natural causal order of the three experimenters. In fact there is no ambiguity if K_1, K_2 and K_3 admit other causal orders (which can happen if K_2 is spacelike from K_1 or K_3) – see Sec. V. As argued above, the expectation

value of Charlie's measurement in step three is given by Eq. (5) for a probe-induced system observable C , which is determined by the interaction between Charlie's probe and the system in coupling zone K_3 and may be localised in O_3 . The superluminal signalling between the spacelike separated experimenters Alice and Charlie in Sorkin's protocol arises if $\text{Tr}(\rho_{AB}C)$ differs from $\text{Tr}(\rho_B C)$, where ρ_B is the updated state in a situation where Alice does not perform an experiment, i.e., where there is *no* coupling between her probe and the system and hence no measurement is made on the system. This corresponds to Eq. (5) in the case where $S_1 = \mathbb{1}$. Hence, there is *no* superluminal signalling if

$$\begin{aligned} & \text{Tr} \left((\rho_S \otimes \rho_{P_1} \otimes \rho_{P_2})(S_1 S_2 (C \otimes \mathbb{1} \otimes \mathbb{1}) S_2^\dagger S_1^\dagger) \right) \\ &= \text{Tr} \left((\rho_S \otimes \rho_{P_2})(S_2 (C \otimes \mathbb{1}) S_2^\dagger) \right) \end{aligned} \quad (6)$$

for system observables C induced by Charlie's probe. The main result of this paper is that (6) holds under very mild technical assumptions. This result is stated and proved as Theorem 2 in Sec. IV and makes essential use of the localisation properties of the scattering map.

In fact, the statement we prove is actually more general as it establishes the desired equality for *all* system observables C localisable in O_3 and not just the ones induced by Charlie's probe (if this class is smaller).

III. TECHNICAL DESCRIPTION

The setting of the FV-framework is algebraic quantum field theory in possibly curved spacetime, which we now briefly recall.

a. Lorentzian geometry We start by fixing notation and recalling standard results of Lorentzian Geometry. Let M be a globally hyperbolic spacetime, i.e., a time-oriented Lorentzian spacetime of dimension at least two that contains a Cauchy surface. For $N \subseteq M$ let $J^+(N)$ and $J^-(N)$ denote its causal future and past respectively and define its causal hull to be $\text{ch}(N) := J^+(N) \cup J^-(N)$; N is called causally convex if it equals its causal hull. Any open causally convex subset of M will be called a *region* and is itself globally hyperbolic when regarded as a spacetime in its own right. Let $D^+(N)$ and $D^-(N)$ denote the future and past Cauchy developments of N , that is the set of points $p \in M$ such that every past-, respectively future-inextendible piecewise smooth causal curve through p intersects N . Then $D(N) := D^+(N) \cup D^-(N)$ is called the Cauchy development or domain of dependence of N . The causal complement of a subset K is defined to be $K^\perp := M \setminus (J^+(K) \cup J^-(K))$. For a compact subset K , the sets $M \setminus J^\mp(K)$ and K^\perp are all open and causally convex and therefore globally hyperbolic. See, for example, the appendix of [12] for details and proofs.

b. Algebraic quantum field theory Let M be a globally hyperbolic spacetime. An algebraic quantum

field theory (AQFT), or simply a *theory*, on M consists of a $*$ -algebra \mathcal{A} with a unit $\mathbb{1}$, together with a family of sub- $*$ -algebras $\mathcal{A}(N)$ of $\mathcal{A}(M) := \mathcal{A}$, each containing $\mathbb{1}$ and labelled by the regions $N \subseteq M$. The elements of $\mathcal{A}(N)$ are considered to be local observables of N , e.g., algebraic combinations of smeared fields $\int_N f(x) \phi(x) dx$ for a quantum field ϕ and a test function f vanishing outside N . This interpretation motivates the following additional assumptions:

Isotony: For regions $N_1 \subseteq N_2$: $\mathcal{A}(N_1) \subseteq \mathcal{A}(N_2)$.

Einstein causality: For spacelike separated regions N_1 and N_2 : the elements of $\mathcal{A}(N_1)$ commute with the elements of $\mathcal{A}(N_2)$.

Time-slice property: For regions $N_1 \subseteq N_2$, so that N_1 contains a Cauchy surface for N_2 : $\mathcal{A}(N_1) = \mathcal{A}(N_2)$.

The time-slice property encodes the existence of a (not further specified) local dynamical law. Morally: A quantum field is determined by its data on a Cauchy surface. We emphasise that the time-slice property is local in the sense that it applies to every region N_2 .

Due to time-slice (and isotony), every observable is localisable in many different, possibly disjoint regions. For example, if an observable A is localisable in a region N_1 and N_2 is a disjoint region containing N_1 in its domain of dependence, i.e., $N_1 \subseteq D(N_2)$, then A is also localisable in N_2 .

One also assumes a *Haag property*, which heuristically guarantees that the theory captures all relevant degrees of freedom. It is used to show that induced observables are localisable in every *connected* region containing the coupling zone – see [7] for details.

Haag property: For every compact set $K \subseteq M$ and every *connected* region L containing K , $\mathcal{A}(L)$ contains every $C \in \mathcal{A}$ that commutes with all elements of $\mathcal{A}(K^\perp)$. [13]

In AQFT, a *state* is a linear map $\omega : \mathcal{A}(M) \rightarrow \mathbb{C}$ which assigns expectation values to algebra elements and is therefore required to be normalised, $\omega(\mathbb{1}) = 1$, and positive, $\omega(A^*A) \geq 0$ for all $A \in \mathcal{A}(M)$.

c. Coupled theories The coupling between probe and system theory and the resulting scattering map arise as follows: Suppose we have three theories on a globally hyperbolic spacetime M : a system-theory \mathcal{S} , a probe-theory \mathcal{P} and a coupled theory \mathcal{C} , which mirrors the crucial assumption that the coupled structure is itself *local*. Let $\mathcal{S} \otimes \mathcal{P}$ denote the tensor-product theory, i.e., the uncoupled combination. As discussed before, \mathcal{S} and \mathcal{P} are coupled together *only* in a compact coupling zone $K \subseteq M$, which is modelled by the existence of a bijective, structure and localisation preserving identification between the coupled and uncoupled theories *outside* (the causal hull) of K , see [7] for the details. For the *in-region* M^- and *out-region* M^+ defined by $M^\pm := M \setminus J^\mp(K)$, this gives us the following maps:

$$\begin{aligned} \mathcal{S} \otimes \mathcal{P} &\rightarrow (\mathcal{S} \otimes \mathcal{P})(M^+) \rightarrow \mathcal{C}(M^+) \rightarrow \mathcal{C}, \\ \mathcal{C} &\rightarrow \mathcal{C}(M^-) \rightarrow (\mathcal{S} \otimes \mathcal{P})(M^-) \rightarrow \mathcal{S} \otimes \mathcal{P}, \end{aligned} \quad (7)$$

each of which is an isomorphism. The first, third, fourth and sixth are given by the time-slice property as M^\pm each contain a Cauchy surface for M [14]. The other arrows are given by the localisation preserving identification map. The overall composition defines the *scattering map* $\Theta : \mathcal{S} \otimes \mathcal{P} \rightarrow \mathcal{S} \otimes \mathcal{P}$, which is an automorphism preserving algebraic relations but *not* localisation. Our earlier discussion in Sec. II implicitly assumed that Θ was implemented as the adjoint action of a unitary scattering operator S , i.e., $\Theta(A) = SAS^\dagger$, but this is neither needed nor assumed in what follows. The localisation properties of Θ are summarised in the following lemma.

Lemma 1 (Proposition 3.1(b),(c) in [7]).

1. For every region $N \subseteq K^\perp : \Theta$ acts trivially on $(\mathcal{S} \otimes \mathcal{P})(N)$.
2. For every region $N \subseteq M^+$ and every region $N^- \subseteq M^-$ with $N \subseteq D(N^-) : \Theta(\mathcal{S} \otimes \mathcal{P})(N) \subseteq (\mathcal{S} \otimes \mathcal{P})(N^-)$.

The first property captures the idea that the coupling has no effect in spacelike separated regions, whereas the second property indicates how Θ changes the localisation of observables.

Now suppose that the system is prepared in state ω and the probe in state σ , and that a measurement of a probe observable is made. The state update rule (without selection) is that $\omega \mapsto \omega'$, where

$$\omega'(C) = (\omega \otimes \sigma)(\Theta(C \otimes \mathbb{1})), \quad (8)$$

which is readily recognised as the analogue of (4).

IV. MAIN RESULT

Let us now discuss the rigorous FV-version of Sorkin's protocol and Eq. (5). Alice, Bob and Charlie each perform actions in the connected *regions* O_1, O_2, O_3 . We assume they fulfill (a) $O_2 \cap J^-(O_1) = \emptyset$; (b) $O_3 \cap J^-(O_2) = \emptyset$; (c) O_3 is spacelike separated from O_1 ; (d) O_3 has compact closure \overline{O}_3 . Note that this covers the situation sketched in Fig 1 but is more general. Let \mathcal{S} be the system theory and let $\mathcal{P}_1, \mathcal{P}_2$ be the two probe theories of Alice and Bob with compact coupling zones K_1, K_2 contained in the regions O_1, O_2 respectively. Denote the corresponding in- and out-regions by M_1^\mp, M_2^\mp , the initial states by σ_1, σ_2 and the associated scattering maps by $\Theta_i : \mathcal{S} \otimes \mathcal{P}_i \rightarrow \mathcal{S} \otimes \mathcal{P}_i$ for $i = 1, 2$. On $\mathcal{S} \otimes \mathcal{P}_1 \otimes \mathcal{P}_2$ define $\hat{\Theta}_1 := \Theta_1 \otimes \mathbb{1}$ and $\hat{\Theta}_2 := \Theta_2 \otimes \mathbb{1}$, where the subscript on the tensor product indicates the slot into which the second factor is inserted.

Let C be a system-observable localisable in O_3 , Charlie's 'region of control'. For example, C could be the induced observable corresponding to any probe observable of Charlie's. Owing to assumptions (a–c), Alice, Bob and Charlie admit a causal order in which Alice's region precedes Bob's, and Bob's region precedes

Charlie's. If Alice and Bob each perform a measurement, the expectation value for Charlie's measurement is therefore given by

$$\omega_{AB}(C) := (\omega \otimes \sigma_1 \otimes \sigma_2)((\hat{\Theta}_1 \circ \hat{\Theta}_2)(C \otimes \mathbb{1} \otimes \mathbb{1})), \quad (9)$$

for initial system state ω . Strictly speaking, when writing down Eq. (9) at this stage, we make the assumption that the effect of two, causally orderable measurements on the initial state ω is given in terms of a composition of individual state updates. However, further below in Sec. VI we show how Eq. (9) can be *derived* in the FV-framework. Furthermore, assumptions (a–c) do not exclude the possibility that the regions controlled by Alice, Bob and Charlie also admit other causal orderings, but Charlie's expectation value is well-defined and independent of any choices made. This will also be discussed in greater depth in Sec. V. On the other hand, if Alice does not perform her experiment, Charlie's expectation value is

$$\omega_B(C) = (\omega \otimes \sigma_2)(\Theta_2(C \otimes \mathbb{1})). \quad (10)$$

The following theorem (the rigorous analogue of Eq. (6)) shows that Sorkin's protocol does *not* signal in the FV-framework. Note that it gives the desired equality without the (possibly restricting) assumption that C is an induced observable.

Theorem 2. *In the notation above, suppose the following assumptions hold: (a) $K_2 \cap J^-(K_1) = \emptyset$; (b) O_3 is a region with compact closure; (c) $O_3 \cap J^-(K_2) = \emptyset$; (d) \overline{O}_3 is spacelike separated from K_1 . Then*

$$\forall C \in \mathcal{S}(O_3) : (\hat{\Theta}_1 \circ \hat{\Theta}_2)(C \otimes \mathbb{1} \otimes \mathbb{1}) = \hat{\Theta}_2(C \otimes \mathbb{1} \otimes \mathbb{1}). \quad (11)$$

This immediately implies

$$\begin{aligned} \omega_{AB}(C) &= (\omega \otimes \sigma_1 \otimes \sigma_2)((\hat{\Theta}_1 \circ \hat{\Theta}_2)(C \otimes \mathbb{1} \otimes \mathbb{1})) \\ &= (\omega \otimes \sigma_2)(\Theta_2(C \otimes \mathbb{1})) = \omega_B(C), \end{aligned} \quad (12)$$

i.e., Charlie's measurement outcome is independent of whether Alice does or does not perform an experiment at all. There is no superluminal signalling.

The proof of Theorem 2 relies on localisation properties of the scattering map, combined with a geometrical lemma.

Lemma 3. *Let K_1, K_2, L be compact subsets of M , let $K_2 \cap J^-(K_1) = \emptyset$ and $L \cap J^-(K_1) = \emptyset$. Then there exists a Cauchy surface Σ of M_1^+ such that $\Sigma \subseteq M \setminus (J^-(K_1) \cup J^+(K_2) \cup J^+(L))$.*

Proof. $M_1^+ = M \setminus J^-(K_1)$ is globally hyperbolic (see Lemma A.4 in [14]). By Proposition 4 in [15] (due to Geroch [16]) there exists a surjective, continuous function $t : M_1^+ \rightarrow \mathbb{R}$, strictly increasing on every future-directed causal curve, whose level sets are Cauchy surfaces for M_1^+ . Since K_2 and L are compact and t is continuous,




FIG. 2. Schematic spacetime diagram of the relative causal position of the compact sets K_1 , K_2 (coupling zones) and L (\bar{O}_3 in Lemma 4) as well as the Cauchy surface Σ in Lemma 3.

$\tilde{\tau} := \min t[K_2 \cup L]$ exists. Choose $\tau < \tilde{\tau}$ and set $\Sigma := t^{-1}[\{\tau\}]$. Σ is a Cauchy surface for M_1^+ and fulfills the desired properties. \square

We apply the lemma for the case where K_1, K_2 are the coupling zones of Alice and Bob and L is the closure of Charlie's region of control, i.e., $L = \bar{O}_3$, which is compact. This allows us to prove that O_3 is contained in the domain of dependence of $K_1^\perp \cap M_2^-$.

Lemma 4. *Let K_1, K_2 be compact subsets of M such that $K_2 \cap J^-(K_1) = \emptyset$. Then for every region O_3 with compact closure such that $O_3 \cap J^-(K_2) = \emptyset$ and $\bar{O}_3 \subseteq K_1^\perp$ it holds that $O_3 \subseteq D(K_1^\perp \cap M_2^-)$.*

Proof. By setting $L := \bar{O}_3$ and using Lemma 3, we can find Σ , a Cauchy surface for M_1^+ which lies in $M \setminus (J^-(K_1) \cup J^+(K_2) \cup J^+(\bar{O}_3))$. Set $T := J^-(\bar{O}_3) \cap \Sigma \subseteq K_1^\perp \cap M_2^-$. (T is spacelike separated from K_1 , because Σ is disjoint from $J^-(K_1)$ and because $J^-(\bar{O}_3)$ is disjoint from $J^+(K_1)$ as $\bar{O}_3 \subseteq K_1^\perp$ by assumption.) Now $O_3 \subseteq D(T)$. $K_1^\perp \cap M_2^- = M_1^+ \cap M_1^- \cap M_2^-$; as M_1^+, M_1^- and M_2^- are open and causally convex (see Lemma A.4 in [14]), so is their intersection, i.e., it is a region, and since it contains T , we have that $O_3 \subseteq D(T) \subseteq D(K_1^\perp \cap M_2^-)$. \square

Theorem 2 now follows by using the localisation properties of Θ and the fact that Charlie's region of control is contained in the domain of dependence of a sub-region of K_1^\perp .

Proof of Theorem 2. Since $C \in \mathcal{S}(O_3)$, $C \otimes \mathbb{1} \otimes \mathbb{1}$ can be localised in O_3 too. According to Lemma 4, $O_3 \subseteq D(K_1^\perp \cap M_2^-)$. According to Lemma 1, we know that $\hat{\Theta}_2(C \otimes \mathbb{1} \otimes \mathbb{1})$ can be localised in the region $K_1^\perp \cap M_2^-$. But since $K_1^\perp \cap M_2^- \subseteq K_1^\perp$, we have by Lemma 1 that $(\hat{\Theta}_1 \circ \hat{\Theta}_2)(C \otimes \mathbb{1} \otimes \mathbb{1}) = \hat{\Theta}_2(C \otimes \mathbb{1} \otimes \mathbb{1})$. \square

V. CAUSAL FACTORISATION

In the previous section we showed that measurements of three observers described in the FV-framework do

not run into the potential superluminal signalling issues associated to Sorkin's *impossible measurements*. To do this, we made the assumption that the effect of causally orderable measurements may be given in terms of a composition of individual state updates as in Eq. (9). In the next section we will show that this assumption can actually be derived in the FV-framework as a result of what is called *causal factorisation*, which we now describe. Our presentation here is certainly not the most general possible but will be sufficient for our current purposes. We intend to report elsewhere on more abstract and general properties of causal factorisation.

To start, let \mathcal{K} be a collection of compact spacetime subsets. A linear order \leq on \mathcal{K} is said to be a *causal linear order* if $K < K'$ implies $J^-(K) \cap J^+(K') = \emptyset$ for every $K, K' \in \mathcal{K}$. It follows that whenever $K < K'$, there is a Cauchy surface of M with K to its past and K' to its future. If \mathcal{K} admits a causal linear order, we say that \mathcal{K} is *causally orderable*. A causally orderable set may admit more than one distinct causal linear order; this happens, for example, in the case of two spacelike separated sets. When the members of a causally orderable set \mathcal{K} are the coupling zones for a collection of observers, we will describe the observers as causally orderable and use any causal linear ordering of \mathcal{K} to induce a linear order on the collection of observers.

Now let \mathcal{S} be a theory of interest and consider two causally orderable observers, A and B, with probe theories \mathcal{P}_A and \mathcal{P}_B . The description of A's measurements in the FV scheme involves *inter alia* the uncoupled combination $\mathcal{S} \otimes \mathcal{P}_A$ and a coupled theory \mathcal{C}_A with a coupling zone K_A , along with a corresponding scattering map Θ_A on $\mathcal{S} \otimes \mathcal{P}_A$; B's measurements are described in a similar way. If both A and B measure independently, they can be considered as a combined "super-observer" whose probe theory $\mathcal{P}_{\{A,B\}}$ is a tensor product of \mathcal{P}_A and \mathcal{P}_B . As the two coupling regions K_A and K_B may be separated by a Cauchy surface, it is reasonable to assume that there is a combined coupled theory $\mathcal{C}_{\{A,B\}}$ with coupling zone $K_A \cup K_B$, and a scattering map $\Theta_{\{A,B\}}$, which can be decomposed as an appropriate composition of the individual scattering maps. Accordingly, we say that the combination of A and B respects *bipartite causal factorisation*, if and only if the coupled theory $\mathcal{C}_{\{A,B\}}$ exists and

$$\Theta_{\{A,B\}} = \begin{cases} \hat{\Theta}_A \circ \hat{\Theta}_B & \text{if } K_B \cap J^-(K_A) = \emptyset \\ \hat{\Theta}_B \circ \hat{\Theta}_A & \text{if } K_A \cap J^-(K_B) = \emptyset, \end{cases} \quad (13)$$

where $\hat{\Theta}_X$ ($X = A, B$) denotes the trivial extension of the scattering map Θ_X from an automorphism of $\mathcal{S} \otimes \mathcal{P}_X$ to an automorphism of $\mathcal{S} \otimes \mathcal{P}_{\{A,B\}}$ by tensoring with a suitable identity map. In particular, if the two coupling regions are spacelike separated then $\Theta_{\{A,B\}}$ may be factored in both ways. The assumption of bipartite causal factorisation is motivated by the expression for the scattering map in terms of time-ordered products in conventional perturbation theory. As a special

case of Bogoliubov's factorisation relation, bipartite causal factorisation is in particular fulfilled by the time-ordered exponential of local coupling terms (with smooth cutoff) in renormalised perturbation theory [17] and by the generators in recent non-perturbative Lagrangian approaches [18]. Moreover it is proved to hold for the probe models considered in [7].

Our treatment of multiple observers is based on three physically motivated assumptions:

1. every finite collection of causally orderable observers can be combined to form a super-observer, whose probe theory is a tensor product of the individual probe theories;
2. the combination process may be achieved in a single step, or equivalently, as the result of successive stages of combination;
3. the combination of any causally orderable pair of observers fulfils *bipartite causal factorisation*.

To illustrate these ideas, let us consider three (distinct) causally orderable observers A, B, C admitting a causal linear order \leq in which $A \leq B \leq C$. The super-observer $\{A, B, C\}$ can be formed in one go, or equivalently by first combining A and B to $\{A, B\}$ and then further combining with C ; alternatively, we can first combine B with C and then combine with A . Understanding 'equivalence' as equality of scattering maps, we have

$$\Theta_{\{\{A,B\},C\}} = \Theta_{\{A,B,C\}} = \Theta_{\{A,\{B,C\}\}} \quad (14)$$

and, on using bipartite causal factorisation, one has

$$\Theta_{\{A,B,C\}} = \hat{\Theta}_{\{A,B\}} \circ \hat{\Theta}_C = \hat{\Theta}_A \circ \hat{\Theta}_B \circ \hat{\Theta}_C, \quad (15)$$

where the hats denote the extension of the scattering maps to $\mathcal{S} \otimes \mathcal{P}_{\{A,B,C\}}$. Moreover the assumptions also imply that whenever there is a choice between different causal orders for fixed A, B, C , the combined scattering map $\Theta_{\{A,B,C\}}$ can be written as a composition of the individual scattering maps in either of these orders.

In general, given any finite set Obs of N causally orderable observers, the super-observer has a combined probe theory

$$\mathcal{P}_{\text{Obs}} = \bigotimes_{X \in \text{Obs}} \mathcal{P}_X \quad (16)$$

and an overall scattering map Θ_{Obs} on $\mathcal{S} \otimes \mathcal{P}_{\text{Obs}}$ that factorises as

$$\Theta_{\text{Obs}} = \hat{\Theta}_{X_1} \circ \hat{\Theta}_{X_2} \circ \dots \circ \hat{\Theta}_{X_N} \quad (17)$$

whenever $X_1 < X_2 < \dots < X_N$ according to some causal linear ordering \leq of Obs ; the hats denote extensions to $\mathcal{S} \otimes \mathcal{P}_{\text{Obs}}$. There are many equivalent formulae for Θ_{Obs} , arising from different ways of successively combining the observers.

VI. MEASUREMENTS BY MULTIPLE OBSERVERS

In this section we demonstrate how multiple successive measurements can be treated in the FV-framework. We start with a discussion of one single observer and a pair of two observers, where we recall results from [7]. We then move on to present the treatment of three observers, which readily generalises to the general $N \in \mathbb{N}$ observer case. We end this section with a discussion of the process of post-selection.

a. Induced observables and effects Let \mathcal{S} be a theory of interest and let A be an observer who wishes to measure (the expectation value) of some local observable of \mathcal{S} in initial state ω . Suppose A has probe theory \mathcal{P}_A , initial probe state σ_A , compact coupling zone K_A , a coupled theory \mathcal{C}_A , identification maps and associated scattering map $\Theta_A : \mathcal{S} \otimes \mathcal{P}_A \rightarrow \mathcal{S} \otimes \mathcal{P}_A$. The prediction for the expectation value of a probe observable $O_A \in \mathcal{P}_A$ in the actual experiment conducted by A , given initial system state ω , is denoted by $E_A(O_A; \omega)$ and given by [7]

$$E_A(O_A; \omega) = (\omega \otimes \sigma_A)(\Theta_A(\mathbb{1} \otimes O_A)). \quad (18)$$

If the observable algebras are represented as operators on a Hilbert space, we can consider the case where ω and σ_A are given by density matrices and where Θ_A is implemented as the adjoint action of a unitary scattering operator. Then the above equation is easily recognised as a straightforward generalisation of Eq. (1).

Returning to the general situation, it was shown in [7] that σ_A and Θ_A give rise to a map $\varepsilon_A : \mathcal{P}_A \rightarrow \mathcal{S}$ such that

$$\forall O_A \in \mathcal{P}_A : \omega(\varepsilon_A(O_A)) = (\omega \otimes \sigma_A)(\Theta_A(\mathbb{1} \otimes O_A)). \quad (19)$$

We call $\varepsilon_A(O_A)$ the *induced (system) observable* corresponding to O_A , as introduced in Eq. (2).

Exploiting the *Haag property* it can be shown that $\varepsilon_A(O_A)$ is a local observable of the system theory, which can be localised in any connected region containing K_A [7, Theorem 3.3].

The interpretation of Eq. (19) is the following: If observer A is interested in the expectation value of a specific local system observable, then she needs to prepare and tune her physical detector, i.e., find O_A , σ_A and Θ_A such that $\varepsilon_A(O_A)$ is the desired system observable. It is an open question whether this is always possible, so we take the pragmatic viewpoint and say the system observables of interest to an observer are those which can be measured using a probe, i.e., those which can be induced by some probe observable upon tuning the probe state and scattering map.

The result of an actual experiment is generically not immediately a sharp numerical outcome but rather an answer to a (finite collection of) yes-no question(s). In quantum theory, this is modelled abstractly by considering a projector or more generally an *effect* P associated to *yes* and $\mathbb{1} - P$ associated to *no* as the main observables of interest. Recall that a $*$ -algebra element

P is an effect if and only if $P^\dagger = P$ and $0 \leq P \leq \mathbb{1}$ where $0 \leq P$ means that P is a convex combination of elements of the form $A^\dagger A$. One frequently calls the expectation value of an effect P its *success probability*.

b. Two observers Consider a set of two observers $\text{Obs} = \{A, B\}$ each of whom wishes to determine the expectation value of a system observable $\varepsilon_A(O_A)$ and $\varepsilon_B(O_B)$ respectively for probe observables $O_A \in \mathcal{P}_A$ and $O_B \in \mathcal{P}_B$. We intend to answer the following question: “What is the expected outcome of observer B’s measurement?”

Similar to before, for every $X \in \text{Obs}$ who interacts with a system-theory \mathcal{S} in initial state ω we have a probe-theory \mathcal{P}_X , initial state σ_X , compact coupling zone K_X , coupled theory \mathcal{C}_X identification maps and associated scattering map $\Theta_X : \mathcal{S} \otimes \mathcal{P}_X \rightarrow \mathcal{S} \otimes \mathcal{P}_X$.

As in the previous section, we may regard the collection of all observers as super-observer in its own right with probe $\mathcal{P} := \bigotimes_{X \in \text{Obs}} \mathcal{P}_X$, coupling zone $K := \bigcup_{X \in \text{Obs}} K_X$ and initial state $\sigma := \bigotimes_{X \in \text{Obs}} \sigma_X$ on \mathcal{P} . Let $O := \bigotimes_{X \in \text{Obs}} O_X$ be the probe observable of interest. We assume the existence of an associated coupled theory \mathcal{C} emerging from coupling \mathcal{S} to \mathcal{P} in K giving rise to a scattering map $\Theta : \mathcal{S} \otimes \mathcal{P} \rightarrow \mathcal{S} \otimes \mathcal{P}$.

Let us now assume that after (an ensemble of) their experiments, the observers meet in their joint future to analyse their experimental data together. Since we consider the two of them to constitute a single super-observer, the expectation value of their joint measurement is, according to Eq. (18),

$$E_{\{A,B\}}(O; \omega) = (\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes O)). \quad (20)$$

If O_A and O_B are effects, O is an effect as well and Eq. (20) can be understood as the success probability of the “combined effect” O corresponding to the success of both O_A and O_B , i.e., $E_{\{A,B\}}(O; \omega) = \text{Prob}_{\{A,B\}}(O_A \& O_B; \omega)$.

In the context of effects it is also immediately possible to give an answer to the posed question. B’s expected outcome is the success probability $\text{Prob}_{\{A,B\}}(O_B; \omega)$ that B observes probe effect O_B given initial system state ω irrespective of A’s outcome. It is given as a marginal probability

$$\begin{aligned} & \text{Prob}_{\{A,B\}}(O_B; \omega) \\ &= \text{Prob}_{\{A,B\}}(O_A \& O_B; \omega) + \text{Prob}_{\{A,B\}}((\neg O_A) \& O_B; \omega) \\ &= (\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes O_A \otimes O_B)) \\ & \quad + (\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes (\mathbb{1} - O_A) \otimes O_B)) \\ &= (\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes \hat{O}_B)), \end{aligned} \quad (21)$$

where we used an explicit order of the tensor product, $\mathcal{P} = \mathcal{P}_A \otimes \mathcal{P}_B$ and where the hat denotes the inclusion of O_B in \mathcal{P} , $\hat{O}_B = \mathbb{1} \otimes O_B$. (The case in which one wishes to perform an analysis post-selected on a specific outcome of observer A is known as *selective* measurement and is discussed further at the end of this section.)

Note that Eq. (21) only depends on A through σ_A and the coupled scattering map Θ and, in particular, is independent of O_A . As a matter of fact, the above discussion in terms of effects can be seen as a motivation for considering Eq. (21) to be the answer to the posed question even in the situation where O_A and O_B are *not* effects, i.e., generally:

$$E_{\{A,B\}}(O_B; \omega) = (\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes \hat{O}_B)). \quad (22)$$

This is useful since, e.g., the field $*$ -algebra of the linear scalar field does not admit any non-trivial effects at all. Moreover, the expression makes a prediction for the *expectation value* of B’s experiment, which can be determined from B’s local experimental data alone. There is no need for B to meet A in their joint future to conduct data analysis together.

Let us continue the investigation of expression (22) in the physically relevant case that the set Obs is *causally orderable*. There are at most two possible linear causal orders on Obs , corresponding to the cases 1. $A \leq B$; or 2. $B \leq A$.

If the combination of the two observers respects bipartite causal factorisation, the super-scattering map decomposes as $\Theta = \hat{\Theta}_A \circ \hat{\Theta}_B$ in the first case, while $\Theta = \hat{\Theta}_B \circ \hat{\Theta}_A$ in the second. Before continuing, it is convenient to observe that, upon writing $\hat{\Theta}_B(\mathbb{1} \otimes O_A \otimes O_B) = \sum_j S_j \otimes O_A \otimes B_j$ and noting that $\varepsilon_B(O_B) = \sum_j \sigma_B(B_j) S_j$, the following holds:

$$\begin{aligned} & (\omega \otimes \sigma_A \otimes \sigma_B)(\hat{\Theta}_A \circ \hat{\Theta}_B(\mathbb{1} \otimes O_A \otimes O_B)) \\ &= (\omega \otimes \sigma_A \otimes \sigma_B)(\hat{\Theta}_A(\sum_j S_j \otimes O_A \otimes B_j)) \\ &= (\omega \otimes \sigma_A)(\Theta_A(\sum_j \sigma_B(B_j) S_j \otimes O_A)) \\ &= (\omega \otimes \sigma_A)(\Theta_A(\varepsilon_B(O_B) \otimes O_A)). \end{aligned} \quad (23)$$

This allows us to simplify Eq. (22) in each case:

1. For $A \leq B$ we order the tensor product of probes as $\mathcal{P} = \mathcal{P}_A \otimes \mathcal{P}_B$ and get

$$\begin{aligned} E_{\{A,B\}}(O_B; \omega) &= (\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes \hat{O}_B)) \\ &= (\omega \otimes \sigma_A \otimes \sigma_B)(\hat{\Theta}_A \circ \hat{\Theta}_B(\mathbb{1} \otimes \hat{O}_B)) \\ &= (\omega \otimes \sigma_A)(\Theta_A(\varepsilon_B(O_B) \otimes \mathbb{1})). \end{aligned} \quad (24)$$

Therefore, if the system state ω_A is defined so that $\omega_A(C) := (\omega \otimes \sigma_A)(\Theta_A(C \otimes \mathbb{1}))$ for all $C \in \mathcal{S}$, B’s expected outcome in this situation is

$$E_{\{A,B\}}(O_B; \omega) = \omega_A(\varepsilon_B(O_B)) = E_B(O_B; \omega_A), \quad (25)$$

which is his expected outcome if A does not measure, but with the system prepared in state ω_A instead of ω . This is the justification for regarding ω_A as the updated system state associated with A’s measurement, as asserted in Eq. (4) and (8). For future reference, let us define the update map $\mathcal{J}_A(\omega) := \omega_A$.

2. For $B \leq A$ and after ordering the tensor product of probes as $\mathcal{P} = \mathcal{P}_B \otimes \mathcal{P}_A$ for convenience,

$$\begin{aligned} E_{\{A,B\}}(O_B; \omega) &= (\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes \hat{O}_B)) \\ &= (\omega \otimes \sigma_B \otimes \sigma_A)(\hat{\Theta}_B \circ \hat{\Theta}_A(\mathbb{1} \otimes O_B)) \\ &= (\omega \otimes \sigma_B)(\Theta_B(\mathbb{1} \otimes O_B)) \\ &= \omega(\varepsilon_B(O_B)) = E_B(O_B; \omega), \end{aligned} \quad (26)$$

where we used that $\hat{\Theta}_A(\mathbb{1} \otimes \hat{O}_B) = (\mathbb{1} \otimes \hat{O}_B)$. This follows from the fact that in the present order of the tensor product of probes we have that $\mathbb{1} \otimes \hat{O}_B = \mathbb{1} \otimes O_B \otimes \mathbb{1}$ and $\hat{\Theta}_A = \Theta_A \otimes_2 \mathbb{1}$, where again the subscript on the tensor product indicates the slot into which the second factor is inserted. Recalling that observer B precedes A (with respect to \leq), the above result shows that there is no influence from the future to the past.

Finally, we remark that if K_A and K_B are spacelike separated, the causal order is not unique: there is an ordering corresponding to case 1 and another corresponding to case 2. However, there is no ambiguity because B's expected outcome is given by (22) in either case. This implies in particular that

$$\omega_A(\varepsilon_B(O_B)) = \omega(\varepsilon_B(O_B)) \quad (27)$$

if K_A, K_B are spacelike separated (as has been observed in [7]).

c. Three observers The obvious next step is to consider three observers and to give an answer to the question: “For a set of three observers $\text{Obs} = \{A, B, C\}$, each of which performs a measurement, what is the expected outcome of observer B's measurement of the induced system observable $\varepsilon_B(O_B)$?”

Following the reasoning and notation of before we can immediately write down the answer as

$$E_{\{A,B,C\}}(O_B; \omega) = (\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes \hat{O}_B)), \quad (28)$$

where we again assumed the existence of an appropriate overall super-observer similar to the bipartite case.

Let us further investigate Eq. (28) under the additional assumption that Obs is causally orderable and that causal factorisation holds. As A and C can be interchanged, there are at most three cases: A, B, C are such that there exists a linear order \leq with 1. $A \leq B \leq C$; 2. $C \leq A \leq B$; or 3. $B \leq C \leq A$.

Choosing a convenient order of tensor products and using results from before yields the following:

1. For $A \leq B \leq C$:

$$\begin{aligned} E_{\{A,B,C\}}(O_B; \omega) &= (\omega \otimes \sigma_A \otimes \sigma_B \otimes \sigma_C)(\hat{\Theta}_A \circ \hat{\Theta}_B \circ \hat{\Theta}_C(\mathbb{1} \otimes \hat{O}_B)) \\ &= (\omega \otimes \sigma_A \otimes \sigma_B)(\check{\Theta}_A \circ \check{\Theta}_B(\mathbb{1} \otimes \check{O}_B)) \\ &= (\omega \otimes \sigma_A)(\Theta_A(\varepsilon_B(O_B) \otimes \mathbb{1})) \\ &= \omega_A(\varepsilon_B(O_B)) \\ &= E_B(O_B; \omega_A), \end{aligned} \quad (29)$$

where the haček denotes the extension to $\mathcal{S} \otimes \mathcal{P}_{\{A,B\}}$ and where, similarly to before, we used that $\hat{\Theta}_C(\mathbb{1} \otimes \hat{O}_B) = \mathbb{1} \otimes \hat{O}_B$ and that $\hat{\Theta}_A$ and $\hat{\Theta}_B$ act trivially on \mathcal{P}_C . The upshot is that observer B's outcome is given by taking the initial system state, updating it according to the map \mathcal{J}_A associated to the observer preceding B (with respect to \leq) and evaluating the updated state on B's induced system observable. The observer succeeding B (with respect to \leq) can be completely ignored.

2. For $C \leq A \leq B$, ordering $\mathcal{P} = \mathcal{P}_C \otimes \mathcal{P}_A \otimes \mathcal{P}_B$,

$$\begin{aligned} E_{\{A,B,C\}}(O_B; \omega) &= (\omega \otimes \sigma_C \otimes \sigma_A \otimes \sigma_B)(\hat{\Theta}_C \circ \hat{\Theta}_A \circ \hat{\Theta}_B(\mathbb{1} \otimes \hat{O}_B)) \\ &= (\omega \otimes \sigma_C \otimes \sigma_A)(\check{\Theta}_C \circ \check{\Theta}_A(\varepsilon_B(O_B \otimes \mathbb{1}))) \\ &= \omega_{AC}(\varepsilon_B(O_B)) \\ &= E_B(O_B; \omega_{AC}), \end{aligned} \quad (30)$$

where $\omega_{AC} := (\mathcal{J}_A \circ \mathcal{J}_C)(\omega)$, cf. [7, Theorem 3.5]. Here, the haček denotes the trivial extension of the scattering maps to $\mathcal{S} \otimes \mathcal{P}_{\{A,C\}}$. The investigation of this case has some interesting consequences. First it provides a proof of the Eq. (9) which we used in the discussion of Sorkin's protocol (where Alice takes the place of C and Bob takes the place of A here). Second, if we regard $X := \{A, C\}$ as a super-observer in its own right, then Eq. (30) can be written as

$$\begin{aligned} E_{\{A,B,C\}}(O_B; \omega) &= (\omega \otimes \sigma_X \otimes \sigma_B)(\hat{\Theta}_X \circ \hat{\Theta}_B(\mathbb{1} \otimes \hat{O}_B)) \\ &= (\omega \otimes \sigma_X)(\Theta_X(\varepsilon_B(O_B) \otimes \mathbb{1})) \\ &= \omega_X(\varepsilon_B(O_B)) \\ &= E_B(O_B; \omega_X), \end{aligned} \quad (31)$$

which is the same calculation as in the case of two observers leading to Eq. (25). This idea will be used in the remaining case, as well as later on to simplify the investigation of $N > 3$ observers. (We will continue to use capital sans-serif Latin letters for individual observers, i.e., elements of Obs , as well as super-observers, i.e., subsets of Obs .)

3. For $B \leq C \leq A$, ordering $\mathcal{P} = \mathcal{P}_B \otimes \mathcal{P}_C \otimes \mathcal{P}_A$ and then regarding $X := \{A, C\}$ as super-probe in its own right enables us to write

$$\begin{aligned} E_{\{A,B,C\}}(O_B; \omega) &= (\omega \otimes \sigma_B \otimes \sigma_X)(\hat{\Theta}_B \circ \hat{\Theta}_X(\mathbb{1} \otimes \hat{O}_B)) \\ &= (\omega \otimes \sigma_B)(\Theta_B(\mathbb{1} \otimes \hat{O}_B)) \\ &= \omega(\varepsilon_B(O_B)) \\ &= E_B(O_B; \omega), \end{aligned} \quad (32)$$

which reinforces the message that there is no signalling from the future to the past.

A given set of observers $\{A, B, C\}$ can admit more than one causal order, however, as for two observers, the answer for the various admissible cases will agree.

d. N observers Let us assume that we have a finite set Obs of N causally orderable observers, each of whom wishes to determine the expectation value of an induced system observable. We fix one observer $B \in \text{Obs}$ and ask: “What is the expected outcome $E_{\text{Obs}}(O_B; \omega)$ of observer B ’s measurement of the induced system observable $\varepsilon_B(O_B)$?” More specifically, we want to know how the general answer $E_{\text{Obs}}(O_B; \omega) = (\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes \hat{O}_B))$ may be simplified in this situation.

Any fixed causal order \leq on Obs gives rise to the following tripartite partition $A := \{X \in \text{Obs} | X < B\}$, $\{B\}$ and $C := \{X \in \text{Obs} | X > B\}$. It immediately follows from Eq. (29) in the analysis of three observers that

$$E_{\text{Obs}}(O_B; \omega) = \omega_A(\varepsilon_B(O_B)) = E_B(O_B; \omega_A). \quad (33)$$

That is, B ’s expected outcome is equal to the one obtained in the absence of the other observers, but in system state ω_A (which is equal to ω if A is empty, i.e., if B is the earliest observer according to \leq). Moreover, suppose that the constituent observers of A are ordered $A_1 < A_2 < \dots < A_{|A|}$ according to \leq . Then causal factorisation gives

$$\Theta_A = \hat{\Theta}_{A_1} \circ \dots \circ \hat{\Theta}_{A_{|A|}}, \quad (34)$$

which implies that we can write the updated state ω_A as the result of $|A|$ many individual updates according to

$$\omega_A = (\mathcal{J}_{A_{|A|}} \circ \dots \circ \mathcal{J}_{A_1})(\omega) = \omega_{A_{|A|}, \dots, A_1}, \quad (35)$$

which follows from the $|A| = 2$ case by induction.

This demonstrates how multiple measurements are modelled in the FV-framework and shows in particular how the familiar concept of successive state-updates is recovered in the situation of causally orderable observers. As we have emphasised, in spite of the possible ambiguity of the causal order, *causal factorisation* ensures that the answer is unique and also free of any influence from the future to the past with respect to any causal order on the set of observers. However, Sorkin’s impossible measurements raise the question of whether any rule for assigning expectation values might be plagued by other acausal influences. It is the purpose of Sec. VII to show that this is not the case in the FV-framework, thus generalising the results of Sec. IV.

Before that, we end the present section with a discussion of *selective* measurements, i.e., the process of post-selection and associated conditional expectation values.

e. Conditional expectation and post-selection Let $\text{Obs} = \{A, B\}$ be a set of two causally orderable observers each of whom performs a measurement of system observable $\varepsilon_X(O_X)$ for $X \in \text{Obs}$. In this context: “What is observer B ’s outcome conditioned on a certain outcome of A ’s measurement?”

To this end let us consider the situation where O_A is an effect and where B is interested in the outcome of

his experiment given that A measures successfully. As before, let us motivate our answer by first considering the case of an *effect* O_B and let us assume that, after conducting (an ensemble of) their experiments, the observers meet in their joint future to analyse their data together. In this case (slightly generalising Sec. 3.3 of [7]) we can write down the success probability of O_B conditioned on (success of) O_A as the conditional probability

$$\begin{aligned} \text{Prob}_{\{A, B\}}(O_B | O_A; \omega) &= \frac{\text{Prob}_{\{A, B\}}(O_B \& O_A; \omega)}{\text{Prob}_{\{A, B\}}(O_A; \omega)} \\ &= \frac{(\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes O_A \otimes O_B))}{(\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes \hat{O}_A))} \\ &= \frac{\omega(\varepsilon_{\{A, B\}}(O_A \otimes O_B))}{\omega(\varepsilon_A(O_A))}, \end{aligned} \quad (36)$$

under the standing assumption that $\text{Prob}_{\{A, B\}}(O_A; \omega) = (\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes \hat{O}_A)) \neq 0$, and where we used the explicit order of tensor products $\mathcal{P} = \mathcal{P}_A \otimes \mathcal{P}_B$. We emphasise that the conditional success probability is operationally determined by means of *post-selection*, i.e., the *selection* of those members of the ensemble of the combined experiment, which yielded a positive answer to A ’s measurement. This requires access to the experimental data of both B and A and can consequently only be performed in their joint future.

Having found the conditional success probability, we again view it as a justification for postulating the following conditional expectation as an answer to the question in the case where O_B is not necessarily an effect but a general observable:

$$E_{\{A, B\}}(O_B | O_A; \omega) = \frac{\omega(\varepsilon_{\{A, B\}}(O_A \otimes O_B))}{\omega(\varepsilon_A(O_A))}. \quad (37)$$

Let us investigate Eq. (37) further in the following cases:

1. $A \leq B$: Using Eq. (23) we observe that

$$\omega(\varepsilon_{\{A, B\}}(O)) = (\omega \otimes \sigma_A)(\Theta_A(\varepsilon_B(O_B) \otimes O_A)), \quad (38)$$

which allows us to write

$$\begin{aligned} E_{\{A, B\}}(O_B | O_A; \omega) &= \frac{(\omega \otimes \sigma_A)(\Theta_A(\varepsilon_B(O_B) \otimes O_A))}{(\omega \otimes \sigma)(\Theta(\mathbb{1} \otimes \hat{O}_A))} \\ &= \frac{(\omega \otimes \sigma_A)(\Theta_A(\varepsilon_B(O_B) \otimes O_A))}{(\omega \otimes \sigma_A)(\Theta_A(\mathbb{1} \otimes O_A))} \\ &=: \omega_{A|O_A}(\varepsilon_B(O_B)) \\ &= E_B(O_B; \omega_{A|O_A}), \end{aligned} \quad (39)$$

in terms of the *selective* update map

$$\mathcal{J}_{A|O_A}(\omega) := \omega_{A|O_A}, \quad (40)$$

which yields a well-defined state provided that O_A has nonzero success probability $(\omega \otimes \sigma_A)(\Theta_A(\mathbb{1} \otimes O_A)) \neq 0$ [7].

Equation (39) shows how Eq. (37) can be understood in terms of an updated state in the case where there exists a causal order such that A precedes B and also constitutes a proof of Eq. (3). Additionally note that $\mathcal{J}_{A|\mathbb{1}} = \mathcal{J}_A$.

2. $B \leq A$: The interpretation of this scenario is that B performs post-selection on a measurement that (at least with respect to one causal order) *succeeds* his own. There is *a priori* no reason to expect this post-selection to be trivial and we have not found any simplified expression for $E_{\{A,B\}}(O_B|O_A;\omega)$ in this case. One might naively think that such a post-selection conflicts causality as there is an *apparent* influence from the future to the past. This issue is resolved by reminding oneself that post-selection can only be performed by all observers together in their joint future.

For completeness we mention that it was shown in Theorem 3.4 in [7] how in the case of spacelike separation, any possible *apparent* acausal behaviour of the selective update map can be attributed to spacelike correlations of the initial state ω . To see this we observe that for spacelike separated A, B

$$\begin{aligned} \hat{\Theta}_A \circ \hat{\Theta}_B(\mathbb{1} \otimes O_A \otimes O_B) \\ &= \hat{\Theta}_A \circ \hat{\Theta}_B(\mathbb{1} \otimes O_A \otimes \mathbb{1} \cdot \mathbb{1} \otimes \mathbb{1} \otimes O_B) \\ &= \left(\hat{\Theta}_A \circ \hat{\Theta}_B(\mathbb{1} \otimes O_A \otimes \mathbb{1}) \right) \cdot \left(\hat{\Theta}_A \circ \hat{\Theta}_B(\mathbb{1} \otimes \mathbb{1} \otimes O_B) \right) \\ &= \left(\hat{\Theta}_A \circ \hat{\Theta}_B(\mathbb{1} \otimes O_A \otimes \mathbb{1}) \right) \cdot \left(\hat{\Theta}_B \circ \hat{\Theta}_A(\mathbb{1} \otimes \mathbb{1} \otimes O_B) \right) \\ &= (\Theta_A(\mathbb{1} \otimes O_A) \otimes \mathbb{1}) \cdot (\Theta_B(\mathbb{1} \otimes O_B) \otimes_2 \mathbb{1}), \end{aligned} \quad (41)$$

which (cf. Sec. 3.2 in [7]) implies that

$$\omega(\varepsilon_{\{A,B\}}(O_A \otimes O_B)) = \omega(\varepsilon_A(O_A)\varepsilon_B(O_B)). \quad (42)$$

This shows that for spacelike separated observers

$$E_{\{A,B\}}(O_B|O_A;\omega) = E_B(O_B;\omega) \quad (43)$$

if and only if $\omega(\varepsilon_A(O_A)\varepsilon_B(O_B)) = \omega(\varepsilon_A(O_A))\omega(\varepsilon_B(O_B))$, i.e., $\varepsilon_A(O_A)$ and $\varepsilon_B(O_B)$ are uncorrelated in state ω .

The generalisation to N observers in the case where Obs admits a causal order \leq such that B is the *latest* observers (with respect to \leq) follows immediately: We look at the partition $A := \{X \in \text{Obs} | X < B\}$ and $\{B\}$ and wish to condition B's expected outcome of a measurement of the probe-effect O_B on the successful measurement of the probe-effects O_X for $X \in A$. Setting $O_A := \bigotimes_{X \in A} O_X$ yields just as before

$$E_{\text{Obs}}(O_B|O_A;\omega) = \omega_{A|O_A}(\varepsilon_B(O_B)) = E_B(O_B;\omega_{A|O_A}). \quad (44)$$

It is noteworthy that the total selective update map $\mathcal{J}_{A|O_A}(\omega)$ can be written as a composition of individual selective update maps

$$\mathcal{J}_{A|O_A}(\omega) = (\mathcal{J}_{|A|} \circ \cdots \circ \mathcal{J}_1)(\omega), \quad (45)$$

where we again ordered the constituent observers of A as $A_1 < \cdots < A_{|A|}$ and used the short-hand notation $\mathcal{J}_r := \mathcal{J}_{A_r|O_{A_r}}$. This follows by induction from the case $|A| = 2$ given by Theorem 3.5 in [7].

VII. ABSENCE OF IMPOSSIBLE MEASUREMENTS FOR MULTIPLE OBSERVERS

In this section we demonstrate the absence of any acausal influence in the measurements of an arbitrary finite number of causally orderable observers in a theory respecting causal factorisation.

To that end let us reconsider the situation of N observers Obs. As in the previous section, we focus our attention on a fixed observer B, taking the role played by Charlie in Sec. IV, and a fixed linear order \leq . As showed before in Eq. (33), B's expected outcome $E_{\text{Obs}}(O_B;\omega)$ equals $\omega_A(\varepsilon_B(O_B))$ in the absence of any post-selection on results of any other observers. Let us assume that there is an observer $Y \in A$ who is spacelike separated from B, i.e., $K_B \subseteq K_Y^\perp$, and will play the role of Alice. This gives rise to the partition $A = X \cup \{Y\} \cup Z$, where $X := \{J \in A | J < Y\}$ and $Z := \{J \in \text{Obs} | J > Y\}$. The super-observer Z will play the role of Bob. We can then write

$$\omega_A(\varepsilon_B(O_B)) = (\omega_X \otimes \sigma_Y \otimes \sigma_Z)(\hat{\Theta}_Y \circ \hat{\Theta}_Z(\varepsilon_B(O_B) \otimes \mathbb{1})). \quad (46)$$

The following holds:

Theorem 5. *If, in the above notation, K_B is connected and spacelike separated from K_Y , then:*

$$\begin{aligned} (\omega_X \otimes \sigma_Y \otimes \sigma_Z)(\hat{\Theta}_Y \circ \hat{\Theta}_Z(\varepsilon_B(O_B) \otimes \mathbb{1})) \\ = (\omega_X \otimes \sigma_Z)(\Theta_Y(\varepsilon_B(O_B) \otimes \mathbb{1})). \end{aligned} \quad (47)$$

A consequence of this theorem is that

$$\omega_A(\varepsilon_B(O_B)) = \omega_{A \setminus \{Y\}}(\varepsilon_B(O_B)), \quad (48)$$

and hence

$$E_{\text{Obs}}(O_B;\omega) = E_{A \setminus \{Y\}}(O_B;\omega), \quad (49)$$

emphasising that we can completely ignore the spacelike separated observer Y as well as the super-observer C succeeding B with respect to \leq .

It follows by successive application of the theorem that no observer that is spacelike separated from B can influence the expected outcome of observer B's measurement. This shows that there is no Sorkin-type (or any other) superluminal signalling between the individuals in the N observer case if each coupling zone is connected. We remark that the need to restrict to connected coupling zones comes from the connectedness condition in the formulation of the Haag property. If connectedness is dropped from the Haag property (cf. footnote [13]), then one can also drop the connectedness

in the above theorem.

The proof of Theorem 5 relies on the geometrical Lemma 6 and an application of Theorem 2.

Lemma 6. *For every connected compact subset $K \subseteq M$ there exists a connected region $N \supseteq K$ with compact closure.*

Proof. A subset with compact closure is called *precompact*. As a smooth manifold, M has an exhaustion by countably many precompact open sets G_α such that $\overline{G_\alpha} \subseteq G_{\alpha+1}$ and $M = \bigcup_\alpha G_\alpha$. Since K is compact, it can be covered by finitely many G_α and since they are nested, there exists β such that $K \subseteq G_\beta$. Since M is globally hyperbolic, the causal hull of G_β , $\text{ch}(G_\beta)$, is open (see Lemma A.8 in [14]) and $\text{ch}(\overline{G_\beta})$ is compact (by definition of global hyperbolicity). Since obviously $G_\beta \subseteq \overline{G_\beta}$, it also follows that $\text{ch}(G_\beta) \subseteq \text{ch}(\overline{G_\beta})$ (as $J^\pm(G_\beta) \subseteq J^\pm(\overline{G_\beta})$) and $\text{ch}(G_\beta) = J^-(G_\beta) \cap J^+(G_\beta)$. So $\text{ch}(G_\beta)$ is a subset of a compact set and hence precompact. As $\text{ch}(G_\beta)$ can be viewed as a globally hyperbolic manifold in its own right, every connected component is hence precompact, connected, open and causally convex. Since K is connected and contained in $\text{ch}(G_\beta)$, it is contained in one connected component, which finishes the proof. \square

Let us now prove Theorem 5.

Proof of Theorem 5. By assumption, $K_B \subseteq K_Y^\perp$ and from the existence of the causal order $K_B \subseteq M_Z^+ = M \setminus J^-(K_Z)$, and since K_B is connected, it is contained in one connected component of the open, causally convex subset $K_Y^\perp \cap M_Z^+$. This connected component can be viewed as a globally hyperbolic manifold in its own right and hence we can apply Lemma 6 to furnish a connected region $N \subseteq K_Y^\perp \cap M_Z^+$ that contains K_B and has compact closure fully contained in K_Y^\perp . Now $\varepsilon_B(O_B)$ is localisable in N , moreover, after identifying $K_1 := K_Y$, $K_2 := K_Z$ and $O_3 := N$, we see that the assumptions of Theorem 2 are fulfilled, thus establishing the desired equality. \square

VIII. CONCLUSIONS AND OUTLOOK

The issue of measurement in QFT has been plagued by acausality exemplified by Sorkin's protocol. Our main result shows that a consistent and fully causal interpretation of tripartite measurement processes in the sense of measurement schemes is possible via the local and covariant proposal in [7], which is applicable

to generic quantum field theories coupled to external forces and on possibly curved spacetimes. The principle of *causal factorisation* of scattering processes for an arbitrary finite number of causally orderable observers allowed us to generalise our result to the N observer case. As opposed to other work, such as [19], our result thereby provides a class of “physically allowed operations” that can be characterised abstractly *as well as* constructed explicitly in specific models, see [7].

The FV-framework may be considered a first important step towards a solution to the problem of delineating *all* ‘physically allowed quantum operations’ raised in [3]; however, whether all of them are induced by FV-measurement schemes is unknown. It is therefore important to more explicitly characterize the system observables associated to measurement schemes. We intend to report on this issue elsewhere. It is also worth noting that local scattering operators, understood as operations reflecting the result of measuring observables, have recently been proposed as a new foundation for AQFT [18] and this viewpoint could be fruitfully combined with ours.

For our key assumption of causal factorization, we have restricted ourselves to a physical motivation and formulation of this assumption. A more rigorous, mathematical investigation would be very interesting and we intend to report on this elsewhere.

Finally it is worth mentioning that *non-relativistic, non-local* particle detector models are a very common tool and widely used for example in quantum field theory in curved spacetime and relativistic quantum information. Other authors have shown that coupling such a detector model to a finite number of field modes [2] or to all but the zero mode [20] leads to superluminal signalling. In view of our result, this is due to the non-locality of such a coupling, whereas the detector model sketched at the end of [7] consequently does not signal superluminally. Applying the FV-framework to questions in which particle detector models have so far been used, for example *entanglement harvesting* [21], promises to yield additional insight both on a conceptual level and with respect to explicitly computable results.

ACKNOWLEDGMENTS

We thank Roger Colbeck for useful comments on the text and Ian Jubb and Leron Borsten for stimulating discussions on the general topic of impossible measurements.

[1] R. D. Sorkin, Impossible measurements on quantum fields, in *Directions in general relativity: Proceedings of the 1993 International Symposium, Maryland*, Vol. 2 (Cambridge University Press, Cambridge, 1993) pp. 293–

305.

[2] D. M. T. Benincasa, L. Borsten, M. Buck, and F. Dowker, Quantum information processing and relativistic quantum fields, *Classical Quantum Gravity*

- 31**, 075007 (2014).
- [3] D. Beckman, D. Gottesman, A. Kitaev, and J. Preskill, Measurability of Wilson loop operators, *Phys. Rev. D* **65**, 065022 (2002).
 - [4] R. D. Sorkin, Forks in the road, on the way to quantum gravity, *Internat. J. Theoret. Phys.* **36**, 2759 (1997).
 - [5] F. Dowker, Useless qubits in “relativistic quantum information” (2011), arXiv:1111.2308 [quant-ph].
 - [6] P. Busch, P. Lahti, J.-P. Pellonpää, and K. Ylinen, *Quantum measurement*, Theoretical and Mathematical Physics (Springer, Cham, 2016).
 - [7] C. J. Fewster and R. Verch, Quantum fields and local measurements, *Commun. Math. Phys.* **378**, 851 (2020), arXiv:1810.06512v3 [math-ph].
 - [8] C. J. Fewster, A generally covariant measurement scheme for quantum field theory in curved spacetimes, in *Progress and Visions in Quantum Theory in View of Gravity*, edited by F. Finster, D. Giulini, J. Kleiner, and J. Tolksdorf (Birkhäuser, Basel, 2020) arXiv:1904.06944 [gr-qc].
 - [9] R. Haag, *Local quantum physics*, 2nd ed., Texts and Monographs in Physics (Springer-Verlag, Berlin, 1996).
 - [10] C. J. Fewster and K. Rejzner, Algebraic Quantum Field Theory – an introduction, in *Progress and Visions in Quantum Theory in View of Gravity*, edited by F. Finster, D. Giulini, J. Kleiner, and J. Tolksdorf (Birkhäuser, Basel, 2020) arXiv:1904.04051 [hep-th].
 - [11] N. Drago and V. Moretti, The notion of observable and the moment problem for $*$ -algebras and their gns representations, *Letters in Mathematical Physics* **110**, 1711–1758 (2020).
 - [12] C. J. Fewster and R. Verch, Dynamical locality and covariance: What makes a physical theory the same in all spacetimes?, *Annales Henri Poincaré* **13**, 1613–1674 (2012).
 - [13] The stated Haag property could be strengthened to incorporate all (possibly disconnected) regions L containing K . However it might then conflict with a further desirable property called *additivity*, see p. 146 in [9].
 - [14] C. J. Fewster and R. Verch, Dynamical locality and covariance: what makes a physical theory the same in all spacetimes?, *Ann. Henri Poincaré* **13**, 1613 (2012).
 - [15] A. N. Bernal and M. Sánchez, On smooth Cauchy hypersurfaces and Geroch’s splitting theorem, *Communications in Mathematical Physics* **243**, 461–470 (2003).
 - [16] R. Geroch, Domain of dependence, *J. Mathematical Phys.* **11**, 437 (1970).
 - [17] K. Rejzner, *Perturbative algebraic quantum field theory*, Mathematical Physics Studies (Springer, Cham, 2016).
 - [18] D. Buchholz and K. Fredenhagen, A C^* -algebraic approach to interacting quantum field theories, *Communications in Mathematical Physics* 10.1007/s00220-020-03700-9 (2020).
 - [19] L. Borsten, I. Jubb, and G. Kells, Impossible measurements revisited (2019), arXiv:1912.06141 [quant-ph].
 - [20] E. Tjoa and E. Martín-Martínez, Zero mode suppression of superluminal signals in light-matter interactions, *Phys. Rev. D* **99**, 065005 (2019).
 - [21] A. Pozas-Kerstjens and E. Martín-Martínez, Harvesting correlations from the quantum vacuum, *Phys. Rev. D* **92**, 064042 (2015).