



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/169765/>

Version: Published Version

---

**Article:**

Gambacurta, Danilo, Colo, Gianluca and Pastore, Alessandro (2020) Nuclear incompressibility from spherical and deformed nuclei. *Journal of Physics: Conference Series*. 012129. ISSN: 1742-6596

<https://doi.org/10.1088/1742-6596/1643/1/012129>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

PAPER • OPEN ACCESS

## Nuclear incompressibility from spherical and deformed nuclei

To cite this article: Danilo Gambacurta *et al* 2020 *J. Phys.: Conf. Ser.* **1643** 012129

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

# Nuclear incompressibility from spherical and deformed nuclei

Danilo Gambacurta<sup>a</sup>, Gianluca Colò<sup>b,c</sup>, Alessandro Pastore<sup>d</sup>

<sup>a</sup>Extreme Light Infrastructure - Nuclear Physics (ELI-NP), Horia Hulubei National Institute for Physics and Nuclear Engineering, 30 Reactorului Street, RO-077125 Magurele, Jud. Ilfov, Romania

<sup>b</sup>Dipartimento di Fisica “Aldo Pontremoli”, Università degli Studi di Milano, 20133 Milano, Italy

<sup>c</sup>INFN, Sezione di Milano, 20133 Milano, Italy

<sup>d</sup>Department of Physics, University of York, Heslington, York, YO10 5DD, UK

E-mail: Gianluca.Colo@mi.infn.it

**Abstract.** We present an analysis based on the deformed Quasi Particle Random Phase Approximation, on top of a deformed Hartree-Fock-Bogoliubov description of the ground state, aimed at studying the isoscalar monopole and quadrupole response in a deformed nucleus. This analysis is motivated by the need of understanding the coupling between the two modes and how it might affect the extraction of the nuclear incompressibility from the monopole distribution. After discussing this motivation, we present the main ingredients of our theoretical framework, and we show some results obtained with the SLy4 and SkM\* interactions for the nucleus <sup>24</sup>Mg.

## 1. Introduction

The nuclear Equation of State (EoS) plays a fundamental role in the description and understanding of several properties of atomic nuclei and astrophysical compact objects [1]. However, its knowledge is, from a quantitative point of view, far from being sufficiently enough well determined. One of the key quantities needed to better constrain the EoS is the nuclear matter incompressibility, *e.g.* the curvature of the symmetric nuclear matter EoS, which is closely related to the compressional modes of atomic nuclei [2].

The emergence of collective excitations is a common feature of many-body systems. In atomic nuclei, the most collective excitations are the Giant Resonances (GRs), described as a coherent motion to which many nucleons participate [3]. The Isoscalar Giant Monopole Resonance (ISGMR), known also as the breathing mode, is a typical example of collective excitation whose centroid energy is related to the nuclear incompressibility. The properties of GRs are also considerably affected by the shape of the nucleus and in particular by its deformation. A very well known example is the Isovector Giant Dipole Resonances, whose strength distribution is split in two peaks, corresponding to a vibration along or perpendicular to the symmetry axis. The distance in energy between these two peaks is related to the corresponding deformation of the nucleus [4].

Deformation also affects the ISGMR and we aim at discussing such effects in view of existing and future experimental results on deformed nuclei. The available measurements of the ISGMR done in closed-shell nuclei (mainly <sup>208</sup>Pb) tend to point toward a value of nuclear compressibility




Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.


of  $K_\infty \approx 240 \pm 20$  MeV [5], while other measurements done in open-shell nuclei point to lower values of the nuclear matter incompressibility and the effect of deformation has still to be investigated.

In the case of the ISGMR, deformation produces a coupling of the  $K = 0$  component of the Isoscalar Giant Quadrupole Resonance (ISGQR) with the ISGMR itself,  $K$  being the projection of the total angular momentum on the symmetry axis [6]. How this coupling might affect the extraction of the nuclear incompressibility is still to be investigated and understood. In this work, we show some preliminary results in this direction.

## 2. Methodology and Results



**Figure 1.** HFB potential energy surface, obtained with the SkM\* and SLy4 interactions, as a function of the axial deformation parameter  $\beta$ , defined in Eq. (2), for  $^{24}\text{Mg}$ .



**Figure 2.** Monopole strength functions obtained for the SLy4 force. Red solid (black dashed) lines correspond to the deformed (spherical) solution.

We briefly summarise the main features of our calculations, and for more details we refer to Ref. [7]. The first step in our methodology consists in solving the Skyrme-Hartree-Fock-Bogoliubov (HFB) equations for an axially deformed system in a finite deformed harmonic oscillator basis by using the HFBTHO code [8]. A density-dependent pairing delta interaction, that is,

$$V_{pair}(\mathbf{r}, \mathbf{r}') = V_0 \left[ 1 + \gamma \frac{\rho(\mathbf{r})}{\rho_c} \right] \delta(\mathbf{r} - \mathbf{r}') , \quad (1)$$

is used in the particle-particle channel. The values of the parameters are  $V_{pair} = -280$  MeV  $\text{fm}^3$ ,  $\rho_c = 0.16 \text{fm}^{-3}$  and  $\gamma = 0.5$ . In the particle-hole channel we use a standard Skyrme functional [9].

The total energy is minimized under the constraint of a fixed quadrupole moment to which we associate the  $\beta$  parameter defined as

$$\beta = \sqrt{\frac{\pi}{5}} \frac{\langle \hat{Q} \rangle_n + \langle \hat{Q} \rangle_p}{\langle r^2 \rangle_n + \langle r^2 \rangle_p} , \quad (2)$$

where  $\langle \hat{Q} \rangle_q$  is the average value of the quadrupole-moment operator for protons ( $p$ ) and neutrons ( $n$ ). This step allows to determine the  $\beta$  value for which the minimal energy of the

system is obtained. The potential energy curves for the SkM\* [10] and SLy4 [11] interactions are shown in Fig. 1 as a function of  $\beta$ . In both cases we see a minimum corresponding to the prolate deformation  $\beta \approx 0.4$ . For this particular choice of interactions, we observe that the solution for the ground state is not superfluid.

The next step consists in calculating the nuclear response by using the Quasi Particle Random Phase Approximation (QRPA) approach [12]. The QRPA calculation is carried out on top of the HFB solution corresponding to the absolute minimum of the potential energy. The QRPA excited states  $|\lambda\rangle$  are described in terms of the excitation operators

$$Q_\lambda^\dagger = \sum_{K < K'} \left( X_{KK'}^\lambda \alpha_K^\dagger \alpha_{K'}^\dagger - Y_{KK'}^\lambda \alpha_{K'} \alpha_K \right), \quad (3)$$

normalized as

$$\sum_{K < K'} \left( (X_{KK'}^\lambda)^2 - (Y_{KK'}^\lambda)^2 \right) = 1, \quad (4)$$

where  $\alpha_K^\dagger$ ,  $\alpha_K$  are the quasiparticle creation and annihilation operators, respectively, and the pairs are such as  $\Omega = \Omega_K + \Omega_{K'}$  and  $\pi = \pi_K \cdot \pi_{K'}$ . Here,  $\Omega$  is the projection of the angular momentum on the symmetry axis  $z$  and  $\pi$  is the parity. The  $X$  and  $Y$  amplitudes together with the energies of the excited states are obtained by solving the QRPA eigenvalue problem in the matrix form

$$\begin{pmatrix} A & B \\ -B^* & -A^* \end{pmatrix} \begin{pmatrix} X^\lambda \\ Y^\lambda \end{pmatrix} = \omega_\lambda \begin{pmatrix} X^\lambda \\ Y^\lambda \end{pmatrix}. \quad (5)$$

We will focus here on the isoscalar quadrupole and monopole strengths whose transition operators are defined as

$$\hat{F}_{J=2,K}^{IS} = \frac{eZ}{A} \sum_{i=1}^A r_i^2 Y_{2K}(\hat{r}_i), \quad \hat{F}_{J=0,K=0}^{IS} = \frac{eZ}{A} \sum_{i=1}^A r_i^2. \quad (6)$$

In Fig. 2 we compare the QRPA results for the SLy4 force, obtained at the deformed minimum  $\beta \approx 0.4$  and at the spherical limit  $\beta = 0$ . For sake of comparison, we plot the folded monopole strength functions

$$S(E) = \sum_\lambda \frac{\Gamma/2}{\pi} \frac{|\langle \lambda | \hat{F} | 0 \rangle|^2}{(E - E_\lambda)^2 + \Gamma^2/4}, \quad (7)$$


where  $\Gamma = 2$  MeV has been used as a smoothing parameter. The effect of the deformation is clearly seen, as it leads to a splitting of the main peak obtained in the spherical solution, and to the two peaks observed in the deformed case.

For the sake of further analysis, we plot in Figs. 3 and 4 the unfolded QRPA strength distributions corresponding to isoscalar monopole and quadrupole operators, in the upper and lower panels, respectively. For both interactions, the QRPA calculations are performed at the deformed minimum shown in Fig. 1. The monopole strength shows, in both cases, two main peaks located at 17.76 and 24.29 MeV for SLy4 and 17.62 and 24.03 MeV for SkM\*. The lower state is also excited by the quadrupole operator as it can be seen in the lower panels of the two figures. The Energy Weighted Sum Rules are preserved within less than 1% [13].


In order to have a deeper insight on the nature of these states, we analyze below the transition densities

$$\rho_\lambda(\vec{r}) = \delta\rho_\lambda(r, z) e^{-iK\phi}, \quad (8)$$

where  $K$  is the total angular momentum projection of the excited state and  $r, z$  and  $\phi$  are the cylindrical coordinates.



**Figure 3.** Monopole (upper panel) and quadrupole (lower panel) strength distribution obtained for the SLy4 interaction.



**Figure 4.** Monopole (upper panel) and quadrupole (lower panel) strength distribution obtained for the SkM\* interaction.


In Figs. 7 and 8, we display the neutron and proton transition densities obtained in the SLy4 case for the state at 17.76 MeV, while in Figs. 5 and 6 we show the neutron and proton transition densities for the state located at 24.29 MeV.

For both states, we observe the expected behaviour for isoscalar excitations, *i.e.* neutrons and protons oscillating in phase. In the case of the monopole excitation (Figs 5 and 6) the oscillation is along the radial axis, while in the the quadrupole case (Figs 7 and 8) the nucleons oscillate along the axial  $z$  axis. For comparison, we plot also, in Figs. 9 and 10, the transition densities of a non collective state ( $E \approx 25.46$  MeV) that clearly shows a different pattern, characterized by several nodes. Similar results have been obtained using the SkM\* interaction.


### Conclusions and Outlook

We have presented some results, based on a deformed QRPA approach, for the monopole and quadrupole strength distributions in the deformed nucleus  $^{24}\text{Mg}$ . Our study is motivated by the need of understanding the compressional properties of deformed nuclei. To this aim, we wish to disentangle the compressional properties associated with the ISGMR, from its coupling with the  $K = 0$  component of the ISGQR.


Our results show clearly the splitting of the monopole strength due to deformation. Moreover, the transition densities of the collective states have been shown to be very instrumental to indicate the character of states (compressional or not). In the near future, we plan to implement




**Figure 5.** Neutron transition density  $\delta\rho_\lambda(r, z)$  for the excited state located at 24.29 MeV (see Fig. 3).



**Figure 6.** Proton transition density  $\delta\rho_\lambda(r, z)$  for the excited state located at 24.29 MeV (see Fig. 3).



**Figure 7.** Neutron transition density  $\delta\rho_\lambda(r, z)$  for the excited state located at 17.76 MeV (see Fig. 3).




**Figure 8.** Proton transition density  $\delta\rho_\lambda(r, z)$  for the excited state located at 17.76 MeV (see Fig. 3).


a projection on the total angular momentum of the QRPA phonons. The projection would in fact allow for a clear decoupling the two modes, providing a cleaner way to extract the monopole peak energy. Work in this direction is on going.

### Acknowledgments

This work has been partially supported by STFC Grant No. ST/P003885/1. Funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 654002 is also acknowledged.



**Figure 9.** Neutron transition density  $\delta\rho_\lambda(r, z)$  for the (non collective) excited state located at 25.46 MeV (see Fig. 3).



**Figure 10.** Proton transition density  $\delta\rho_\lambda(r, z)$  for the (non collective) excited state located at 25.46 MeV (see Fig. 3).

## References

- [1] Oertel M, Hempel M, Klähn T and Typel S 2017 *Rev. Mod. Phys.* **89**(1) 015007
- [2] Blaizot J P 1980 *Physics Reports* **64** 171–248
- [3] Harakeh M N and van der Woude A 2001 *Giant Resonances Fundamental High-Frequency Modes of Nuclear Excitation* (Oxford University Press, New York)
- [4] Martini M, Péru S, Hilaire S, Goriely S and Lechaftois F 2016 *Physical Review C* **94** 014304
- [5] Garg U and Colò G 2018 *Progress in Particle and Nuclear Physics* **101** 55–95
- [6] Ragnarsson I and Nilsson S G 2005 *Shapes and shells in nuclear structure* (Cambridge University Press)
- [7] Losa C, Pastore A, Døssing T, Vigezzi E and Broglia R A 2010 *Physical Review C* **81** 064307
- [8] Stoitsov M, Dobaczewski J, Nazarewicz W and Ring P 2005 *Computer physics communications* **167** 43–63
- [9] Bender M, Heenen P H and Reinhard P G 2003 *Reviews of Modern Physics* **75** 121
- [10] Bartel J, Quentin P, Brack M, Guet C and Håkansson H B 1982 *Nuclear Physics A* **386** 79–100
- [11] Chabanat E, Bonche P, Haensel P, Meyer J and Schaeffer R 1997 *Nuclear Physics A* **627** 710–746
- [12] Suhonen J 2007 *From nucleons to nucleus: concepts of microscopic nuclear theory* (Springer Science & Business Media)
- [13] Lipparini E and Stringari S 1989 *Physics Reports* **175** 103–261