



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/170339/>

Version: Accepted Version

Article:

Saegeman, V., Cohen, M.C., Burton, J.L. et al. (2021) Microbiology in minimally invasive autopsy: best techniques to detect infection. ESGFOR (ESCMID study group of forensic and post-mortem microbiology) guidelines. *Forensic Science, Medicine and Pathology*, 17 (1). pp. 87-100. ISSN: 1547-769X

<https://doi.org/10.1007/s12024-020-00337-x>

This is a post-peer-review, pre-copyedit version of an article published in *Forensic Science, Medicine, and Pathology*. The final authenticated version is available online at: <http://dx.doi.org/10.1007/s12024-020-00337-x>.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Microbiology in minimally invasive autopsy: Best techniques to detect infection. ESGFOR (ESCMID Study Group of Forensic and Post-mortem Microbiology) guidelines

Veroniek Saegeman¹, Marta C. Cohen², Julian L. Burton³, Miguel J. Martinez^{4,5}, Natalia Rakislova^{4,6}, Amaka C Offiah⁷, Amparo Fernandez-Rodriguez⁸ #.

On behalf of ESGFOR (working group of the ESCMID –European Society of Clinical Microbiology and Infectious Diseases- Study Group of Forensic and Post-mortem Microbiology)

1. Clinical Laboratory, AZ Nikolaas, Moerlandstraat 1, 9100 Sint-Niklaas, and Infection Control Department, University Hospitals Leuven, Herestraat 49, 3000 Leuven, Belgium. ORCID ID: 0000-0003-0439-0205

2. Sheffield Children’s Hospital NHS FT. Histopathology Department. Western Bank S10 2TH. Sheffield UK. #: Link person between ESCMID and the European Society of

Pathology (ESP). ORCID ID: orcid.org/0000-0001-5534-444X

3. The Medico-Legal Centre, Watery Street, Sheffield, UK.

4. ISGlobal, Barcelona Centre for International Health Research (CRESIB),

5. Department of Microbiology, Hospital Clínic, Universitat de Barcelona, Barcelona, Spain.

6. Department of Pathology, Hospital Clínic, Universitat de Barcelona, Barcelona, Spain.

7. Academic Unit of Child Health, Department of Oncology and Metabolism, University of Sheffield, Sheffield, UK and Department of Radiology, Sheffield Children’s NHS FT, Sheffield, UK ORCID ID: 0000-0001-8991-5036

8. Microbiology Laboratory, Biology Department, Instituto Nacional de Toxicología y Ciencias Forenses, Las Rozas de Madrid, Madrid, Spain. ORCID ID: 0000-0002-2992-3226

*Corresponding Author: Amparo Fernandez-Rodriguez.

Telephone: 34-91-768-89-44

E mail: amparo.fernandezrodriguez@justicia.es

Word count: 3226

Article classification:

10: Forensics

10.020: Medicine

10.190: Autopsy

1 **Microbiology in minimally invasive autopsy: Best techniques to detect infection.**
2 **ESGFOR (ESCMID Study Group of Forensic and Post-mortem Microbiology)**
3 **guidelines**

4
5
6
7 **Abstract**

8
9 *Purpose*

10 This manuscript aims to: 1) provide specific guidelines on PMM techniques in the
11 setting of Minimally Invasive Autopsy (MIA), both for pathologists collecting samples
12 and for microbiologists advising pathologists and interpreting the results and 2)
13 introduce standardization in PMM sampling at MIA
14
15

16 *Methods*

17
18 Post-mortem microbiology (PMM) is crucial to identify the causative organism in
19 deaths due to infection. MIA including the use of post-mortem (PM) computed
20 tomography (CT) and PM magnetic resonance imaging (MRI), is increasingly carried
21 out as a complement or replacement for the traditional PM. In this setting, mirroring the
22 traditional autopsy, PMM aims to: detect infectious organisms causing sudden
23 unexpected deaths; confirm clinically suspected but unproven infection; evaluate the
24 efficacy of antimicrobial therapy; identify emergent pathogens; and recognize medical
25 diagnostic errors. Meaningful interpretation of PMM results requires careful evaluation
26 in the context of the clinical history, macroscopic and microscopic findings.
27
28

29 This guidelines were developed by a multidisciplinary team with experts in various
30 fields of microbiology and pathology on behalf of the ESGFOR (ESCMID – European
31 Society of Clinical Microbiology and Infectious Diseases - Study Group of Forensic and
32 Post-mortem Microbiology, in collaboration with the ESP -European Society of
33 Pathology-) based on literature search and the author's expertise.
34
35

36 *Results*

37
38 Microbiological sampling methods for MIA are presented for various scenarios: adults,
39 children, developed and developing countries. Concordance between MIA and
40 conventional invasive autopsy is substantial for children and adults and moderate for
41 neonates and maternal deaths.
42
43

44 *Conclusion*

45 Networking and closer collaboration among microbiologists and pathologists is vital to
46 maximize the yield of PMM in MIA.
47
48

49 **Keywords:** traditional autopsy, forensic sampling, infection, forensic microbiology.
50 minimally invasive autopsy. post-mortem microbiology
51
52
53
54

55 **Introduction**

56
57
58 Post-mortem microbiology (PMM) is crucial to identify the causative organism in
59 deaths due to infection. Although the traditional autopsy remains the gold standard
60
61
62
63
64
65

1 procedure to obtain samples for PMM, the lack of trained human resources, cultural and
2 religious constraints amongst others, have led to the introduction of Minimally Invasive
3 Autopsy (MIA) techniques as an alternative [1]. In some cases, MIA includes the
4 procurement of tissue samples, also known as Minimally Invasive Tissue Sampling
5 (MITS), using fine needles to collect small amounts of tissue from key organs and body
6 fluids. Currently, MIA is increasingly carried out as a complement or replacement for
7 the traditional complete autopsy, mostly in low- and middle-income countries (LMIC).
8 The dead human body is a complex microbial ecological system that varies significantly
10 between individuals as a result of diet, lifestyle and geographical factors [2,3]. It is
11 therefore important to highlight that a positive result from PMM may reflect a genuine
12 infection but may also reflect sample contamination, commensal organisms and/or
13 post-mortem bacterial translocation (PMBT). The incidence of post-mortem
14 contamination and PMBT has been estimated to approximate 20% in traditional
15 autopsies where PMM has been conducted [4,5]. However, when standardized
16 procedures are adhered to, post-mortem contamination can be kept below 10% [6].

17
18
19
20
21
22 PMBT is a natural phenomenon in which endogenous commensal gut bacteria, most
23 commonly *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*,
24 *Enterococcus spp.*, Clostridia and streptococci multiply and migrate into the blood and
25 tissues. PMBT is a driver of putrefactive decomposition, which begins approximately
26 four minutes after death [2, 7-9]. Although PMBT begins within eight hours of death, it
27 is not solely dependent on the post-mortem interval (PMI). Agonal spread of
28 microorganisms throughout the body during the process of dying and resuscitation has
29 been suggested but is not universally accepted [6, 10,11]. Aerobic bacteria proliferate
30 early during the putrefactive process, to be later overtaken by anaerobes [2].

31
32
33
34
35
36 Sample contamination during collection of peripheral blood is most commonly due to
37 the introduction of coagulase-negative staphylococci from the skin [10]. In general, a
38 mixed bacterial growth likely represents sample contamination whilst single isolates are
39 generally interpreted to represent true positives [6, 11, 12]. Despite close adherence to
40 sampling protocols, a pure growth of a single microorganism from multiple sites may
41 still represent contamination rather than infection [4,10]. Yet, isolation of a pathogenic
42 organism from multiple sites at autopsy, most likely represents a true ante-mortem
43 bacteremia [4].

44
45
46
47
48
49 A prolonged PMI and putrefaction are usually perceived to impair the yield of PMM
50 [13]. Tuomisto *et al.* [14], using real-time quantitative PCR to investigate bacterial
51 migration from the gut into the blood, liver, portal vein, mesenteric lymph node, and
52 pericardial fluid of cadavers stored at 4°C, were able to demonstrate that the relative
53 amounts of intestinal bacterial DNA (bifidobacteria, *Bacteroides*, *Enterobacter* and
54 Clostridia) increased with time. However, a prolonged PMI does not uniformly increase
55 the risk of PMBT or the yield in obtaining positive post-mortem cultures [6, 15].

56
57
58 Differences in detection techniques (routine bacteriology cultures versus molecular
59 techniques) could be part of the explanation in contradictory study results.

1 Palmiere *et al.* [4] showed that bacterial cultures and real-time polymerase chain
2 reaction (qPCR) analyses were reliable, not yielding false-positive results in the
3 detection of *Neisseria meningitidis*, *Listeria monocytogenes*, *Streptococcus pneumoniae*
4 or *Haemophilus influenzae* in CSF obtained from the lateral ventricles and cisterna
5 magna in severely decomposing adults known not to have died infectious deaths.
6
7 Positive results were obtained in both bacterial culture and qPCR analysis of CSF from
8 decomposing controls with known meningitis. Also, reliable results were obtained in
9
10 other *N. meningitidis* qPCR validation studies performed in post-mortem samples [16].
11 Overall, PMBT is a frustrating “side-effect” of death. From a practical point of view,
12 refrigeration of the corpse minimizes this phenomenon [6]. There is increasing interest
13 in the possibility that exploration of successional changes in the human microbiome
14 after death using metagenomics may have a role to play in determining the PMI [3,8],
15 particularly in decomposing bodies [2, 17,18].
16
17

18
19 As PMM is not yet an extended practice in MIA, the contaminating effects of PMBT
20 haven’t been evaluated yet. In the following paragraphs, we develop in more detail
21 microbiological sampling in MIA. With careful interpretation, PMM is a useful tool to
22 identify correctly an infectious cause of death (COD). Our study aims to provide useful
23 guidelines on PMM sampling techniques in MIA, both to the pathologist requesting the
24 analysis and to the microbiologist advising the pathologist and interpreting the results.
25
26
27
28
29
30
31

32 **General post-mortem microbiology sampling technique**

33 PMM samples should be collected as soon as possible and ideally within 24 hours of
34 death [19], but it is often impossible to perform MIA in such a short time frame,
35 especially in poor settings. The collection of samples for PMM can however still be
36 fruitful [6]. However, the PMI, should always be considered along with the imaging
37 and/or macroscopic and microscopic features of the organs when interpreting the results
38 (See Table 1).
39
40
41

42 The nature of the samples to be collected varies depending on the clinical history.
43 Specific examples are considered below. In infant and childhood death without
44 symptoms, a standard set of samples is recommended in the traditional PM:
45 nasopharyngeal swab, CSF, blood, lung, spleen, heart, and bowel content. If sepsis,
46 bacteremia or other conditions are suspected, it is recommended to add samples from
47 other tissues (Table 1). [19-21]. According to the method of MIA used, the aim would
48 be to replicate these samples as much as possible. Some centers complement the
49 traditional full autopsy or the MIA MSCT (Multi-Slice-Computed Tomography), MRI
50 (Magnetic Resonance Imaging), using MITS (tissue sampling through fine needle) or
51 laparoscopic/thoracoscopic approaches which allow direct visualization and sampling of
52 organs [22, 23].
53
54
55
56
57
58

59 A variety of other conditions may cause systemic inflammatory response syndrome and
60
61
62
63
64
65

1 mimic sepsis in the absence of infection [24]. According to the context of the MIA, the
2 provision of non-microbiological samples, such as those referred for toxicological
3 screening, metabolic analysis and histopathological examination may be required.

4
5 Where samples are obtained by percutaneous needle puncture/aspiration in MITS, the
6 body must first be appropriately positioned, and the skin should be cleaned with sterile
7 water and disinfected either with alcohol-based solution containing chlorhexidine or
8 iodine. Isopropyl alcohol should only be used after samples for toxicological analysis
9 have been obtained (or the toxicologist must be notified of its use) [19]. Prior
10 post-mortem angiography via femoral cannulation does not impede the collection of
11 peripheral blood for PMM provided that the skin is disinfected prior to cannulation for
12 angiography [5].
13
14
15
16
17

18 Ideally, a separate set of sterile instruments should be used to collect each sample, in
19 order to avoid cross-contamination [13].
20
21

22 Tissue samples for microbiology should be stored in sterile containers or bottles without
23 additives, refrigerated until transport and sent to the microbiology laboratory within
24 24-48 hours after the autopsy [13,18]. It is recommended to cryopreserve additional
25 samples at -80°C for future molecular analyses in the event they are required.
26
27
28

29 Consideration must be given to the health and safety of mortuary workers during the
30 collection of PMM samples, as the decedent's microbiome may pose a risk. Specific
31 measures to protect mortuary staff from potential infectious hazards have been
32 described elsewhere [13,25]. Direct contact and puncture wounds are the most
33 important routes for transmission of blood-borne infections, and aerosol transmission is
34 mainly a risk for airborne pathogens such as *Mycobacterium tuberculosis complex* or
35 the MERS, SARS or COVID-19 coronaviruses [26]. Universal safety precautions to be
36 applied are personal protective equipment including eye protection, a surgical mask,
37 surgical gown, waterproof apron, gauntlets, surgical gloves and, optionally, a
38 cut-resistant glove on the non-dominant hand. However, when airborne pathogens are
39 anticipated, additional respiratory protection, such as a Free Flight Phase Two (FFP2) or
40 even FFP3 masks for highly contagious pathogens such as COVID-19 is required [27].
41 Where fungal growth is evident macroscopically either during the crime scene review,
42 exhumation or at autopsy, respiratory protection should also be worn due to the risk of
43 inhaling spores from *Aspergillus fumigatus* complex and *Candida albicans* [18].
44
45
46
47
48
49
50

51 **Different settings for microbiology in MIA**

52 **The adult MIA**

53 Changing societal, cultural, religious and political attitudes around the world have seen
54 the conventional invasive autopsy become less acceptable and frequent [1,28,29]. MIA
55 has gained acceptability and is a valid alternative to the traditional autopsy in the
56 investigation of natural adult deaths, especially in low-income countries [28,30,31,32,
57
58
59
60
61
62
63
64
65

33].

1
2
3
4 As in the conventional PM, MIA protocol includes a review of the available history of
5 events leading to death and a thorough external examination. Unlike the conventional
6 complete autopsy, in MIA the body cavities are not opened. Instead, key organs may be
7 sampled through percutaneous puncture using specific needles (see below under 3.3.
8 MIA in developing countries). Swabs represent another alternative to be used in
9 noninvasive post mortems. Where facilities and resources permit, non-invasive or MIAs
10 using either multi-slice computed tomography (MSCT) or MRI scans with or without
11 angiography may be used to replace or limit the conventional autopsy [31,34,35]. In
12 adults, MSCT has been shown to be superior to MRI in detecting the COD [34]. The
13 combination of post-mortem whole body MSCT scan, CT angiography and tissue
14 biopsy has been shown to be the most sensitive technique in detecting the COD when
15 MIAs are undertaken and surpass non-invasive techniques alone [35].
16
17
18
19
20
21
22

23 *Microbiology sampling during the adult MIA*

24 While MIA can reduce the need to undertake invasive autopsies in patients with known
25 or suspected Hazard Group 3 infections such as HIV or COVID-19 [31,35,36,37], very
26 few recent articles detail the collection of samples for microbiological examination in
27 adults during MIA. Previous studies have focused on the collection of biopsies for
28 histopathological examination [38,39]. In some studies, patients with known high-risk
29 infections have been actively excluded and/or tissue sampling limited to
30 histopathological examination [30,40]. Previous research has focused on the role of
31 MIA to assess the prevalence of tuberculosis and other infections in patients with known
32 HIV infection [41]. Collection of blood, CSF and samples from the lungs, liver and at
33 MIA conducted 3-6 days after death have demonstrated that tuberculosis and other
34 bacterial infections are the commonest COD in patients with advanced HIV infection).
35 Such findings concur with those from conventional invasive autopsies performed in
36 other studies [42].
37

38 Swab samples – either nasopharyngeal or oropharyngeal - for SARS-CoV-2 can be
39 taken without damage to the body, this before performing the MIA protocol [43].

40 Swabbing the upper respiratory tract, particularly the nasopharyngeal site, is relevant for
41 postmortem diagnostic of viral respiratory pathogens [44] while the throat is relevant
42 for measles [45].
43
44
45
46
47
48
49
50

51 Prior to procuring needle tissue samples, the skin should be cleaned with sterile water
52 and disinfected either with alcohol-based solution containing chlorhexidine or iodine.
53 Under the suspicion of legionellosis, it is particularly important avoiding contact with
54 tap water to prevent contamination [41]. Blood samples are collected into an
55 EDTA-containing tube, and an aerobic blood culture bottle. CSF is collected into a
56 sterile tube for analysis and stored at -80°C in Eppendorf Safe-Lock Tubes. Tissue
57 samples obtained by needle biopsy are placed both in thioglycolate broth and in lysis
58 buffer for microbiological analyses. Similarly, samples from these sites and from the
59
60

heart, spleen, kidney, uterus and skin can be collected for histopathological examination. MIA is ideally performed within 24 hours of death to minimize the effects of PMBT. MIAs performed after 24h after death can still render reliable diagnostic results, although in this situation the contribution of some microorganisms such as *Enterobacteriaceae* and *Pseudomonas* spp. might be overestimated [46] and may reflect bacterial translocation rather than true infections. This translocation may become a larger problem given the delay in performing MIA while waiting for the result of a SARS-CoV-2 PCR taken after admission of a body in the forensic pathology laboratory.

Castillo *et al.* [32] compared MIA to conventional invasive autopsy. Within 24 hours of death, the authors disinfected and sampled the skin, blood and CSF. This was followed by targeted collection of tissue using needle biopsies for microbiological and histopathological analysis from the liver, lungs, bone marrow and central nervous system [47]. The conventional autopsy was performed immediately after MIA. MIA sensitivity was highest for disseminated infections (98%, 95% Confidence Interval 87-100%) and lowest for pulmonary infections (79%, 95% CI: 58-93%) due to difficulties in targeting lesions within the lung on blind needle biopsy. MIA specificity was highest for gastrointestinal infections (100%, 95% CI 97-100%) and lowest for disseminated infections (99%, 95% CI 92-100%). The single most frequent infectious COD identified in this study was *M. tuberculosis* complex. [32,48].

In an innovative study, Van der Linden *et al.* [49] used needle biopsies under imaging guidance during MIA to collect samples from the heart, liver and kidney. The authors found that the integrity of the patient's RNA in these samples was larger than in samples subsequently obtained by conventional autopsy, though glyceraldehyde-3-phosphate dehydrogenase (GAPDH) expression was lower in samples with a PMI greater than 15 hours. Needle biopsies had also been used during MIA with CT-guidance to perform PCR analysis for the presence of bacterioplankton DNA in lungs, spleen, kidney and brain [50] as a diagnostic marker in drowning six days after the discovery of the body. Samples were collected, each using a new sterile cutting needle after cleaning the skin with alcohol and stored at 4°C in sterile containers without preservative until processing.

Protocol for the collection of microbiology samples at MIA in adults

At present, the literature reflects that there is insufficient worldwide experience to allow a definitive recommendation on how and when samples should be collected for microbiological investigation during MIA. Relatively few studies have addressed this issue, and almost all have been undertaken in areas of high HIV and tuberculosis prevalence [47,51]. Microbiological investigation during adult MIA is likely to have a lower diagnostic yield with longer PMIs and in areas of low HIV prevalence. It is clear that samples can be successfully obtained for microbiology at MIA with or without imaging-guidance. Provided these have been obtained after first appropriately disinfecting the skin, these samples can be submitted to a wide variety of microbiological investigations, including serology, culture and PCR-based analyses

(Table 2). What remains unclear is how many biopsies should be obtained from each site, whether all samples should be collected in every MIA or whether sample collection should be directed by the clinical history and what impact the PMI has on the sensitivity and specificity of such investigations. These are areas for future research.

For several BSL3 pathogens, such as HIV, *M. tuberculosis* complex and SARS-CoV-2, it is difficult to determine the duration of infectiousness of the body. Tuberculosis could be infectious for more than 12 months [52], HIV and SARS-CoV-2 may be detected by PCR for several days postmortem [53,54]. The duration of their infectiousness i.e. isolation in viral culture in dead human bodies is not known to the best of our knowledge. Therefore, it is worthwhile taking swabs or punctures for molecular analyses for these BSL3 pathogens, following stringent safety measures [26] even with postmortem intervals of more than 24 h.

The MIA in stillbirth, neonates and children

Microbiology sampling in pediatric MIA can be obtained using the protocol described above for adult patients (Table 3) [47,51,55,56]. Successful implementation of pediatric MIA into routine clinical practice requires further development and has many issues to consider in order to ensure that the best possible service is provided [22,23]. However, provided that MIA is undertaken jointly by pediatric pathologists and radiologists, it could be an acceptable alternative to conventional autopsy in selected cases [50].

Currently, the pediatric MIA uses MRI rather than CT [57, 58]. The examination of the placenta becomes an invaluable tool in the post-mortem examination of stillbirth, and in neonatal deaths the study of the placenta becomes key, providing significant information in more than 50% of the MIA cases [59].

In low/medium income countries, where resources are not available, some studies suggest using Minimally Invasive Tissue Sampling (MITS) with available clinical data, for attributing underlying and immediate causes of neonatal deaths [60].

In pediatric post-mortems swab samples for SARS-CoV-2 should add a rectal swab in addition to the nasopharyngeal swab used in adults [61].

MIA in developing countries

Most of the world's population dies without qualified medical personnel either certifying their death or identifying a specific cause. The conventional invasive autopsy is indisputably the gold standard for COD determination. However, it is rarely performed in low-income countries due to the lack of resources, trained available pathologists, poor acceptability and the fact that in these countries most of the deaths occur outside the hospital. Minimally invasive techniques have been proposed as an alternative to the traditional complete autopsy to better refine the mortality statistics in these settings and therefore contribute to effective public health interventions.

In 2013, the Barcelona's Institute for Global Health (ISGlobal) started the CaDMIA research project (validation of the MIA tool for COD investigation in developing

1 countries), funded by the Bill & Melinda Gates Foundation, aiming to validate the use
2 of a less invasive methodology for COD ascertainment. The MIA protocol is based on
3 the collection of post-mortem biopsies from key organs (central nervous system,
4 lungs, heart, liver, kidney, bone marrow, spleen and skin) and fluids (CSF and blood).
5 MIA and conventional invasive autopsy samples are studied using traditional and
6
7 advanced histopathological and microbiological techniques. Pathology and
8 microbiology results are analyzed by a multidisciplinary team in order to reach a
9 consensus regarding the most plausible COD, both for the MIA and the conventional
10 autopsy. The procedure involves the collection of 20 mL of blood and CSF and
11 puncture of liver, lungs, heart, spleen, kidneys, bone marrow and brain in all cases
12 plus the uterus in women of childbearing age, using biopsy needles [22,55,456]. The
13 type and main characteristics of the different needles used in the MIA procedure for
14 each particular biopsy, the sites of puncture and the number of samples to be obtained
15 are summarized in Table 4 [47]. The organs are presented in the order in which the
16 samples are collected.
17

18 Extensive microbiological investigations are included to identify infectious COD [51].
19 The sampling and testing scheme for microbiology includes a universal screening for
20 several key pathogens (i.e. HIV, HBV, malaria) and bacterial/fungal culture of
21 biological fluids and tissue samples. In addition to universal screening, frozen aliquots
22 of plasma, CSF or other biological fluids and tissue samples in nucleic
23 acid-preserving lysis buffer are collected for molecular assays (i.e. broad-spectrum
24 PCR assays and specific qPCR tests) guided by histopathological findings.
25 Preliminary results suggest that CSF and lung tissue are among the samples offering a
26 better performance for microbiological analysis to detect an infectious COD. The
27 MIA tool has recently been validated for neonates, children, maternal deaths and other
28 adult deaths from Mozambique [32,55,56,62]. In these validation studies, the
29 concordance between MIA and conventional complete autopsy is moderate for
30 neonates and maternal deaths ($\kappa = 0.404$ and 0.485 respectively) and substantial
31 for children and adults ($\kappa = 0.704$ and 0.732 respectively). The overall
32 concordance between the MIA diagnosis and conventional invasive autopsy was 75.9%
33 (85/112). The concordance was higher for infectious diseases and malignant tumors
34 (63/80 [78.8%] and 13/16 [81.3%], respectively) than for other diseases (9/16; 56.2%).
35 The specific microorganisms causing death were identified in the MIA in 62/74
36 (83.8%) of the infectious disease deaths with a recognized cause.
37
38
39
40
41
42
43
44
45
46
47
48
49

50 Discussion

51 PMM has shown to be useful to detect an infectious COD, though it must be interpreted
52 with caution, and in the context of the clinical history (including ante-mortem
53 microbiology) as well as imaging, macroscopic and microscopic findings [9,10]. Where
54 there is concordance between PMM, the history and the autopsy findings, it is
55 reasonable to attribute the identified microorganism as a causative factor. Where
56 concordance is lacking, it is likely that the microbiological findings are spurious and
57
58
59
60
61
62
63
64
65

represent contamination or PMBT [4].

Laboratory interpretative criteria for post-mortem bacterial culturing have been previously defined by many authors [11,12,63-68]. The possibility of sample degradation affecting the detection of nucleic acids and therefore compromising the molecular detection of pathogens should also be considered when dealing with post-mortem samples [65]. However, these recommendations are generally addressed to the interpretation of the results obtained in samples taken during a traditional autopsy.

The PMM protocols using MIA offer an alternative to better ascertain infection under specific settings. As mentioned above, some validation studies have already been performed, showing their value [32,55,56,62]. However, additional investigations in a variety of scenarios and the evaluation of how the PMI affects MIA microbiological results compared to those obtained by a traditional autopsy could be of help for a better estimation of its possible utility under different circumstances. Also, in the MIA setting, more interpretation criteria based on prospective studies including not only post-mortem cultures, but also molecular analyses are required for a better comprehension of PMM.

Acknowledgments

Conflict of Interest

There is no conflict of interest to be reported.

Word count: 3435

Key Points

- The study provides specific guidelines on post mortem microbiology (PMM) techniques in the setting of Minimally Invasive Autopsy (MIA)
- In low and middle -income countries MIA combines clinical history, external examination, Minimally Invasive Tissue Sampling (MITS) and microbiology.
- In developed countries MIA adds to the above multi slice computed tomography (MSCT) and/or Magnetic Resonance Imaging (MRI)
- Swabbing is an additional approach to be considered as part of MIA, as it is useful for SARS-CoV-2 and other respiratory viruses.
- Where there is concordance between PMM, the history, histology and/or imaging findings, it is reasonable to attribute the identified microorganism as a causative factor. Where concordance is lacking, it is likely that the microbiological findings are spurious and represent contamination or postmortem bacterial translocation

References

1. Bassat Q, Ordi J, Vila J, Ismail MR, Carrilho C, et al. Development of a post-mortem procedure to reduce the uncertainty regarding causes of death in developing countries. *The Lancet*. 2013;Sep;1(3):e125-6.
2. Finley SJ, Benbow ME, Javan GT. Microbial communities associated with human decomposition and their potential use as postmortem clocks. *Int J Legal Med*. 2015;129:623-32.

-
- 1 3. Hauther KA, Cobaugh KL, Meadows Jantz L, Sparer TE, DeBruyn JM. Estimating
2 time since death from postmortem human gut microbial communities. *J Forensic Sci.*
3 2015;60:1234-40.
 - 4 4. Palmiere C, Vanhaebost J, Ventura F, Bonsignore A, Bonetti LR. Cerebrospinal fluid
5 PCR analysis and biochemistry in bodies with severe decomposition. *J Forensic Leg*
6 *Med.* 2015;30:21-24.
 - 7 5. Palmiere C, Egger C, Grabherr S, Jatton-Ogay K, Greub G. Postmortem angiography
8 using femoral cannulation and postmortem microbiology. *Int J Legal*
9 *Med.* 2015;129:861-7.
 - 10 6. Morris JA, Harrison LM, Partridge SM. Practical and theoretical aspects of
11 postmortem bacteriology. *Curr Diagn Pathol.* 2007;13,65–74.
 - 12 7. Buchelo SR, Lynne AM. The microbiome of human decomposition. *Microbe*
13 *Magazine.* 2016;11: 165-71.
 - 14 8. Burcham ZM, Hood JA, Pechal JL, Krausz KL, Schmidt CJ et al. Fluorescently
15 labeled bacteria provide insight on post-mortem microbial transmigration. *Forensic*
16 *Sci Int.* 2016;264:63-9.
 - 17 9. Christoffersen S. The importance of microbiological testing for establishing cause of
18 death in 42 forensic autopsies. *Forensic Sci Int.* 2015;250:27-32.
 - 19 10. Palmiere C, Egger C, Prod'Hom G, Greub G. Bacterial translocation and sample
20 contamination in postmortem microbiological analyses. *J Forensic Sci.*
21 2016;61:367-74.
 - 22 11. Weber MA, Sebire NJ. Postmortem investigation of sudden unexpected death in
23 infancy: current issues and autopsy protocol. *Diagn Histopathol.*
24 2009;15(11):510-23.
 - 25 12. Lobmaier IV, Vege A, Gaustad P, Rognum TO. Bacteriological
26 investigation-significance of time lapse after death. *Eur J Clin Microbiol Infect Dis.*
27 2009;28:1191-8.
 - 28 13. Saegeman V, Cohen MC, Alberola J, Ziyade N, Farina C et al. How is post-mortem
29 microbiology appraised by pathologists? Results from a practice survey conducted
30 by ESGFOR. *Eur J Clin Microbiol Infect Dis.* 2017;Aug;36(8):1381-85.
 - 31 14. Tuomisto S, Karhunen PJ, Vuento R, Aittoniemi J, Pessi T. Evaluation of
32 postmortem bacterial migration using culturing and real-time quantitative PCR. *J*
33 *Forensic Sci.* 2013;58:910-6.
 - 34 15. Weber MA, Hartley JC, Brooke I, Lock PE, Klein NJ et al. Post-mortem interval
35 and bacteriological culture yield in sudden unexpected death in infancy (SUDI).
36 *Forensic Sci Int.* 2010, May 20;198(1-3):121-5.
 - 37 16. Fernández-Rodríguez A, Alcalá B, Alvarez-Lafuente R. Real-time PCR detection of
38 *Neisseria meningitidis* in formalin-fixed tissues from sudden deaths. *Diagn*
39 *Microbiol Infect Dis.* 2008;60:339–46
 - 40 17. Damann FE, Williams DE, Layton AC. Potential use of bacterial community
41 succession in decaying human bone for estimating postmortem interval. *J Forensic*
42 *Sci.* 2015;60:844-50.
 - 43 18. Schwarz P, Dannaoui E, Gehl A, Felske-Zech H, Birngruber CG et al. Molecular
44 identification of fungi found on decomposed human bodies in forensic autopsy cases.

Int J Legal Med. 2015;129:785-91.

19. Fernández-Rodríguez A, Cohen MC, Lucena J, Van de Voorde W, Angelini A et al. How to optimise the yield of forensic and clinical post-mortem microbiology with an adequate sampling: a proposal for standardisation. *Eur J Clin Microbiol Infect Dis.* 2015;34:1045-57.
20. Fernández-Rodríguez A, Burton J, Andreoletti, L, Alberola J, Fornes P et al. Post-mortem microbiology in Sudden Death: Sampling protocols proposed in different clinical settings. *Clin Microb Infect.* 2019; 25 (5): 570-9.
21. Burton JL, Saegeman V, Arribi A, Rello J, Andreoletti L, et al. Post-mortem microbiology sampling following death in hospital: An ESGFOR Task Force Consensus Statement. *J Clin Pathol.* 2019; 0:1–8. doi:10.1136/jclinpath
22. Addison S, Arthurs OJ, Thayyil S. Post-mortem MRI as an alternative to non-forensic autopsy in fetuses and children: from research into clinical practice. *Br J Radiol.* 2014;Apr:87(1036):20130621.
23. Cohen MC, Paley MN, Griffiths PD, Whitby EH. Less invasive autopsy: benefits and limitations of the use of magnetic resonance imaging in the perinatal postmortem. *Pediatr Dev Pathol.* 2008;11:1-9.
24. Tsokos M. Postmortem diagnosis of sepsis. *Forensic Sci Int.* 2007;165:155-64.
25. Burton JL. Health and safety at autopsy. *J Clin Pathol.* 2003;56:254-60.
26. Hanley B, Lucas SB, Youd E, et al. Autopsy in suspected COVID-19 cases. *J Clin Pathol* Epub ahead of print: [please include Day Month Year]. doi:10.1136/jclinpath-2020-206522 *Clin Pathol.* 2020;0:1–4. doi:10.1136/jclinpath-2020-206522
27. Transmission-based precautions. Guidance for care of deceased. during COVID-19 pandemic. Issued: 19 March 2020. The Royal College of Pathologists and the Association for Anatomical Pathology Technology in conjunction with Public Health England.
28. Burton JL, Underwood JCE. Clinical, educational and epidemiological value of autopsy. *Lancet.* 2007;369:1471-80.
29. Fan JK, Tong DK, Poon JT, Lo OS, Beh PS, et al. Multimodality minimally invasive autopsy – a feasible and accurate approach to post-mortem examination. *Forensic Sci Int.* 2010;195:93-8.
30. Weustinik AC, Hunink MGM, van Dijke CF, Renken CF, Krestin GP, et al. Minimally invasive autopsy: an alternative to conventional autopsy? *Radiology.* 2009;250: 897-904.
31. Roberts ISD, Traill ZC. Minimally invasive autopsy employing post-mortem CT and targeted coronary angiography: evaluation of its application to a routine coronial service. *Histopathology.* 2014;64:211-7.
32. Castillo P, Martínez MJ, Ussene E, Jordao D, Lovane L, et al. Validity of a minimally invasive autopsy for cause of death determination in adults in Mozambique: an observational study. *PLoS Med.* 2016;13(11):e1002171.
33. Maixenchs M, Anselma R, Zielinski-Gutiérrez E, Odiambo FO, Akello C, et al. Willingness to know the cause of death and hypothetical acceptability of the minimally invasive autopsy in six diverse African and Asian settings: a mixed

-
- methods socio-behavioural study. *PLoS Med.* 2016;13(11):e1002172.
- 1 34. Roberts ISD, Benamore RE, Benbow EW, Lee SH, Harris JN, et al. Post-mortem
2 imaging as an alternative to autopsy in the diagnosis of adult deaths: a validation
3 study. *The Lancet.* 2012;379:136–42.
- 4 35. Blokker BM, Wagenveld IM, Weustink AC, Oosterhuis JW, Hunink MG.
5
6 Non-invasive or minimally invasive autopsy compared to conventional autopsy of
7 suspected natural deaths in adults: a systematic review. *Eur Radiol.*
8 2016;26:1159-79.
- 9 36. Fryer EP, Traill ZC, Benamore RE, Roberts IS. High risk medicolegal autopsies: is a
10 full postmortem examination necessary?. *J Clin Pathol.* 2012;66:1-7.
- 11 37. Gonzalez-Arnay E, Martin-Olivera R, Quintero-Quintero YC, Hernandez-Guerra AI.
12 Proposal for a harmonized protocol for COVID-19 screening and necropsy in
13 forensic sciences facilities. *J Forensic Leg Med* 2020:Oct 3;76:102067, doi:
14 10.1016/j.jjflm.2020.102067
- 15 38. Bolliger SA, Filograna L, Spendlove D, Thali MJ, Sirnhofer S, et al. (2010).
16 Postmortem imaging-guided biopsy as an adjuvant to minimally invasive autopsy
17 with CT and postmortem angiography: a feasibility study. *Am J Roentgenol.*
18 2010;195:1051-6.
- 19 39. Ebert LC, Ptacek W, Breitbeck R, Fürst M, Kronreif G, et al. (2014). Virtobot 2.0:
20 the future of automated surface documentation and CT-guided needle placement in
21 forensic medicine. *Forensic Sci Med Pathol.* 2014;10:179-86.
- 22 40. Ross SG, Thali MJ, Bollinger S, Germerott T, Ruder TD, et al. Sudden death after
23 chest pain: feasibility of virtual autopsy with post-mortem CT angiography and
24 biopsy. *Radiology.* 2012;264:250-9.
- 25 41. Karat AS, Omar T, von Gottberg A, Tlali M, Chihota VN, et al. Autopsy prevalence
26 of tuberculosis and other potentially treatable infections among adults with
27 advanced HIV enrolled in out-patient care in South Africa. *PLOS One.*
28 2016;11(11):e0166158.
- 29 42. Cox JA, Lukande RL, Lucas S, Nelson AM, Van Marck E, et al. Autopsy causes of
30 death in HIV-positive individuals in sub-Saharan Africa and correlation with clinical
31 diagnoses. *AIDS Rev.* 2010;12:183-94
- 32 43. Santurro A, Scopetti M, D’Errico S, Fineschi V. A technical report from the Italian
33 SARS-CoV-2 outbreak. Postmortem sampling and autopsy investigation in cases of
34 suspected or probable COVID-19 *Forensic Science, Medicine and Pathology.*
35 Epub:2020 May 12 <https://doi.org/10.1007/s12024-020-00258-9>
- 36 44. Moore C, Jones R. The use of coroner's autopsy reports to validate the use of
37 targeted swabbing rather than tissue collection for rapid confirmation of virological
38 causes of sudden death in the community. *J Clin Virol* 2015 Feb;63:59-62. doi:
39 10.1016/j.jcv.2014.11.031. Epub 2014 Dec 3
- 40 45. Wise J. Largest group of children affected by measles outbreak in Wales is 10–18
41 year olds. *BMJ* 2013;346:f2545.
- 42 46. Hurtado JC, Quintó L, Castillo P, Carrilho C, Fernandes F, et al. Postmortem
43 Interval and Diagnostic Performance of the Autopsy Methods. *Sci Rep.*
44 2018 ;8(1):16112.
- 45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

-
- 1 47. Castillo P, Ussene E, Ismail MR, Jordao D, Iovane L, et al. Pathological methods
2 applied to the investigation of causes of death in developing countries: minimally
3 invasive autopsy approach. *PLoS ONE*.2015;10(6):e0132057.
- 4 48. Palhares AEM, Ferreira L, Freire M, Castillo P, Martínez MJ, et al. Performance of
5 the minimally invasive autopsy tool for cause of death determination in adult deaths
6 from the Brazilian Amazon: an observational study. *Virchows Arch*. 2019 doi:
7 10.1007/s00428-019-02602-z. [Epub ahead of print]
- 8 49. Van der Linden A, Blokker BM, Kap M, Weustink AC, Riegman PHJ, et al.
9 Post-mortem tissue biopsies obtained at minimally invasive autopsy: an
10 RNA-quality analysis. *PLOS One* .2014;9(12):e115675.
- 11 50. Ruddy GN, Johnson C, Amoroso J, Robinson C, Bradley CJ, et al. Post-mortem
12 computed tomography coaxial cutting needle biopsy to facilitate the detection of
13 bacterioplankton using PCR probes as a diagnostic indicator for drowning. *Int J*
14 *Legal Med*. 2017;131:211-6.
- 15 51. Martínez MJ, Massora S, Mandomando I, Ussene E, Jordao D, et al. Infectious
16 cause of death determination using minimally invasive autopsies in developing
17 countries. *Diagn Microbiol Infect Dis*. 2016;84:80-6.
- 18 52. Smith R. Survival of tubercle bacilli: *American Review of Tuberculosis*, 45(3), pp.
19 334–345
- 20 53. *Zz Guide to infection control in the healthcare setting, The infection hazards of*
21 *human cadavers. International Society of Infectious Diseases. Hoffman PN, Healing*
22 *TD*
- 23 54. Lu L, Liu D, Zhang H et al, Air and surface contamination in non-health care
24 settings among 641 environmental specimens of 39 COVID-19 cases. *PLoS Negl*
25 *Trop Dis* 2020; 9;14(10):e0008570, doi: 10.1371/journal.pntd.0008570
- 26 55. Bassat Q, Castillo P, Martínez MJ, Jordao D, Iovane L, Hurtado JC, et al. Validity
27 of a minimally invasive autopsy tool for cause of death determination in pediatric
28 deaths in Mozambique: An observational study. *PLoS Med*. 2017;14(6): e1002317.
29 <https://doi.org/10.1371/journal.pmed.1002317>
- 30 56. Menendez C, Castillo P, Martínez MJ, Jordao D, Iovane L, Ismail MR, et al.
31 Validity of a minimally invasive autopsy for cause of death determination in
32 stillborn babies and neonates in Mozambique: An observational study. *PLoS Med*.
33 2017;14(6): e1002318. <https://doi.org/10.1371/journal.pmed.1002318>
- 34 57. Thayyil S, Sebire NJ, Chitty LS, Wade A, Chong W, et al. Post-mortem MRI versus
35 conventional autopsy in fetuses and children: a prospective validation study. *The*
36 *Lancet*. 2013;Jul 20;382(9888):223-33.
- 37 58. Sethuraman C, Cohen M, Whitby E, Marven S, Lindley R. Sheffield minimally
38 invasive autopsy research technique (smart)- a technical report on feasibility of
39 keyhole autopsy and magnetic resonance imaging. Platform presentation at the 50th
40 Meeting of the Pediatric Pathology Society. 21-22 March 2015, Boston, US. 2015
41 DOI: 10.2350/15-02-1606-MISC.1
- 42 59. Rummery R, DiGiovanni J, Raghavan A, Whitby E, Cohen M. Benefits and
43 limitations of the minimally invasive perinatal autopsy: A review of the first 5 years
- 44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

-
- of an innovative service development. Society of Pediatric Pathology Spring Meeting. 16-18 March 2018. Pinnacle Hotel Harbourfront. Vancouver, BC. Canada
60. Madhi SA, Pathirana J, Baillie V, Izu A, Bassat Q et al. Unraveling Specific Causes of Neonatal Mortality Using Minimally Invasive Tissue Sampling: An Observational Study, *Clinical Infectious Diseases*, Volume 69, Issue Supplement_4, 15 October 2019, Pages S351–S360, <https://doi.org/10.1093/cid/ciz574>
61. Xu, Y., Li, X., Zhu, B. et al. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat Med* 26, 502–505 (2020). <https://doi.org/10.1038/s41591-020-0817-4>
62. Castillo P, Hurtado JC, Martínez MJ, Jordao D, Lovane L, et al. Validity of a minimally invasive autopsy for cause of death determination in maternal deaths in Mozambique: an observational study. *PLOS Medicine*. 2017;Nov 8;14(11):e1002431.
63. Prtak L, Al-Adnani M, Fenton P, Kudesia G, Cohen MC. Contribution of bacteriology and virology in sudden unexpected death in infancy. *Arch Dis Child*. 2010;95:371-6.
64. Roberts FJ. A review of postmortem bacteriological cultures. *Can Med Assoc J*. 1969;Jan 11;100(2):70-4.
65. Alberola J, Cohen MC, Fernandez Rodriguez A. Análisis microbiológico post mortem. Recomendaciones de la Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. In: *Procedimientos de Microbiología Clínica*. SEIMC 2012:43. Eds: Cercenado E, Canton R. 2012 ISBN- 978-84-616-0446-3 Available at: www.seimc.org/documentos/protocolos/microbiologia/cap43.asp
66. Sunagawa K, Sugitani M. Post-mortem detection of bacteremia using pairs of blood culture samples. *Leg Med*. 2017;Jan;24:92-7.
67. Weber M, Klein N, Hartley J, Lock P, Malone M, et al. Infection and sudden unexpected death in infancy: a systematic retrospective case review. *The Lancet*. 2008;May 31;371(9627):1848-53.
68. Fernández-Rodríguez A, Alberola J, Cohen MC. Análisis microbiológico post-mortem. *Enferm Infecc Microbiol Clin*. 2013;31(10):685-691 <http://dx.doi.org/10.1016/j.eimc.2012.09.019>.
69. Zhao W, Zhong Z, Xie X, Yu Q, Liu J, et al. Relation between chest CT findings and clinical conditions of coronavirus disease (COVID-19) pneumonia: a multicenter study. *Am J Roentgenol*. 2020:1–6.
70. Yao XH, Li TY, He ZC, Ping YF, Liu HW, et al. A pathological report of three COVID-19 cases by minimally invasive autopsies, *Chinese Journal of Pathology*. 49 (2020) E009. DOI:10.3760/cma.j.cn112151-20200312-00193.
71. Driggin E, Madhavan MV, Bikdeli B, Chuich T, Laracy J, et al. Cardiovascular Considerations for Patients, Health Care Workers, and Health Systems During the Coronavirus Disease 2019 (COVID-19) Pandemic, *Journal of the American College of Cardiology*. 2020; doi: <https://doi.org/10.1016/j.jacc.2020.03.031>.
72. Dubey SP, Larawin V, Molumi CP. Intracranial spread of chronic middle ear suppuration. *Am J Otolaryngol*. 2010;Mar-Apr;31(2):73–7.
73. Geyik MF, Kokoglu OF, Hosoglu S, Ayaz C. Acute bacterial meningitis as a

complication of otitis media and related mortality factors. *Yonsei Med J.* 2002;Oct;43(5):573–8.

- 1
2
3 74. Saegeman V, Bammens B, De Munter P, Neyt J, Cossey V, et al. How to isolate a
4 patient with *Cladophialophora bantiana* infection? An opinion. *J Hosp Infect.*
5 2015;Sep;91(1):89-90.
6
7 75. Johnson JA, Boyce TG, Cetta F, Steckelberg JM, Johnson JN. Infective Endocarditis
8 in the Pediatric Patient: A 60-Year Single-Institution Review. *Mayo Clin Proc.*
9 2012;87(7):629-35.
10
11 76. Achermann Y, Rossle M, Hoffmann M, Deggim V, Kister S et al. Prosthetic valve
12 endocarditis and bloodstream infection due to *Mycobacterium chimaera*. *J Clin*
13 *Microbiol.* 2013;51:1769-73.
14
15 77. Williamson D, Howden B, Stinear T. *Mycobacterium chimaera* spread from heating
16 and cooling units in heart surgery. *N Engl J Med.* 2017;376:600-2.
17
18 78. Satoh F, Tachibana H, Hasegawa I, Osawa M. Sudden death caused by chronic
19 Chagas disease in a non-endemic country: Autopsy report. *Pathol Int.* 2010;60:235–
20 240.
21
22 79. Liu A, Hu Y, Coates A. Sudden cardiac death and tuberculosis - How much do we
23 know? *Tuberculosis.* 2012;92:307–31
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1: Macroscopic and imaging features in infectious conditions

Condition	Macroscopic appearance	Imaging*	Etiological agents	Remarks and references
Respiratory infections				
Tonsillitis, laryngitis, tracheitis, bronchitis	Congested mucosa +/- mucopurulent secretions	<i>Radiographs:</i> Enlarged tonsils <i>CT:</i> Enlarged “kissing” tonsils, parapharyngeal fat stranding <i>MRI:</i> Enlarged tonsils, low to isointense on T1, high signal on T2	Bacteria Viruses	
Pneumonia	-Heterogeneous, multifocal affecting several lobes (bronchopneumonia) or -a homogenous consolidation affecting one or more lobes (lobar pneumonia) 4 histopathological stages: congestive phase: enlarged and congested lung – red hepatisation: a reddish firm and consolidated lung – grey hepatisation: an opaque lung with grey discoloration and a purulent exudate on the cut surface - resolution	Air space shadowing with air bronchograms in a rounded (rounded pneumonia), lobar (lobar pneumonia) or patchy (bronchopneumonia) distribution	Bacteria Viruses Fungi	Clinical classification: community acquired Hospital acquired Ventilator associated Aspiration pneumonia More information on etiological agents in [21]
Tuberculosis	-Disseminated Tuberculosis: multiple small whitish/yellowish, well circumscribed, firm nodules +/- a central caseous necrosis. -miliary tuberculosis: numerous whitish nodules (2mm diameter) invariably involving the pleural surface, may also be present in vertebra and other organs	Perihilar/paratracheal lymphadenopathy (primary MTB) Calcification of Ghon focus and lymph nodes plus solitary or multiple granulomas (latent MTB) Consolidation, obstructive atelectasis, cavitation (progressive primary MTB) Innumerable small (≤ 2 mm) calcified nodules best seen on CT (miliary MTB) Pleural/pericardial thickening and effusion (pleural/pericardial MTB)	<i>Mycobacterium tuberculosis</i> complex (MTB)	
Abscess	A cavity filled with necrotic and purulent material.	Rounded cavity with air/fluid level. There may be surrounding consolidation	Bacteria	May be associated to an adjacent area of bronchopneumonia
<i>Empyema thoracis</i>	A fibrinopurulent exudate on the pleural surfaces or frank pus in the pleural cavity.	<i>Radiographs:</i> Lenticular-shaped fluid density in the pleural space <i>CT:</i> May show locules of gas. Displacement and compression of adjacent lung. “Split pleura” sign		A possible complication of lobar bronchopneumonia or pneumonia, lymphatic or vascular dissemination of a distant infection, or from a hepatic or subphrenic abscess
COVID-19	Pleurisy, pericarditis,	<i>Radiographs:</i> multiple ground glass opacities	SARS-CoV-2	Although mainly affecting the lung (diffuse alveolar damage, hyaline

	lung consolidation and pulmonary oedema. Possibility of lung weight increased and also of purulent inflammation due bacterial co-infection	<i>CT</i> : GGO, mixed GGO, consolidation, reticulation. [69]		membranes, macrophages and CD4+ T cell lymphocytes, and microthrombus) other organs can also be damaged: heart (myocarditis), spleen (decreased numbers of lymphocyte, cell degeneration and necrosis), vessels, liver, kidney and others, which should be sampled in the MIA [70,71],
Central nervous system infections				
Meningo-encephalitis	Non-specific aspect of brain: pale, swollen, with flattened gyri and devoid of any exudate	Unenhanced post-mortem imaging may be normal <i>CT</i> : Mild hydrocephalus. Hyperdensity surrounding basal cisterns <i>MRI</i> : Sulci less hypointense than normal (T1), hyperintense signal in sulci (FLAIR)	Viruses	More information on etiological agents in [21]
	<i>N. meningitidis</i> : no exudate. Other bacteria: a purulent exudate, regularly associated with middle ear infection (eg. <i>S. pneumoniae</i>)		Bacteria	Post-mortem culture often negative for <i>N. meningitidis</i> since fastidious organism. PCR from skin petechiae or CSF gives highest yield. Swab of the middle ear is important in PMM [63-68,72,73]
	- Dense gelatinous inflammatory exudate, mainly in the basal cisterns - meningeal or parenchymatous location - release of <i>M. tuberculosis</i> complex and dissemination and causing encephalitis		<i>Mycobacterium tuberculosis</i> complex	
			<i>Candida</i> spp., Mucorales spp., <i>Aspergillus fumigatus</i> complex, <i>Cryptococcus</i> spp.	Most often in immunocompromised patients
Cerebral phaeohyphomycosis		Single or multiple brain abscesses seen on CT or MRI as solid round/oval space occupying lesions with central necrosis/fluid and perilesional edema. Central diffusion restriction on MRI highly suggestive of an abscess rather than tumour	<i>Histoplasma capsulatum</i> , <i>Coccidioides immitis</i> , <i>Blastomyces dermatitidis</i> , <i>Cladophialophora bantiana</i> [74]	Healthy individuals endemic regions contact and airborne precautions
Cerebral abscess in the deep grey matter and in the cortex		Solid round/oval space occupying lesion with central necrosis/fluid and perilesional edema. Central diffusion restriction on MRI highly suggestive of an abscess rather than tumour	Parasites (eg. Toxoplasmosis)	Immunocompromised patients

Cardiovascular infection				
Endocarditis	Friable vegetations containing fibrin, cellular debris, bacteria. Vegetations may erode the adjacent valve ring and cause annular abscesses	<i>Radiographs/CT:</i> Non-specific, patchy consolidation, cardiomegaly with pericardial effusion, pleural effusion (on occasion) <i>MRI:</i> May show valvular vegetations	<i>Streptococcus viridans</i> , <i>Staphylococcus aureus</i> , <i>Enterococcus</i> spp., <i>Streptococcus gallolyticus</i> , HACEK group bacteria*, atypical mycobacteria	- In children with repaired complex heart diseases: mean age of 12.3+/-5.5 yrs, viridans streptococci and <i>S. aureus</i> [75] - <i>S. aureus</i> most prevalent in IV drug users - <i>M. chimaera</i> endocarditis reports after open heart surgery linked to contamination of heater cooler units [76,77]
Myocarditis	Heart is floppy and dilated. Myocardium may appear macroscopically normal, although frequently pale with punctate haemorrhagic areas.	Non-specific post-mortem imaging findings	Viruses: Enteroviruses, <i>Parvovirus B19</i> , Human Herpesvirus 6 and 8, Adenoviruses; <i>Trypanosoma cruzi</i> ** [78]	Human Immunodeficiency virus, <i>M. tuberculosis</i> complex may cause myocarditis [79] More information on etiological agents in [20,21]
Pericarditis	Pericardium opaque with a purulent exudate. May be complicated by fibrosis leading to constrictive pericarditis.	Cardiomegaly with pericardial effusion <i>MRI:</i> May demonstrate thickened pericardium ($\geq 4\text{mm}$)	Bacteria	By direct or indirect invasion of bacteria (eg. bronchopneumonia)
Septicaemia	Generalized bacterial infection with bacterial replication in the circulation. Gram-negative sepsis may result in endotoxin shock. May be complicated by diffuse intravascular coagulation with petechiae in skin and mucous membranes.	Non-specific post-mortem imaging findings		Different agents depending on community acquired or nosocomial. More information of pathogens involved in nosocomial septicaemia in [21]
Gastro-intestinal infections				More information on etiological agents in [21]
Hepatitis	Enlarged and congested liver, with green discoloration if cholestasis, +/- pale areas of hepatocellular necrosis. Chronic viral hepatitis may result in fibrosis and cirrhosis. A cirrhotic liver is firm, with a reduced volume and multinodular appearance on the external and cut surfaces.	Non-specific post-mortem imaging findings <i>CT:</i> May show decreased parenchymal attenuation (representing oedema)	Viruses: hepatitis A, B, C, D, E virus, Epstein-Barr virus, Cytomegalovirus	Virus detection via PCR on tissue samples, complementary to viral serology
Liver abscess		<i>Radiographs:</i> May show air in the biliary tree or within an abscess, right pleural effusion <i>CT:</i> Solid round/oval space occupying lesion with central necrosis/fluid. May contain gas	Gastro-intestinal bacteria: <i>Clostridium</i> spp., <i>Peptostreptococcus</i> spp., <i>S. aureus</i> , gram-negative bacteria -Parasites: <i>Entamoeba histolytica</i>	

	cavity filled with a purulent exudate,	<i>MRI</i> : Solid round/oval space occupying lesion with central necrosis/fluid with perilesional edema		After/during amoebic colitis or as a complication of bacterial peritonitis
Peritonitis	Peritoneal surface appears opaque and erythematous with an associated purulent exudate. Amount of exudate related to the magnitude and duration of the infectious process.	<i>CT/MRI</i> : Thickened peritoneum (smooth, irregular or nodular appearance); free or loculated ascites	Gram-negatives, <i>Enterococcus</i> spp., <i>S. aureus</i> , alpha- and beta haemolytic streptococci, <i>Clostridium perfringens</i> , yeasts	Underlying infectious condition (eg. Acute appendicitis, cholecystitis, diverticulitis, salpingitis) or other abdominal condition (trauma, volvulus, cirrhosis, nephrotic syndrome)
Colitis	Oedematous mucosa, haemorrhagic and ulcerated. Shigella usually affects the distal colon; hyperaemic mucosa, oedematous, haemorrhagic with prominence of Peyer patches and associated purulent exudates. <i>Salmonella</i> is prevalent in ileum and proximal colon. Mucosa is oedematous, haemorrhagic and associated with linear ulcers. <i>Clostridium</i> spp. May cause haemorrhage and necrosis of the colon with risk of perforation. <i>C. difficile</i> causes pseudomembranous colitis with a firmly adherent fibrinopurulent mucosal exudate. Colon mucosa is slightly affected	<i>CT/MRI</i> : Multiple fluid-filled bowel loops with thickened walls, ascites, inflammation/stranding of pericolonic fat	Bacteria: <i>E. coli</i> , <i>S. aureus</i> , <i>Salmonella</i> , <i>Shigella</i> , <i>Campylobacter</i> spp., <i>Yersinia</i> spp., <i>Clostridium</i> spp. Viruses: Adenovirus type 40 and 41, Astroviruses, Noroviruses, Enterovirus, Rotavirus	Bloody diarrhoea
Urogenital Infections				
Pyelonephritis and pyonephrosis	White exudate in pelvis	Non-specific post-mortem imaging findings <i>CT</i> : Low attenuation (edema) of renal parenchyma, renal calculus, hydronephrosis, abscess/perinephric collection <i>MRI</i> : Affected regions are hypointense on T1 and hyperintense on T2 compared to normal renal tissue	Gram-negatives	
Pelvic inflammatory disease	Yellowish clumps or sulfur granules	<i>CT/MRI</i> : Smooth peritoneal thickening, stranding/haziness of pelvic fat, fluid-filled fallopian tubes with thickened walls, tubo-ovarian or pelvic abscesses	<i>Actinomyces</i> spp. <i>Neisseria gonorrhoeae</i> , <i>Chlamydia trachomatis</i> , <i>Trichomonas vaginalis</i>	Associated with intra-uterine devices

Notes: *HACEK group: *Haemophilus parainfluenzae*, *Aggregatibacter aphrophilus*, *Aggregatibacter actinomycetemcomitans*, *Cardiobacterium hominis*, *Eikenella corrodens* and *Kingella kingae*

** Other less frequent pathogens causing myocarditis are: *Epstein-Barr virus*, *Cytomegalovirus*, *Chlamydia psittaci*, *Candida albicans*, *Toxoplasma gondii*, *Trichinella* spp.

Table 2. Sampling collection during adult MIA for microbiological investigation.

Tissue/Body fluid	Site of collection	Means of collection	Type of analysis
Blood	Subclavian vein or heart	Needle aspiration	Direct bacterial culture Virology/serology for viral hepatitis, HIV Molecular analyses*
Cerebrospinal fluid	Occipital approach to the cisterna magna	Needle aspiration	Direct bacterial culture Molecular analyses Antigenic analyses
Urine	Suprapubic aspiration	Needle aspiration	Direct bacterial culture
Pleural effusion	Aspirate via chest wall	Needle aspiration	Direct bacterial culture
Ascites	Aspirate via abdominal wall	Needle aspiration	Direct bacterial culture
Brain	Trans-ethmoidal approach	Cutting needle biopsy	Direct bacterial culture Molecular analyses
Bone marrow	Anterior superior iliac crest	Bone marrow aspiration	Direct bacterial culture
Nasopharyngeal	Nasopharynx	Swabs (Amies/viral transport media/sodium chloride)	Molecular analyses Antigenic analyses
Lungs	Via chest wall	Cutting needle biopsy	Direct bacterial culture Molecular analyses
Spleen	Via left lateral abdominal wall	Cutting needle biopsy	Direct bacterial culture Molecular analyses
Liver	Via right lateral abdominal wall	Cutting needle biopsy	Direct bacterial culture Molecular analyses
Kidney	Via posterior abdominal wall	Cutting needle biopsy	Direct bacterial culture

Notes:

Samples that can be collected during adult MIA for microbiological investigation. The range of samples collected will depend on the population being served, the clinical history and the resources available. A new sterile needle should be used for each biopsy. [49,51]

*Molecular analyses either for bacteria or viruses are included in the table in those samples in which they are most frequently useful.

Table 3. Standard sampling collection during pediatric MIA for microbiological investigation [49, 51, 55-56]

Tissue/Body fluid	Site of collection	Means of collection	Type of analysis
Nasopharynx/ Throat	Nasopharynx Throat	Swab with transport media (Amies/viral)	Direct bacterial culture Molecular analyses* Antigenic analyses
Blood	Heart	Needle aspiration Through chest wall or during thoracoscopy (according to procedure)	Bacterial culture Molecular analyses Antigenic analyses
Cerebrospinal fluid	Occipital approach to the cisterna magna or a lumbar puncture	Needle aspiration	Direct bacterial culture Molecular analyses Antigenic analyses
Urine	Suprapubic aspiration	Needle aspiration	Direct bacterial culture
Lungs	Through thoracoscopy	Tissue sample	Direct bacterial culture Molecular analyses
Additional tissues: -Liver, central nervous system) - Bone marrow		-Biopsy needles (14G+16G)	Direct bacterial culture Molecular analyses
Bowel content	Through laparoscopy	Incision of bowel, which is sutured thereafter.	Direct bacterial culture Molecular analyses Antigenic analyses
Rectal swab	Rectum	Swab (viral transport media)	PCR SARS-CoV-2

Note: *Molecular analyses either for bacteria or viruses are included in the table in those samples in which they are most frequently useful.

Table 4. Type and main characteristics of the different needles used in the MIA procedure for each particular biopsy, sites of puncture and number of samples to be obtained [47].

	Needle	Type	Gauge	Needle length (mm)	Site of Puncture
Cerebrospinal fluid	Quincke Spinal [#]	Manual	20	100	Occipital puncture
Blood	Quincke Spinal	Manual	20	100	Supra/infra-clavicular or left ventricle
Liver	Monopty*	Automatic	14-16	115	Anterior right axillar line, 11 th -12 th intercostal space
Lungs and heart	Monopty*	Automatic	14-16	100	Right and left clavicular region down to the diaphragm for microbiology samples. Multiple random thoracic punctures for pathology
Bone Marrow	T-Lok TM Trepine**	Manual	8	100	Anterior iliac crest
Central nervous system	Monopty*	Automatic	16	200	Occipital puncture Trans ethmoidal puncture. Perforation of the cribriform plate with the bone marrow trephine to reach the cranial cavity
Skin	Biopsy punch ^{##}	Manual	5 mm	-	Macroscopically detected lesions

(# Becton Dickinson, Franklin Lakes, NJ, USA, ##KAI Europe GMBH, Solingen, Germany, * BARD Biopsy Systems, Tempe, AZ; USA, **Mana-Tech Ltd, Staffordshire, UK