UNIVERSITY OF LEEDS

This is a repository copy of *An evaluation of changes to protein supply in Zambia between 1961 and 2013 shows consistent dominance of plant-based sources*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/171045/>

Version: Accepted Version

Proceedings Paper:

Kapulu, NP, Clark, H, Orfila, C orcid.org/0000-0003-2564-8068 et al. (1 more author) (2020) An evaluation of changes to protein supply in Zambia between 1961 and 2013 shows consistent dominance of plant-based sources. In: Proceedings of the Nutrition Society. Nutrition Society Live 2020, 14 Jul 2021 - 15 Jul 2020, Online. Cambridge University Press (CUP) .

<https://doi.org/10.1017/s002966512000748x>

© The Authors, 2020. This article has been published in a revised form in Proceedings of the Nutrition Society, <https://doi.org/10.1017/S002966512000748X>. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

An evaluation of changes to protein supply in Zambia between 1961 and 2013 shows consistent dominance of plant-based sources

N.P. Kapulu,
H. Clark,
C. Orfila and
J.I. Macdiarmid

Dietary protein is essential to maintain optimum growth and maintenance of body muscles in humans ⁽¹⁾. Although animal sources of protein have a higher biological value and protein digestibility than most foods ⁽²⁾, they are beyond the economic reach of many, especially those living in the global south. However, protein from animal sources is environmentally unsustainable, and excess consumption of red and processed meats are associated with increased risk of non-communicable diseases ⁽³⁾. Using Zambia as a case study, this study aimed to assess the supply of protein from various food sources to meet population-level requirements, including changes that have occurred over time.

A list of 264 food items for 13 commodity groups in the FAO food balance sheets was developed using information from the FAO handbook and local knowledge of commonly consumed foods. The protein content for each item was estimated from West African food composition tables and where the food was not in these tables data from United States Department of Agriculture composition tables were used. The primary sources of protein and supply trends to meet age-and-gender specific population-level nutrition requirements for the period 1961–2013, were assessed.

The results showed that in Zambia, 74% of the protein was supplied from plant-based sources, including cereals (62%), oil crops (5%), starchy roots (3%), vegetables (2%) and pulses (2%), while about 24% came from animal foods comprising meat (17%), fish (4%), dairy (2%), and eggs (1%). Zambia was able to meet the protein population requirements from plant-based sources alone sufficiently. Although the supply of protein from 1961–2013 exceeded population-level nutrient requirements, its availability gradually declined from about 76 g/capita/day to approximately 63 g/capita/day. Total protein supply, especially between the late 1970s and early 2000 was affected by a decline in supply from cereals and oil crops, which may be due to reduced wheat imports, maize and groundnut production, respectively. Beyond 2005, contributions to total protein from oil crops improved exponentially because of increased soybean production, while amounts from animal sources were notably unchanged between 1961 and 2013. The trend in protein supply follows the same pattern in the total energy supply.

This study shows that population-level nutritional requirements for total protein is adequate and can be met from plant-based sources in developing countries where animal source foods are unaffordable for the majority of poor people. However, the diversity of diets and food sources needs to increase to ensure the quality of the protein is sufficient and nutrition security can be achieved. Dietary changes to achieve this must be carefully considered so as not to create a negative impact on health and the environment.

References

- 1 Bauer, J, Biolo, G, Cederholm, T et al. (2013) *J Am Med Dir Assoc* 14, 542–559. [10.1016/j.jamda.2013.05.021](https://doi.org/10.1016/j.jamda.2013.05.021) [CrossRef](#) [Google Scholar](#)
- 2 Hoffman, JR & Falvo, MJ (2004) *J Sports Sci Med* 3, 118–130. [Google Scholar](#)
- 3 Nijdam, D, Rood, T & Westhoek, H (2012) *Food Policy* 37, 760–770. [10.1016/j.foodpol.2012.08.002](https://doi.org/10.1016/j.foodpol.2012.08.002) [CrossRef](#) [Google Scholar](#)