UNIVERSITY OF LEEDS

This is a repository copy of *Application of pharmaceutical waste sludge compost alters the antibiotic resistome in soil under the Chinese cabbage system.*

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/171344/>

Version: Accepted Version

Article:

Xiao, Z, Zhang, Y, Li, G et al. (6 more authors) (2020) Application of pharmaceutical waste sludge compost alters the antibiotic resistome in soil under the Chinese cabbage system. Journal of Cleaner Production. 125229. ISSN 0959-6526

<https://doi.org/10.1016/j.jclepro.2020.125229>

© 2020, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

1 There are 8848 words in this paper.

2 **Application of pharmaceutical waste sludge compost alters the**
3 **antibiotic resistome in soil under the Chinese cabbage system**

4 Zufe Xiao^{a, b}, Yuan Zhang^{a, *}, Gang Li^{b, c, d, *}, Laura J. Carter^e, Hongtao Wang^{b, c}, Jing Ding^f, Faith
5 Ka Shun Chan^{e, g, h}, Yilong Hao^{b, d}, Yaoyang Xu^{b, c, d}

6

7 ^a School of Environmental Science and Engineering, Suzhou University of Science and Technology,
8 99 Xuefu Road, Suzhou, 215009, China.

9 ^b Key Lab of Urban Environment and Health, Fujian Key Lab of Watershed Ecology, Institute of
10 Urban Environment, Chinese Academy of Sciences, 1799 Jimei Road, Xiamen, 361021, China.

11 ^c University of Chinese Academy of Sciences, 19A Yuquan Road, Beijing, 100049, China.

12 ^d Ningbo Urban Environmental Observatory and Research Station, Institute of Urban Environment,
13 Chinese Academy of Science, Ningbo 315830, China.

14 ^e School of Geography, Faculty of Environment, University of Leeds, Leeds, LS2 9JT, UK.

15 ^f School of Environmental and Material Engineering, Yantai University, 30 Qingquan Road, Yantai,
16 264005, China

17 ^g School of Geographical Sciences, Faculty of Science and Engineering, University of Nottingham
18 Ningbo China, 315100, Ningbo, China

19 ^h Water@Leeds Research Institute, University of Leeds, Leeds, LS2 9JT, UK

20

21 * Joint corresponding authors: Yuan Zhang (yuanzhang_1001@mail.usts.edu.cn) & Gang Li
22 (gli@iue.ac.cn)

Highlights

- Cabbage associated with resistome was measured in amended soils.
- ARG profiles in amended soil and phyllosphere of cabbage were closely clustered.
- The amendment of compost might improve the accessibility of resistome in the soil especially for the Chinese cabbage system.

ABSTRACT

Antibiotic resistance is a global threat posing risks to public health. China, as the largest consumer and producer of antibiotics, is generating a large amount of pharmaceutical waste sludge from the antibiotic manufacturing industry, which has the potential to be released into the environments by anthropogenic activities. Land application of pharmaceutical waste sludge compost (PWSC) is a popular way of PWSC disposal, with large amount of antibiotics might hence be introduced into the soil environments and result in the development of antibiotic resistance genes (ARGs). ARGs in PWSC amended soil-plant systems, their transmission routes and potential ecological risks are still unknown. A high-throughput qPCR chip was used to profile ARGs in Chinese cabbage (Shanghaiqing, *Brassica chinensis* L.) and soils (including phyllosphere, root endosphere, rhizosphere soil, and bulk soil) with PWSC amendment, aiming to study the effect of PWSC application on ARGs soil-plant systems. A total of 249 ARGs and 12 mobile genetic elements (MGEs) were detected in all collected samples. The highest number of detected ARGs was in the soil samples (up to 181) compared to the above and below-ground components of Chinese cabbage. Our results demonstrated the PWSC amendment increased the diversity and normalized abundance of ARGs in the amended soil-Chinese cabbage system. Mantel test and Procrustes analysis revealed

the connection between ARG profiles and microbial communities (fungal and bacterial communities). Shared ARGs were identified among the Chinese cabbage phyllosphere, Chinese cabbage root, and rhizosphere soils, demonstrating a potential link between antibiotic resistome in Chinese cabbage and soils, with the amended soil as a key source of ARGs in the phyllosphere of Chinese cabbage. In summary, our findings provided novel evidence for a transmission route shared Zero-radius operational taxonomic units (ZOTUs) and ARGs were passed between the soil-Chinese cabbage system elements (i.e., amended soil, root, and phyllosphere) of ARGs from the PWSC amended soils to common green vegetables, highlighting a potential safety hazard of antibiotic resistome transfer from soils to the human food chain.

Keywords: Chinese cabbage; antibiotic resistome; pharmaceutical sludge; compost-amended soil; composting.

1. INTRODUCTION

Antibiotic resistance has become notorious in human pathogens (Allen et al., 2010). The spread of antibiotic-resistant bacteria (ARB) in the environment is contributing to the global crisis of clinically relevant antimicrobial resistance (AMR). Environmental pollution of ARB and antibiotic resistance genes (ARGs) can be attributed to commonly use of antibiotics in the animal industry (farm and agricultural industries), which are linked to anthropogenic activities such as intensive use of bio-organic fertilizers (e.g. animal manures and sewage sludge) in the agriculture (Chen et al., 2016; Chen et al., 2017; Cheng et al., 2013; Liu et al., 2019).

Appropriate usage of organic fertilizers on farmland can increase soil organic matter content,

but offering a pathway to recycle nutrients in the soil (Martínez Salgado et al., 2019). Studies have proved that the unregulated use of untreated organic fertilizers can increase levels of antibiotic resistome in farm settings (Chen et al., 2017; Su et al., 2015; Xie et al., 2016a). These antibiotics and enriched ARGs in farm settings could further transfer from untreated organic wastes to soils and gradually accumulate in vegetation (by absorption from the soil). For instance, supererogatory ARGs were detected in tomato and Chinese cabbage after the use of swine excrement and struvite as a soil amendment (An et al., 2018; Romain et al., 2013). Consumption of green products, particularly raw greens, represents a typical pathway of direct human exposure to the soil microbiome (Chen et al., 2019). ARGs can enter the human food chain or human pathogens via the contaminated plants (e.g. fruits and vegetables) or via horizontal gene transfer (HGT), posing a risk to human health (Amy et al., 2006; Gillings, 2017). China is one of the world's largest producers and consumers of antibiotics (Qiao et al., 2018; Zhang et al., 2015; Zhu et al., 2013). Vast quantities of pharmaceutical waste sludge (PWS) are produced from antibiotic manufacturing and wastewater treatment processes. PWS harbors large amounts of mobile genetic elements (MGEs) and ARGs (Tao et al., 2016), with a number of studies suggesting that PWS harbors more MGEs and ARGs compared with the municipal waste sludge (Liu et al., 2014; Liu et al., 2012; Tong et al., 2017).

Waste sludge is often pre-treated before the land application (e.g. via aerobic composting, anaerobic digestion, or sludge carbonization), with aerobic composting widely regarded as an efficient way to minimize chemical and biological hazards from organic wastes (Bernal et al., 2009; Liu et al., 2019; Su et al., 2015).

Previous studies have demonstrated that aerobic composting was promoted as an effective means to reduce the concentration of antibiotics and ARGs in waste sludge (Selvam and Wong,

2017; Xie et al., 2016b; Zhang et al., 2019). Antibiotic resistome is a resistance reservoir of all the ARGs and precursor genes in the pathogen and nonpathogen. It includes resistance elements carried by both antibiotic-producing bacteria and pathogenic bacteria, and cryptic resistance genes (Wright, 2007). Whilst, ARGs can be effectively reduced through composting, the antibiotic resistome will not be eliminated. Using Pharmaceutical Waste Sludge Compost (PWSC) as an organic fertilizer, therefore, represents an underlying origin of human exposure to antibiotic resistome during subsequent consumption of vegetables grown in compost-amended soils.

Our understanding of the fate of antibiotic resistome by following the application of composted waste, has seldom been evaluated. In particular, the effect on indigenous soil, root, and phyllosphere antibiotic resistome is a key knowledge gap. It is therefore imperative to elucidate the factors influencing the fate of antibiotic resistome, which follows PWSC application to agricultural land in order to develop effective tactics and technologies to alleviate the rapid spread of antibiotic resistome.

High-throughput quantitative PCR (HT-qPCR) with 296 recognized primer sets covering common 12 MGEs and 283 ARGs marker genes in the existing research (Zhu et al., 2013), was employed together with Illumina sequencing of fungi ITS genes and bacterial 16S rRNA genes. Analysis was targeted to: (1) assess the fate of four types of waste compost on the molecular structure of ARGs, bacterial and fungal communities in the soil-Chinese cabbage system; (2) explore the potential migration route of ARGs from the amended soil to Chinese cabbage, and (3) identify the shared MGEs and ARGs between the Chinese cabbage and amended soil by bipartite network analysis. The pot experiment aimed to assess the potential ecological risk of the use of pharmaceutical waste sludge compost. We explored the impacting factors of ARGs in the crops and

proposed suggestions for agricultural utilization of pharmaceutical waste sludge compost.

2. MATERIALS AND METHODS

2.1 Properties and Materials

Four types of compost (refer to T1-T3 and S) with a range of characteristics by following production under different composting treatments were used in this pot experiment. Three types of PWSCs (coded T1-T3) were derived from carbon-rich amendments of pharmaceutical sludge compost. Sewage sludge compost (S) was obtained from an online sales platform (Ningbo, China) (Table 1). The PWSCs were air-dried, prior to being used in the experiments. Soils were collected from a vegetable plot (Ningbo, China), which had not received sewage sludge or been amended with other organic manure in the past five years. Prior to use, soils were sieved (mesh size: 2mm) and air-dried. Chemical and physical properties are described in Table 1, including the contents of heavy metals, antibiotics, total carbon, and total nitrogen in the initial samples. Total nitrogen and total carbon were evaluated via dry combustion in a CNS element analyzer (*Vario MAX, Elementar*) (Chen et al., 2017). Concentrations of antibiotics were analyzed by liquid chromatography-mass spectrometry (*LC-MS/MS. XevoG2-SQTOF, Waters*) (details are provided in Supplementary file). The content of heavy metals was determined with inductively coupled plasma- mass spectrometer (*ICP-MS, iCAP Qnova, Thermo Scientific*) after oxidative digestion of samples in sealed microwave digestion tubes (Zhu et al., 2013). The sample pH was determined by preparing a 1:5 solid sample/water (w/v) suspension (*Basic pH Meter, PB-10, Sartorius*). Determination of nitrate and ammonium in the soil was performed using an AA3 Continuous Flow Analytical System (*AA3, SEAL Analytical GmbH*) (Wang et al., 2019).

2.2 Experiment Design

Figure S1A shows a schematic of the experimental design. Each pot consisted of 1kg of soil amended with or without 5% PWSC (w/w, dry weight) (Luo et al., 2016). Five treatments were arranged: untreated soil without fertilizer application (hereinafter called CK), PWSC-T1 amended-soil (hereinafter called T1), PWSC-T2 amended-soil (hereinafter called T2), PWSC-T3 amended-soil (hereinafter called T3), and S amended-soil (hereinafter called S).

Each treatment was replicated for three times, giving 15 pots in total (Fig S1A). Soil samples were then planted with Chinese cabbage (Shanghaiqing, *Brassica chinensis* L.) in rhizo-bags to separate the bulk soil from rhizosphere soil. Each rhizo-bag contained 160g soil (30 μ m nylon mesh, 4cm diameters, 6cm height), which only allowed the penetration of small molecular weight compounds and prohibited the penetration of plant roots.

In this experiment, 14-day-old Chinese cabbage seedlings were potted in the soil and treated with different fertilizer regimes in a greenhouse experiment located at the Institute of Urban Environment, Chinese Academy of Sciences. Each pot was transplanted with 3 plants. Only one plant was kept after survival. The experiment was conducted at room temperature (25-30 °C) with the relative humidity maintained at 70%. On a daily basis, deionized water was used for each pot to maintain the soil water content throughout the experiment. All samples, including bulk soil, rhizosphere soil, root, and Chinese cabbage leaves were collected after 60 days. The plants were harvested at full maturity.

All harvested samples were placed into aseptic bags and delivered on ice to the lab promptly after collection. A portion of the soil was sieved (mesh size: 2mm) to determine the selected soil properties and the rest was stored at -80 °C for DNA extraction. Plant samples were brought back

to the laboratory immediately for processing (Fig S1B).

2.3 Extraction and Purification of DNA Samples

DNA was extracted from soil (bulk and rhizosphere soil) according to the operation guide (*FastDNA Spin Kit for Soil* (MP Biomedical, CA)). Phyllosphere DNA was extracted according to previously published methods with slight modifications (Chen et al., 2017; Duran et al., 2018; Marasco et al., 2018).

Within 8 hours after harvesting, epiphytic microbes were isolated from the phyllosphere using extensive shaking in sterile water and phosphate buffer (8.5 g L⁻¹ Na₂HPO₄ anhydrous, 6.33 g L⁻¹ NaH₂PO₄, and pH = 6.5) added with 0.1% Triton X-100. Flushing liquid was stored as the epiphytic fraction after filtering through 0.22 µm pore size membranes. The extraction for DNA in the filtrate was processed using the *FastDNA Spin Kit for Soil*.

Root samples were rinsed with 80% ethanol and 0.25% NaClO to wash living microorganisms of the root surfaces and followed by a sterile water-rinse step three times (at 1 min each). The efficiency of the sterilization was evaluated through culturing on agar medium plates (15 g L⁻¹ agar, 3 g L⁻¹ MgSO₄, 10 mL L⁻¹ glycerol, 1.5 g L⁻¹ K₂HPO₄, 10 g L⁻¹ hydrolyzed casein, and 10 g L⁻¹ protease peptone for solid medium). The root was treated and homogenized by liquid nitrogen grinding plus. Then DNA was extracted with a *FastDNA Spin Kit for Soil* (Figure S1B).

2.4 Microbial Community Profiling

The DNA concentration of samples was determined through fluorescence quantitative analysis. Samples were diluted to 20 ng µL⁻¹ for downstream a two-step PCR amplification (Chen et al., 2017). In the first step, the V4-V5 variable region of bacterial *16S rRNA* (515F-907R) (Jing et al., 2015) and *fungus primers* (*gITS7* - *ITS4*) (Ihrmark et al., 2012) were amplified (Table S1).

PCR amplification was performed in sterile conditions with technical triplicates. Details of the PCR amplification program and reaction system were presented in Table S2. PCR products were checked on a 1% agarose gel with negative control. Purification was followed for replicated reactions: 1) the bacterial amplicons recovery of purified products was carried out following previous methods (Chen et al., 2019; Zhu et al., 2018); and 2) fungal amplicons were purified using the *Universal DNA Purification kit DP214*.

The DNA concentration was measured again with fluorescence quantitation analysis, and pooling 300 ng DNA of each barcoded amplicon in one library for the relevant microbial group. Barcoded amplicons sequencing was implemented with the *Illumina Hiseq2500* platform (*MICROANALY, Hefei, China*).

2.5 16S rRNA Gene and ITS Read Processing

Raw data were pre-processed in *VSEARCH v2.12.0* (Rognes et al., 2016) and *USEARCH v10.0* (Edgar, 2013), Zero-radius operational taxonomic units (ZOTUs) were denoised using the *UNOISE3* (Edgar, 2016) algorithm with a 100% sequence similarity. Double-end amplicon sequences were merged (minimum 50 bp overlap). After removing primers and barcode, chimeras, and quality filtered (largest prospective error threshold of 1.0) were removed with *USEARCH v10.0*, the rest of the high-quality amplicon reads were devoted to downstream analysis. Taxonomical classification based on ZOTUs was performed using *VSEARCH v2.12.0* and the *Silva/UNITE (Silva 123/UNITE 8.0)* database (Quast et al., 2013).

ZOTUs of samples that were in low mass number (< 10 sequence compositions) generally represent a PCR or sequencing error and were removed from the samples. All raw sequences were stored in the public databases (*National Center for Biotechnology Information Sequence Read*

Archive). That session specification is numbered SRP227163. Alpha-diversity for each sample was profiled through a measurement of Chao1 index, Buzas_gibson index and Shannon_e index, and box plots were described to compare the diversity level of fungal and bacterial ZOTU.

2.6 HT-qPCR and Data Analysis

SmartChip Real-time PCR system (Wafergen Inc., USA) was employed with HT-qPCR, which is specifically used for large gene expression studies (Wang et al., 2014). Compared with traditional qPCR, this system is a new dedicated high-throughput processing chip, which could greatly accelerate the performance of all the levels. Compared with shotgun metagenomic sequencing, the proposed system is more advantageous than shotgun metagenomic sequencing in operation, analytical method, and absolute quantification analysis, although it might not have a big database (Xiang et al., 2020). Primer sets targeting 283 ARGs, 12 marker genes for mobile genetic elements (MGEs) (Ouyang et al., 2015) including of four universal integron-integrase genes (*intI-1(clinic)*, *intI-1LC*, *intI2*, and *intI3*) and eight transposase genes, and one 16S rRNA gene (Table S5) were included in recent research work (Zheng et al., 2019).

The program and reaction system of HT-qPCR amplification was performed as outlined in Table S2. The qPCR results were examined employing SmartChip qPCR Software. Wells with PCR efficiencies are not in the normal range (1.8-2.2) or with multiple melting peaks were removed. Then, according to eq 1, the amplification efficiency was converted to 2. Each experiment was conducted and repeated three times and a detectable threshold cycle (CT) limit value was set to 31 (Zhu et al., 2013).

In the end, a valid value was determined only if all the repeated trials had amplification. The normalized copy number of MGEs and ARGs was measured and transformed to absolute gene copy

numbers and normalized to 16S rRNA gene copy numbers, which were quantified respectively from the Wafergen platform (eq 2) (Chen et al., 2016). To minimize error as a result of differences in 16S rRNA gene abundance between samples, a normalized copy number of ARGs per bacterial cell were used and calculated as follows (eq 3). (Klappenbach et al., 2001; Stalder et al., 2014; Zhu et al., 2018)

$$C_T = C_T' \log_2 E \quad (\text{eq1})$$

$$\text{Relative Gene Copy Number} = 10^{(31 - C_T)/(10/3)} \quad (\text{eq2})$$

$$\text{Normalized copy number of ARG gene} = (\text{Relative ARG gene copy number} / \text{Relative 16S rRNA gene copy number}) \times 4.1 \quad (\text{eq3})$$

Where C_T means the threshold value, 4.1 represents the average number of 16S rRNA gene per bacterium, the related calculation is applied according to the Ribosomal RNA Operon Copy Number Database

2.7 Statistical Analysis

All data analyses in the plots that follow (e.g. box plots) were computed with the default settings of ggplot2 (Hadley Wickham et al., 2016) for the software R (R Team, 2010). Statistical correlation analysis (Spearman's correlation) was undertaken through SPSS software (PASW Statistics 18.0). A co-association network of ARGs was performed using R with the psych packages and igraph (Adair et al., 2018; Csardi and Nepusz, 2006). Visualized bipartite network analysis was based on Gephi.

Based on Bray-Curtis distance, the paper makes a PCoA analysis of the ARGs, bacterial, and fungi community profiles. Procrustes analysis and Mantel analysis of correlation between ARGs, bacterial, and fungal communities were used in R package, vegan (Oksanen et al., 2011).

3. Results and Discussion

3.1 Distribution of MGEs and ARGs in the Soil-Chinese Cabbage System

A total of 12 MGEs and 249 ARGs were detected among all samples, which was higher than previously reported for ARGs in soil samples (Chen et al., 2017; Zhang et al., 2019). The diversity of MGEs and ARGs ranged from 25 to 181 in all samples, with phyllosphere (CK) and rhizosphere soil (soil amended with PWSC T3/S) harboring the lowest (25) and the highest (181) detected number of ARGs, respectively. One possible explanation of this result is that the rhizosphere is usually considered as the most active area in plant microbiome and rhizospheric microorganism has the potential of increasing the spread of ARB and ARG around the rhizosphere.

A total of 249 ARGs were identified, representing nine types, of which the majority belonged to vancomycin resistance genes, beta-lactams, tetracycline, and MLSB resistance genes (Fig 1). The detection frequency of the initial soil was similar to the CK at the end of the experiment, which was statistically lower than in the soil samples amended with compost (Adonis, $P < 0.05$). After compost incorporated into the soil, diversity of MGEs and ARGs was raised in each sample, including soil, root endophytes, and phyllospheric samples.

ARGs were detected by HT-qPCR in all samples. This proved that ARGs generally exist in the environment (D'Costa et al., 2006). The number of ARGs detected in phyllospheric samples was lower than those in the soil and root endophyte samples following amendment with different fertilizer treatments (Fig 1). This could be a result of the fact that the soil bacterial community was in direct contact with the composted material, which contains high levels of antibiotics and ARGs (Zhu et al., 2017). The number of MGEs and ARGs detected in the initial soil and compost samples combined was less than the detected number of MGEs and ARGs in the soil amended with compost.

This phenomenon can be well explained by the following two reasons: (1) the application of organic fertilizer effectively facilitated microorganism activity in the soil and then promoted the spread of antibiotic resistance (Zhu et al., 2013), and (2) the antibiotic residues in PWSCs acted as selective agents and enriched ARGs in the soil (Baym et al., 2016; Nolivos et al., 2019).

The change of normalized abundance and the absolute abundance of MGEs and ARGs were different in soils amended with different composts. Figure S2 showed the absolute abundance of ARGs in all soil and phyllospheric samples ranged from 6.16×10^8 to 1.68×10^{10} and from 3.06×10^7 to 1.38×10^8 copies g^{-1} solid (dry weight), respectively. It is clear that following compost (T1, T2, T3, and S) amendment, the absolute abundance of ARGs in soil, rhizosphere, and phyllosphere were raised (Fig S2). Compared with CK, the T1 amendment significantly increased ($p < 0.05$) the absolute abundance of ARGs in rhizosphere soil samples, while S amendment significantly raised ($p < 0.05$) ARGs in soil and phyllospheric samples, suggesting that S compost may also have a strong effect on the Chinese cabbage antibiotic resistome (Fig S2). Our results showed that the S-compost consistently resulted in the highest concentration of antibiotic residues as well as the largest number of ARGs (Fig 1 and Table S1), which infers that the antibiotic residues might be responsible for the elevated number of ARGs.

Based on the comparison of normalized results of MGEs and ARGs in bacterial cell numbers, we evaluated normalized abundance levels of MGEs and ARGs in the total microbial community (Fig S3). In the bulk soil, compost (T1, T2, T3, and S) applications consistently led to increases in the normalized abundance of MGEs and ARGs (Fig S3). In the rhizosphere soil, root endophytes, and phyllospheric samples, whilst there were differences in the normalized abundance of MGEs and ARGs, the increase in abundance did not always coincide compared with CK (Fig S3). In all samples,

the bulk soil harbored a greater resistome than the rhizosphere soil and the absolute abundance of MGEs and ARGs in all soil samples were greater than that in the phyllosphere (Fig 2).

Our results were consistent with a previously published research confirming that the rhizosphere was a key area for propagation and spread of antibiotic resistome and horizontal transfer of ARGs (Zhang et al., 2019). Previous studies have postulated that the rhizosphere microbes provide a range of ecosystem services for plants, such as nutrient acquisition, and abiotic stress tolerance (Meena et al., 2017; Mendes et al., 2014).

Hence, this study further highlights the importance of studying ARGs and ARB in the rhizosphere. Diverse and abundant ARGs were detected in different environments using different new molecular tools. The metagenomic approach is considered as a feasible method for studying the environmental dimension of antibiotic resistance. For example, Li et al. (2015) identified 260 ARG from 6 different typical environment samples including soil, water, sediment, excrement, wastewater biofilm, and sludge. Moreover, diverse ARGs (381 ARG subtypes) were also detected in urban sewage samples from 32 municipal wastewater treatment plants by Su et al. (2017). High-throughput qPCR based ARG chip as another popular approach for studying ARGs in environment, has the advantage of measuring the absolute abundance of ARGs, which provide a more intuitive way to describe the potential risk. We propose combining various alternative methods might have great potentials in providing more comprehensive understanding of ARGs in environments and their dissemination.

In addition to the changes in the diversity and abundance of MGEs and ARGs, even ARG subtypes and MGEs turned out to have more variants (interactions) in network analysis (correlation coefficient $> |0.6|$ and P-value < 0.05), mostly in large proportions (Fig S4). Co-occurrence styles

among MGEs and ARG subtypes in the soil-Chinese cabbage system were evaluated using network analysis. Network analysis results suggested that the use of composts (T1, T2, T3, and S) might promote the propagation and spread of ARGs by way of HGT (Fig S4, Table S1). Of all networks, the node was considered the “hub”, chiefly meaning the most continually connected or the most abundant node in each network. For example, *vanC2/vanC3* was the hub gene for the network in the control soil-Chinese cabbage system (CK) (Fig S4A), *ermX* was a “hub” gene for the network in T1 amended-soil (Fig S4B), *bla_{CTX-M-01}* was a “hub” gene for the network in T2 amended-soil (Fig S4C), *tetG-01* was a “hub” gene for the network in T3 amended-soil (Fig S4D) and *bla_{IMP-02}* was a “hub” gene for the network in S amended-soil (Fig S4E).

Our network analyses also indicated that MGEs co-occurred with diverse ARG subtypes in cabbage-soil systems. Among the detected MGEs, the transposase genes (e.g., *tnpA-01*, *tnpA-02*, *tnpA-04* and *tnpA-05*) and the integrase genes (e.g., *intI-ILC* and *intI-1(clinic)*) showed significant co-occurrence with related ARGs in the S amended soil-Chinese cabbage system, with matching degrees of 30, 33, 32, 32, 30 and 30 (Fig S4E). Bray-Curtis distance and PCoA analysis (Fig S5) indicated that the structure and distribution of ARGs in root endophytes and phyllospheric samples were significantly different from soil samples (Adonis, $P < 0.01$).

Network analysis is deemed to be one of the most effective methods to uncover the co-occurrence patterns of MGEs and ARGs (Forsberg et al., 2014; Li et al., 2015). Network analysis carried out in this study, supported previous observations regarding the relationship between ARGs and MGEs (An et al., 2018; Li et al., 2015; Quintela-Baluja et al., 2019). Network analysis also demonstrated that the relationship between ARGs and MGEs became more intertwined and complex following the introduction of compost into the soil eco-system. The ARGs exhibited a complex

interaction with the MGEs, resulting in a more uniform deformation compared with the previous study (Liu et al., 2019). Co-occurrence networks represented that there was an "easy correlation" between two genes, but without unequivocal demonstrable evidence of direct association and interaction. It still needs further identifying specific ARGs-MGEs interactions and associates them with microbial community structures and ecological functions.

3.2 Microbiota in Soil and Chinese Cabbage Samples

A total of 3,924-69,847 and 2,525-93,570 effective tags were acquired for the bacterial and fungal communities, respectively, through 16S rRNA and ITS amplicon sequencing of all samples.

3.2.1 Bacterial Community

Bacterial community data were further evaluated using the Buzas_gibson and Shannon_e diversity index to explore changes in bacterial diversity. Compared with those of the CK, the Shannon_e and Buzas_gibson indices increased in the soil amended with fertilizer (Fig S6). The Shannon_e and Buzas_gibson indices of the bacterial community were more sensitive to the T2 than that of the other groups (Fig S6). In bulk and rhizosphere soil but not showed the cabbage phyllosphere, alpha diversity of bacterial after application of compost was raised significantly with obvious features ($P < 0.05$) (Fig S6). Calculated alpha diversity revealed samples from the phyllosphere had the lowest bacterial alpha diversity, followed by bulk soil and then rhizosphere soil (Fig S6).

The bacterial community structure of the soil (bulk soil and rhizosphere soil) samples at the phylum level was shown in Fig S7A. Firmicutes, Actinobacteria, Acidobacteria, and Proteobacteria were the four dominant phyla in all soil (bulk soil and rhizosphere soil) samples, with their relative

abundance accounted for 68.2-79% totally (Fig S7A). At the class level, Bacilli (ranging from 7.88 to 18.5%), Acidobacteria (ranging from 8.28 to 16.2%), and Alphaproteobacteria (ranging from 7.4 to 15.2%) were the dominant classes of all soil (bulk soil and rhizosphere soil) samples (Fig S7B).

3.2.2 Fungal Community

Compared with the bulk soil, Chao1 and Richness indices increased in the rhizosphere soil. Comparatively, when amended with T1, T2, T3, and S, the Chao1 index declined by 7.36%-30.60% and 7.49%-15.90% in rhizosphere soil (except for T1) and bulk soil, respectively (Fig S9).

In the bulk soil, at the phylum level, the abundance of Ascomycota accounted for 73.93-92.13% of the total composition of all libraries, followed by Chytridiomycota (0.11-20.78%), Basidiomycota (1.00-7.00%), and Zygomycota (0.34-5.58%) (Except for CK). (Fig S10 A) In rhizosphere soil, at the phylum level, the abundance of Ascomycota accounted for 88.9-98.1% of the total composition of all libraries, followed by Zygomycota (0.32-6.57%), Basidiomycota (0.63-4.57%), and Chytridiomycota (0.01-2.02%) (Fig S10 A).

In all soil samples, in comparison to the CK, the abundance of Ascomycota increased following the addition of compost, the highest value reached at 36.03% (Fig S10 A). The abundance of Ascomycota in samples from bulk soil had the lowest value, followed by rhizosphere soil (Fig S10 A). Meanwhile, those abundance of Basidiomycota and Zygomycota increased by 0.24-0.83% (except for treatment S) and 0.04-6.25%, respectively.

In the phyllosphere, compared with the CK, the Chao1 index and Richness increased by 28.42% and 30% respectively in treatments spiked with T3, with minimal changes after the addition of T1, T2, and S (Fig S9 C and D). Ascomycota and Basidiomycota were the dominant fungal community in the phyllosphere and accounted between 68.94- 90.61% of the total composition in all libraries.

In comparison to the control treatments, species of Ascomycota were more sensitive to the addition of T1. Consequently, a decrease ranging between 34.51 and 95.38% was observed for Basidiomycota (except for S), while increases between 4.28-41.49% were found for Ascomycota (Fig S10 E).

Our findings support previous research, which observed that the addition of fertilizer can affect the composition and structure of the plant rhizosphere fungal and bacterial communities. The additional compost has been shown to positively influence bacterial richness and diversity in soil, and that over time, the bacterial community structure becomes more stable. (Kavamura et al., 2018)

Soil microbial communities affect plant growth, resource use efficiency, and health, especially the microbial communities that are picked up according to vegetation to form the rhizosphere microbiome (Berendsen et al., 2012; Rodrigo et al., 2013). The bacterial and fungal community compositions in phyllospheric samples were similar to the rhizosphere soil samples, particularly at the phylum level (Fig S7, S8, and S10). We can conclude that a certain amount of fungal and bacterial communities, derived from amended soil, subsequently migrated to the vegetable tissue and colonized the endophyte.

However, phyllospheric samples, principal fungal and bacterial families were different from the soil samples (Fig S7, S8, and S10). It has been suggested that in comparison to endophyte samples, phyllospheric samples, are more sensitive to the airborne fungal and bacterial community than the soil fungal and bacterial communities, which would also support our findings (Yan et al., 2019). We observed differences in phyllospheric samples and bulk soil samples for the major ARGs detected (aminoglycoside, beta-lactams, and multidrug resistance genes) (Fig 1 and 2). This finding may be ascribed to the unique microbial community composition in the phyllosphere, where particular

microbial ZOTUs are able to port only specific types of ARGs (Li et al., 2015).

3.3 Pathway of ARGs from PWSC to Phyllosphere

Diverse shared ARGs were found among rhizosphere soil (compost amended-soil), root, and Chinese cabbage phyllosphere, and those only detected in individual samples were represented utilizing a bipartite network analysis (Fig 2). Seven categories of ARGs (I-VII) were identified based on the detected sample types, with two categories identified, where the antibiotic resistome was shared between the Chinese cabbage phyllosphere and environment following compost treatment (namely, IV and VI).

Cluster IV contained 20 (Fig 2a), 12 (Fig 2b), 1 (Fig 2c), and 18 (Fig 2d) ARGs, respectively, which were detected simultaneously in the Chinese cabbage phyllosphere and root samples. Cluster VI contained 19 (Fig 2a), 15 (Fig 2b), 12 (Fig 2c), and 21 (Fig 2d) ARGs, respectively, which were detected simultaneously in the Chinese cabbage phyllosphere and the amended rhizosphere soil. The genes in these two categories conferred resistance to multidrug, beta-lactamase, aminoglycoside, MLSB, vancomycin, and tetracycline, and were possible candidates for moving resistance from the soil, to the endophyte and phyllosphere. The ARGs in Cluster VII were shared in the middle of three compartments (Chinese cabbage phyllosphere, rhizosphere soil, and root). In PWSC-T1 amended soils, these genes included 69 ARGs and 5 MGEs (*tnpA*-02, *tnpA*-04, *tnpA*-05, *intI*-1 (*clinic*), *intI*-1(*ILC*), most of which conferred multidrug resistance as well as resistance to beta-lactamase, aminoglycoside, MLSB, and tetracycline (Fig 2a).

In PWSC-T2 amended soils, these genes included 53 ARGs and 6 MGEs (*tnpA*-01, *intI*3, *tnpA*-02, *intI*-1(*ILC*), *intI*-1(*clinic*), *tnpA*-05), most of which conferred multidrug resistance and resistance to beta-lactamase, sulfonamide, aminoglycoside, and MLSB (Fig 2b). In PWSC-T3 amended soil,

these genes included 60 ARGs and 3 MGEs (*intl-ILC*, *intl-1(clinic)*, *tnpA-01*), the majority of which conferred resistance to vancomycin, sulfonamide, aminoglycoside, multidrug, beta-lactamase, and MLSB (Fig 2c). In S amended-soil, these genes included 60 ARGs and 5 MGEs (*tnpA-02*, *tnpA-04*, *intl-ILC*, *intl-1(clinic)*, *tnpA-05*), most of which delivered multidrug resistance as well to aminoglycoside, beta-lactamase and MLSB (Fig 2d). The bipartite network graph of PWSC treatment (T1-T3) revealed a similar layout to the S treatment. However, there were generally fewer shared ARGs clusters between the PWSC amended rhizosphere soil, Chinese cabbage phyllosphere, and root endophyte samples (Fig 2).

Our findings provide further evidence that compost amended soil-derived ARGs are a key origin of the antibiotic resistome found in plants (Chen et al., 2017; Zhang et al., 2019; Zhu et al., 2017). To understand further the migration pattern of the PWSC-derived antibiotic resistome to Chinese cabbage tissues and soil. We have illustrated the possible routes in the soil-plant system (Fig 2). The migration of ARGs from amended soil to plant through the plant tissues was accounted for as an inherent route.

All categories in the same soil-Chinese cabbage system shared and exchanged ARGs effortlessly across categories, especially the exchange of shared ARGs. For example, in the PWSC-T1 treatment, we found that 121 ARGs were shared between root endophyte and rhizosphere soil. Of most interest were the 74 genes shared solely by rhizosphere soil and Chinese cabbage phyllosphere, which provided clear evidence of ARGs migrating from amended soil to the Chinese cabbage phyllosphere. The specific concern is infections caused by multidrug-resistant gram-positive pathogens. Vancomycin, which is an important medical measure to assure the safe running of the hospital system, is the last line of defense to avoid the hospital infection from complete

collapse. Agent resistant genes to vancomycin, except for T1, were not detected in the phyllosphere in this study. A study on biosafety is an important aspect among studies on applying fertilizer of PWSC.

3.4 Shared Microbiota and Antibiotic Resistome among Chinese Cabbage and Amended Soil Samples

Procrustes analysis and Mantel analysis were used to explore shared ZOTUs associated with the antibiotic resistome composition from Chinese cabbage phyllosphere, root, and rhizosphere soil samples. Our study indicated that ARG profiles were associated with the shared bacterial structures and composition in both the control group and treatment groups, especially for T1 (Fig 3). Further Procrustes analyzes of the ARG profiles and bacterial 16S rRNA gene clustered the samples automatically according to the type of sample, and the foundation of Bray-Curtis dissimilarity metrics, the results offered high accuracy. (Table S3, Fig 3)

We also described the fungal communities' roles in the dissemination of ARGs. Our findings, in agreement with previous studies, show that ARG profiles were remarkably associated with the shared fungal structures and composition in both the control group and treatment group. Procrustes analyzes of ARG profiles and fungal ITS gene clustered the samples automatically according to the type of sample, and the foundation of Bray-Curtis dissimilarity metrics fitting the results offered high accuracy (Table S4, Fig 4).

Studies have suggested that shared ARGs were found in both the endophytes and phyllosphere of plants cultured in soil amended with manure. For instance, Bai et al. (2015) observed that in terms of taxonomic structure, leaf and root microbiota, were extensively overlapped whilst. Knief et al.

(2012) observed that the dinitrogen reductase gene and dinitrogenase gene, were in both the rhizosphere and phyllosphere metagenome, indicating that at a functional gene level, these genes were partially overlapping in the leaf and root microbiota.

Shared ZOTUs and shared ARGs imply that phyllosphere and root microbiota might interconnect fungal and bacterial communities including antibiotic-resistant microbiota which have the potential to move between the rhizosphere and phyllosphere (Beattie and Lindow, 1999; Ruiz-Pérez et al., 2016). A study has indicated that roots can effectively recruit soil bacteria or fungal communities to colonize the root, allow antibiotic resistome to migrate from the soil into the phyllosphere and roots (Bulgarelli et al., 2012). This might be a key migration path for the antibiotic resistome in organic fertilizer to move into plants, whereby in this process, the roots can act as a bridge. The results showed that shared ZOTUs support the origin of ARGs by a host in the bacterial or fungal communities. Ultimately, the phyllosphere and root endophyte in Chinese cabbage forms their own ARG compositions.

4. Conclusions

In summary, this research showed that the application of both PWSCs (T1-T3) and S increased the diversity and absolute abundance of the ARGs in the phyllosphere, root, rhizosphere soil, and bulk soil of Chinese cabbage. ARGs were examined in the Chinese cabbage microbiome and disproportionally distributed in distinguishing parts (e.g. Phyllosphere and root) of Chinese cabbage. Enrichment of Chinese cabbage antibiotic resistome was associated with fungal and bacterial taxa. Based on these findings, we suggested that rhizosphere soil, root, and phyllosphere microbiota might interconnect with microbial communities (fungal and bacterial communities), including

antibiotic-resistant microbiota, potentially moving between the rhizosphere and phyllosphere. In this process, we propose that the roots play the role of a bridge.

Nevertheless, shared antibiotic resistome in the amended soil-plant system indicates a feasible pathway of ARGs migration into human pathogens and the microbiome via the food chain (e.g. Vegetable and salad). These observations are critical for the assessment of public health risks on Chinese cabbage grown in the soil amendment dealt with PWSCs. Soil utilization of sewage sludge is an important universal method of treating sewage sludge. How to prevent the biological pollution is always a much-discussed problem, especially on the pharmaceutical waste sludge. This research has significant implications for the recycling of pharmaceutical waste sludge, to improve the sustainable agricultural practices and the human health protection.

Acknowledgments

This research is based upon the fund supports provided by Ministry of Science and Technology of China (MSTC) with National Key Research and Development Program (2017YFE0119000), National Natural Science Foundation of China (41701564, 41850410497, 41571130063), Science and Technology Project in Suzhou (SNG201613), Natural Science Fund for Colleges and Universities in Jiangsu Province (17KJB610010), Graduate Student Scientific Research Innovation Projects in Jiangsu Province (SJCX17_0674), and Xiamen Bureau of Science and Technology of China (3502Z20193075). Finally, we thank Yi Zhao for her contributions to constructive comments.

References

- Adair, K.L., Wilson, M., Bost, A., Douglas, A.E., 2018. Microbial community assembly in wild populations of the fruit fly *Drosophila melanogaster*. *ISME J.* 12(4), 959-972. doi:10.1038/s41396-017-0020-x.
- Allen, H.K., Donato, J., Wang, H.H., Cloud-Hansen, K.A., Davies, J., Handelsman, J., 2010. Call of the

509 wild: antibiotic resistance genes in natural environments. *Nature Reviews Microbiology*. 8(4), 251-259.
510 doi:10.1038/nrmicro2312.

511 Amy, P., Ruoting, P., Heather, S., Carlson, K.H., 2006. Antibiotic resistance genes as emerging
512 contaminants: studies in northern Colorado. *Environ Sci Technol*. 40(23), 7445. doi:10.1021/es0680156.

513 An, X.L., Chen, Q.L., Zhu, D., Su, J.Q., 2018. Distinct effects of struvite and biochar amendment on the
514 class 1 integron antibiotic resistance gene cassettes in phyllosphere and rhizosphere. *Sci Total Environ*.
515 631-632, 668-676. doi:10.1016/j.scitotenv.2018.03.034.

516 Bai, Y., Muller, D.B., Srinivas, G., Garrido-Oter, R., Potthoff, E., Rott, M., Dombrowski, N., Munch,
517 P.C., Spaepen, S., Remus-Emsermann, M., Huttel, B., McHardy, A.C., Vorholt, J.A., Schulze-Lefert, P.,
518 2015. Functional overlap of the Arabidopsis leaf and root microbiota. *Nature*. 528(7582), 364-369.
519 doi:10.1038/nature16192.

520 Baym, M., Lieberman, T.D., Kelsic, E.D., Chait, R., Gross, R., Yelin, I., Kishony, R., 2016.
521 Spatiotemporal microbial evolution on antibiotic landscapes. *Science*. 353(6304), 1147-1151.
522 doi:10.1126/science.aag0822.

523 Beattie, G.A., Lindow, S.E., 1999. Bacterial colonization of leaves: a spectrum of strategies.
524 *Phytopathology*. 89(5), 353-359. doi:10.1094/PHYTO.1999.89.5.353.

525 Berendsen, R.L., Pieterse, C.M., Bakker, P.A., 2012. The rhizosphere microbiome and plant health.
526 *Trends Plant Sci*. 17(8), 478-486. doi:10.1016/j.tplants.2012.04.001.

527 Bernal, M.P., Alburquerque, J.A., Moral, R., 2009. Composting of animal manures and chemical criteria
528 for compost maturity assessment. A review. *Bioresour Technol*. 100(22), 5444-5453.
529 doi:10.1016/j.biortech.2008.11.027.

530 Bulgarelli, D., Rott, M., Schlaeppi, K., Ver Loren van Themaat, E., Ahmadinejad, N., Assenza, F., Rauf,
531 P., Huettel, B., Reinhardt, R., Schmelzer, E., Peplies, J., Gloeckner, F.O., Amann, R., Eickhorst, T.,
532 Schulze-Lefert, P., 2012. Revealing structure and assembly cues for Arabidopsis root-inhabiting bacterial
533 microbiota. *Nature*. 488(7409), 91-95. doi:10.1038/nature11336.

534 Chen, Q.-L., An, X.-L., Li, H., Zhu, Y.-G., Su, J.-Q., Cui, L., 2017. Do manure-borne or indigenous soil
535 microorganisms influence the spread of antibiotic resistance genes in manured soil? *Soil Biology and*
536 *Biochemistry*. 114, 229-237. doi:10.1016/j.soilbio.2017.07.022.

537 Chen, Q.-L., Ding, J., Zhu, D., Hu, H.-W., Delgado-Baquerizo, M., Ma, Y.-B., He, J.-Z., Zhu, Y.-G., 2019.
538 Rare microbial taxa as the major drivers of ecosystem multifunctionality in long-term fertilized soils.
539 *Soil Biology and Biochemistry*. doi:10.1016/j.soilbio.2019.107686.

540 Chen, Q., An, X., Li, H., Su, J., Ma, Y., Zhu, Y.G., 2016. Long-term field application of sewage sludge
541 increases the abundance of antibiotic resistance genes in soil. *Environ Int*. 92-93, 1-10.

542 Chen, Q.L., An, X.L., Zhu, Y.G., Su, J.Q., Gillings, M.R., Ye, Z.L., Cui, L., 2017. Application of Struvite
543 Alters the Antibiotic Resistome in Soil, Rhizosphere, and Phyllosphere. *Environ Sci Technol*. 51(14),
544 8149-8157. doi:10.1021/acs.est.7b01420.

545 Chen, Q.L., Cui, H.L., Su, J.Q., Penuelas, J., Zhu, Y.G., 2019. Antibiotic Resistomes in Plant
546 Microbiomes. *Trends Plant Sci*. doi:10.1016/j.tplants.2019.02.010.

547 Cheng, W., Chen, H., Su, C., Yan, S., 2013. Abundance and persistence of antibiotic resistance genes in
548 livestock farms: A comprehensive investigation in eastern China. *Environ Int*. 61(4), 1-7.

549 Csardi, G., Nepusz, T., 2006. The Igraph Software Package for Complex Network Research. *Interjournal*
550 *Complex Systems*. 1695.

551 D'Costa, V.M., McGrann, K.M., Hughes, D.W., Wright, G.D., 2006. Sampling the Antibiotic Resistome.
552 *Science*. 311(5759), 374-377. doi:10.1126/science.1120800.

553 Edgar, R.C., 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nature*
554 *Methods*. 10(10), 996-998. doi:10.1038/nmeth.2604.

555 Edgar, R.C., 2016. UNOISE2: improved error-correction for Illumina 16S and ITS amplicon sequencing.
556 *bioRxiv*. 081257. doi:10.1101/081257.

557 Forsberg, K.J., Patel, S., Gibson, M.K., Lauber, C.L., Knight, R., Fierer, N., Dantas, G., 2014. Bacterial
558 phylogeny structures soil resistomes across habitats. *Nature*. 509(7502), 612-616.
559 doi:10.1038/nature13377.

560 Gillings, M.R., 2017. Lateral gene transfer, bacterial genome evolution, and the Anthropocene. *Ann N*
561 *Y Acad Sci*. 1389(1), 20-36. doi:10.1111/nyas.13213.

562 Hadley Wickham, Winston Chang, Lionel Henry, Thomas Lin Pedersen, Kohske Takahashi, Claus
563 Wilke, ., K.W., 2016. ggplot2: Create Elegant Data Visualisations Using the Grammar of Graphics. *Book*
564 *of Abstracts*.

565 Ihrmark, K., Bodeker, I.T., Cruz-Martinez, K., Friberg, H., Kubartova, A., Schenck, J., Strid, Y., Stenlid,
566 J., Brandstrom-Durling, M., Clemmensen, K.E., Lindahl, B.D., 2012. New primers to amplify the fungal
567 ITS2 region--evaluation by 454-sequencing of artificial and natural communities. *FEMS Microbiol Ecol*.
568 82(3), 666-677. doi:10.1111/j.1574-6941.2012.01437.x.

569 Jing, X., Sanders, N.J., Shi, Y., Chu, H., Classen, A.T., Zhao, K., Chen, L., Shi, Y., Jiang, Y., He, J.-S.,
570 2015. The links between ecosystem multifunctionality and above- and belowground biodiversity are
571 mediated by climate. *Nature Communications*. 6(1). doi:10.1038/ncomms9159.

572 Kavamura, V.N., Hayat, R., Clark, I.M., Rossmann, M., Mendes, R., Hirsch, P.R., Mauchline, T.H., 2018.
573 Inorganic Nitrogen Application Affects Both Taxonomical and Predicted Functional Structure of Wheat
574 Rhizosphere Bacterial Communities. *Front Microbiol*. 9, 1074. doi:10.3389/fmicb.2018.01074.

575 Klappenbach, J.A., Saxman, P.R., Cole, J.R., Schmidt, T.M., 2001. rrndb: the Ribosomal RNA Operon
576 Copy Number Database. *Nucleic Acids Res*. 29(1), 181-184. doi:10.1093/nar/29.1.181.

577 Knief, C., Delmotte, N., Chaffron, S., Stark, M., Innerebner, G., Wassmann, R., Von, M.C., Vorholt, J.A.,
578 2012. Metaproteogenomic analysis of microbial communities in the phyllosphere and rhizosphere of rice.
579 *ISME J*. 6(7), 1378. doi:10.1038/ismej.2011.192.

580 Li, B., Yang, Y., Ma, L., Ju, F., Guo, F., Tiedje, J.M., Zhang, T., 2015. Metagenomic and network analysis
581 reveal wide distribution and co-occurrence of environmental antibiotic resistance genes. *ISME J*. 9(11),
582 2490-2502. doi:10.1038/ismej.2015.59.

583 Liu, J., Meng, Z., Liu, X., Zhang, X.-H., 2019. Microbial assembly, interaction, functioning, activity and
584 diversification: a review derived from community compositional data. *Marine Life Science &*
585 *Technology*. 1(1), 112-128. doi:10.1007/s42995-019-00004-3.

586 Liu, M., Ding, R., Zhang, Y., Gao, Y., Tian, Z., Zhang, T., Yang, M., 2014. Abundance and distribution
587 of Macrolide-Lincosamide-Streptogramin resistance genes in an anaerobic-aerobic system treating
588 spiramycin production wastewater. *Water Res*. 63, 33-41. doi:10.1016/j.watres.2014.05.045.

589 Liu, M., Ni, H., Yang, L., Chen, G., Yan, X., Leng, X., Liu, P., Li, X., 2019. Pretreatment of swine manure
590 containing β -lactam antibiotics with whole-cell biocatalyst to improve biogas production. *Journal of*
591 *Cleaner Production*. 240. doi:10.1016/j.jclepro.2019.118070.

592 Liu, M., Zhang, Y., Yang, M., Tian, Z., Ren, L., Zhang, S., 2012. Abundance and Distribution of
593 Tetracycline Resistance Genes and Mobile Elements in an Oxytetracycline Production Wastewater
594 Treatment System. *Environ Sci Technol*. 46(14), 7551-7557. doi:10.1021/es301145m.

595 Liu, X., Lendormi, T., Lanoisell, J.-L., 2019. Overview of hygienization pretreatment for pasteurization
596 and methane potential enhancement of biowaste: Challenges, state of the art and alternative technologies.

Journal of Cleaner Production. 236. doi:10.1016/j.jclepro.2019.06.356.

Luo, X., Liu, G., Xia, Y., Chen, L., Jiang, Z., Zheng, H., Wang, Z., 2016. Use of biochar-compost to improve properties and productivity of the degraded coastal soil in the Yellow River Delta, China. *Journal of Soils and Sediments*. 17(3), 780-789. doi:10.1007/s11368-016-1361-1.

Martínez Salgado, M.M., Ortega Blu, R., Janssens, M., Fincheira, P., 2019. Grape pomace compost as a source of organic matter: Evolution of quality parameters to evaluate maturity and stability. *Journal of Cleaner Production*. 216, 56-63. doi:10.1016/j.jclepro.2019.01.156.

Meena, K.K., Sorty, A.M., Bitla, U.M., Choudhary, K., Gupta, P., Pareek, A., Singh, D.P., Prabha, R., Sahu, P.K., Gupta, V.K., Singh, H.B., Krishanani, K.K., Minhas, P.S., 2017. Abiotic Stress Responses and Microbe-Mediated Mitigation in Plants: The Omics Strategies. *Front Plant Sci*. 8, 172. doi:10.3389/fpls.2017.00172.

Mendes, L.W., Kuramae, E.E., Navarrete, A.A., van Veen, J.A., Tsai, S.M., 2014. Taxonomical and functional microbial community selection in soybean rhizosphere. *ISME J*. 8(8), 1577-1587. doi:10.1038/ismej.2014.17.

Nolivos, S., Cayron, J., Dedieu, A., Page, A., Delolme, F., Lesterlin, C., 2019. Role of AcrAB-TolC multidrug efflux pump in drug-resistance acquisition by plasmid transfer. *Science*. 364(6442), 778-782. doi:10.1126/science.aav6390.

Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P., 2011. The vegan package: Community Ecology Package. R package version 2.0--2.

Ouyang, W.Y., Huang, F.Y., Zhao, Y., Li, H., Su, J.Q., 2015. Increased levels of antibiotic resistance in urban stream of Jiulongjiang River, China. *Appl Microbiol Biotechnol*. 99(13), 5697-5707. doi:10.1007/s00253-015-6416-5.

Qiao, M., Ying, G.G., Singer, A.C., Zhu, Y.G., 2018. Review of antibiotic resistance in China and its environment. *Environ Int*. 110, 160-172. doi:10.1016/j.envint.2017.10.016.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glockner, F.O., 2013. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res*. 41(Database issue), D590-596. doi:10.1093/nar/gks1219.

Quintela-Baluja, M., Abouelnaga, M., Romalde, J., Su, J.Q., Yu, Y., Gomez-Lopez, M., Smets, B., Zhu, Y.G., Graham, D.W., 2019. Spatial ecology of a wastewater network defines the antibiotic resistance genes in downstream receiving waters. *Water Res*. 162, 347-357. doi:10.1016/j.watres.2019.06.075.

Rodrigo, M., Paolina, G., Raaijmakers, J.M., 2013. The rhizosphere microbiome: significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. *Fems Microbiology Reviews*. 37(5), 634-663. doi:10.1111/1574-6976.12028.

Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahe, F., 2016. VSEARCH: a versatile open source tool for metagenomics. *PeerJ*. 4, e2584. doi:10.7717/peerj.2584.

Romain, M., Andrew, S., Yuan-Ching, T., Roger, M., Lyne, S., Yun, Z., Edward, T., 2013. Impact of manure fertilization on the abundance of antibiotic-resistant bacteria and frequency of detection of antibiotic resistance genes in soil and on vegetables at harvest. *Applied & Environmental Microbiology*. 79(18), 5701-5709. doi:10.1128/AEM.01682-13.

Ruiz-Pérez, C.A., Restrepo, S., Zambrano, M.M., 2016. Microbial and Functional Diversity within the Phyllosphere of *Espeletia* sp. in an Andean High Mountain Ecosystem. *Applied & Environmental Microbiology*. 82(6), 1807. doi:10.1128/AEM.02781-15.

Selvam, A., Wong, J.W.C., 2017. Degradation of Antibiotics in Livestock Manure During Composting. 267-292. doi:10.1016/b978-0-444-63664-5.00012-5.

Stalder, T., Barraud, O., Jove, T., Casellas, M., Gaschet, M., Dagot, C., Ploy, M.C., 2014. Quantitative and qualitative impact of hospital effluent on dissemination of the integron pool. *ISME J.* 8(4), 768-777. doi:10.1038/ismej.2013.189.

Su, J.Q., An, X.L., Li, B., Chen, Q.L., Gillings, M.R., Chen, H., Zhang, T., Zhu, Y.G., 2017. Metagenomics of urban sewage identifies an extensively shared antibiotic resistome in China. *Microbiome*. 5(1), 84. doi:10.1186/s40168-017-0298-y.

Su, J.Q., Wei, B., Ou-Yang, W.Y., Huang, F.Y., Zhao, Y., Xu, H.J., Zhu, Y.G., 2015. Antibiotic resistome and its association with bacterial communities during sewage sludge composting. *Environ Sci Technol.* 49(12), 7356-7363. doi:10.1021/acs.est.5b01012.

Tao, W., Zhang, X.X., Zhao, F., Huang, K., Ma, H., Wang, Z., Ye, L., Ren, H., 2016. High Levels of Antibiotic Resistance Genes and Their Correlations with Bacterial Community and Mobile Genetic Elements in Pharmaceutical Wastewater Treatment Bioreactors. *Plos One*. 11(6), e0156854. doi:10.1371/journal.pone.0156854.

Team, C.R., 2010. Team RDC.R: A Language And Environment For Statistical Computing. R Foundation for Statistical Computing: Vienna, Austria. *Computing*. 14, 12-21.

Tong, J., Lu, X., Zhang, J., Sui, Q., Wang, R., Chen, M., Wei, Y., 2017. Occurrence of antibiotic resistance genes and mobile genetic elements in enterococci and genomic DNA during anaerobic digestion of pharmaceutical waste sludge with different pretreatments. *Bioresour Technol.* 235, 316-324. doi:10.1016/j.biortech.2017.03.104.

Wang, F.H., Qiao, M., Su, J.Q., Chen, Z., Zhou, X., Zhu, Y.G., 2014. High throughput profiling of antibiotic resistance genes in urban park soils with reclaimed water irrigation. *Environ Sci Technol.* 48(16), 9079-9085. doi:10.1021/es502615e.

Wang, W., Yang, M., Shen, P., Zhang, R., Qin, X., Han, J., Li, Y., Wen, X., Liao, Y., 2019. Conservation tillage reduces nitrous oxide emissions by regulating functional genes for ammonia oxidation and denitrification in a winter wheat ecosystem. *Soil and Tillage Research*. 194. doi:10.1016/j.still.2019.104347.

Wright, G.D., 2007. The antibiotic resistome: the nexus of chemical and genetic diversity. *Nat Rev Microbiol.* 5(3), 175-186. doi:10.1038/nrmicro1614.

Xiang, Q., Zhu, D., Giles, M., Neilson, R., Yang, X.-R., Qiao, M., Chen, Q.-L., 2020. Agricultural activities affect the pattern of the resistome within the phyllosphere microbiome in peri-urban environments. *J Hazard Mater.* 382, 121068. doi:10.1016/j.jhazmat.2019.121068.

Xie, W.Y., McGrath, S.P., Su, J.Q., Hirsch, P.R., Clark, I.M., Shen, Q., Zhu, Y.G., Zhao, F.J., 2016a. Long-Term Impact of Field Applications of Sewage Sludge on Soil Antibiotic Resistome. *Environ Sci Technol.* 50(23), 12602-12611. doi:10.1021/acs.est.6b02138.

Xie, W.Y., Yang, X.P., Li, Q., Wu, L.H., Shen, Q.R., Zhao, F.J., 2016b. Changes in antibiotic concentrations and antibiotic resistome during commercial composting of animal manures. *Environ Pollut.* 219, 182-190. doi:10.1016/j.envpol.2016.10.044.

Yan, Z.Z., Chen, Q.L., Zhang, Y.J., He, J.Z., Hu, H.W., 2019. Antibiotic resistance in urban green spaces mirrors the pattern of industrial distribution. *Environ Int.* 132, 105106. doi:10.1016/j.envint.2019.105106.

Zhang, J., Lin, H., Ma, J., Sun, W., Yang, Y., Zhang, X., 2019. Compost-bulking agents reduce the reservoir of antibiotics and antibiotic resistance genes in manures by modifying bacterial microbiota. *Sci Total Environ.* 649, 396-404. doi:10.1016/j.scitotenv.2018.08.212.

Zhang, Q.Q., Ying, G.G., Pan, C.G., Liu, Y.S., Zhao, J.L., 2015. Comprehensive evaluation of antibiotics emission and fate in the river basins of China: source analysis, multimedia modeling, and linkage to

bacterial resistance. *Environ Sci Technol.* 49(11), 6772-6782. doi:10.1021/acs.est.5b00729.

Zhang, Y.J., Hu, H.W., Chen, Q.L., Singh, B.K., Yan, H., Chen, D., He, J.Z., 2019. Transfer of antibiotic resistance from manure-amended soils to vegetable microbiomes. *Environ Int.* 130, 104912. doi:10.1016/j.envint.2019.104912.

Zheng, F., Zhu, D., Giles, M., Daniell, T., Neilson, R., Zhu, Y.G., Yang, X.R., 2019. Mineral and organic fertilization alters the microbiome of a soil nematode *Dorylaimus stagnalis* and its resistome. *Sci Total Environ.* 680, 70-78. doi:10.1016/j.scitotenv.2019.04.384.

Zhu, B., Chen, Q., Chen, S., Zhu, Y.G., 2017. Does organically produced lettuce harbor higher abundance of antibiotic resistance genes than conventionally produced? *Environ Int.* 98, 152-159. doi:10.1016/j.envint.2016.11.001.

Zhu, D., An, X.L., Chen, Q.L., Yang, X.R., Christie, P., Ke, X., Wu, L.H., Zhu, Y.G., 2018. Antibiotics Disturb the Microbiome and Increase the Incidence of Resistance Genes in the Gut of a Common Soil Collembolan. *Environ Sci Technol.* 52(5), 3081-3090. doi:10.1021/acs.est.7b04292.

Zhu, Y.G., Johnson, T.A., Su, J.Q., Qiao, M., Guo, G.X., Stedtfeld, R.D., Hashsham, S.A., Tiedje, J.M., 2013. Diverse and abundant antibiotic resistance genes in Chinese swine farms. *Proc Natl Acad Sci U S A.* 110(9), 3435-3440. doi:10.1073/pnas.1222743110.

703 Table 1. Chemical properties of fertilizers and soil.

Items	Soil	PWSC T1	PWSC T2	PWSC T3	S
pH	7.52	7.43	8.06	6.94	8.73
TC	17.74	283.1	242.2	280.5	146.2
TN	2.71	9.45	12.95	11.27	9.59
Cr	78.18	47.73	226.58	36.43	13.35
Cu	34.81	65.53	164.41	229.65	95.38
Pb	38.45	5.87	14.35	1.91	4.08
Zn	132.37	148.85	256.64	320.68	290.66
Cd	0.37	0.48	0.51	0.45	0.45
Tetracycline	-	4.96	3.86	4.06	8.79
Oxytetracycline	-	2.33	1.12	1.97	4.68
Doxycycline	-	2.56	2..13	1.86	3.33
Sulfadiazine	-	0.81	0.31	0.54	0.34
Sulfamethazine	-	0.23	0.12	0.10	0.05
Ciprofloxacin	-	4.32	4.12	3.24	5.66
Enrofloxacin	-	0.32	0.29	0.31	0.51
Chlorotetracycline	-	0.54	0.59	0.48	0.46

704

705

706

707

708

709

710

711

712

713

Figure 1. The detected MGEs and ARGs in soil samples (including initial soil, rhizosphere soil and bulk soil), Chinese cabbage phyllosphere and compost (including T1, T2, T3 and S). ARGs were classified according to the antibiotic they resisted. Control treatments (CK), PWSCs of T1 treatments (T1), PWSCs of T2 treatments (T2), PWSCs of T3 treatments (T3), sewage sludge compost treatments (S) respectively. The mean values of three replicates, data as means $\pm$ SD.

Figure 2. Bipartite network illustrating the shared genes between Chinese cabbage phyllosphere, root, and rhizosphere soil (compost amended soil) cultured in soil amended with compost (T1 (a), T2 (b), T3 (c) and S (d)).

Figure 3. Procrustes analysis is unveiling the significant association between the absolute abundance of ARGs and shared bacterial taxa composition (16S rRNA gene ZOTUs) relied on the Bray-Curtis distance. a, without the application of manure, b, with application of T1 composting, c, with application of T2 composting, d, with application of T3 composting, e, with application of S composting.

Figure 4. Procrustes analysis unveiling the significant association between the absolute abundance of ARGs and shared fungal taxa composition (ITS read ZOTUs) relied on the Bray-Curtis distance. a, without application of composting, b, with application of T1 composting, c, with application of T2 composting, d, with application of T3 composting, e, with application of S composting.