



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/172007/>

Version: Accepted Version

Article:

Halliwell, J., Barbaric, I. and Andrews, P.W. (2020) Acquired genetic changes in human pluripotent stem cells : origins and consequences. *Nature Reviews Molecular Cell Biology*, 21 (12). pp. 715-728. ISSN: 1471-0072

<https://doi.org/10.1038/s41580-020-00292-z>

This is a post-peer-review, pre-copyedit version of an article published in *Nature Reviews Molecular Cell Biology*. The final authenticated version is available online at:
<https://doi.org/10.1038/s41580-020-00292-z>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1
2 **ACQUIRED GENETIC CHANGES IN PLURIPOTENT STEM CELLS: ORIGINS AND**
3 **CONSEQUENCES**
4
5
6

7 **Jason Halliwell, Ivana Barbaric*, Peter W Andrews***
8
9

10
11
12 *The Centre for Stem Cell Biology, The Department of Biomedical Science, The University*
13 *of Sheffield, Western Bank, Sheffield S10 2TN, UK*
14
15
16

17 **Correspondence to:** p.w.andrews@sheffield.ac.uk and i.barbaric@sheffield.ac.uk
18
19

20 **Acknowledgements:** This work was funded in part by grants from the European Union's
21 Horizon 2020 research and innovation program under grant agreement No. 668724 and
22 from the UK Regenerative Medicine Platform, MRC reference MR/R015724/1.
23
24
25

26 **ABSTRACT (200 words)**

27 In the twenty years since human embryonic stem cells, and subsequently induced pluripotent stem
28 cells (collectively, pluripotent stem cells), were first described, it has become apparent that these
29 cells may acquire genetic changes during long term culture, commonly manifest by gains or losses
30 of particular chromosomal regions, or by mutations in certain cancer associated genes, especially
31 *TP53*. Such changes raise concerns for the safety of products destined for clinical applications in
32 regenerative medicine. Although acquired changes may not be present in a cell line at the start of a
33 research program, the low sensitivity of current detection methods means that mutations may be
34 difficult to detect if they arise but are only present in a small proportion of the cells. Nevertheless,
35 recent work suggests that the underlying mutation rate in pluripotent stem cells is low, though they
36 also seem to be particularly susceptible to genomic damage. This apparent contradiction can be
37 reconciled by the observations that, in contrast to somatic cells, pluripotent stem cells are
38 programmed to die in response to genomic damage, which may reflect the requirements of early
39 embryogenesis. Thus, the common variants that do occur are likely rare events that offer the cells
40 a selective growth advantage.

41

42 **INTRODUCTION**

43 Although little more than 20 years has passed since the first human embryonic stem cells (ESC)
44 were reported (Thomson et al., 1998, Reubinoff et al., 2000), and less than 14 years since human
45 induced pluripotent cells (iPSC) were described (Takahashi et al., 2007, Yu et al., 2007), clinical
46 trials for regenerative medicine using derivatives of these cells are already underway or on the
47 horizon (da Cruz et al., 2018, Schwartz et al., 2012, Song et al., 2015, Mandai et al., 2017, Barker
48 et al., 2017). Yet over this period it has become evident that both human ESC and iPSC (collectively
49 here denoted as PSC), although mostly diploid when first derived, may acquire genetic alterations,
50 ranging from large scale structural modifications readily recognized as karyotypic variants through
51 to single base pair changes on subsequent passage (**Figure 1**). Although not the focus of this
52 review, it is worth noting that epigenetic changes encompassing aberrations in DNA methylation,
53 imprinting and X-chromosome inactivation have also been reported in PSC (reviewed in Bar and
54 Benvenisty, 2019). The observation of genetic and epigenetic changes in PSC has triggered worries
55 about the significance of such variants for the safety of PSC-based regenerative medicine (Yasuda
56 et al., 2018, Sato et al., 2019). In particular, some recurrent genetic changes, for example mutations
57 in *TP53*, or gains of chromosome 12p, 17q, and 20q have been associated with various cancers,
58 notably the association of gain of chromosome 12p with embryonal carcinoma (EC) cells, the
59 malignant counterpart of PSC (Andrews, 2002). Indeed, a planned trial of iPSC-derived retinal
60 pigment cells to treat age related macular degeneration was halted when a point mutation was
61 detected, although whether this particular mutation may have caused a problem was unknown
62 (Mandai et al., 2017, Garber, 2015).

63 Some of the genetic variants found in PSC may well have been present in the embryos or somatic
64 cells from which they were derived, or may have been induced during derivation (Hussein et al.,
65 2011, Rouhani et al., 2016). The outcomes of the currently pursued in-depth studies into the effect
66 of the choice of a starting cell type and a method of reprogramming on the overall mutational burden
67 in iPSC (reviewed in Steichen et al., 2019) will have an important bearing on the practical applications
68 of PSC. However, regardless of their ultimate findings, a key feature of such ‘Mutations of Origin’ is
69 that they will be present in all the cells of a given PSC line and so should be readily detectable and
70 assessed for their significance before research with a particular line is initiated. Much more
71 problematic, and the subject of this review, are mutations that were not present initially but arise,
72 often recurrently, during culture of the cells – ‘Acquired Mutations’. Nevertheless, it is important to
73 view these variants in perspective: many PSC lines do not acquire the commonly observed variants,
74 or only in late passage. In a study by the International Stem Cell Initiative, 79 lines out of 122 retained
75 a normal karyotype (Amps et al., 2011). In another study, recurrent mutants of *TP53* were detected
76 in only five out of 140 human PSC lines (Merkle et al., 2017). On the other hand, since the sensitivity
77 of detecting mutant cells in a mosaic culture is low (see **BOX 1**), such variant cells may lurk in
78 cultures for a considerable time until they take over due to a selective growth advantage, or the line
79 is subject to a population bottleneck. Further, some variants, for example gains of a small region of
80 the long arm of chromosome 20, are particularly difficult to detect by G-banding karyotyping and may
81 go unnoticed, even when present in a substantial proportion of the cells (Amps et al., 2011).

82 Over the years, many studies have sought to find ways to minimize the appearance of genetic
83 variants in PSC. However, in such quests, it is important to recall that the appearance of the common
84 recurrent variants depends upon two independent events, mutation followed by subsequent
85 selection (**Figure 1**), and these can be experimentally difficult to disentangle. The mechanisms of
86 selective growth advantage, due to the altered expression or activity of one or more genes (‘Driver’
87 genes), or to the effects of culture conditions on selection, can be relatively easily analysed by spiking
88 wild type cultures with variant cells and monitoring their subsequent growth patterns (Olariu et al.,
89 2010). By contrast, mutation occurs at a very low frequency so that mutations are difficult to monitor
90 directly without expansion of the variant cells, in which case the estimates of mutation rate may be
91 compromised by selection unless this is avoided by often cumbersome clonogenic strategies
92 (Thompson et al., 2020).

93 In this review we discuss the nature of the acquired genetic variants that commonly arise during
94 culture of human PSC cultures and consider the potential consequences of acquired genetic variants
95 in human PSC, both for research and for clinical applications. We then discuss the mechanisms of
96 selective growth advantage that lead to the recurrent appearance of particular variants. Finally, we
97 focus on the underlying mechanisms of mutation in PSC; it seems that PSC differ substantially from
98 somatic cells in both their susceptibility and response to DNA damage, which may reflect the
99 exigencies of cell proliferation in the early embryo.

100

101 **ACQUIRED GENETIC VARIATION**

102 **Karyotypic abnormalities**

103 Traditionally, routine screening of PSC lines for the identification of genetic changes has been
104 performed mainly by cytogenetic and molecular methods that are capable of detecting numerical
105 and structural aneuploidies rather than DNA sequence changes (Draper et al., 2004, Amps et al.,
106 2011). Consequently, karyotypic abnormalities are the most comprehensively catalogued genetic
107 changes in PSC to date. In taking stock of the reports of karyotypic abnormalities in PSC over the
108 last two decades, it is clear that the aberrant PSC karyotypes can encompass virtually any type of
109 an abnormality, including numerical aneuploidies, such as a whole chromosome gain (trisomy) or
110 loss (monosomy), as well as structural aneuploidies, including interstitial duplications, deletions,
111 inversions, amplifications and translocations (Draper et al., 2004, Amps et al., 2011, Taapken et al.,
112 2011). That said, the distribution of chromosomal aberrations appears to be non-random and certain
113 types of variants are more commonly seen (Taapken et al., 2011, Draper et al., 2004, Amps et al.,
114 2011, Baker et al., 2016). The first apparent bias is towards gains rather than losses of chromosomal
115 material. Indeed, it is estimated that over 70% of all karyotypic abnormalities reported in ESC
116 represent whole or partial chromosome gains, whereas only around 20% of reported abnormalities
117 are losses of chromosomes or chromosomal material (Baker et al., 2016). Losses of entire
118 chromosomes are particularly rare, representing only around 2% of reported abnormalities in PSC
119 cultures (Baker et al., 2016). The under representation of monosomies in PSC concurs with the
120 observation that cells in general tolerate gains of genetic material more readily than losses (Torres
121 et al., 2008). Both unbalanced and balanced translocations, i.e. with or without the overt net gain or
122 loss of chromosomal material, respectively, have also been reported, but unlike in certain
123 haematological malignancies, for example, no common recurrent translocations or fusion genes
124 have so far been associated with variant PSC (Draper et al., 2004, Amps et al., 2011, Baker et al.,
125 2016, Assou et al., 2020). On the other hand, some chromosomes are rarely, if ever, reported as
126 gained or lost in PSC, including chromosomes 2, 4, 19 and 21 (Amps et al., 2011, Baker et al., 2016).
127 Finally, most striking is the observation of a consistent pattern of chromosomes affected by
128 aneuploidy in PSC, with the majority of detected aberrations in PSC karyotypes representing gains
129 of the whole or fragments of chromosomes 1, 12, 17, 20 and X (Baker et al., 2007, Amps et al., 2011,
130 Nguyen et al., 2013, Baker et al., 2016, Assou et al., 2020) (**Box 2**). Of these, a particularly insidious
131 change is the frequent gain of a small, variable region of located near the centromere of the long
132 arm of chromosome 20 (Lefort et al., 2008, Spits et al., 2008). Often this gain is below the resolution
133 of G-band karyotyping but, for example, it was noted by SNP array analysis in over 20% of the cell
134 lines in the ISCI study (Amps et al., 2011, Baker et al., 2016). That the same repertoire of
135 aneuploidies is observed across different ESC and iPSC lines and across different laboratories
136 world-wide (Amps et al., 2011) points to the enhanced fitness of such variant cells, likely due to an

137 increased expression of one or more of the genes located on amplified chromosomes (Enver et al.,
138 2005, Ben-David et al., 2014).

139

140 **Point mutations**

141 Point mutation screening has not yet become routine in PSC maintenance and, therefore, culture
142 acquired nucleotide changes in these cells remain largely unexplored. A couple of, recent studies
143 have investigated the presence and potential recurrence of cancer-related point mutations in PSC.
144 Merkle *et al.* (Merkle et al., 2017) subjected the DNA from 140 ESC lines provided by different
145 laboratories world-wide to whole exome sequencing. After filtering out inherited polymorphisms and
146 focusing only on variants that were present in a subset of cells, as suspected culture-acquired
147 mutations, 28 of the 263 mosaic variants detected across the 140 lines were predicted to alter gene
148 function. Of these, the tumor suppressor gene *TP53* was the only gene in which mutations were
149 detected in multiple cell lines, with six different missense mutations in five independent ESC lines.
150 All of the identified *TP53* missense mutations affected cytosines of highly mutable CpG dinucleotides
151 within four of the residues encoding the DNA-binding domain of the TP53 protein, therefore rendering
152 the mutant TP53 protein inactive. Further *TP53* mutations in both ESC and iPSC were uncovered
153 by leveraging RNA-sequencing datasets from public repositories (Merkle et al., 2017, Avior et al.,
154 2019), thereby establishing *TP53* as a recurrently mutated gene in PSC. As the observed *TP53*
155 mutations represent some of the most frequent mutations in cancer (Kandoth et al., 2013) and are
156 also known to cause a familial cancer predisposition disorder, the Li-Fraumeni syndrome (Malkin et
157 al., 1990, Srivastava et al., 1990), these findings have brought into focus the need for monitoring of
158 PSC for culture-acquired *TP53* mutations.

159 Despite being the most prevalent, *TP53* mutations are not the only recurrent point mutations arising
160 in cancer-related genes upon PSC expansion (Avior et al., 2019). Recently, recurrent point
161 mutations were detected in at least 22 other genes that were previously classified within the COSMIC
162 Census database as genes with a documented cancer-related activity, including *CCND2*, *PCM1*,
163 *MYH9*, *HIF1A*, *BCL9* and *VHL* (Avior et al., 2019). Intriguingly, the mutational burden seems to differ
164 between different pluripotent states; human PSC in the naïve state, representing the pluripotent state
165 of the pre-implantation epiblast cells (Nichols and Smith, 2009), were estimated to carry four times
166 more mutations than their primed counterparts, which correspond to the pluripotent state of the post-
167 implantation epiblast (Avior et al., 2019). As the naïve samples analysed in this study were reset
168 from primed PSC, rather than derived directly from embryos, the mutational load differences may
169 not be intrinsic to different cell states, but may in fact reflect a substantial selection pressure imposed
170 on the cells during resetting to naïve pluripotency. Supporting the latter view, the genes found to
171 be mutated in the naïve cells were in pathways affected by chemical inhibitors used in resetting
172 primed cells to the naïve state (Avior et al., 2019).

173 The studies by Merkle et al (2017) and Avior et al (2019) offered an important insight into the
174 mutational landscape of PSC, albeit they focused on analysing a relatively small portion of the
175 PSC genome (i.e. the exome, which represents only about 1% of the genome). Undoubtedly,
176 uncovering the true extent and pattern of point mutations arising in PSC will require much larger,
177 ideally longitudinal datasets, and scrutiny of PSC sequence changes at a genome-wide level.
178 Important for this endeavour will be the implementation of next generation sequencing as a
179 component of routine monitoring of PSC genomes. Currently, the turnaround time and cost of
180 sequencing preclude its use as a routine screening method, but the ongoing technological
181 developments, which are driving down the cost and the data processing time, make this a feasible
182 prospect for the coming decade. Nonetheless, we must remain cognizant of the fact that the
183 reliable detection of mutations is only the first step in handling culture-acquired genetic changes.
184 A far more difficult hurdle is ascribing the functional meaning to the detected mutations, and
185 predicting their potential impact for applications of human PSC.

186

187 **Consequences for applications of human PSC**

188 The close relationship between experimentally-derived PSC and EC cells, the malignant stem cells
189 of teratocarcinomas, which occur predominantly as testicular germ cell tumors in young men
190 (Andrews, 2002, Mostofi and Price, 1973, Damjanov and Solter, 1974), and the ability of PSC to
191 produce teratomas when grown in immunodeficient mice, has always provoked concerns that cancer
192 presents a significant safety hazard for PSC based regenerative medicine. However, it is important
193 to recognize the distinction between teratomas and teratocarcinomas. Teratomas are tumors
194 containing differentiated cells without any persisting PSC. By contrast, teratocarcinomas are tumors
195 with the characteristics of teratomas that also contain undifferentiated PSC (**Figure 2**) (Damjanov
196 and Andrews, 2007b, Damjanov and Andrews, 2016). Clinically, teratocarcinomas are highly
197 malignant cancers, but they can also be effectively treated because PSC are exceptionally sensitive
198 to the chemotherapeutic agent, Cis-Platinum (Einhorn and Donohue, 1977, Oosterhuis et al., 1984,
199 Einhorn et al., 1981) as part of a standard treatment that also includes Bleomycin and Etoposide
200 (Williams et al 1987). Although some PSC do produce teratocarcinomas in which undifferentiated
201 PSC can be recognized histologically, or by outgrowths of PSC from explanted tumors (Andrews et
202 al., 2005), many of the xenograft tumors derived from PSC are better classified as teratomas (Allison
203 et al., 2018). It might be anticipated that variant PSC carrying mutations that enhance their
204 proliferative potential and, perhaps, reduce their propensity to differentiate would be more likely to
205 generate teratocarcinomas. Certainly, aneuploid PSC can produce teratocarcinomas (Andrews et
206 al., 2005). Further, the transcriptomes of ESC carrying an extra copy of chromosome 12 clustered
207 more closely with EC cells from germ cell tumors, which almost always exhibit a gain of the short
208 arm of chromosome 12, while ESC with a gain of chromosome 12 were more likely to produce
209 teratocarcinomas than the parent diploid cells from which they were derived (Ben-David et al., 2014).

210 On the other hand, in a recent ISCI study, albeit limited in scope, teratocarcinomas were produced
211 by PSC without overt karyotypic abnormalities, whereas PSC with such variants, including gains of
212 chromosome 12, produced teratomas, indicating no clear correlation between the formation of
213 teratocarcinomas and the presence of overt karyotypic changes (Allison et al., 2018). These
214 discrepancies point to the need for a more systematic study of the relationship between genotype
215 and the ability of PSC to form teratocarcinomas rather than teratomas.

216 On the other hand, regenerative medicine applications depend upon transplantation of specific
217 differentiated derivatives, not undifferentiated cells, so it is the possibility that genetic variants of PSC
218 may cause a neoplastic transformation of their derivative differentiated cells that is the greater
219 concern (**Figure 2**). Unfortunately, there is very little direct evidence upon which to draw any definite
220 conclusions about the extent of the risks. The somatic elements in teratomas of the laboratory
221 mouse are almost always benign and non-tumorigenic (Damjanov and Solter, 1974) and this may
222 be generally true of human teratomas. However, pathologists with expertise in clinical gonadal
223 teratocarcinomas do have concerns since in the human tumors, in contrast to those of the laboratory
224 mouse, many of the differentiated elements such as neural tubes exhibit features of immaturity that
225 may be regarded as potentially neoplastic (Damjanov and Andrews, 2016). Certainly, secondary
226 somatic tumors derived from primary germ cell tumors have been found clinically, although they are
227 very rare (Mostofi and Price, 1973). Experimental, PSC-derived teratomas often also contain
228 primitive endodermal elements, which is a further concern since yolk sac carcinoma, representing
229 malignant primitive endoderm, is a well-known clinical form of germ cell tumors of the newborn
230 (Cunningham et al., 2012).

231 Although the relationship of malignant transformation of teratoma elements to particular genetic
232 variants has not been established, some of the common karyotypic variants occurring in human PSC
233 are also associated with other types of somatic cancer - for example, gains of the long arm of
234 chromosome 17 with neuroblastoma (Plantaz et al., 1997) (**BOX 2**). Further, two of the genes
235 associated with recurrent variants in human PSC, *TP53* (Merkle et al., 2017) and *BCL2L1*, the driver
236 gene of the chromosome 20q amplicon (see below) (Avery et al., 2013), which derive their selective
237 advantage for PSC from their anti-apoptotic functions, are associated with many cancers (Beroukhim
238 et al., 2010, Hainaut and Hollstein, 1999). Although the driver genes of the other common recurrent
239 variants of human PSC have yet to be identified, they provide a selective advantage because of their
240 specific effects on the undifferentiated PSC and it is entirely possible that their effects on specific
241 differentiated derivatives may be quite different. However, since many of the recurrent variants
242 involve gains or losses of large chromosomal regions, it is also possible that other 'hitch-hiker' genes
243 linked to the driver gene may also cause effects in the differentiated derivatives, separately from the
244 effects of the driver genes on the undifferentiated PSC themselves.

245 Apart from cancer, the genetic variants of PSC have the potential to cause a wide range of effects
246 on cellular physiology that could compromise the efficacy of derivative cells used in clinical

247 applications, or the production of such cells, or indeed the use of PSC in research, for example into
248 disease mechanisms. Nevertheless, there has been very little systematic consideration of these
249 issues. They were discussed by international key opinion leaders at a meeting of the International
250 Stem Cell Initiative (ISCI), at the Jackson Laboratory in 2016 (Andrews et al., 2017), and again at a
251 meeting hosted by Nature in London in 2018 (Technologies, 2018), while the Japanese regulatory
252 authorities have issued some guidelines (Research and Development Division, Health Policy
253 Bureau, The Ministry of Health, Labour and Welfare (MHLW),
254 http://www.nihs.go.jp/cbtp/sispsc/pdf/Eg.ver.Annex_0613-3_2016.pdf). However, there is no
255 international consensus about potential risk assessment and the ISCI meeting in 2016 suggested
256 the establishment of an advisory group to collate information about the common genetic variants of
257 PSC, including any evidence of their effects on cell behaviour, and linking that information to other
258 cancer and disease related genomic databases. Meanwhile, the Nature meeting strongly
259 recommended that researchers clearly document any genetic variants that may have been present
260 in cells used for particular research, so providing the data for future retrospective analysis of their
261 potential consequences. Certainly, as a minimum, the documentation should include appropriate
262 characterisation of the karyotype of the cells, and also assessment of the chromosome 20 amplicon,
263 given that these represent the most commonly observed genomic changes seen in these cells.

264

265 **SELECTION DRIVES THE APPEARANCE OF RECURRENT GENETIC VARIANTS**

266 **Growth advantage**

267 Although genetic variants may be occasionally fixed when PSC cultures are passed through a
268 population bottleneck, such as cloning, the recurrence of specific mutations within PSC populations
269 suggests that such genetic changes endow the variant cells with a selective growth advantage.
270 Consistent with this, the proportions of variant cells in a culture typically increase over time from
271 when they are first detected (Draper et al., 2004, Catalina et al., 2008, Imreh et al., 2006). Similarly,
272 in experiments designed to recapitulate the takeover of cultures by variant clones, co-mixing a small
273 proportion of commonly occurring variants with their wild-type counterparts led to a gradually
274 increased representation of variant cells in subsequent passages until they eventually dominated
275 the cultures (Olariu et al., 2010, Avery et al., 2013). Commonly, a variant may be first detected when
276 it constitutes around 5 – 10% of the cells in a culture, rising rapidly to 100% in as few as 5 passages.
277 Based on these longitudinal evaluations, the takeover of PSC cultures by variant cells has been
278 likened to Darwin's principle of natural selection, whereby the variant PSC that are best adapted to
279 particular selective conditions outcompete their neighbours and populate cultures with their own
280 progeny. The specific phenotypic features associated with genetic variants hold clues as to the
281 selective pressures operating in PSC cultures, the reduction of which is key to minimising the
282 appearance of genetic variants in expanding PSC populations.

283 In principle, genetically variant PSC could gain a selective advantage by acquiring one or several of
284 the following features: a proliferative advantage underpinned by faster cell cycle time, a decreased
285 rate of differentiation, or altered pattern of differentiation, or an increased rate of survival (**Figure 3**).
286 Indeed, a number of studies have reported that such traits typify variant cells harbouring the
287 commonly acquired aneuploidies. For example, the growth advantage of trisomy 12 PSC was
288 attributed mainly to their significantly reduced cell cycle time, although the variant cells also displayed
289 an increased resistance to apoptosis and a reduced tendency for differentiation (Ben-David et al.,
290 2014).

291 With regard to the reduced propensity for differentiation, no studies have so far reported a total block
292 of variant ESC or iPSC to differentiation, although nullipotent EC cells are well known in the context
293 of testicular germ cell tumors (Andrews et al., 1980, Andrews et al., 1982). Rather, either a reduced
294 differentiation capacity (Fazeli et al., 2011) or a delayed differentiation dynamic (Werbowski-
295 Ogilvie et al., 2009) compared with wild-type cells has been observed. In some instances,
296 genetically variant PSC appeared to yield alternative cell types to wild-type cells exposed to the
297 same set of differentiation conditions. For example, the same differentiation protocol applied to wild-
298 type PSC and variants with a gain of the long arm of chromosome 17 resulted in mesodiencephalic
299 dopaminergic neurons or dorsal telencephalic neurons, respectively (Lee et al., 2015). Given that
300 this gain entails amplification of a large chromosomal region and, hence, increased expression of
301 most of the genes in that region (Enver et al., 2005), it is easy to envision that such extensively
302 altered gene and protein expression profiles could include changes that skew the differentiation
303 trajectory of cells. In this case, the skewed differentiation was attributed to an increased expression
304 of *WNT3* and *WNT9B* genes localized in the amplified part of chromosome 17 (Lee et al., 2015). In
305 another case, it has been reported that BCL-XL over expression perturbs SMAD and TGF β signalling
306 in PSC with the chromosome 20q11.21 gain, resulting in impaired neurectoderm differentiation
307 (Markouli et al., 2019). Although the altered propensity for differentiation of variant cells may be a
308 mere consequence of hitch-hiker genes rather than the driver of their growth advantage, the
309 converse may also hold true if the differentiation process itself exerts selection on the differentiating
310 cells. For example, in one study, cardiac differentiation favoured cells with a gain of the long arm of
311 chromosome 20 (Laurent et al., 2011) whereas in another report Merkle et al (Merkle et al., 2017)
312 noted an enrichment of mutant *TP53* cells upon PSC differentiation. In both cases, a variant PSC
313 population was already present in the starting cultures prior to differentiation, but it is also possible
314 that variant cells may arise and be selected during the differentiation process itself.

315 Although faster cell cycle and altered differentiation have been associated with some of the recurrent
316 variants, resistance to apoptosis seems to be a frequent feature of variants commonly detected in
317 PSC cultures. This is, perhaps, not surprising given that marked sensitivity to apoptosis represents
318 one of the notable features of early-passage diploid PSC. Excessive cell death is particularly
319 prominent when PSC are grown at a low cell density (Ohgushi et al., 2010), a condition under which

320 single PSC are confronted with a series of bottlenecks preventing their clonal growth (Barbaric et al.,
321 2014). At the molecular level, propensity for apoptosis has been explained by a low apoptotic
322 threshold of PSC, governed by low expression levels of anti-apoptotic proteins and high expression
323 levels of pro-apoptotic proteins (Liu et al., 2013). In addition to preferential expression of pro-
324 apoptotic factors, PSC store a constitutively active pro-apoptotic factor BAX in the Golgi (Dumitru et
325 al., 2012). This effectively primes PSC for a rapid apoptotic response to appropriate cues. Apart
326 from hampering the efficient scale up of PSC, the severe reduction in cell numbers during culture
327 clearly creates conditions for selection of genetically variant cells capable of blunting the apoptotic
328 pathways (Avery et al., 2013, Merkle et al., 2017).

329 The emergence of variant cells in PSC cultures inevitably entails interactions of variants with their
330 wild-type counterparts, as the two populations share their environment and some of their cell-cell
331 contacts. The nature of these interactions can determine the fate of wild-type cells in a non-cell
332 autonomous manner, thereby impacting on the dynamics of the variant's overtake of cultures (**Figure**
333 **3**). Some of the commonly occurring PSC variants were shown to suppress the growth of wild-type
334 populations by inducing apoptosis in their neighbouring wild-type cells (Price et al., 2019), in a
335 manner similar to the phenomenon of cell competition described in other model systems (Bowling et
336 al., 2019). In PSC cultures, a differential sensitivity of wild-type and variant PSC to mechanical
337 pressures imposed by cell crowding allowed variants to effectively eliminate wild-type cells from
338 mixed cultures, therefore enhancing the ability of variants to rapidly achieve the clonal dominance
339 (Price et al., 2019). Therefore, consideration of cell interactions, in addition to cell autonomous
340 mechanisms, is needed in developing effective strategies for prevention of growth supremacy of
341 variant cells.

342

343 **Driver genes**

344 The simplest working hypothesis to account for the recurrent selection of a particular chromosomal
345 variant is that it is the altered expression of a single 'driver' gene located in the variant region that
346 provides a growth advantage by altering a cell's behaviour in response to proliferation, differentiation
347 or cell death cues (**Figure 3**). It is, of course, possible that interaction of multiple linked genes in a
348 particular chromosomal rearrangement, or indeed alterations to the chromatin architecture itself,
349 may be responsible. Nevertheless, most studies have focused on seeking a single driver gene.

350 Often the size of the genomic region affected is too large to home in on a likely candidate, but in the
351 case of amplifications affecting chromosome 20, a common minimal amplicon of 0.55Mb was
352 identified in the pericentromeric region of the long arm in all reported examples (Amps et al., 2011).
353 Within this minimal amplicon, containing only thirteen annotated genes, *BCL2L1* was a likely
354 candidate driver gene as its anti-apoptotic splice variant, BCL-XL, is expressed in human PSC (Amps
355 et al., 2011). Experiments in which cells carrying a gain chromosome 20, or that had been
356 transfected with a *BCL2L1*-over expressing vector, were mixed with diploid cells, confirmed that

357 *BCL2L1* and its BCL-XL product was indeed the driver providing a selective growth advantage by
358 blocking apoptosis (Nguyen et al., 2013, Avery et al., 2013).

359 Like chromosome 20, a common minimal amplicon has also been identified on the long arm of
360 chromosome 1 (Baker et al., 2016)(E. McIntire et al, International Society for Stem Cell Research
361 Meeting abstract). A likely candidate driver gene located in this region is *MDM4*, which regulates
362 p53 by suppressing its response to cellular stresses and increasing the threshold to apoptosis (Haupt
363 et al., 2019). Since recurrent dominant negative mutations of *TP53* provide a growth advantage to
364 human PSC (Merkle et al., 2017) it is likely that dysregulation of other genes, such as *MDM4*, that
365 affect apoptosis through p53 would confer a similar growth advantage. On the long arm of
366 chromosome 17, another anti-apoptotic gene, *SURVIVIN (BIRC5)*, encoded in the chromosome
367 17q25.3 region, has been proposed since its inhibition leads to apoptosis of human PSC and cancer
368 cells (Blum et al., 2009, Mesri et al., 2001, Ma et al., 2006, Yang et al., 2004). On the other hand,
369 other candidate genes encoded on chromosome 17 include *WNT3* and *WNT9B*, suggested by their
370 involvement in the altered patterns of differentiation of cells carrying a gain of the long arm of
371 chromosome 17 (Lee et al., 2015).

372 Interest in gains of chromosome 12 has a long history since testicular germ cell tumors almost always
373 have a gain of the short arm, mostly as an isochromosome (Atkin and Baker, 1982), or more rarely
374 as an interstitial amplification (Rodriguez et al., 2003, Korkola et al., 2006). However, there is no
375 definitive evidence to identify the specific driver gene, either for the progression of germ cell tumors,
376 or for the appearance of variant human PSC with a gain of chromosome 12. An obvious candidate
377 driver gene on the short arm of chromosome 12 is *NANOG*, given its central role in maintaining
378 pluripotency, and that its over-expression inhibits differentiation (Chambers et al., 2007). Also, over
379 expression of *NANOG* does allow human ESC to efficiently form colonies at low density, which is
380 normally associated with extensive apoptosis, perhaps mediated by downregulating *LECTIN1*, which
381 normally promotes apoptosis, and upregulating *HSPA1A*, which inhibits apoptosis (Darr et al., 2006).
382 Indeed, *NANOG* is located in a minimal amplicon that has been identified in germ cell tumors,
383 chromosome 12p13.31, but so are two other genes, *DPPA3* and *GDF3*, that also may affect the
384 behaviour of human PSC (Korkola et al., 2006). However, a different minimal amplicon has also
385 been reported in human germ cell tumors, at 12p11.2–p12.1, in which a number of other genes have
386 been highlighted, such as the oncogene, *KRAS* (Rodriguez et al., 2003).

387 Although less frequent, deletions may promote enhanced survival through copy number loss of pro-
388 apoptotic genes. The BCL-2 apoptotic pathway is controlled by interactions between pro- and anti-
389 apoptotic protein family members. Human ESC show elevated expression of the pro-apoptotic
390 genes *NOXA*, *BIK*, *BIM*, *BMF* and *PUMA*, which may contribute to their low apoptotic threshold
391 (Madden et al., 2011, Liu et al., 2013, Dumitru et al., 2012). The two of these most highly expressed
392 in human PSC, *NOXA* and *BIK*, are located in chromosomal regions, 18q21.32 and 22q13.2, that do
393 undergo recurrent deletion. Deletion of *NOXA* by genetic manipulation decreases the sensitivity of

394 human PSC to mitotic errors, thereby increasing the survival of aneuploidy cells (Zhang et al., 2019),
395 and improves survival during cell dissociation, similar to the overexpression of the anti-apoptotic
396 proteins, BCL-2 and BCL-XL (Ohgushi et al., 2010, Ardehali et al., 2011).

397

398 **ACQUISITION OF MUTATIONS**

399 **Mutation Rate in PSC**

400 Whereas the mechanisms by which genetic variants offer cells a selective growth advantage are
401 relatively easy to assess and have been extensively studied, addressing the mechanisms that drive
402 the appearance of the variants in the first instance is more problematic: Mutations occur
403 stochastically and at low frequency in single cells within much larger populations, so that by the time
404 they become detectable the frequency of mutation may have been grossly distorted by the effects of
405 selection. To overcome this problem, we recently adopted a clonogenic strategy in which a single
406 cell was isolated and allowed to expand as a clonal colony for a fixed time, after which the clone was
407 subcloned, with about 20 subclones being isolated and, after expansion, subjected to whole genome
408 sequencing (Thompson et al., 2020). Using this approach, in which most of the mutants that arose
409 were in genes and locations unlikely to result in growth advantage or disadvantage, we estimated
410 the mutation rate of two human, clinical grade ESC lines, MShef4 and MShef11, as 0.37×10^{-9} and
411 0.28×10^{-9} SNVs per base pair, per day, respectively, equating to approximately 0.30×10^{-9} and
412 0.23×10^{-9} SNVs per cell division, respectively, given that the cell cycle time of human PSC, in our
413 experience, approximates 20 hours (+/- 2 hours) (Barbaric et al., 2014). This rate was not affected
414 by the use of the Rho associated coiled coil containing protein kinase (ROCK) inhibitor, Y-27632,
415 commonly used in human PSC culture (Watanabe et al., 2007) The frequency of INDELS was 10-
416 fold lower. These low rates are comparable with another study of human iPSC in which the mutation
417 rate was estimated to be 0.18×10^{-9} SNV per base-pair, per cell division which was considerably
418 lower than in the endothelial progenitor cells (Rouhani et al., 2016). These mutation rates in human
419 PSC contrast with an estimated rate of 2.66×10^{-9} mutations per base pair, per mitosis in somatic
420 cells (Milholland et al., 2017). In another more limited study of a human ESC line, a slightly higher
421 mutation rate of 1×10^{-9} SNV per base-pair, per cell division, but again this was much lower than an
422 estimate of a corresponding somatic cell in the same study (Kuijk et al., 2018), while in a study of a
423 single locus, *Aprt*, in mouse ESC, the mutation rate was estimated to be 10 fold lower than in
424 corresponding somatic cells (Cervantes et al., 2002). These low rates are consistent with the
425 infrequency of recurrent point mutations observed in PSC lines: for example Merkle et al (2017)
426 (Merkle et al., 2017) only observed mutations in TP53 in five out of 140 human PSC lines.

427 In our study of the MShef4 and MShef11 human ESC lines (Thompson et al., 2020), the mutation
428 rate was similar across all chromosomes, with no obvious hotspots, with the exception of a slightly
429 raised rate on the X chromosome, which might have been a consequence of both lines being male.
430 Nevertheless, the mutation rate was significantly higher in intergenic regions than in exons and

431 introns, suggesting an influence of chromatin structure on mutation. Further, the predominant
432 mutation signatures that we detected were consistent with oxidative damage being the predominant
433 cause of mutation and, indeed, the mutation rate for both SNV and INDELS was reduced by about
434 50% when the cells were maintained under low (5%) oxygen atmospheres.

435

436 **DNA Replication Stress and Mitotic Errors**

437 While many SNV in PSC, as in other cultured cells (Petljak et al., 2019, Kucab et al., 2019, Viel et
438 al., 2017), are caused by misincorporation of bases due to oxidative stress, the relatively rapid cell
439 cycle of PSC might also expose them to high levels of DNA replication stress, characterised by
440 reduced rates of DNA replication together with stalling and collapse of replication forks (Bartkova et
441 al., 2005, Gorgoulis et al., 2005). Errors in the repair of resulting double stranded DNA breaks could
442 then lead to chromosomal rearrangements (Cannan and Pederson, 2016). Self-renewal of human
443 PSC is characterised by an abbreviated G1 phase that bypasses the RB1-E2F checkpoint due to
444 the high expression of cyclin D2 and its CDK4 partner together with the constitutive expression of
445 cyclin E, which together maintain RB1 in a hyperphosphorylated and inactive state (Becker et al.,
446 2006, Becker et al., 2010, Filipczyk et al., 2007). Using DNA fibre assays we have recently found
447 that, in comparison to isogenic somatic cells, human PSC do exhibit the features of DNA replication
448 stress, including slower DNA replication speeds with evidence of stalled replication forks, and
449 replication initiating from quiescent replication origins (**Figure 4**) (Halliwell et al., 2019). In parallel
450 we also observed more extensive replication-associated DNA damage in the PSC compared to
451 somatic cells, as has also been reported by others (Simara et al., 2017, Vallabhaneni et al., 2018).

452 A similar situation pertains in many cancers where cyclin E is frequently over-expressed and RB1-
453 E2F is constitutively activated (Akli and Keyomarsi, 2003). One of the consequences of this is
454 replication stress, double stranded breaks and genetic instability (Bester et al., 2011, Burrell et al.,
455 2013, Frame et al., 2006, Pickering and Kowalik, 2006). In mouse ESC, also, molecular hallmarks
456 of replication stress are almost identical to those observed when oncogenes, such as cyclin E, are
457 dysregulated in somatic cells (Ahuja et al., 2016), suggesting that atypical cell cycle control with
458 consequent susceptibility to DNA replication stress and genomic damage in PSC parallels the
459 oncogene-induced DNA damage model for cancer development and progression (Halazonetis et al.,
460 2008).

461 Replication stress induced from oncogene expression can lead to nucleotide deficiency and collision
462 of replication forks with transcription complexes (Jones et al., 2013, Bester et al., 2011).
463 Supplementing cancer cells or primary cell lines that overexpress oncogenes, such as cyclin E, with
464 nucleosides has been found to alleviate replication stress and its associated DNA damage and
465 genetic instability in these cases (Bester et al., 2011, Burrell et al., 2013). In a similar manner, we

466 have recently found that exogenous nucleosides increase the rate of replication fork progression and
467 decrease DNA damage in human PSC cultures (**Figure 4**) (Halliwell et al., 2019).

468 While chromosomal non-dysjunction and numerical instabilities may be the product of merotelic
469 kinetochore attachment, in which the microtubules from both poles bind to the same sister chromatid,
470 leading to lagging and potential mis-segregation of chromosomes (Cimini et al., 2001), the
471 persistence of DNA replication defects from S phase into mitosis can also result in the formation of
472 mitotic errors that are a source of chromosomal instabilities (Burrell et al., 2013). Under-replicated
473 regions can interlink sister chromatids during segregation forming anaphase bridges that are prone
474 to breakage forming double stranded breaks (Chan et al., 2009). Often, to prevent anaphase
475 bridges, nucleases cleave the DNA that again generates double stranded breaks (Naim et al., 2013).
476 Further, the condensation of chromosomes that harbour replication intermediates are particularly
477 prone to breakage (Lukas et al., 2011). These double stranded breaks that result from replication
478 intermediates in mitosis are the substrates for genetic instability caused by error induced repair.

479 By fluorescently labelling human PSC with histone H2B-mCherry it was observed that 30% of
480 mitoses were abnormal including a high proportion with lagging chromosomes and anaphase bridges
481 (Zhang et al., 2019), a level substantially higher than that observed in somatic cell lines (Lamm et
482 al., 2016). In comparison to somatic cell lines, diploid human PSC show condensation defects that
483 result in partially condensed and entangled chromosomes (Lamm et al., 2016). Supplementing
484 cultures with exogenous nucleosides alleviated replication stress and decreased the frequency of
485 mitotic errors, providing further evidence that these are linked in human PSC (Halliwell et al., 2019),
486 as well as providing an approach to reducing their appearance in PSC cultures. However, it should
487 be noted that the continued occurrence of mitotic errors, even with the addition of nucleosides
488 suggests that there are other factors driving their occurrence.

489

490 **Response to genomic damage**

491 Human PSC deploy a number of mechanisms to minimise the effective mutation rate that otherwise
492 might be anticipated from their high susceptibility to DNA damage. Genes involved in various repair
493 pathways show increased expression compared to somatic cells (Maynard et al., 2008, Momcilović
494 et al., 2009), and nucleotide excision repair, base excision repair, and the resolution of inter-strand
495 crosslinks caused by ionising radiation all have all been reported to be faster in human PSC than
496 somatic cell lines (Luo et al., 2012, Hyka-Nouspikel et al., 2012, Maynard et al., 2008). PSC also
497 tend to repair double strand break using homologous recombination, which is prone to less errors
498 than non-homologous end joining (Adams et al., 2010a), although they do also utilise a higher fidelity
499 system of non-homologous end joining that is independent of DNA-PKc and ATM (Adams et al.,
500 2010b). Further, in response to the formation of reactive oxygen species as a by-product of
501 respiration, and consequent oxidative stress, PSC express higher levels of SOD2 and GPX2 anti-
502 oxidant enzymes compared to differentiated lines (Saretzki et al., 2008).

503 Nevertheless, human PSC generally activate apoptosis when exposed to lower doses of genotoxic
504 insults than do somatic cells, suggesting that a low apoptotic threshold is the key element in their
505 response to genomic damage. After human PSC are exposed to ultraviolet C radiation to induce
506 nucleotide base adducts or DNA breaks they respond with extensive apoptosis even at mild doses
507 that have little effect on somatic cell lines (Hyka-Nouspikel et al., 2012, Luo et al., 2012, Simara et
508 al., 2017). Similarly, the treatment of human PSC with cis-platinum or thymidine to initiate replication
509 block (Desmarais et al., 2012, Desmarais et al., 2016), or with nocadazole to induce mitotic block
510 (Zhang et al., 2019) also elicits an extensive apoptotic response in contrast to the response by
511 somatic cells, while PSC also efficiently activate apoptosis in response to oxidative stress (Saretzki
512 et al., 2008). The particular sensitivity of embryonal carcinoma cells, the malignant PSC of
513 teratocarcinomas, to drugs such as cis-platinum (Oosterhuis et al., 1984, Einhorn and Donohue,
514 1977), a DNA cross linking agent, makes germ cell tumors one of the most treatable forms of solid
515 cancer, most likely reflecting this particular low apoptotic threshold.

516 An atypical cell cycle checkpoint control mechanism most likely underlies the low apoptotic threshold
517 of human PSC. In response to DNA damage, human PSC fail to activate p21, which is normally
518 required to execute the G1/S checkpoint, providing less time for repair before apoptosis is initiated
519 in a p53 dependant manner (Hyka-Nouspikel et al., 2012, Hong and Stambrook, 2004, Momcilović
520 et al., 2009). Further, in response to DNA replication stress caused by high levels of thymidine, or
521 the presence of cis-platinum, human PSC, unlike somatic cells, do not activate ATR-CHK1, while
522 foci of RPA, which binds to single stranded DNA at stalled replication forks, are not formed: instead
523 the cells commit to apoptosis (Desmarais et al., 2012, Desmarais et al., 2016). Human PSC also
524 undergo extensive apoptosis in response to mitotic stress which may safeguard the genome from
525 abnormal mitosis by clearing the effected cell from the cell pool (Zhang et al., 2019). Collectively,
526 these studies support a model in which genomic stability, and the particularly low observed mutation
527 rate of PSC is primarily maintained by a low apoptotic threshold. Consequently, blocking apoptosis
528 seems to be the most likely mechanism that provides selective growth advantage for the common
529 genetic variants found in human PSC: the two driver genes so far identified, *TP53* and *BCL2L1*, both
530 act to inhibit apoptosis, while other proposed candidates, *MDM4* and *SURVIVIN*, are also anti-
531 apoptotic.

532 This low apoptotic threshold of human PSC may reflect their relationship to the early embryo in which
533 the need for rapid cell doublings is accomplished by the lack of cell cycle checkpoints, rendering the
534 cells particularly susceptible to errors in DNA synthesis and mitosis, which could be catastrophic for
535 subsequent embryonic development. Indeed, almost half of human embryos fail to survive due to
536 chromosomal instability, which does not seem to be a mere artefact of *in vitro* fertilisation (van
537 Echten-Arends et al., 2011, Munné et al., 2019, Starostik et al., 2020). It has been observed that
538 the mosaic embryos that survive to the blastocyst stage undergo “genetic normalization” when
539 cultured under routine IVF conditions (Brezina et al., 2011). The mechanism of genetic normalization

540 is still widely debated although, in the mouse, activation of apoptosis during the later pre-implantation
541 stages may allow for the removal of aneuploid cells from the developing embryo (Kops et al., 2004,
542 Bolton et al., 2016). This model is supported by observations that the proportion of aneuploidy in
543 the inner cell mass is reduced, whereas in the trophectoderm it is enriched as development proceeds
544 (Hardy, 1997). However, this is still widely debated as apoptosis is a feature of all embryos, and
545 may be a mechanism for maintaining cellular homeostasis regardless of their genomic state (Haouzi
546 and Hamamah, 2009).

547 Another context in which the apoptotic response of human PSC to DSB induction is of central
548 importance, is the process of genome editing. The rising prominence of genome editing
549 technologies, in particular CRISPR/Cas9-based methods, has fuelled efforts aimed at, for example,
550 correcting germline mutations in PSC to allow autologous cell therapy, or removing HLA antigens to
551 reduce the need for immunosuppressants in transplanted patients. Crucially, as gene editing relies
552 on the induction of DSB by nucleases, the edited PSC undergo high levels of apoptosis in
553 comparison to their unedited counterparts (Ihry et al., 2018). The rate of cell death was shown to be
554 similar between different, edited PSC, regardless of whether the targeted gene was expressed in
555 PSC or whether it was non-expressed and dispensable for PSC maintenance. This observation
556 supports the view that the induction of DSB during the gene editing process commits PSC to
557 apoptosis (Ihry et al., 2018). Mechanistically, DSB induction by Cas9 was shown to trigger
558 differential gene expression in edited cells, most notably by promoting the TP53 transcriptional
559 response (Ihry et al., 2018). In line with the importance of TP53 activation during the genome editing
560 process, performing the PSC genome editing in cells with genetically inactivated *TP53* reduced the
561 levels of cell death and improved the efficiency of PSC genome editing (Ihry et al., 2018). While
562 genetic inactivation of TP53 is deemed too risky for editing of cells destined for clinical use, transient
563 TP53 inactivation has been proposed as a possible alternative (Schiroli et al., 2019). Further work
564 will need to carefully address this possibility, to ensure that the transient TP53 inactivation does not
565 inadvertently select for *TP53* genetic mutants. Relevant to this notion is the recent data
566 demonstrating the emergence of *TP53*-inactivating mutations in cancer cell lines during the
567 CRISPR/Cas9 genome editing process (Enache et al., 2020).

568

569 **CONCLUSIONS**

570 Efforts to collect and catalogue genetic variation in PSC over the last two decades demonstrated
571 that particular variants do arise in cultures and are sometimes difficult to detect because of the
572 limitation of sensitivity of detection methods. Nonetheless, it is reassuring that the rate of mutation
573 in PSC cultures is low compared with somatic cells. Indeed, based on the data from large-scale
574 retrospective studies, such as the ISCI (Amps et al., 2011), and on the direct measurements of the
575 mutation rates in PSC (Thompson et al., 2020), there is no evidence to suggest that PSC genomes
576 are particularly unstable. Rather, the clonal expansion of genetic variants against the backdrop of

577 low mutation rates, can be explained by the effect of selective pressures operating in PSC cultures.
578 Optimizing culture conditions and protocols to minimize the growth advantages of the common
579 variants is, therefore, a key route to maintaining the genetic integrity of PSC lines. However, the
580 predominant selective force dominating PSC cultures appears to be a high rate of apoptosis (Dumitru
581 et al., 2012, Barbaric et al., 2014). Apoptosis seems to be a default fate choice of PSC in many
582 different scenarios, including a response of cells to genome damage or mitotic stress (Desmarais et
583 al., 2012, Desmarais et al., 2016, Dumitru et al., 2012, Zhang et al., 2019), most likely reflecting its
584 function of maintaining the genetic integrity of the early embryo. Consequently, optimising culture
585 conditions should entail removing the apoptotic stimuli, but not blocking apoptosis *per se* which
586 would be counterproductive.

587 Armed with a knowledge of recurrent karyotypic and sequence changes, our attention now needs to
588 turn to finding ways of minimizing their occurrence by lowering the genome damage and reducing
589 the selective pressures. In that respect, the observations that mutation rates can be decreased by
590 growing cells under low oxygen (Thompson et al., 2020) and that replication-stress induced genome
591 damage can be alleviated by addition of exogenous nucleosides (Halliwell et al., 2019) provide
592 foundations for optimised culture conditions of PSC. Further work should also address the
593 contribution of epigenetic variants to aberrant PSC phenotypes, as the understanding of epigenetic
594 variation in PSC cultures remains limited.

595 Finally, the field awaits deciphering of the functional consequences of the karyotype and sequence
596 changes on PSC traits and on the behaviour of their differentiated derivatives. Interpreting the role
597 of specific variants is complicated by the fact that their consequences are likely to be context-
598 dependent. For example, a mutation in a gene expressed specifically in an endodermal lineage may
599 have little impact on the clinical application of neuronal cells. To aid these analyses, ISCI is
600 proposing an international study group to collate and monitor evidence of genetic variants in PSC
601 and their potential consequences (Andrews et al., 2017). However, it should be noted that the
602 success of this approach requires a concerted effort within the field to perform routine monitoring
603 and report the presence of genetic variants, thereby allowing retrospective analyses of their effects.
604 We envisage that these initiatives, in a synergy with cancer genome efforts, would provide a rational
605 strategy to assess the potential risk of different mutations, a necessary requirement for routine, safe
606 clinical implementation of cellular therapies.

607

608 REFERENCES

- 609 ADAMS, B. R., GOLDING, S. E., RAO, R. R. & VALERIE, K. 2010a. Dynamic dependence on
610 ATR and ATM for double-strand break repair in human embryonic stem cells and neural
611 descendants. *PLoS One*, 5, e10001.
- 612 ADAMS, B. R., HAWKINS, A. J., POVIRK, L. F. & VALERIE, K. 2010b. ATM-independent, high-
613 fidelity nonhomologous end joining predominates in human embryonic stem cells. *Aging*
614 (*Albany NY*), 2, 582-96.

615 AHUJA, A. K., JODKOWSKA, K., TELONI, F., BIZARD, A. H., ZELLWEGER, R., HERRADOR,
616 R., ORTEGA, S., HICKSON, I. D., ALTMAYER, M., MENDEZ, J. & LOPES, M. 2016. A
617 short G1 phase imposes constitutive replication stress and fork remodelling in mouse
618 embryonic stem cells. *Nat Commun*, 7, 10660.

619 AKLI, S. & KEYOMARSI, K. 2003. Cyclin E and its low molecular weight forms in human cancer
620 and as targets for cancer therapy. *Cancer Biol Ther*, 2, S38-47.

621 ALLISON, T. F., ANDREWS, P. W., AVIOR, Y., BARBARIC, I., BENVENISTY, N., BOCK, C.,
622 BREHM, J., BRÜSTLE, O., DAMJANOV, I., ELEFANTY, A., FELKNE, D., GOKHALE,
623 P. J., HALBRITTER, F., HEALY, L. E., HU, T. X., KNOWLES, B. B., LORING, J. F.,
624 LUDWIG, T. E., MAYBERRY, R., MICALLEF, S., MOHAMED, J. S., MÜLLER, F.-J.,
625 MUMMERY, C. L., NAKATSUJI, N., NG, E. S., OH, S. K. W., O'SHEA, O., PERA, M. F.,
626 REUBINOFF, B., ROBSON, P., ROSSANT, J., SCHULDT, B. M., SOLTER, D., SOURRIS,
627 K., STACEY, G., STANLEY, E. G., SUEMORI, H., TAKAHASHI, K. & YAMANAKA, S.
628 2018. Assessment of established techniques to determine developmental and malignant
629 potential of human pluripotent stem cells. *Nat Commun*, 9, 1925.

630 AMPS, K., ANDREWS, P. W., ANYFANTIS, G., ARMSTRONG, L., AVERY, S., BAHARVAND,
631 H., BAKER, J., BAKER, D., MUNOZ, M. B., BEIL, S., BENVENISTY, N., BEN-YOSEF,
632 D., BIANCOTTI, J. C., BOSMAN, A., BRENA, R. M., BRISON, D., CAISANDER, G.,
633 CAMARASA, M. V., CHEN, J., CHIAO, E., CHOI, Y. M., CHOO, A. B., COLLINS, D.,
634 COLMAN, A., CROOK, J. M., DALEY, G. Q., DALTON, A., DE SOUSA, P. A.,
635 DENNING, C., DOWNIE, J., DVORAK, P., MONTGOMERY, K. D., FEKI, A., FORD, A.,
636 FOX, V., FRAGA, A. M., FRUMKIN, T., GE, L., GOKHALE, P. J., GOLAN-LEV, T.,
637 GOURABI, H., GROPP, M., LU, G., HAMPL, A., HARRON, K., HEALY, L., HERATH,
638 W., HOLM, F., HOVATTA, O., HYLLNER, J., INAMDAR, M. S., IRWANTO, A. K., ISHII,
639 T., JACONI, M., JIN, Y., KIMBER, S., KISELEV, S., KNOWLES, B. B., KOPPER, O.,
640 KUKHARENKO, V., KULIEV, A., LAGARKOVA, M. A., LAIRD, P. W., LAKO, M.,
641 LASLETT, A. L., LAVON, N., LEE, D. R., LEE, J. E., LI, C., LIM, L. S., LUDWIG, T. E.,
642 MA, Y., MALTBY, E., MATEIZEL, I., MAYSHAR, Y., MILEIKOVSKY, M., MINGER,
643 S. L., MIYAZAKI, T., MOON, S. Y., MOORE, H., MUMMERY, C., NAGY, A.,
644 NAKATSUJI, N., NARWANI, K., OH, S. K., OLSON, C., OTONKOSKI, T., PAN, F.,
645 PARK, I. H., PELLIS, S., PERA, M. F., PEREIRA, L. V., QI, O., RAJ, G. S., REUBINOFF,
646 B., ROBINS, A., ROBSON, P., ROSSANT, J., SALEKDEH, G. H., SCHULZ, T. C., et al.
647 2011. Screening ethnically diverse human embryonic stem cells identifies a chromosome 20
648 minimal amplicon conferring growth advantage. *Nat Biotechnol*, 29, 1132-44.

649 ANDREWS, P. W. 2002. From teratocarcinomas to embryonic stem cells. *Philos Trans R Soc Lond*
650 *B Biol Sci*, 357, 405-17.

651 ANDREWS, P. W., BEN-DAVID, U., BENVENISTY, N., COFFEY, P., EGGAN, K., KNOWLES,
652 B. B., NAGY, A., PERA, M., REUBINOFF, B., RUGG-GUNN, P. J. & STACEY, G. N.
653 2017. Assessing the Safety of Human Pluripotent Stem Cells and Their Derivatives for
654 Clinical Applications. *Stem Cell Reports*, 9, 1-4.

655 ANDREWS, P. W., BRONSON, D. L., BENHAM, F., STRICKLAND, S. & KNOWLES, B. B.
656 1980. A comparative study of eight cell lines derived from human testicular teratocarcinoma.
657 *Int J Cancer*, 26, 269-80.

658 ANDREWS, P. W., GOODFELLOW, P. N., SHEVINSKY, L. H., BRONSON, D. L. & KNOWLES,
659 B. B. 1982. Cell-surface antigens of a clonal human embryonal carcinoma cell line:
660 morphological and antigenic differentiation in culture. *Int J Cancer*, 29, 523-31.

661 ANDREWS, P. W., MATIN, M. M., BAHRAMI, A. R., DAMJANOV, I., GOKHALE, P. &
662 DRAPER, J. S. 2005. Embryonic stem (ES) cells and embryonal carcinoma (EC) cells:
663 opposite sides of the same coin. *Biochem Soc Trans*, 33, 1526-30.

664 ARDEHALI, R., INLAY, M. A., ALI, S. R., TANG, C., DRUKKER, M. & WEISSMAN, I. L. 2011.
665 Overexpression of BCL2 enhances survival of human embryonic stem cells during stress and
666 obviates the requirement for serum factors. *Proc Natl Acad Sci U S A*, 108, 3282-7.

667 ASSOU, S., GIRAULT, N., PLINET, M., BOUCKENHEIMER, J., SANSAC, C., COMBE, M.,
668 MIANNÉ, J., BOURGUIGNON, C., FIELDS, M., AHMED, E., COMMES, T.,
669 BOUREUX, A., LEMAÎTRE, J.-M. & DE VOS, J. 2020. Recurrent Genetic Abnormalities
670 in Human Pluripotent Stem Cells: Definition and Routine Detection in Culture Supernatant
671 by Targeted Droplet Digital PCR. *Stem Cell Reports*, 14, 1-8.

672 ATKIN, N. B. & BAKER, M. C. 1982. Specific chromosome change, i(12p), in testicular tumours?
673 *Lancet*, 2, 1349.

674 AVERY, S., HIRST, A. J., BAKER, D., LIM, C. Y., ALAGARATNAM, S., SKOTHEIM, R. I.,
675 LOTHE, R. A., PERA, M. F., COLMAN, A., ROBSON, P., ANDREWS, P. W. &
676 KNOWLES, B. B. 2013. BCL-XL mediates the strong selective advantage of a 20q11.21
677 amplification commonly found in human embryonic stem cell cultures. *Stem Cell Reports*, 1,
678 379-86.

679 AVIOR, Y., EGGAN, K. & BENVENISTY, N. 2019. Cancer-Related Mutations Identified in Primed
680 and Naive Human Pluripotent Stem Cells. *Cell Stem Cell*, 25, 456-461.

681 BAKER, D., HIRST, A. J., GOKHALE, P. J., JUAREZ, M. A., WILLIAMS, S., WHEELER, M.,
682 BEAN, K., ALLISON, T. F., MOORE, H. D., ANDREWS, P. W. & BARBARIC, I. 2016.
683 Detecting Genetic Mosaicism in Cultures of Human Pluripotent Stem Cells. *Stem Cell*
684 *Reports*, 7, 998-1012.

685 BAKER, D. E., HARRISON, N. J., MALTBY, E., SMITH, K., MOORE, H. D., SHAW, P. J.,
686 HEATH, P. R., HOLDEN, H. & ANDREWS, P. W. 2007. Adaptation to culture of human
687 embryonic stem cells and oncogenesis in vivo. *Nat Biotechnol*, 25, 207-15.

688 BARBARIC, I., BIGA, V., GOKHALE, P. J., JONES, M., STAVISH, D., GLEN, A., COCA, D. &
689 ANDREWS, P. W. 2014. Time-lapse analysis of human embryonic stem cells reveals multiple
690 bottlenecks restricting colony formation and their relief upon culture adaptation. *Stem Cell*
691 *Reports*, 3, 142-55.

692 BARKER, R. A., PARMAR, M., STUDER, L. & TAKAHASHI, J. 2017. Human Trials of Stem
693 Cell-Derived Dopamine Neurons for Parkinson's Disease: Dawn of a New Era. *Cell Stem Cell*,
694 21, 569-573.

695 BARTKOVA, J., HOREJSÍ, Z., KOED, K., KRÄMER, A., TORT, F., ZIEGER, K., GULDBERG,
696 P., SEHESTED, M., NESLAND, J. M., LUKAS, C., ØRNTOFT, T., LUKAS, J. & BARTEK,
697 J. 2005. DNA damage response as a candidate anti-cancer barrier in early human
698 tumorigenesis. *Nature*, 434, 864-70.

699 BECKER, K. A., GHULE, P. N., LIAN, J. B., STEIN, J. L., VAN WIJNEN, A. J. & STEIN, G. S.
700 2010. Cyclin D2 and the CDK substrate p220(NPAT) are required for self-renewal of human
701 embryonic stem cells. *J Cell Physiol*, 222, 456-64.

702 BECKER, K. A., GHULE, P. N., THERRIEN, J. A., LIAN, J. B., STEIN, J. L., VAN WIJNEN, A.
703 J. & STEIN, G. S. 2006. Self-renewal of human embryonic stem cells is supported by a
704 shortened G1 cell cycle phase. *J Cell Physiol*, 209, 883-93.

705 BEN-DAVID, U., ARAD, G., WEISSBEIN, U., MANDEFRO, B., MAIMON, A., GOLAN-LEV,
706 T., NARWANI, K., CLARK, A. T., ANDREWS, P. W., BENVENISTY, N. & CARLOS
707 BIANCOTTI, J. 2014. Aneuploidy induces profound changes in gene expression,
708 proliferation and tumorigenicity of human pluripotent stem cells. *Nat Commun*, 5, 4825.

709 BEROUKHIM, R., MERMEL, C. H., PORTER, D., WEI, G., RAYCHAUDHURI, S., DONOVAN,
710 J., BARRETINA, J., BOEHM, J. S., DOBSON, J., URASHIMA, M., MC HENRY, K. T.,
711 PINCHBACK, R. M., LIGON, A. H., CHO, Y. J., HAERY, L., GREULICH, H., REICH, M.,
712 WINCKLER, W., LAWRENCE, M. S., WEIR, B. A., TANAKA, K. E., CHIANG, D. Y.,
713 BASS, A. J., LOO, A., HOFFMAN, C., PRENSNER, J., LIEFELD, T., GAO, Q., YECIES,
714 D., SIGNORETTI, S., MAHER, E., KAYE, F. J., SASAKI, H., TEPPER, J. E., FLETCHER,
715 J. A., TABERNERO, J., BASELGA, J., TSAO, M. S., DEMICHELIS, F., RUBIN, M. A.,
716 JANNE, P. A., DALY, M. J., NUCERA, C., LEVINE, R. L., EBERT, B. L., GABRIEL, S.,
717 RUSTGI, A. K., ANTONESCU, C. R., LADANYI, M., LETAI, A., GARRAWAY, L. A.,
718 LODA, M., BEER, D. G., TRUE, L. D., OKAMOTO, A., POMEROY, S. L., SINGER, S.,

719 GOLUB, T. R., LANDER, E. S., GETZ, G., SELLERS, W. R. & MEYERSON, M. 2010. The
720 landscape of somatic copy-number alteration across human cancers. *Nature*, 463, 899-905.

721 BESTER, A. C., RONIGER, M., OREN, Y. S., IM, M. M., SARNI, D., CHAOAT, M., BENSIMON,
722 A., ZAMIR, G., SHEWACH, D. S. & KEREM, B. 2011. Nucleotide deficiency promotes
723 genomic instability in early stages of cancer development. *Cell*, 145, 435-46.

724 BLUM, B., BAR-NUR, O., GOLAN-LEV, T. & BENVENISTY, N. 2009. The anti-apoptotic gene
725 survivin contributes to teratoma formation by human embryonic stem cells. *Nat Biotechnol*,
726 27, 281-7.

727 BOLTON, H., GRAHAM, S. J. L., VAN DER AA, N., KUMAR, P., THEUNIS, K., FERNANDEZ
728 GALLARDO, E., VOET, T. & ZERNICKA-GOETZ, M. 2016. Mouse model of chromosome
729 mosaicism reveals lineage-specific depletion of aneuploid cells and normal developmental
730 potential. *Nat Commun*, 7, 11165.

731 BOWLING, S., LAWLOR, K. & RODRÍGUEZ, T. A. 2019. Cell competition: the winners and losers
732 of fitness selection. *Development*, 146.

733 BREZINA, P., BARKER, A., BENNER, A., ROSS, R., NGUYEN, K.-H., ANCHAN, R., RICHTER,
734 K., CUTTING, G. & KEARNS, W. 2011. Genetic Normalization of Differentiating
735 Aneuploid Human Embryos. *Nature Precedings*.

736 BURRELL, R. A., MCCLELLAND, S. E., ENDEFELDER, D., GROTH, P., WELLER, M. C.,
737 SHAIKH, N., DOMINGO, E., KANU, N., DEWHURST, S. M., GRONROOS, E., CHEW,
738 S. K., ROWAN, A. J., SCHENK, A., SHEFFER, M., HOWELL, M., KSCHISCHO, M.,
739 BEHRENS, A., HELLEDAY, T., BARTEK, J., TOMLINSON, I. P. & SWANTON, C. 2013.
740 Replication stress links structural and numerical cancer chromosomal instability. *Nature*, 494,
741 492-496.

742 CANNAN, W. J. & PEDERSON, D. S. 2016. Mechanisms and Consequences of Double-Strand DNA
743 Break Formation in Chromatin. *J Cell Physiol*, 231, 3-14.

744 CATALINA, P., MONTES, R., LIGERO, G., SANCHEZ, L., DE LA CUEVA, T., BUENO, C.,
745 LEONE, P. E. & MENENDEZ, P. 2008. Human ESCs predisposition to karyotypic
746 instability: Is a matter of culture adaptation or differential vulnerability among hESC lines
747 due to inherent properties? *Mol Cancer*, 7, 76.

748 CERVANTES, R. B., STRINGER, J. R., SHAO, C., TISCHFIELD, J. A. & STAMBROOK, P. J.
749 2002. Embryonic stem cells and somatic cells differ in mutation frequency and type. *Proc*
750 *Natl Acad Sci U S A*, 99, 3586-90.

751 CHAMBERS, I., SILVA, J., COLBY, D., NICHOLS, J., NIJMEIJER, B., ROBERTSON, M.,
752 VRANA, J., JONES, K., GROTEWOLD, L. & SMITH, A. 2007. Nanog safeguards
753 pluripotency and mediates germline development. *Nature*, 450, 1230-4.

754 CHAN, K. L., PALMAI-PALLAG, T., YING, S. & HICKSON, I. D. 2009. Replication stress induces
755 sister-chromatid bridging at fragile site loci in mitosis. *Nat Cell Biol*, 11, 753-60.

756 CIMINI, D., HOWELL, B., MADDOX, P., KHODJAKOV, A., DEGRASSI, F. & SALMON, E. D.
757 2001. Merotelic kinetochore orientation is a major mechanism of aneuploidy in mitotic
758 mammalian tissue cells. *J Cell Biol*, 153, 517-27.

759 CUNNINGHAM, J. J., ULBRIGHT, T. M., PERA, M. F. & LOOIJENGA, L. H. 2012. Lessons from
760 human teratomas to guide development of safe stem cell therapies. *Nat Biotechnol*, 30, 849-
761 57.

762 DA CRUZ, L., FYNES, K., GEORGIADIS, O., KERBY, J., LUO, Y. H., AHMADO, A., VERNON,
763 A., DANIELS, J. T., NOMMISTE, B., HASAN, S. M., GOOLJAR, S. B., CARR, A. F.,
764 VUGLER, A., RAMSDEN, C. M., BICTASH, M., FENSTER, M., STEER, J.,
765 HARBINSON, T., WILBREY, A., TUFAIL, A., FENG, G., WHITLOCK, M., ROBSON, A.
766 G., HOLDER, G. E., SAGOO, M. S., LOUDON, P. T., WHITING, P. & COFFEY, P. J. 2018.
767 Phase 1 clinical study of an embryonic stem cell-derived retinal pigment epithelium patch in
768 age-related macular degeneration. *Nat Biotechnol*, 36, 328-337.

769 DAMJANOV, I. & ANDREWS, P. W. 2007a. The terminology of teratocarcinomas and teratomas.
770 *Nat Biotechnol*, 25, 1212; discussion.

771 DAMJANOV, I. & ANDREWS, P. W. 2007b. The terminology of teratocarcinomas and teratomas.
772 *Nat Biotechnol*, 25, 1212; discussion 1212.

773 DAMJANOV, I. & ANDREWS, P. W. 2016. Teratomas produced from human pluripotent stem cells
774 xenografted into immunodeficient mice - a histopathology atlas. *Int J Dev Biol*, 60, 337-419.

775 DAMJANOV, I. & SOLTER, D. 1974. Experimental teratoma. *Curr Top Pathol*, 59, 69-130.

776 DARR, H., MAYSHAR, Y. & BENVENISTY, N. 2006. Overexpression of NANOG in human ES
777 cells enables feeder-free growth while inducing primitive ectoderm features. *Development*,
778 133, 1193-201.

779 DESMARAIS, J. A., HOFFMANN, M. J., BINGHAM, G., GAGOU, M. E., MEUTH, M. &
780 ANDREWS, P. W. 2012. Human embryonic stem cells fail to activate CHK1 and commit to
781 apoptosis in response to DNA replication stress. *Stem Cells*, 30, 1385-93.

782 DESMARAIS, J. A., UNGER, C., DAMJANOV, I., MEUTH, M. & ANDREWS, P. 2016. Apoptosis
783 and failure of checkpoint kinase 1 activation in human induced pluripotent stem cells under
784 replication stress. *Stem Cell Res Ther*, 7, 17.

785 DRAPER, J. S., SMITH, K., GOKHALE, P., MOORE, H. D., MALTBY, E., JOHNSON, J.,
786 MEISNER, L., ZWAKA, T. P., THOMSON, J. A. & ANDREWS, P. W. 2004. Recurrent gain
787 of chromosomes 17q and 12 in cultured human embryonic stem cells. *Nat Biotechnol*, 22, 53-
788 4.

789 DUMITRU, R., GAMA, V., FAGAN, B. M., BOWER, J. J., SWAHARI, V., PEVNY, L. H. &
790 DESHMUKH, M. 2012. Human embryonic stem cells have constitutively active Bax at the
791 Golgi and are primed to undergo rapid apoptosis. *Mol Cell*, 46, 573-83.

792 EINHORN, L. H. & DONOHUE, J. 1977. Cis-diamminedichloroplatinum, vinblastine, and
793 bleomycin combination chemotherapy in disseminated testicular cancer. *Ann Intern Med*, 87,
794 293-8.

795 EINHORN, L. H., NAGY, C., FURNAS, B. & WILLIAMS, S. D. 1981. Nabilone: an effective
796 antiemetic in patients receiving cancer chemotherapy. *J Clin Pharmacol*, 21, 64S-69S.

797 ENVER, T., SONEJI, S., JOSHI, C., BROWN, J., IBORRA, F., ORNTOFT, T., THYKJAER, T.,
798 MALTBY, E., SMITH, K., ABU DAWUD, R., JONES, M., MATIN, M., GOKHALE, P.,
799 DRAPER, J. & ANDREWS, P. W. 2005. Cellular differentiation hierarchies in normal and
800 culture-adapted human embryonic stem cells. *Hum Mol Genet*, 14, 3129-40.

801 FAZELI, A., LIEW, C. G., MATIN, M. M., ELLIOTT, S., JEANMEURE, L. F., WRIGHT, P. C.,
802 MOORE, H. & ANDREWS, P. W. 2011. Altered patterns of differentiation in karyotypically
803 abnormal human embryonic stem cells. *Int J Dev Biol*, 55, 175-80.

804 FILIPCZYK, A. A., LASLETT, A. L., MUMMERY, C. & PERA, M. F. 2007. Differentiation is
805 coupled to changes in the cell cycle regulatory apparatus of human embryonic stem cells. *Stem*
806 *Cell Res*, 1, 45-60.

807 FRAME, F. M., ROGOFF, H. A., PICKERING, M. T., CRESS, W. D. & KOWALIK, T. F. 2006.
808 E2F1 induces MRN foci formation and a cell cycle checkpoint response in human fibroblasts.
809 *Oncogene*, 25, 3258-66.

810 GARBER, K. 2015. RIKEN suspends first clinical trial involving induced pluripotent stem cells. *Nat*
811 *Biotechnol*, 33, 890-1.

812 GORGOULIS, V. G., VASSILIOU, L. V., KARAKAIDOS, P., ZACHARATOS, P., KOTSINAS,
813 A., LILOGLOU, T., VENERE, M., DITULLIO, R. A., KASTRINAKIS, N. G., LEVY, B.,
814 KLETSAS, D., YONETA, A., HERLYN, M., KITTAS, C. & HALAZONETIS, T. D. 2005.
815 Activation of the DNA damage checkpoint and genomic instability in human precancerous
816 lesions. *Nature*, 434, 907-13.

817 HAINAUT, P. & HOLLSTEIN, M. 1999. p53 and Human Cancer: The First Ten Thousand
818 Mutations. In: VANDE WOUDE, G. F. & KLEIN, G. (eds.) *Advances in Cancer Research*.
819 Academic Press.

820 HALAZONETIS, T. D., GORGOULIS, V. G. & BARTEK, J. 2008. An oncogene-induced DNA
821 damage model for cancer development. *Science*, 319, 1352-5.

822 HAUPT, S., MEJÍA-HERNÁNDEZ, J. O., VIJAYAKUMARAN, R., KEAM, S. P. & HAUPT, Y.
823 2019. The long and the short of it: the MDM4 tail so far. *Journal of Molecular Cell Biology*,
824 11, 231-244.

825 HONG, Y. & STAMBROOK, P. J. 2004. Restoration of an absent G1 arrest and protection from
826 apoptosis in embryonic stem cells after ionizing radiation. *Proc Natl Acad Sci U S A*, 101,
827 14443-8.

828 HUSSEIN, S. M., BATADA, N. N., VUORISTO, S., CHING, R. W., AUTIO, R., NÄRVÄ, E., NG,
829 S., SOUROUR, M., HÄMÄLÄINEN, R., OLSSON, C., LUNDIN, K., MIKKOLA, M.,
830 TROKOVIC, R., PEITZ, M., BRÜSTLE, O., BAZETT-JONES, D. P., ALITALO, K.,
831 LAHESMAA, R., NAGY, A. & OTONKOSKI, T. 2011. Copy number variation and selection
832 during reprogramming to pluripotency. *Nature*, 471, 58-62.

833 HYKA-NOUSPIKEL, N., DESMARAIS, J., GOKHALE, P. J., JONES, M., MEUTH, M.,
834 ANDREWS, P. W. & NOUSPIKEL, T. 2012. Deficient DNA damage response and cell cycle
835 checkpoints lead to accumulation of point mutations in human embryonic stem cells. *Stem*
836 *Cells*, 30, 1901-10.

837 IMREH, M. P., GERTOW, K., CEDERVALL, J., UNGER, C., HOLMBERG, K., SZÖKE, K.,
838 CSÖREGH, L., FRIED, G., DILBER, S., BLENNOW, E. & AHRLUND-RICHTER, L. 2006.
839 In vitro culture conditions favoring selection of chromosomal abnormalities in human ES
840 cells. *J Cell Biochem*, 99, 508-16.

841 JONES, R. M., MORTUSEWICZ, O., AFZAL, I., LORVELLEC, M., GARCÍA, P., HELLEDAY,
842 T. & PETERMANN, E. 2013. Increased replication initiation and conflicts with transcription
843 underlie Cyclin E-induced replication stress. *Oncogene*, 32, 3744-53.

844 KANDOTH, C., MCLELLAN, M. D., VANDIN, F., YE, K., NIU, B., LU, C., XIE, M., ZHANG,
845 Q., MCMICHAEL, J. F., WYCZALKOWSKI, M. A., LEISERSON, M. D. M., MILLER, C.
846 A., WELCH, J. S., WALTER, M. J., WENDL, M. C., LEY, T. J., WILSON, R. K.,
847 RAPHAEL, B. J. & DING, L. 2013. Mutational landscape and significance across 12 major
848 cancer types. *Nature*, 502, 333-339.

849 KOPS, G. J., FOLTZ, D. R. & CLEVELAND, D. W. 2004. Lethality to human cancer cells through
850 massive chromosome loss by inhibition of the mitotic checkpoint. *Proc Natl Acad Sci U S A*,
851 101, 8699-704.

852 KORKOLA, J. E., HOULDSWORTH, J., CHADALAVADA, R. S., OLSHEN, A. B.,
853 DOBRZYNSKI, D., REUTER, V. E., BOSL, G. J. & CHAGANTI, R. S. 2006. Down-
854 regulation of stem cell genes, including those in a 200-kb gene cluster at 12p13.31, is
855 associated with in vivo differentiation of human male germ cell tumors. *Cancer Res*, 66, 820-
856 7.

857 KUCAB, J. E., ZOU, X., MORGANELLA, S., JOEL, M., NANDA, A. S., NAGY, E., GOMEZ, C.,
858 DEGASPERI, A., HARRIS, R., JACKSON, S. P., ARLT, V. M., PHILLIPS, D. H. & NIK-
859 ZAINAL, S. 2019. A Compendium of Mutational Signatures of Environmental Agents. *Cell*,
860 177, 821-836.e16.

861 KUIJK, E., JAGER, M., ROEST, B. V. D., LOCATI, M., HOECK, A. V., KORZELIUS, J.,
862 JANSSEN, R., BESSELINK, N., BOYMANS, S., BOXTEL, R. V. & CUPPEN, E. 2018.
863 Mutational impact of culturing human pluripotent and adult stem cells. *bioRxiv*, 430165.

864 LAMM, N., BEN-DAVID, U., GOLAN-LEV, T., STORCHOVÁ, Z., BENVENISTY, N. &
865 KEREM, B. 2016. Genomic Instability in Human Pluripotent Stem Cells Arises from
866 Replicative Stress and Chromosome Condensation Defects. *Cell Stem Cell*, 18, 253-61.

867 LAURENT, L. C., ULITSKY, I., SLAVIN, I., TRAN, H., SCHORK, A., MOREY, R., LYNCH, C.,
868 HARNESS, J. V., LEE, S., BARRERO, M. J., KU, S., MARTYNOVA, M., SEMECHKIN,
869 R., GALAT, V., GOTTESFELD, J., IZPISUA BELMONTE, J. C., MURRY, C.,
870 KEIRSTEAD, H. S., PARK, H. S., SCHMIDT, U., LASLETT, A. L., MULLER, F. J.,
871 NIEVERGELT, C. M., SHAMIR, R. & LORING, J. F. 2011. Dynamic changes in the copy
872 number of pluripotency and cell proliferation genes in human ESCs and iPSCs during
873 reprogramming and time in culture. *Cell Stem Cell*, 8, 106-18.

874 LEE, C. T., BENDRIEM, R. M., KINDBERG, A. A., WORDEN, L. T., WILLIAMS, M. P.,
875 DRGON, T., MALLON, B. S., HARVEY, B. K., RICHIE, C. T., HAMILTON, R. S., CHEN,
876 J., ERRICO, S. L., TSAI, S. Y., UHL, G. R. & FREED, W. J. 2015. Functional consequences
877 of 17q21.31/WNT3-WNT9B amplification in hPSCs with respect to neural differentiation.
878 *Cell Rep*, 10, 616-32.

879 LEFORT, N., FEYEUX, M., BAS, C., FÉRAUD, O., BENNACEUR-GRISCELLI, A.,
880 TACHDJIAN, G., PESCHANSKI, M. & PERRIER, A. L. 2008. Human embryonic stem cells
881 reveal recurrent genomic instability at 20q11.21. *Nat Biotechnol*, 26, 1364-6.

882 LIU, J. C., GUAN, X., RYAN, J. A., RIVERA, A. G., MOCK, C., AGRAWAL, V., AGARWAL,
883 V., LETAI, A., LEROU, P. H. & LAHAV, G. 2013. High mitochondrial priming sensitizes
884 hESCs to DNA-damage-induced apoptosis. *Cell Stem Cell*, 13, 483-91.

885 LUKAS, C., SAVIC, V., BEKKER-JENSEN, S., DOIL, C., NEUMANN, B., PEDERSEN, R. S.,
886 GRØFTE, M., CHAN, K. L., HICKSON, I. D., BARTEK, J. & LUKAS, J. 2011. 53BP1
887 nuclear bodies form around DNA lesions generated by mitotic transmission of chromosomes
888 under replication stress. *Nat Cell Biol*, 13, 243-53.

889 LUO, L. Z., GOPALAKRISHNA-PILLAI, S., NAY, S. L., PARK, S. W., BATES, S. E., ZENG, X.,
890 IVERSON, L. E. & O'CONNOR, T. R. 2012. DNA repair in human pluripotent stem cells is
891 distinct from that in non-pluripotent human cells. *PLoS One*, 7, e30541.

892 MA, X., ZHENG, W., WEI, D., MA, Y., WANG, T., WANG, J., LIU, Q. & YANG, S. 2006. High-
893 level expression, purification and pro-apoptosis activity of HIV-TAT-survivin (T34A) mutant
894 to cancer cells in vitro. *J Biotechnol*, 123, 367-78.

895 MADDEN, D. T., DAVILA-KRUGER, D., MELOV, S. & BREDESEN, D. E. 2011. Human
896 embryonic stem cells express elevated levels of multiple pro-apoptotic BCL-2 family
897 members. *PLoS One*, 6, e28530.

898 MALKIN, D., LI, F. P., STRONG, L. C., FRAUMENI, J. F., NELSON, C. E., KIM, D. H., KASSEL,
899 J., GRYKA, M. A., BISCHOFF, F. Z. & TAINSKY, M. A. 1990. Germ line p53 mutations
900 in a familial syndrome of breast cancer, sarcomas, and other neoplasms. *Science*, 250, 1233-
901 8.

902 MANDAI, M., KURIMOTO, Y. & TAKAHASHI, M. 2017. Autologous Induced Stem-Cell-Derived
903 Retinal Cells for Macular Degeneration. *N Engl J Med*, 377, 792-793.

904 MARKOULI, C., COUVREU DE DECKERSBERG, E., REGIN, M., NGUYEN, H. T.,
905 ZAMBELLI, F., KELLER, A., DZIEDZICKA, D., DE KOCK, J., TILLEMANN, L., VAN
906 NIEUWERBURGH, F., FRANCESCHINI, L., SERMON, K., GEENS, M. & SPITS, C.
907 2019. Gain of 20q11.21 in Human Pluripotent Stem Cells Impairs TGF- β -Dependent
908 Neuroectodermal Commitment. *Stem Cell Reports*, 13, 163-176.

909 MAYNARD, S., SWISTOWSKA, A. M., LEE, J. W., LIU, Y., LIU, S. T., DA CRUZ, A. B., RAO,
910 M., DE SOUZA-PINTO, N. C., ZENG, X. & BOHR, V. A. 2008. Human embryonic stem
911 cells have enhanced repair of multiple forms of DNA damage. *Stem Cells*, 26, 2266-74.

912 MERKLE, F. T., GHOSH, S., KAMITAKI, N., MITCHELL, J., AVIOR, Y., MELLO, C., KASHIN,
913 S., MEKHOUBAD, S., ILIC, D., CHARLTON, M., SAPHIER, G., HANDSAKER, R. E.,
914 GENOVESE, G., BAR, S., BENVENISTY, N., MCCARROLL, S. A. & EGGAN, K. 2017.
915 Human pluripotent stem cells recurrently acquire and expand dominant negative P53
916 mutations. *Nature*, 545, 229-233.

917 MESRI, M., WALL, N. R., LI, J., KIM, R. W. & ALTIERI, D. C. 2001. Cancer gene therapy using
918 a survivin mutant adenovirus. *J Clin Invest*, 108, 981-90.

919 MILHOLLAND, B., DONG, X., ZHANG, L., HAO, X., SUH, Y. & VIJG, J. 2017. Differences
920 between germline and somatic mutation rates in humans and mice. *Nat Commun*, 8, 15183.

921 MOMCILOVIĆ, O., CHOI, S., VARUM, S., BAKKENIST, C., SCHATTEEN, G. & NAVARA, C.
922 2009. Ionizing radiation induces ataxia telangiectasia mutated-dependent checkpoint
923 signaling and G(2) but not G(1) cell cycle arrest in pluripotent human embryonic stem cells.
924 *Stem Cells*, 27, 1822-35.

925 MOSTOFI, F. K. & PRICE, E. B. 1973. Tumors of the male genital system. A.F.I.P. Atlas of Tumor

926 Pathology. 2nd series, Fascicle 8, Washington D.C.

927 NAIM, V., WILHELM, T., DEBATISSE, M. & ROSSELLI, F. 2013. ERCC1 and MUS81-EME1
928 promote sister chromatid separation by processing late replication intermediates at common
929 fragile sites during mitosis. *Nat Cell Biol*, 15, 1008-15.

930 NGUYEN, H. T., GEENS, M. & SPITS, C. 2013. Genetic and epigenetic instability in human
931 pluripotent stem cells. *Hum Reprod Update*, 19, 187-205.

932 NICHOLS, J. & SMITH, A. 2009. Naive and primed pluripotent states. *Cell Stem Cell*, 4, 487-92.

933 OHGUSHI, M., MATSUMURA, M., EIRAKU, M., MURAKAMI, K., ARAMAKI, T.,
934 NISHIYAMA, A., MUGURUMA, K., NAKANO, T., SUGA, H., UENO, M., ISHIZAKI, T.,
935 SUEMORI, H., NARUMIYA, S., NIWA, H. & SASAI, Y. 2010. Molecular pathway and cell
936 state responsible for dissociation-induced apoptosis in human pluripotent stem cells. *Cell Stem*
937 *Cell*, 7, 225-39.

938 OLARIU, V., HARRISON, N. J., COCA, D., GOKHALE, P. J., BAKER, D., BILLINGS, S.,
939 KADIRKAMANATHAN, V. & ANDREWS, P. W. 2010. Modeling the evolution of culture-
940 adapted human embryonic stem cells. *Stem Cell Res*, 4, 50-6.

941 OOSTERHUIS, J. W., ANDREWS, P. W., KNOWLES, B. B. & DAMJANOV, I. 1984. Effects of
942 cis-platinum on embryonal carcinoma cell lines in vitro. *Int J Cancer*, 34, 133-9.

943 PETLJAK, M., ALEXANDROV, L. B., BRAMMELD, J. S., PRICE, S., WEDGE, D. C.,
944 GROSSMANN, S., DAWSON, K. J., JU, Y. S., IORIO, F., TUBIO, J. M. C., KOH, C. C.,
945 GEORGAKOPOULOS-SOARES, I., RODRÍGUEZ-MARTÍN, B., OTLU, B., O'MEARA,
946 S., BUTLER, A. P., MENZIES, A., BHOSLE, S. G., RAINE, K., JONES, D. R., TEAGUE,
947 J. W., BEAL, K., LATIMER, C., O'NEILL, L., ZAMORA, J., ANDERSON, E., PATEL, N.,
948 MADDISON, M., NG, B. L., GRAHAM, J., GARNETT, M. J., MCDERMOTT, U., NIK-
949 ZAINAL, S., CAMPBELL, P. J. & STRATTON, M. R. 2019. Characterizing Mutational
950 Signatures in Human Cancer Cell Lines Reveals Episodic APOBEC Mutagenesis. *Cell*, 176,
951 1282-1294.e20.

952 PICKERING, M. T. & KOWALIK, T. F. 2006. Rb inactivation leads to E2F1-mediated DNA double-
953 strand break accumulation. *Oncogene*, 25, 746-55.

954 PLANTAZ, D., MOHAPATRA, G., MATTHAY, K. K., PELLARIN, M., SEEGER, R. C. &
955 FEUERSTEIN, B. G. 1997. Gain of chromosome 17 is the most frequent abnormality detected
956 in neuroblastoma by comparative genomic hybridization. *Am J Pathol*, 150, 81-9.

957 PRICE, C. J., STAVISH, D., GOKHALE, P. J., SARGEANT, S., LACEY, J., RODRIGUEZ, T. A.
958 & BARBARIC, I. 2019. Genetically variant human pluripotent stem cells selectively
959 eliminate wild-type counterparts through YAP-mediated cell competition. *bioRxiv*, 854430.

960 REUBINOFF, B. E., PERA, M. F., FONG, C. Y., TROUNSON, A. & BONGSO, A. 2000.
961 Embryonic stem cell lines from human blastocysts: somatic differentiation in vitro. *Nat*
962 *Biotechnol*, 18, 399-404.

963 RODRIGUEZ, S., JAFER, O., GOKER, H., SUMMERSGILL, B. M., ZAFARANA, G., GILLIS, A.
964 J., VAN GURP, R. J., OOSTERHUIS, J. W., LU, Y. J., HUDDART, R., COOPER, C. S.,
965 CLARK, J., LOOIJENGA, L. H. & SHIPLEY, J. M. 2003. Expression profile of genes from
966 12p in testicular germ cell tumors of adolescents and adults associated with i(12p) and
967 amplification at 12p11.2-p12.1. *Oncogene*, 22, 1880-91.

968 ROUHANI, F. J., NIK-ZAINAL, S., WUSTER, A., LI, Y., CONTE, N., KOIKE-YUSA, H.,
969 KUMASAKA, N., VALLIER, L., YUSA, K. & BRADLEY, A. 2016. Mutational History of
970 a Human Cell Lineage from Somatic to Induced Pluripotent Stem Cells. *PLoS Genet*, 12,
971 e1005932.

972 SARETZKI, G., WALTER, T., ATKINSON, S., PASSOS, J. F., BARETH, B., KEITH, W. N.,
973 STEWART, R., HOARE, S., STOJKOVIC, M., ARMSTRONG, L., VON ZGLINICKI, T. &
974 LAKO, M. 2008. Downregulation of multiple stress defense mechanisms during
975 differentiation of human embryonic stem cells. *Stem Cells*, 26, 455-64.

976 SATO, Y., BANDO, H., DI PIAZZA, M., GOWING, G., HERBERTS, C., JACKMAN, S., LEONI,
977 G., LIBERTINI, S., MACLACHLAN, T., MCBLANE, J. W., PEREIRA MOURIÈS, L.,

978 SHARPE, M., SHINGLETON, W., SURMACZ-CORDLE, B., YAMAMOTO, K. & VAN
979 DER LAAN, J. W. 2019. Tumorigenicity assessment of cell therapy products: The need for
980 global consensus and points to consider. *Cytotherapy*, 21, 1095-1111.

981 SCHWARTZ, S. D., HUBSCHMAN, J. P., HEILWELL, G., FRANCO-CARDENAS, V., PAN, C.
982 K., OSTRICK, R. M., MICKUNAS, E., GAY, R., KLIMANSKAYA, I. & LANZA, R. 2012.
983 Embryonic stem cell trials for macular degeneration: a preliminary report. *Lancet*, 379, 713-
984 20.

985 SIMARA, P., TESAROVA, L., REHAKOVA, D., MATULA, P., STEJSKAL, S., HAMPL, A. &
986 KOUTNA, I. 2017. DNA double-strand breaks in human induced pluripotent stem cell
987 reprogramming and long-term in vitro culturing. *Stem Cell Res Ther*, 8, 73.

988 SONG, W. K., PARK, K. M., KIM, H. J., LEE, J. H., CHOI, J., CHONG, S. Y., SHIM, S. H., DEL
989 PRIORE, L. V. & LANZA, R. 2015. Treatment of macular degeneration using embryonic
990 stem cell-derived retinal pigment epithelium: preliminary results in Asian patients. *Stem Cell*
991 *Reports*, 4, 860-72.

992 SPITS, C., MATEIZEL, I., GEENS, M., MERTZANIDOU, A., STAESSEN, C., VANDESKELDE,
993 Y., VAN DER ELST, J., LIEBAERS, I. & SERMON, K. 2008. Recurrent chromosomal
994 abnormalities in human embryonic stem cells. *Nat Biotechnol*, 26, 1361-3.

995 SRIVASTAVA, S., ZOU, Z. Q., PIROLLO, K., BLATTNER, W. & CHANG, E. H. 1990. Germ-
996 line transmission of a mutated p53 gene in a cancer-prone family with Li-Fraumeni syndrome.
997 *Nature*, 348, 747-9.

998 TAAPKEN, S. M., NISLER, B. S., NEWTON, M. A., SAMPSELL-BARRON, T. L., LEONHARD,
999 K. A., MCINTIRE, E. M. & MONTGOMERY, K. D. 2011. Karotypic abnormalities in human
1000 induced pluripotent stem cells and embryonic stem cells. *Nat Biotechnol*, 29, 313-4.

1001 TAKAHASHI, K., TANABE, K., OHNUKI, M., NARITA, M., ICHISAKA, T., TOMODA, K. &
1002 YAMANAKA, S. 2007. Induction of pluripotent stem cells from adult human fibroblasts by
1003 defined factors. *Cell*, 131, 861-72.

1004 TECHNOLOGIES, S. 2018. *Challenges in Ensuring hPSC Quality*. *Stemcell.com* [Online].
1005 Available: <https://www.stemcell.com/nature-research-roundtable-hPSC-quality> [Accessed].

1006 THOMPSON, O., VON MEYENN, F., HEWITT, Z., ALEXANDER, J., WOOD, A.,
1007 WEIGHTMAN, R., GREGORY, S., KRUEGER, F., ANDREWS, S., BARBARIC, I.,
1008 GOKHALE, P. J., MOORE, H. D., REIK, W., MILO, M., NIK-ZAINAL, S., YUSA, K. &
1009 ANDREWS, P. W. 2020. Low rates of mutation in clinical grade human pluripotent stem cells
1010 under different culture conditions. *Nature Communications*, 11, 1528.

1011 THOMSON, J. A., ITSKOVITZ-ELDOR, J., SHAPIRO, S. S., WAKNITZ, M. A., SWIERGIEL, J.
1012 J., MARSHALL, V. S. & JONES, J. M. 1998. Embryonic stem cell lines derived from human
1013 blastocysts. *Science*, 282, 1145-7.

1014 TORRES, E. M., WILLIAMS, B. R. & AMON, A. 2008. Aneuploidy: cells losing their balance.
1015 *Genetics*, 179, 737-46.

1016 VALLABHANENI, H., LYNCH, P. J., CHEN, G., PARK, K., LIU, Y., GOEHE, R., MALLON, B.
1017 S., BOEHM, M. & HURSH, D. A. 2018. High Basal Levels of γ H2AX in Human Induced
1018 Pluripotent Stem Cells Are Linked to Replication-Associated DNA Damage and Repair. *Stem*
1019 *Cells*, 36, 1501-1513.

1020 VAN ECHTEN-ARENDS, J., MASTENBROEK, S., SIKKEMA-RADDATZ, B., KOREVAAR, J.
1021 C., HEINEMAN, M. J., VAN DER VEEN, F. & REPPING, S. 2011. Chromosomal
1022 mosaicism in human preimplantation embryos: a systematic review. *Hum Reprod Update*, 17,
1023 620-7.

1024 VIEL, A., BRUSELLES, A., MECCIA, E., FORNASARIG, M., QUAIA, M., CANZONIERI, V.,
1025 POLICICCHIO, E., URSO, E. D., AGOSTINI, M., GENUARDI, M., LUCCI-CORDISCO,
1026 E., VENESIO, T., MARTAYAN, A., DIODORO, M. G., SANCHEZ-METE, L.,
1027 STIGLIANO, V., MAZZEI, F., GRASSO, F., GIULIANI, A., BAIOCCHI, M., MAESTRO,
1028 R., GIANNINI, G., TARTAGLIA, M., ALEXANDROV, L. B. & BIGNAMI, M. 2017. A

1029 Specific Mutational Signature Associated with DNA 8-Oxoguanine Persistence in MUTYH-
1030 defective Colorectal Cancer. *EBioMedicine*, 20, 39-49.

1031 WATANABE, K., UENO, M., KAMIYA, D., NISHIYAMA, A., MATSUMURA, M., WATAYA,
1032 T., TAKAHASHI, J. B., NISHIKAWA, S., MUGURUMA, K. & SASAI, Y. 2007. A ROCK
1033 inhibitor permits survival of dissociated human embryonic stem cells. *Nat Biotechnol*, 25,
1034 681-6.

1035 WERBOWETSKI-OGILVIE, T. E., BOSSÉ, M., STEWART, M., SCHNERCH, A., RAMOS-
1036 MEJIA, V., ROULEAU, A., WYNDER, T., SMITH, M. J., DINGWALL, S., CARTER, T.,
1037 WILLIAMS, C., HARRIS, C., DOLLING, J., WYNDER, C., BOREHAM, D. & BHATIA,
1038 M. 2009. Characterization of human embryonic stem cells with features of neoplastic
1039 progression. *Nat Biotechnol*, 27, 91-7.

1040 YANG, D., WELM, A. & BISHOP, J. M. 2004. Cell division and cell survival in the absence of
1041 survivin. *Proc Natl Acad Sci U S A*, 101, 15100-5.

1042 YASUDA, S., KUSAKAWA, S., KURODA, T., MIURA, T., TANO, K., TAKADA, N.,
1043 MATSUYAMA, S., MATSUYAMA, A., NASU, M., UMEZAWA, A., HAYAKAWA, T.,
1044 TSUTSUMI, H. & SATO, Y. 2018. Tumorigenicity-associated characteristics of human iPS
1045 cell lines. *PLoS One*, 13, e0205022.

1046 YU, J., VODYANIK, M. A., SMUGA-OTTO, K., ANTOSIEWICZ-BOURGET, J., FRANE, J. L.,
1047 TIAN, S., NIE, J., JONSDOTTIR, G. A., RUOTTI, V., STEWART, R., SLUKVIN, I. I. &
1048 THOMSON, J. A. 2007. Induced pluripotent stem cell lines derived from human somatic
1049 cells. *Science*, 318, 1917-20.

1050 ZHANG, J., HIRST, A. J., DUAN, F., QIU, H., HUANG, R., JI, Y., BAI, L., ZHANG, F.,
1051 ROBINSON, D., JONES, M., LI, L., WANG, P., JIANG, P., ANDREWS, P. W.,
1052 BARBARIC, I. & NA, J. 2019. Anti-apoptotic Mutations Desensitize Human Pluripotent
1053 Stem Cells to Mitotic Stress and Enable Aneuploid Cell Survival. *Stem Cell Reports*, 12, 557-
1054 571.

1055

1056 **FIGURE LEGENDS**

1057

1058 **Figure 1: Appearance of mutations in human PSC involves both mutation and**
1059 **selection.**

1060 Human PSC are subject to the full range of mutations seen in other systems, including single
1061 base changes and small insertions and deletions (INDELS) (Thompson et al., 2020), as well
1062 as larger scale genomic rearrangements with gains or losses of whole chromosomes or
1063 chromosomal fragments that alter the number of copies of whole sets of genes, and
1064 consequently their levels of expression (Enver et al., 2005). The most commonly seen
1065 chromosomal rearrangements are illustrated. Generally, in the absence of a clonogenic
1066 bottleneck, these mutations will never be detected unless they offer the mutant cell a
1067 selective growth advantage (Olariu et al., 2010).

1068

1069 **Figure 2: Nature of tumors derived from pluripotent stem cells**

1070 PSC produce tumours that may contain both undifferentiated PSC together with their
1071 differentiated derivatives, in which case the tumor is termed a 'Teratocarcinoma' and is
1072 regarded as highly malignant. If the PSC fully differentiate, so that the tumor contains only
1073 differentiated derivatives, it is termed a 'Teratoma' and is generally regarded as benign and
1074 not malignant (Damjanov and Andrews, 2007a, Damjanov and Andrews, 2016). However,
1075 a caveat is that some of the differentiated derivatives may develop into a secondary
1076 malignancy corresponding to their particular cell type, and this may be driven by a mutation
1077 present in the parent PSC. In addition to somatic derivatives, PSC may also generate
1078 primitive endoderm elements which are precursors of highly malignant yolk sac carcinomas

1079

1080 **Figure 3: Mechanisms of Variant Growth Advantage**

1081 A variety of mechanisms can be envisaged by which variant cells could gain a growth
1082 advantage over wild type cells. **Cell autonomous mechanisms:** mutation **(a)** drives faster
1083 cell cycle, or **(b)** blocks differentiation, or **(c)** blocks apoptosis. **Cell interactive**
1084 **mechanisms:** mutation **(d)** causes variant cell to inhibit the growth of its wild-type
1085 counterpart (Price et al., 2019) or **(e)** alters the patterns of differentiation. The latter could
1086 generate a selective advantage either if certain differentiated derivatives produced factors
1087 that promote differentiation (advantage would derive from blocking such lineages) or
1088 produced factors that blocked general differentiation (advantage would derive from
1089 enhancing differentiation to such lineages).

1090

1091 **Figure 4: Overview of the origins of mutation in human PSC. a)** Somatic cellular
1092 proliferation occurs predominantly without replication stress, allowing for faithful cell division.
1093 However, should replication stress occur, somatic cells can respond with cellular
1094 senescence, apoptosis or DNA repair. **b)** Human pluripotent stem cells proliferate rapidly
1095 and are susceptible to replication stress, DNA damage and mitotic errors. However, they
1096 display a low mutation rate, which is reconciled by responding to genetic stresses with
1097 apoptosis and efficient DNA repair. **c)** Culture of human PSC with exogenous nucleoside
1098 alleviates replication stress, DNA damage and mitotic errors. Alleviating genetic stress
1099 minimises apoptosis, enhancing growth rate and removing the selective advantage of anti-
1100 apoptotic mutations.

1101

1102

1103

1104

1105 **BOX 1. Detection of Genetically Variant Cells**

1106 Routine monitoring of PSC cultures for the appearance of genetic variants is essential: the proportion
1107 of some common variants in a culture can rise from undetectable to 100% within 5 -10 passages
1108 (Olariu et al., 2010). Different screening methods have different advantages and limitations,
1109 including the minimum proportions of variant cells that can be detected in mosaic populations, and
1110 whether prior knowledge of the variant to be detected is required. Other factors to consider are the
1111 types of variant detectable, cost, speed and whether specialist facilities and expertise are required.
1112 At a minimum we would recommend regular screening of cultures for common variants using the
1113 PCR method, which is rapid and can easily be carried out in most laboratories, backed up at key
1114 decision points in specific experiments with more in-depth analysis using other methods as
1115 appropriate.

- 1116 • **Karyotyping by G banding:** Indiscriminate view of structural and numerical chromosomal
1117 abnormalities greater than 5Mb (Steinemann et al., 2013). Limited to analysis of proliferating
1118 cells, but the only convenient method to detect balanced genomic rearrangements.
1119 Sensitivity and precision can be enhanced by use of fluorescent hybridization probes.
1120 Minimum mosaicism detected: 18% (routine analysis of 30 metaphase spreads), 6%
1121 (analysis of 100 metaphase spreads) (Baker et al., 2016).
- 1122 • **Array Comparative Genome Hybridisation (aCGH) and SNP array:** Global screening
1123 detects copy number variants down to 1Kb. Minimum mosaicism detected: 10-15% for small
1124 copy number variants. (Valli et al., 2011).
- 1125 • **Interphase Fluorescent In Situ Hybridisation (FISH):** Hybridisation of fluorescent probes
1126 to highly complementary nucleic acid sequences in interphase cells. Requires prior
1127 knowledge of the copy number variant to be detected. The presence of false negatives
1128 dependent upon the type of variant restricts the sensitivity of detection: Minimum mosaicism
1129 detected depends upon the number of nuclei assessed and the size of the CNV: ~5% is
1130 possible for gains of whole chromosome arms, with 100 nuclei assessed, but this falls
1131 markedly for smaller CNV (Baker et al., 2016).
- 1132 • **qPCR based assays:** Rapid method suitable for any well-equipped molecular cell biology
1133 laboratories. Requires prior knowledge of the CNV to be detected. Minimum mosaicism
1134 detected: ~5% (Baker et al., 2016).
- 1135 • **Expression karyotyping:** Regions of CNV are detected based on their global gene
1136 expression from RNA-seq data in comparison to a diploid calibrator sample. Requires no
1137 prior knowledge of the CNV to be detected, although it is only capable of detecting
1138 chromosomal aberrations greater than ~10Mb with a sensitivity of ~30% (Maysnar et al.,
1139 2010).

- 1140 • **eSNP Karyotyping:** An adaptation of expression karyotyping with allele bias. This approach
1141 does not require a diploid calibrator sample for CNV detection. Resolution of eSNP
1142 karyotyping depends on the genomic region and sequencing depth, yet it comfortably
1143 detected amplifications to entire chromosomes or chromosome arms. The eSNP
1144 karyotyping approach shows comparable sensitivity to other DNA based array approaches
1145 (~30%) (Weissbein et al., 2016).
- 1146
- 1147 BAKER, D., HIRST, A. J., GOKHALE, P. J., JUAREZ, M. A., WILLIAMS, S., WHEELER, M., BEAN,
1148 K., ALLISON, T. F., MOORE, H. D., ANDREWS, P. W. & BARBARIC, I. 2016. Detecting
1149 Genetic Mosaicism in Cultures of Human Pluripotent Stem Cells. *Stem Cell Reports*, 7, 998-
1150 1012.
- 1151 MAYSHAR, Y., BEN-DAVID, U., LAVON, N., BIANCOTTI, J. C., YAKIR, B., CLARK, A. T., PLATH,
1152 K., LOWRY, W. E. & BENVENISTY, N. 2010. Identification and classification of chromosomal
1153 aberrations in human induced pluripotent stem cells. *Cell Stem Cell*, 7, 521-31.
- 1154 OLARIU, V., HARRISON, N. J., COCA, D., GOKHALE, P. J., BAKER, D., BILLINGS, S.,
1155 KADIRKAMANATHAN, V. & ANDREWS, P. W. 2010. Modeling the evolution of culture-
1156 adapted human embryonic stem cells. *Stem Cell Res*, 4, 50-6.
- 1157 STEINEMANN, D., GÖHRING, G. & SCHLEGELBERGER, B. 2013. Genetic instability of modified
1158 stem cells - a first step towards malignant transformation? *Am J Stem Cells*, 2, 39-51.
- 1159 VALLI, R., MARLETTA, C., PRESSATO, B., MONTALBANO, G., LO CURTO, F., PASQUALI, F. &
1160 MASERATI, E. 2011. Comparative genomic hybridization on microarray (a-CGH) in
1161 constitutional and acquired mosaicism may detect as low as 8% abnormal cells. *Mol*
1162 *Cytogenet*, 4, 13.
- 1163 WEISSBEIN, U., SCHACHTER, M., EGLI, D. & BENVENISTY, N. 2016. Analysis of chromosomal
1164 aberrations and recombination by allelic bias in RNA-Seq. *Nat Commun*, 7, 12144.
- 1165
- 1166
- 1167

BOX 2: Features of common genetic changes in PSC.

Gains:

Chromosome 1q

- Typically acquired as a structural chromosome rearrangement (unbalanced translocation or interstitial duplication) (Amps et al., 2011, Baker et al., 2016)
- Minimal region: 1q25-q41 (Baker et al., 2016)
- Potential driver gene: *MDM4*
- Association with cancer: Wilms' tumor (Chagtai et al., 2016), multiple myeloma (Walker et al., 2010) and intracranial ependymomas (Kilday et al., 2012)

Chromosome 12p

- Typically acquired as a whole chromosome trisomy, although a number of studies have also reported a partial trisomy of the short arm of chromosome 12 (Amps et al., 2011, Baker et al., 2016)
- Minimal region: 12p11-pter (Baker et al., 2016)
- Potential driver gene: *NANOG*
- Functional consequences observed to date: increased proliferation rate due to increased replication, reduced propensity for spontaneous differentiation and apoptosis (Ben-David et al., 2014)
- Association with cancer: testicular germ cell tumors (Atkin and Baker, 1982)

Chromosome 17q

- Typically acquired as a structural chromosome rearrangement (unbalanced translocation or interstitial duplication) (Amps et al., 2011, Baker et al., 2016)
- Minimal region: 17q25-qter (Baker et al., 2016)
- Potential driver gene: *SURVIVIN (BIRC5)*
- Functional consequences observed to date: growth advantage in undifferentiated cultures (Olariu et al., 2010) due to enhanced proliferation (Lee et al., 2015); faster differentiation to mesodiencephalic dopaminergic neural cells compared with wild-type cells (Lee et al., 2015)
- Association with cancer: testicular germ cell tumors (Kraggerud et al., 2002) and neuroblastoma (Bown et al., 2001)

Chromosome 20q

- Typically acquired as an interstitial duplication (Amps et al., 2011)
- Minimal region: 20q11.21 (Amps et al., 2011)
- Driver gene: *BCL2L1* (Avery et al., 2013, Nguyen et al., 2013)
- Functional consequences observed to date: growth advantage due to reduced propensity for apoptosis (Avery et al., 2013, Nguyen et al., 2013), reduced dependence on bFGF (Werbowski-Ogilvie et al., 2009), delayed differentiation to neural lineages (Werbowski-Ogilvie et al., 2009) and reduced efficiency of neuroectodermal lineage commitment (Markouli et al., 2019)
- Association with cancer: colorectal cancer (Nguyen and Duong, 2018)

Losses:

Chromosome 10p

- Typically acquired as an interstitial deletion (Amps et al., 2011)
- Minimal region: 10p13-pter (Amps et al., 2011, Baker et al., 2016)
- Association with cancer: melanoma (Robertson et al., 1999) and glioblastoma (Kimmelman et al., 1996)

1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274

Chromosome 18q

- Typically acquired as an interstitial deletion (Amps et al., 2011)
- Minimal region: 18q21-qter (Amps et al., 2011, Baker et al., 2016)
- Association with cancer: colorectal carcinoma (Popat and Houlston, 2005)

AMPS, K., ANDREWS, P. W., ANYFANTIS, G., ARMSTRONG, L., AVERY, S.,
BAHARVAND, H., BAKER, J., BAKER, D., MUNOZ, M. B., BEIL, S., BENVENISTY,
N., BEN-YOSEF, D., BIANCOTTI, J. C., BOSMAN, A., BRENA, R. M., BRISON, D.,
CAISANDER, G., CAMARASA, M. V., CHEN, J., CHIAO, E., CHOI, Y. M., CHOO, A.
B., COLLINS, D., COLMAN, A., CROOK, J. M., DALEY, G. Q., DALTON, A., DE
SOUSA, P. A., DENNING, C., DOWNIE, J., DVORAK, P., MONTGOMERY, K. D.,
FEKI, A., FORD, A., FOX, V., FRAGA, A. M., FRUMKIN, T., GE, L., GOKHALE, P. J.,
GOLAN-LEV, T., GOURABI, H., GROPP, M., LU, G., HAMPL, A., HARRON, K.,
HEALY, L., HERATH, W., HOLM, F., HOVATTA, O., HYLLNER, J., INAMDAR, M. S.,
IRWANTO, A. K., ISHII, T., JACONI, M., JIN, Y., KIMBER, S., KISELEV, S.,
KNOWLES, B. B., KOPPER, O., KUKHARENKO, V., KULIEV, A., LAGARKOVA, M.
A., LAIRD, P. W., LAKO, M., LASLETT, A. L., LAVON, N., LEE, D. R., LEE, J. E., LI,
C., LIM, L. S., LUDWIG, T. E., MA, Y., MALTBY, E., MATEIZEL, I., MAYSHAR, Y.,
MILEIKOVSKY, M., MINGER, S. L., MIYAZAKI, T., MOON, S. Y., MOORE, H.,
MUMMERY, C., NAGY, A., NAKATSUJI, N., NARWANI, K., OH, S. K., OLSON, C.,
OTONKOSKI, T., PAN, F., PARK, I. H., PELLIS, S., PERA, M. F., PEREIRA, L. V., QI,
O., RAJ, G. S., REUBINOFF, B., ROBINS, A., ROBSON, P., ROSSANT, J., SALEKDEH,
G. H., SCHULZ, T. C., et al. 2011. Screening ethnically diverse human embryonic stem
cells identifies a chromosome 20 minimal amplicon conferring growth advantage. *Nat
Biotechnol*, 29, 1132-44.

ATKIN, N. B. & BAKER, M. C. 1982. Specific chromosome change, i(12p), in testicular tumours?
Lancet, 2, 1349.

AVERY, S., HIRST, A. J., BAKER, D., LIM, C. Y., ALAGARATNAM, S., SKOTHEIM, R. I.,
LOTHE, R. A., PERA, M. F., COLMAN, A., ROBSON, P., ANDREWS, P. W. &
KNOWLES, B. B. 2013. BCL-XL mediates the strong selective advantage of a 20q11.21
amplification commonly found in human embryonic stem cell cultures. *Stem Cell Reports*,
1, 379-86.

BAKER, D., HIRST, A. J., GOKHALE, P. J., JUAREZ, M. A., WILLIAMS, S., WHEELER, M.,
BEAN, K., ALLISON, T. F., MOORE, H. D., ANDREWS, P. W. & BARBARIC, I. 2016.
Detecting Genetic Mosaicism in Cultures of Human Pluripotent Stem Cells. *Stem Cell
Reports*, 7, 998-1012.

BEN-DAVID, U., ARAD, G., WEISSBEIN, U., MANDEFRO, B., MAIMON, A., GOLAN-LEV,
T., NARWANI, K., CLARK, A. T., ANDREWS, P. W., BENVENISTY, N. & CARLOS
BIANCOTTI, J. 2014. Aneuploidy induces profound changes in gene expression,
proliferation and tumorigenicity of human pluripotent stem cells. *Nat Commun*, 5, 4825.

BOWN, N., LASTOWSKA, M., COTTERILL, S., O'NEILL, S., ELLERSHAW, C., ROBERTS,
P., LEWIS, I., PEARSON, A. D. & GROUP, U. K. C. C. G. A. T. U. K. C. S. C. S. 2001.
17q gain in neuroblastoma predicts adverse clinical outcome. U.K. Cancer Cytogenetics
Group and the U.K. Children's Cancer Study Group. *Med Pediatr Oncol*, 36, 14-9.

CHAGTAI, T., ZILL, C., DAINESE, L., WEGERT, J., SAVOLA, S., POPOV, S., MIFSUD, W.,
VUJANIĆ, G., SEBIRE, N., LE BOUC, Y., AMBROS, P. F., KAGER, L., O'SULLIVAN,
M. J., BLAISE, A., BERGERON, C., MENGELBIER, L. H., GISSELSSON, D., KOOL,
M., TYTGAT, G. A., VAN DEN HEUVEL-EIBRINK, M. M., GRAF, N., VAN
TINTEREN, H., COULOMB, A., GESSLER, M., WILLIAMS, R. D. & PRITCHARD-
JONES, K. 2016. Gain of 1q As a Prognostic Biomarker in Wilms Tumors (WTs) Treated

1275 With Preoperative Chemotherapy in the International Society of Paediatric Oncology (SIOP)
1276 WT 2001 Trial: A SIOP Renal Tumours Biology Consortium Study. *J Clin Oncol*, 34, 3195-
1277 203.

1278 KILDAY, J. P., MITRA, B., DOMERG, C., WARD, J., ANDREIUOLO, F., OSTESO-IBANEZ,
1279 T., MAUGUEN, A., VARLET, P., LE DELEY, M. C., LOWE, J., ELLISON, D. W.,
1280 GILBERTSON, R. J., COYLE, B., GRILL, J. & GRUNDY, R. G. 2012. Copy number gain
1281 of 1q25 predicts poor progression-free survival for pediatric intracranial ependymomas and
1282 enables patient risk stratification: a prospective European clinical trial cohort analysis on
1283 behalf of the Children's Cancer Leukaemia Group (CCLG), Societe Francaise d'Oncologie
1284 Pediatrique (SFOP), and International Society for Pediatric Oncology (SIOP). *Clin Cancer*
1285 *Res*, 18, 2001-11.

1286 KIMMELMAN, A. C., ROSS, D. A. & LIANG, B. C. 1996. Loss of heterozygosity of chromosome
1287 10p in human gliomas. *Genomics*, 34, 250-4.

1288 KRAGGERUD, S. M., SKOTHEIM, R. I., SZYMANSKA, J., EKNAES, M., FOSSÅ, S. D.,
1289 STENWIG, A. E., PELTOMÄKI, P. & LOTHE, R. A. 2002. Genome profiles of
1290 familial/bilateral and sporadic testicular germ cell tumors. *Genes Chromosomes Cancer*, 34,
1291 168-74.

1292 LEE, C. T., BENDRIEM, R. M., KINDBERG, A. A., WORDEN, L. T., WILLIAMS, M. P.,
1293 DRGON, T., MALLON, B. S., HARVEY, B. K., RICHIE, C. T., HAMILTON, R. S.,
1294 CHEN, J., ERRICO, S. L., TSAI, S. Y., UHL, G. R. & FREED, W. J. 2015. Functional
1295 consequences of 17q21.31/WNT3-WNT9B amplification in hPSCs with respect to neural
1296 differentiation. *Cell Rep*, 10, 616-32.

1297 MARKOULI, C., COUVREU DE DECKERSBERG, E., REGIN, M., NGUYEN, H. T.,
1298 ZAMBELLI, F., KELLER, A., DZIEDZICKA, D., DE KOCK, J., TILLEMANN, L., VAN
1299 NIEUWERBURGH, F., FRANCESCHINI, L., SERMON, K., GEENS, M. & SPITS, C.
1300 2019. Gain of 20q11.21 in Human Pluripotent Stem Cells Impairs TGF- β -Dependent
1301 Neuroectodermal Commitment. *Stem Cell Reports*, 13, 163-176.

1302 NGUYEN, H. T. & DUONG, H. Q. 2018. The molecular characteristics of colorectal cancer:
1303 Implications for diagnosis and therapy. *Oncol Lett*, 16, 9-18.

1304 NGUYEN, H. T., GEENS, M. & SPITS, C. 2013. Genetic and epigenetic instability in human
1305 pluripotent stem cells. *Hum Reprod Update*, 19, 187-205.

1306 OLARIU, V., HARRISON, N. J., COCA, D., GOKHALE, P. J., BAKER, D., BILLINGS, S.,
1307 KADIRKAMANATHAN, V. & ANDREWS, P. W. 2010. Modeling the evolution of
1308 culture-adapted human embryonic stem cells. *Stem Cell Res*, 4, 50-6.

1309 POPAT, S. & HOULSTON, R. S. 2005. A systematic review and meta-analysis of the relationship
1310 between chromosome 18q genotype, DCC status and colorectal cancer prognosis. *Eur J*
1311 *Cancer*, 41, 2060-70.

1312 ROBERTSON, G. P., HERBST, R. A., NAGANE, M., HUANG, H. J. & CAVENEE, W. K. 1999.
1313 The chromosome 10 monosomy common in human melanomas results from loss of two
1314 separate tumor suppressor loci. *Cancer Res*, 59, 3596-601.

1315 WALKER, B. A., LEONE, P. E., CHIECCHIO, L., DICKENS, N. J., JENNER, M. W., BOYD, K.
1316 D., JOHNSON, D. C., GONZALEZ, D., DAGRADA, G. P., PROTHEROE, R. K., KONN,
1317 Z. J., STOCKLEY, D. M., GREGORY, W. M., DAVIES, F. E., ROSS, F. M. & MORGAN,
1318 G. J. 2010. A compendium of myeloma-associated chromosomal copy number
1319 abnormalities and their prognostic value. *Blood*, 116, e56-65.

1320 WERBOWETSKI-OGILVIE, T. E., BOSSÉ, M., STEWART, M., SCHNERCH, A., RAMOS-
1321 MEJIA, V., ROULEAU, A., WYNDER, T., SMITH, M. J., DINGWALL, S., CARTER, T.,
1322 WILLIAMS, C., HARRIS, C., DOLLING, J., WYNDER, C., BOREHAM, D. & BHATIA,
1323 M. 2009. Characterization of human embryonic stem cells with features of neoplastic
1324 progression. *Nat Biotechnol*, 27, 91-7.

1325
1326