UNIVERSITY of York

This is a repository copy of Steric and Electronic Effects on the 1H Hyperpolarisation of Substituted Pyridazines by Signal Amplification by Reversible Exchange.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/172122/</u>

Version: Published Version

Article:

Rayner, Peter John orcid.org/0000-0002-6577-4117, Burns, Michael orcid.org/0000-0002-5569-1270, Fear, Elizabeth et al. (1 more author) (2021) Steric and Electronic Effects on the 1H Hyperpolarisation of Substituted Pyridazines by Signal Amplification by Reversible Exchange. MAGNETIC RESONANCE IN CHEMISTRY. pp. 1187-1198. ISSN 0749-1581

https://doi.org/10.1002/mrc.5152

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/ SPECIAL ISSUE RESEARCH ARTICLE

WILEY

Steric and electronic effects on the ¹H hyperpolarisation of substituted pyridazines by signal amplification by reversible exchange

Peter J. Rayner 💿 | Michael J. Burns 💿 | Elizabeth J. Fear 💿 | Simon B. Duckett 💿

Centre for Hyperpolarisation in Magnetic Resonance, Department of Chemistry, University of York, York, UK

Correspondence

Simon B. Duckett, Centre for Hyperpolarisation in Magnetic Resonance, Department of Chemistry, University of York, York YO10 5NY, UK. Email: simon.duckett@york.ac.uk

Funding information

University of York; EPSRC, Grant/Award Number: EP/R51181X/1; Wellcome Trust, Grant/Award Numbers: 092506, 098335

Abstract

Utility of the pyridazine motif is growing in popularity as pharmaceutical and agrochemical agents. The detection and structural characterisation of such materials is therefore imperative for the successful development of new products. Signal amplification by reversible exchange (SABRE) offers a route to dramatically improve the sensitivity of magnetic resonance methods, and we apply it here to the rapid and cost-effective hyperpolarisation of substituted pyridazines. The 33 substrates investigated cover a range of steric and electronic properties and their capacity to perform highly effective SABRE is assessed. We find the method to be tolerant to a broad range of electron donating and withdrawing groups; however, good sensitivity is evident when steric bulk is added to the 3- and 6-positions of the pyridazine ring. We optimise the method by reference to a disubstituted ester that yields signal gains of >9000-fold at 9.4 T (>28% spin polarisation).

KEYWORDS

hyperpolarisation, NMR, parahydrogen, SABRE

1 | INTRODUCTION

Pyridazine is a dinitrogen containing aromatic heterocycle that contains two sp^2 -hybridised nitrogen atoms within a six-membered ring. Whilst being isosteric with benzene, it offers increased possibilities for molecular interactions, higher electron density, lower LogP values and improved crystallinity. This has resulted in its use increasing in both pharmaceutical and agrochemical industries.^[1] Indeed, the replacement of phenyl rings within drug molecules has led to several hundred articles dealing with bioactive pyridazines that belong to almost all therapeutic classes in the last 15 years alone.^[2] As such, pyridazine has been described as a 'privileged structure' within the field of medicinal chemistry.^[3] Most recent examples have focused on the synthesis and evaluation of fused *bi*- and *tri*-cyclic pyridazines such as imidazo,^[2] pyrrolo^[4] and thienopyridazines.^[5]

With such prevalence of pyridazines in medicinal and agricultural chemistry, routes to their rapid and accurate detection and structural determination are necessary. Nuclear magnetic resonance (NMR) is one of the most widely used methods in the analytic sciences and offers unique insight into molecular structure, activity and environment. However, when compared with other analytical techniques, such as mass spectrometry or infrared spectroscopy, the required sample size and concentration are much higher. This is due to the lower sensitivity of

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. Magnetic Resonance in Chemistry published by John Wiley & Sons Ltd.

1

NMR that is caused by very small population differences between the magnetic state energy levels that the method probes. Classically, this insensitivity is overcome by signal averaging and increased concentrations, which has time and cost implications. However, this sensitivity challenge will hinder the ability of NMR to rapidly detect transient intermediates or trace impurities.

NMR hyperpolarisation offers an attractive solution by creating magnetic state energy level populations that deviate from the Boltzmann distribution.^[6] This has led to the development of methods with sensitivity gains that are many orders of magnitude higher than could be achieved using conventional NMR. These large signals have been widely exploited for probing reaction mechanisms,^[7] detection of low concentration species^[8] and the in vivo quantification of anatomy^[9] and metabolism^[10] amongst others. A number of methods for creating the hyperpolarised species are known with spin exchange optical pumping (SEOP),^[11] dynamic nuclear polarisation (DNP)^[6a,12] and *para*hydrogen induced polarisation (PHIP)^[6a,13] being the most common.

Signal amplification by reversible exchange $(SABRE)^{[14]}$ is a PHIP method that derives its hyperpolarised spin order from the singlet state of hydrogen, *para*hydrogen (*p*-H₂). SABRE typically utilises an iridium *N*-heterocyclic catalyst^[15] that reversible binds *p*-H₂, as hydride ligands, and a substrate molecule such that spin polarisation is transferred through the temporarily formed *J*-coupling network.^[16] Since its first report in 2009, SABRE has received wide attention from the research community, and the method and its applications have been recently reviewed in detail.^[17]

Pyridazine has previously been shown to be SABRE hyperpolarisation, and the amenable to mechanism for the formation of the active SABRE catalyst is well defined.^[18] When [IrCl (COD)(IMes)] (where COD = 1.5-cyclooctadience and IMes = 1.3-bis (2,4,6-trimethylphenyl)imidazol-2-ylidene) is reacted with H₂ in the presence of pyridazine at 298 K, [Ir (H)₂(IMes)(pyridazine)₃]Cl is formed which undergoes reversible ligand exchange in methanol- d_4 . Additionally, a hapotropic shift between the two nitrogen atoms in the ring is also observed using EXSY. When this sample was exposed to p-H₂ (3 bar), a ca. 400-fold ¹H signal gain was observed at 9.4 T for its two proton resonances after polarisation transfer at 65 G. The use of selective deuteration enabled the SABRE polarisation level to increase such that $3,5-d_2$ -pyridazine yields polarisation levels of greater than 22% (>7000-fold signal gain at 9.4 T) in conjunction with T_1 lifetimes that approach 3 min when a deuterated catalyst and 5 bar pressure of p-H₂ were employed.^[19] The high symmetry of the pyridazine motif has also been exploited for the creation of long-lived

magnetic states that allow detection of a ¹H NMR signal 15 min after its creation.^[20] Additionally, SABRE hyperpolarisation of its ¹³C and ¹⁵N nuclei with in pyridazine has also been reported.^[21]

Here, we wish to report the effects of pyridazine functional group substitution on the SABRE derived ¹H NMR enhancements and T_1 relaxation lifetimes. Over 30 mono and disubstituted pyridazines are investigated that encompass a range of steric and electronic properties in this study. This allows us to draw conclusions as to what are the optimum substitution patterns, in addition to defining the functional group tolerance on the SABRE effect.

2 | RESULTS AND DISCUSSION

Our examination of the effects of pyridazine substitution began with the study of mono substituted substrates prior to building in complexity. The 3-Substituted pyridazines 2-7 shown in Table 1 were purchased or synthesised as described in the supporting information. NMR tubes equipped with a J. Young's Tap were filled with a solution containing [IrCl (COD)(IMes)] (5 mM), substrate (20 mM) in methanol- d_4 and exposed to H₂ (3 bar). After 1 h at room temperature, the samples were analysed by ¹H NMR spectroscopy for the formation of new hydride containing complexes. For comparison, we also recorded the ¹H NMR spectrum of a sample containing pyridazine 1, which showed a single hydride containing complex with a resonance at $\delta_{\rm H}$ –21.49 that is consistent with [Ir(H)₂(IMes)(pyridazine)₃]Cl as previously reported in the literature.^[18] The headspace of this sample is then replaced with 3 bar p-H₂ (>99% enrichment) and manually shaken in a 70 G polarisation transfer field, prior to rapid insertion into a 9.4 T NMR spectrometer and a ¹H NMR spectrum was immediately recorded (full experimental details are available in the supporting information). This ¹H NMR spectrum showed per proton signal enhancements of 316- and 472-fold for the ortho and meta resonances of pyridazine after SABRE polarisation transfer at 70 G after quantification. T_1 relaxation values of 29.1 s for the ortho (H3/6) resonance and 26.6 s for the meta (H4/5) resonance were observed for this sample.

Our first step in adding complexity to the pyridazine substrate was through the introduction of a methyl group adjacent to the nitrogen atom in order to investigate steric influence on SABRE hyperpolarisation. In analogous 2-picoline, the methyl group inhibits SABRE when using the [IrCl (COD)(IMes)] precatalyst^[22]; however, this has been recently overcome through the use of a bidentate PHOX ligand and up to 132-fold signal gains are reported.^[23] In contrast, 3-methylpyridazine **2**, performed

Substrate	Hydride resonance for [Ir(H)2(IMes)(sub)3]cl at 298 K/ppm	¹ H SABRE signal gain at 9.4 T (% polarisation)	<i>T</i> ¹ relaxation time at 298 K measured at 9.4 T/s
	-21.49	H3/6-472 ± 24 (1.5%)	H3/6-29.1
1 (N-N)		H4/5-316 ± 42 (1.0%)	H4/5-26.6
	-21.59	H4-758 ± 48 (2.4%)	H4-21.4
		H5-1161 ± 101 (3.6%)	Н5-23.0
2 — () / N-N		H6-1177 ± 76 (3.7%)	Н6-29.4
	-21.63	H4-648 ± 68 (2.0%)	H4-28.2
		H5-884 ± 87 (2.8%)	H5-24.5
3 MeO一\\// N-N		H6-759 ± 64 (2.4%)	H6-31.0
	-21.54	H4-99 ± 11 (0.3%)	H4-12.1
но —		H5-260 ± 32 (0.8%)	H5-16.3
4N_N		H6-273 ± 54 (0.9%)	Н6-17.6
	-21.52	H4-1266 ± 105 (4.0%)	H4-15.4
		H5-553 ± 86 (1.7%)	H5-20.8
5 CI—(\//) N—N		H6-936 ± 65 (2.9%)	Н6-27.9
	-21.42	H4-307 ± 31 (1.0%)	H4-22.2
MeO		H5-84 ± 14 (0.3%)	H5-13.6
6 O N-N		H6-252 ± 18 (0.8%)	H6-21.7
	$-20.68, -23.03 (J_{\rm HH} = 7.3 \text{ Hz})$	H4-3 ± 1 (0.01%)	H4-17.4
		H5-6 ± 2 (0.02%)	H5-25.8
7 NC-\/ N-N		H6-6 ± 1 (0.02%)	Н6-27.2
	_[a]	H3-613 ± 31 (1.9%)	Н3-27.2
Δ.		H5-354 ± 19 (1.1%)	H5-20.8
8		H6-541 ± 12 (1.7%)	H6-24.2
	_[a]	H3-564 ± 35 (1.8%)	H3-45.4
/		H5-374 ± 48 (1.2%)	H5-15.0
9 O		H6-652 ± 52 (2.0%)	Н6-17.4
	_[a]	H3-862 ± 65 (2.7%)	H3-57.4
OMe		H5-589 ± 71 (1.8%)	H5-18.4
10 O=		H6-778 ± 32 (2.4%)	H6-18.0

-WILEY 3

TABLE 1 (Continued)

Substrate	Hydride resonance for [Ir(H)2(IMes)(sub)3]cl at 298 K/ppm	¹ H SABRE signal gain at 9.4 T (% polarisation)	<i>T</i> ¹ relaxation time at 298 K measured at 9.4 T/s
	_[a]	H3-1204 ± 86 (3.8%)	H3-69.6
<i>d</i> ₃ -10 OCD ₃ N-N		H5-754 ± 75 (2.4%)	H5-12.2
		H6-897 ± 101 (2.8%)	H6-24.2

Note: Superscript [a] indicates a complex hydride region in the ¹H NMR spectrum that contains a minimum of five products.

well under SABRE conditions and the signal gains were improved when compared with **1**. Indeed, ¹H signal enhancements for the H4, H5 and H6 resonances were 758-, 1161- and 1177-fold, respectively. A 145-fold signal gain was also observed for the resonance of the methyl proton. Only a single hydride containing species could be detected under both PHIP and low temperature NMR studies with a hydride resonance at $\delta_{\rm H}$ –21.59. Therefore, we suggest that each of the three 2 ligands in $[Ir(H)_2(IMes)(2)_3]$ Cl bind through the N1 atom and other potential regioisomers either do not form or are present at too low concentrations to be detected. This is likely to be due to the steric influence of the methyl group which may hinder binding through the N2 atom. 3-Methoxypyridazine 3 also formed a single hydride containing complex and its hydride resonance shifted to $\delta_{\rm H}$ -21.63 in line with the increased electron donating capacity of this ligand. After SABRE polarisation transfer under analogous conditions, a 759-fold signal gain was observed for the H6 resonance. This is >2 times the analogous signal gain for 1, although it is a reduction in that observed relative to 2. Its T_1 relaxation times are comparable with that of **1**.

Further complexity was introduced by utilising the hydroxymethyl functionality in 4, and this resulted in a significantly reduced SABRE signal enhancements. Now, the maximum signal gain was 273-fold for the H6 resonance. Although the major hydride containing species remains that of $[Ir(H)_2(IMes)(4)_3]Cl (\delta_H - 21.54)$, an additional complex with broad hydride resonances at $\delta_{\rm H}$ -20.68 and -30.69 also appears, which could be due to bidentate binding of 4 in the equatorial plane through its N and O donor atoms. The broadness of these resonances indicates rapid hydride loss to form H₂, a process which would quickly consume the available $p-H_2$ thereby reducing the observed signal gains. Additionally, its T_1 values for H4, H5 and H6 were 12.1, 16.3 and 17.6 s, respectively, and therefore shorter than those for 3-methylpyridazine 2.

Next, we probed the effect of introducing electronwithdrawing substituents into the three position with substrates 5-7. 3-Chloropyridazine 5 forms a single hydride containing complex with a resonance $\delta_{\rm H}$ –21.52 and performs well under SABRE. A signal gain for its H4 proton of 1266-fold was observed, and its H5 and H6 resonances showed signal enhancements of 552- and 935-fold, respectively. T_1 values of 15.4, 20.8 and 27.9 s were also measured at 9.4 T. This indicates that the SABRE of monosubstituted pyridazines is tolerant to both electron-rich and electron-deficient ring systems. Methyl-3-pyridazinecarboxylate 6 did not yield as high signal enhancements as 5. Instead, a maximum of 307-fold signal gain was quantified for its H4 resonance. The dominant hydride containing complex gives a resonance at $\delta_{\rm H}$ –21.52; however, a minor product that has hydride resonances at $\delta_{\rm H}$ –20.12 and –29.42 also appears under PHIP conditions.

In contrast, 3-pyridazinecarbonitrile 7 does not form a dihydride-symmetric active catalyst and instead yields a product with inequivalent hydride resonances at $\delta_{\rm H}$ -20.68 and -23.03 that share a J-coupling of 7.3 Hz. As the binding of nitrile ligands to form an active SABRE catalyst is well established,^[24] we attribute these hydride resonances to the complex whereby one molecule of 7 is bound through the N1 atom and the other is bound through its nitrile group in the equatorial plane. Additionally, a minor product with hydride resonances at $\delta_{\rm H}$ –20.8 and –21.9 appears under PHIP conditions, which is likely to be due to the regioisomeric binding of 7 in the axial position. In this case, poor SABRE enhancement results and a maximum 6-fold ¹H NMR signal enhancement was quantified at 9.4 T. The T_1 values for 7 are comparable with that of pyridazine 1.

Next, we considered the effects of 4-substituted pyridazines **8–11**. 4-Methylpyridazine **8** showed good polarisation levels of 613-, 354- and 541-fold for its H3, H5 and H6 resonances, respectively. In contrast to 3-methylpyridazine **2**, 4-methylpyridazine **8** yields

multiple hydride-containing complexes on reaction with [IrCl (COD)(IMes)] in the presence of H₂. Cooling the sample to 243 K reveals >5 hydride containing complexes with resonances between $\delta_{\rm H}$ –21.25 and –21.42. Because the methyl group of 8 has a reduced steric influence on the ligation to the metal centre when compared with 2, it is now possible that the regioisomeric complexes, where binding can occur through either the substrates N1 or N2 atoms, lie much closer in energy than for the case of 2. Statistically, this would lead to the possibility of six

TABLE 2 Summary of SABRE hyperpolarisation and T_1 relaxation times of 3,6-disubstituted pyridazines

Substrate	Hydride resonance for [Ir (H)2(IMes)(sub)3]cl at 298 K	¹ H SABRE signal gain at 9.4 T (% polarisation)	<i>T</i> ₁ relaxation time at 298 K measured at 9.4 T
3,6-d₂-1 D-√N−N	-21.50	H4/5-492 ± 35 (1.5%)	H4/5-58.2
11 — — — — — — — — — — — — — — — — — —	-21.61	H4/5–129 ± 14 (0.4%)	H4/5–33.4
12 CI-CI	-24.45 ^[a]	_	H4/5-49.4
13 MeO-	_	_	H4/5-24.1
	-	_	H4-26.9
14 CI-			H5-18.2
	-	-	H4-30.9
15 CI-OMe			пэ-29.0
	_	-	H4-22.3
16 CI-CF ₃ -CF ₃			H3-20.0
	_[b]	H4-117 ± 21 (0.4%)	H4–16.2
17 $CI \longrightarrow NH_2$		H5-31 ± (0.1%)	H5–15.2
	_[b]	H4-12 ± 3 (0.03%)	H4–26.7
		H5-3 ± 1 (0.01%)	Н5–25.8

Note: Superscript [a] indicates chemical shift measured at 243 K. Superscript [b] indicates a complex hydride region in the ¹H NMR spectrum that contains multiple products.

regioisomers of $[Ir(H)_2(IMes)(8)_3]Cl$ being present in solution.

A complex hydride region is also present after formation of the active SABRE catalysts of 4-acetylpyridazine 9. Strong SABRE performance is still present and a signal gain of 564-fold for its H3 resonance is detected. Additionally, the T_1 value of the H3 resonance is increased to 45.7 s, which can be attributed to the lack of spin-spin routes to relaxation for this isolated proton. When a methyl ester is introduced into the 4-position to give 11 an improvement in signal gain is noted. The H3, H5 and H6 resonances of 11 exhibit signal gains of 862-, 589- and 778-fold, respectively. Additionally, a 57.4 s T_1 relaxation time is quantified for its H3 resonance, which is again isolated from other spins in the substrate due to the presence of the ester. Deuteration of the methyl group to form d_3 -11 significantly improved its SABRE polarisation levels. Now 1204-, 754- and 897-fold signal gains were quantified for the H3, H4 and H6 resonances, respectively. The T_1 relaxation values also increase, and the isolated H3 proton now has a value that is 69.6 s.

In order to increase the substrate complexity, a second substituent was added onto the pyridazine ring. First, symmetrical 3,6-substituted pyridazines 11-13 were investigated followed by unsymmetrical variants 14-18 as shown in Table 2. $3,6-d_2$ -Pyridazine (3,6-d_2-1) gave higher signal enhancements than its protio counterpart (1) with a 492-fold per proton signal gain for the remaining H4/5 position. Additionally, the T_1 relaxation time is also increased to 58.2 s from 26.6 s due to a reduction in spin-spin interactions within the substrate. 3,6-Dimethylpyridazine (11) presents an increased steric challenge when compared with its mono-substituted counterpart 2. Although an active SABRE complex forms with a hydride resonance at $\delta_{\rm H}$ –21.61, a significant reduction in polarisation transfer results. After SABRE transfer at 70 G, a 129-fold signal gain is observed for its H4/5 proton resonance which is ca. 50% of that which was quantified for 1. Similarly, 3,6-dichloropyridazine 12 also forms a SABRE complex; however, its hydride resonance only becomes visible at 243 K and gives a peak at $\delta_{\rm H}$ –24.45. At 298 K, a very broad hydride resonance is detected that indicates rapid ligand loss. Now, no ¹H SABRE polarisation is detected after polarisation transfer over a range of magnetic fields (0.5-120 G). This is in contrast to the reported ¹⁵N signal gain of 12 000-fold that is achieved at 8.5 T.^[21c] We suggest that the faster ligand dissociation is better matched for the strong ${}^{2}J_{HN}$ coupling between the substrate and hydride ligands in the active complex when compared with the weaker ${}^{5}J_{\rm HH}$ coupling.

No SABRE transfer is observed for 3,6dimethoxypyridazine (13) after SABRE polarisation transfer, and no evidence of any hydride containing complexes could be obtained at either low temperature or under PHIP conditions. Unsymmetrical substrates 14-16 also yielded no SABRE enhancements in their ¹H NMR spectra after polarisation transfer. This indicates that the steric influence of the 3,6-disubstituted substrates is not conducive to polarisation transfer. Finally, we were able to obtain weak SABRE enhancements for 3-chloro-6-aminopyridazine 17 and 3-chloro-6-cyanopyridazine 18. In both cases, multiple hydride resonances are present in the ¹H NMR spectrum, which reflects their multiple binding modalities. 3-Chloro-6-aminopyridazine 17 showed SABRE derived signal gains of 112-fold for the H5 signal and 61-fold for its H4 resonance. Its ¹H NMR spectrum reveals the presence of two dominant hydride containing complexes: first, a symmetric complex with a hydride resonance at $\delta_{\rm H}$ –22.1 and second with broad resonances at $\delta_{\rm H}$ –21.8 and –29.7. These complexes are formed in a 1:5 ratio and can be accounted for by binding through both the pyridazine ring nitrogen and also the aromatic amine.^[25] For **18**, the hydride signals are much more complex with ca. 4 hydride containing complexes being formed, which may arise due to the ability for binding through both the ring nitrogen and nitrile group. The SABRE-derived polarisation transfer to 18 was significantly lower, and a maximum 12-fold signal gain was quantified.

Overall, the signal enhancements for the 3,6disubstituted pyridazines are significantly lower than those of the monosubstituted pyridazines. This is like to be due to a combination of effects. Firstly, the occupation of the *ortho* position increases the steric demand around the metal centre, which will either inhibit the formation of the active SABRE complex or significantly reduce the lifetime of the active catalyst. Secondly, as the *ortho* sites are occupied, the SABRE process utilises the weaker ⁵J coupling from the hydride to the *meta* proton which may lead to less efficacious polarisation transfer.

The investigation was then extended to 4,5disubstituted pyridazines which were synthesised by the well-established^[26] reverse electron demand Diels-Alder reaction as detailed in the supporting information. Exposure of a methanol- d_4 solution containing [IrCl(COD) (IMes)] (5 mM), substrate **19–29** (20 mM) to H_2 (3 bar) led to the formation of single hydride containing complexes in all cases with their resonances appearing between $\delta_{\rm H}$ –20.89 and –22.12 for each of the substrates. We attribute these resonances to the respective product of type [Ir(H)₂(IMes)(sub)₃)]Cl.^[18] When the sample was exposed to p-H₂ (3 bar) in a ca. 70 G polarisation transfer field, significant improvements in their ¹H NMR signal enhancements were observed when compared with the corresponding thermally polarised spectra.

Hydride resonance for [Ir(H)₂(IMes)(sub)₃]cl ¹H SABRE signal gain T_1 relaxation time Substrate at 298 K at 9.4 T (% polarisation) at 298 K measured at 9.4 T -21.59 H3/6-1871 ± 201 (5.8%) H3/6-85.8 $\langle \rangle = \langle \rangle$ 4,5-*d*₂-1 -22.12 H3/6-998 ± 78 (3.1%) H3/6-7.1 OMe 19 -21.00H3/6-128 ± 18 (0.4%) H3/6-8.2 Me₃Si SiMe₃ 20 -21.15H3/6-823 ± (2.6%) H3/6-13.8 21 -21.15 H3/6-1365 ± 145 (4.3%) H3/6-33.2 d₁₀-21 -21.03 H3/6-878 ± 76 (2.7%) H3/6-84.3 22 -20.89H3/6-1647 ± 79 (5.1%) H3/6-120.8 DМе 23 H3/6-83.4 -20.84 H3/6-1134 ± 124 (3.5%) OEt 24

TABLE 3 Summary of SABRE hyperpolarisation and T_1 relaxation times of 4,5-disubstituted pyridazines

WILEY 7

(Continues)

TABLE 3 (Continued)

WILEY.



4,5- d_2 -Pyridazine 4,5- d_2 -1 gave a 3480-fold per proton signal enhancement for its ortho resonance. Additionally, a T_1 relaxation time of 85.8 s, which is longer than that of the corresponding $3,6-d_2$ -pyridazine $3,6-d_2-1$, however, is shorter than that of $3,5-d_2$ -pyridazine isotopologue which has been reported previously.^[19] Introduction of electron donating methoxy groups (19) into the 4,5-positions gave signal enhancements of 998-fold at 9.4 T for the ortho protons of **19**. Disappointingly, its T_1 relation time significantly shortens to just 7.1 s. We propose this could be due to intramolecular dipole-dipole coupling between protons. Trimethylsilyl derivative 20 shows significantly reduced SABRE activity and just a 129-fold ¹H NMR signal gain was quantified at 9.4 T. Polarisation transfer was also attempted into its ²⁹Si nuclei; however, no signal gain was observed after polarisation transfer at a range of fields from 0.5 G to 120 G. This is in contrast to other silyl-derived substrates that showed SABRE polarisation transfer to ²⁹Si is possible via the adjacent ¹H nuclei at 25 G.^[27] The ¹H T_1 value for the *ortho* resonances was measured to be just 11.1 s. Comparatively, 4,5-Diphenylpyridazine **21** showed good SABRE transfer, and a 823-fold signal gain was quantified at 9.4 T. However, its T_1 value was just 13.8 s. Synthesis of its selectively deuterated isotopologue yielded an improvement in both SABRE polarisation transfer and the T_1 value of the remaining proton resonance. Indeed, d_{10} -**21** gave a 1365-fold signal gain at 9.4 T in conjunction with a relaxation time of 33.2 s were quantified.

When electronegative atoms are placed in the 4,5-positions, good SABRE enhancement is still evident. Thus, when 4,5-dichloropyridazine **22** reduced the signal gain was 878-fold and its T_1 value increased to 84.3 s. Other electron-withdrawing groups performed better

8

with dimethyl ester **23** yielding a 1647-fold signal gain for its *ortho* resonance, and the relaxation time was to 120.1 s. This is equivalent to 5.1% polarisation at 9.4 T.

As this diester derivative showed strong polarisation transfer and long magnetic lifetimes, we wished to further probe the effect of the ester moiety on the SABRE effect. Therefore, diethyl ester 24 was synthesised and subject to analogous SABRE testing. After interrogation at 9.4 T a 1134-fold signal enhancement resulted. This is a reduction when compared with dimethyl ester 23 and a reduction in T_1 value from 120.8 to 83.4 s is also noted. Introduction of isopropyl esters (25) further reduces both the SABRE signal gains and the T_1 values. These T_1 effects can be quantified by working out an enhancement value at the point of creation of the hyperpolarised state. Assuming this T_1 value remains constant for 5 s during the moving into the measurement field where the polarisation level is quantified, we estimate that initial polarisation levels of ca. 1700-, 1250- and 500-fold for 23, 24 and 25 respectively result. This indicates that there is a steric effect taking place during the SABRE process even in the remote 4,5-positions. Breaking the symmetry of the molecule through the use of one methyl ester and one ethyl ester in 26 returns the substrate to high average SABRE enhancements of 1256. The individual enhancements for H3 and H6 could not be determined due to peak overlap. The average T_1 on the remaining proton sites is now 94.6 s.

The introduction of amide groups into the 4,5-positions reduces the SABRE performance of the pyridazine molecules 27-29 when compared with dimethyl ester 23. Additionally, the T_1 values for the amide derivatives are all significantly smaller than the ester counterparts, which may reduce the detected polarisation level. A similar T_1 analysis reveals that at the point of creation, the polarisation levels for 27, 28 and 29 would be ca. 500-, 500- and 550-fold, respectively. These values are all comparable with one another and yet three times smaller than those calculated for dimethyl ester 23. Such an effect is likely to be due to an increase in the ligand exchange rate relative to that of the ester derivate and away from that which has been predicted to be optimal by Barskiy.^[28] Given the previously reported studies on SABRE as a function of electronic and steric NHC impact, such behaviour is not unexpected.^[15] Analysis of the reported values for nicotinamide and methyl nicotinate reveals a similar 3-fold reduction in signal enhancement where electronic effects are clearly dominant.^[19]

With the success of SABRE hyperpolarisation of dimethyl ester **23**, we began to explore methods to further optimise its hyperpolarisation and explore the effects of other carbonyl containing substrates. First, we investigated the effect of increasing the concentration of **23** on the resulting SABRE signal gains (Figure 1). We found



FIGURE 1 Optimisation of SABRE hyperpolarisation of **23**. (a) Effect of increasing the equivalents of substrate on the SABRE hyperpolarisation of **23**. (b) Effect of changing the *N*-heterocyclic carbene (NHC) ligand in the precatalyst of type [IrCl (COD)(NHC)] on the SABRE hyperpolarisation of **23**. (c) Structures of NHC ligands used in the study



FIGURE 2 Single scan ¹H NMR spectra of d_6 -23 under (top) thermally polarised conditions that has been expanded 512× in the vertical direction and (bottom) after SABRE polarisation transfer at 70 G using [IrCl (COD)(IMes)] (5 mM), d_6 -23 (20 mM) in methanol- d_4

that four equivalents of **23** was optimal and increasing beyond this catalyst loading led to a reduction in polarisation efficacy. This is consistent with the effects of diluting the finite available polarisation from p-H₂ through an increased number of substrate spins.

The nature of the N-heterocyclic carbene (NHC) in the SABRE precatalyst can have a profound effect on the signal enhancement.^[15b,29] Therefore, we screened a series of precatalysts of type [IrCl(COD)(NHC)], which have been shown to modulate the rate of ligand loss through increased steric effects.^[29b] When using [IrCl (COD)(SIMes)] as the precatalyst an increased signal enhancement of 3788 ± 132 -fold was observed for 23, which is over two times larger than that seen with [IrCl (COD)(IMes)]. However, further utilisation of the larger IPr ligand led to a reduction in signal enhancement to 1542 ± 154 -fold. Therefore, we can conclude that the optimal conditions for the polarisation of 23 are [IrCl (COD)(SIMes)] (5 mM), 23 (20 mM) in methanol- d_4 , and these conditions would be applied to further studies on the 4,5-dicarbonyl pyridazines.

We began with the synthesis of its isotopologue, d_6 -23, with a view to extending its relaxation lifetime and increasing its SABRE efficacy. After subjection of d_6 -23 to the optimised SABRE conditions ([IrCl(COD) (SIMes)] (5 mM), d_6 -23 (20 mM) in methanol- d_4), a 9177-fold per proton signal gain was observed for its

ortho resonances at 9.4 T which is equivalent to 28.7% polarisation (Figure 2). Additionally, a T_1 relaxation time of 138.4 s was recorded.

3 | CONCLUSIONS

In summary, we present a general method for the SABRE hyperpolarisation of over 30 substituted pyridazines and signal enhancements of up to 9177-fold at 9.4 T. This is equivalent to 28.7% polarisation and shows the potential of this method to dramatically improve signal strength in magnetic resonance techniques. Monosubstituted pyridazines perform well and yield good signal enhancements. The steric effect of adding groups into the 3-position is minimised through the substrates' ability to bind through the N1 atom. The method is able to tolerate not only electron-rich pyridazines such as 3-methoxypyridazine 3 but also electron poor substrates such as 3-chloropyridazine 4. Similarly, 4-substituted pyridazines perform equally well. Interestingly, when the substrate contains the possibility for multiple binding modalities, through additional groups such as alcohols or nitriles, the active catalyst differs from that of the typical symmetric [Ir(H)₂(IMes)(sub)₃]Cl. For example, 3-cyanopyridazine 7 formed a catalyst where one of the equatorial ligands binds through the N1 atom and the

second ligates via the nitrile moiety. This results in lower SABRE signal enhancements when compared with pyridazine **1**.

The hyperpolarisation of pyridazines becomes more sensitive when increased steric bulk is placed into the 3- and 6-positions on the aromatic ring. Indeed, when 3,6-dimethylpyridazine **11** is utilised, a 129-fold signal gain is observed which is a significant reduction when compared with 3-methylpyridazine **2** (up to 1177-fold signal gain). Increasing the steric bulk further leads to no signal enhancements being observed at 298 K, and no evidence of any hydride containing species could be detected. This suggests that the NHC-derived catalysts commonly utilised for SABRE cannot form under the conditions used here, and therefore, the exploration of other catalysts could be fruitful.^[23,30]

In contrast, 4,5-disubstituted pyridazines all form the expected symmetric $[Ir(H)_2(IMes)(sub)_3]Cl$ catalyst and thus undergo efficacious SABRE. The electronic effect of changing the functionality does not appear to significantly impact the resulting signal enhancements after polarisation transfer. For example, 4,5-dimethoxypyridazine **19** shows an excellent per proton signal gain for its H3/H6 resonance of 998-fold, whereas for 4,5-dichloropyridazine, **22** yielded a signal gain of 878-fold. The best performing substrate was dimethylester **23** which gave a signal gain of 1647-fold in conjunction with a T_1 relaxation time of over 2 min at 9.4 T.

Finally, we exemplify the possibility to further increase the SABRE transfer to substituted pyridazines by optimisation of the polarisation level of **23**. We found that changing the precatalyst to [IrCl(COD)(SIMes)] a higher signal gain could be achieved and deuteration of the methyl esters to give d_6 -23 ultimately yielded a polarisation level of >28%. Deuteration of the methyl esters also extended the T_1 relaxation time to 138 s. The data reported here will allow for the development of methods to hyperpolarise and detect a range of pyridazine containing pharmaceutical and agrochemicals such as the antidepressant minaprine or anticonvulent SR41378.^[3] Next stages will see us consider the effect of fused ring systems and seek to overcome the challenges posed by the steric influence of 3,6-disubstituted systems.

ACKNOWLEDGMENTS

The authors would like to acknowledge the following for supporting this work: The Wellcome Trust (grants 092506 and 098335), the EPSRC (EP/R51181X/1) and the University of York.

PEER REVIEW

The peer review history for this article is available at https://publons.com/publon/10.1002/mrc.5152.

ORCID

Peter J. Rayner https://orcid.org/0000-0002-6577-4117 Michael J. Burns https://orcid.org/0000-0002-5569-1270 Elizabeth J. Fear https://orcid.org/0000-0002-0578-900X

Simon B. Duckett ^b https://orcid.org/0000-0002-9788-6615

REFERENCES

- a] C. Lamberth, *J.Heterocycl. Chem.* 2017, 54, 2974. b] I. Mohd,
 A. Mohammad, *J. Bioorg. Chem.* 2020, 46, 745. c] T. J. Ritchie,
 S. J. F. Macdonald, S. Peace, S. D. Pickett, C. N. Luscombe,
 MedChemComm 2012, (3), 1062.
- [2] C. Hamdouchi, C. Sanchez-Martinez, J. Gruber, M. del Prado, J. Lopez, A. Rubio, B. A. Heinz, J. Med. Chem. 2003, 46, 4333.
- [3] C. G. Wermuth, MedChemComm 2011, 2, 935.
- [4] B. M. Fox, K. Iio, K. Li, R. Choi, T. Inaba, S. Jackson, S. Sagawa, B. Shan, M. Tanaka, A. Yoshida, F. Kayser, *Bioorg. Med. Chem. Lett.* 2010, 20, 6030.
- [5] B. Dyck, S. Markison, L. Zhao, J. Tamiya, J. Grey, M. W. Rowbottom, M. Zhang, T. Vickers, K. Sorensen, C. Norton, J. Wen, C. E. Heise, J. Saunders, P. Conlon, A. Madan, D. Schwarz, V. S. Goodfellow, *J. Med. Chem.* **2006**, *49*, 3753.
- [6] a] K. V. Kovtunov, E. V. Pokochueva, O. G. Salnikov, S. F. Cousin, D. Kurzbach, B. Vuichoud, S. Jannin, E. Y. Chekmenev, B. M. Goodson, D. A. Barskiy, I. V. Koptyug, *Chem. Asian J.* **2018**, *13*, 1857. b] P. Nikolaou, B. M. Goodson, E. Y. Chekmenev, *Chemistry* **2015**, *21*, 3156.
- [7] a] S. A. Colebrooke, S. B. Duckett, J. A. B. Lohman, R. Eisenberg, *Chem. Eur. J.* 2004, 10, 2459. b] C. Godard, S. B. Duckett, S. Polas, R. Tooze, A. C. Whitwood, *Dalton Trans.* 2009, 38, 2496. c] C. Godard, J. Lopez-Serrano, M. D. Galvez-Lopez, M. Rosello-Merino, S. B. Duckett, I. Khazal, A. Lledos, A. C. Whitwood, *Magn. Reson. Chem.* 2008, 46, S107. d] D. J. Fox, S. B. Duckett, C. Flaschenriem, W. W. Brennessel, J. Schneider, A. Gunay, R. Eisenberg, *Inorg. Chem.* 2006, 45, 7197. e] J. López-Serrano, S. B. Duckett, A. Lledós, *J. Am. Chem. Soc.* 2006, 128, 9596. f] I. Atheaux, F. Delpech, B. Donnadieu, S. Sabo-Etienne, B. Chaudret, K. Hussein, J.-C. Barthelat, T. Braun, S. B. Duckett, R. N. Perutz, *Organometallics* 2002, 21, 5347. g] P. D. Morran, S. A. Colebrooke, S. B. Duckett, J. A. B. Lohman, R. Eisenberg, *J. Chem. Soc., Dalton Trans.* 1998, 22, 3363.
- [8] a] P. M. Richardson, W. Iali, S. S. Roy, P. J. Rayner, M. E. Halse, S. B. Duckett, *Chem. Sci.* 2019, *10*, 10607. b] N. Eshuis, B. J. A. van Weerdenburg, M. C. Feiters, F. P. J. T. Rutjes, S. S. Wijmenga, M. Tessari, *Angew. Chem., Int. Ed.* 2015, *54*, 1481. c] L. S. Lloyd, R. W. Adams, M. Bernstein, S. Coombes, S. B. Duckett, G. G. R. Green, R. J. Lewis, R. E. Mewis, C. J. Sleigh, *J. Am. Chem. Soc.* 2012, *134*, 12904.
- [9] S. Fain, M. L. Schiebler, D. G. McCormack, G. Parraga, J. Magn. Reson. Imaging 2010, 32, 1398.
- [10] a] E. Cavallari, C. Carrera, M. Sorge, G. Bonne, A. Muchir, S. Aime, F. Reineri, *Sci. Rep.* **2018**, *8*, 8366. b] H. Gutte, A. E. Hansen, H. H. Johannesen, A. E. Clemmensen, J. H. Ardenkjar-Larsen, C. H. Nielsen, A. Kjar, *Am J Nucl Med Mol Imaging* **2015**, *5*, 548. c] T. Harris, H. Degani, L. Frydman,

12 \perp Wiley-

NMR Biomed. 2013, 26, 1831. d] T. H. Witney, M. I. Kettunen, K. M. Brindle, J. Biol. Chem. 2011, 286, 24572. e] C. von Morze, P. E. Z. Larson, S. Hu, K. Keshari, D. M. Wilson, J. H. Ardenkjaer-Larsen, A. Goga, R. Bok, J. Kurhanewicz, D. B. Vigneron, J. Magn. Reson. Imaging 2011, 33, 692. f] Y.-F. Yen, S. J. Kohler, A. P. Chen, J. Tropp, R. Bok, J. Wolber, M. J. Albers, K. A. Gram, M. L. Zierhut, I. Park, V. Zhang, S. Hu, S. J. Nelson, D. B. Vigneron, J. Kurhanewicz, H. A. A. M. Dirven, R. E. Hurd, Magn. Res. Med. 2009, 62, 1.

- [11] a] T. G. Walker, W. Happer, Rev. Mod. Phys. 1997, 69, 629. b] S. Appelt, A. B.-A. Baranga, C. Erickson, M. Romalis, A. Young, W. Happer, Phys. Rev. a 1998, 58, 1412.
- [12] a] G. Zhang, C. Hilty, Magn. Reson. Chem. 2018, 56, 566. b] T. Maly, G. T. Debelouchina, V. S. Bajaj, K.-N. Hu, C.-G. Joo, M. L. Mak-Jurkauskas, J. R. Sirigiri, P. C. van der Wel, J. Herzfeld, R. J. Temkin, J. Chem. Phys. 2008, 128, 02B611. c] K. R. Keshari, D. M. Wilson, Chem. Soc. Rev. 2014, 43, 1627.
- [13] a] C. R. Bowers, D. P. Weitekamp, J. Am, Chem. Soc. 1987, 109, 5541. [b]C. R. Bowers, D. P. Weitekamp, Phys. Rev. Lett. 1986, 57, 2645.
- [14] R. W. Adams, J. A. Aguilar, K. D. Atkinson, M. J. Cowley, P. I. P. Elliott, S. B. Duckett, G. G. R. Green, I. G. Khazal, J. López-Serrano, D. C. Williamson, Science 2009, 323, 1708.
- [15] a] M. J. Cowley, R. W. Adams, K. D. Atkinson, M. C. R. Cockett, S. B. Duckett, G. G. R. Green, J. A. B. Lohman, R. Kerssebaum, D. Kilgour, R. E. Mewis, J. Am, Chem. Soc. 2011, 133, 6134. b] P. J. Rayner, P. Norcott, K. M. Appleby, W. Iali, R. O. John, S. J. Hart, A. C. Whitwood, S. B. Duckett, Nat. Commun. 2018, 9, 4251.
- [16] R. W. Adams, S. B. Duckett, R. A. Green, D. C. Williamson, G. G. R. Green, J. Chem. Phys. 2009, 131, 194505.
- [17] a] P. J. Rayner, S. Duckett, Angew. Chem., Int. Ed. 2018, 57, 6742. b] J.-B. Hövener, A. N. Pravdivtsev, B. Kidd, C. R. Bowers, S. Glöggler, K. V. Kovtunov, M. Plaumann, R. Katz-Brull, K. Buckenmaier, A. Jerschow, F. Reineri, T. Theis, R. V. Shchepin, S. Wagner, P. Bhattacharya, N. M. Zacharias, E. Y. Chekmenev, Angew. Chem., Int. Ed. 2018, 57, 11140.
- [18] K. M. Appleby, R. E. Mewis, A. M. Olaru, G. G. R. Green, I. J. S. Fairlamb, S. B. Duckett, Chem. Sci. 2015, 6, 3981.
- [19] P. J. Ravner, M. J. Burns, A. M. Olaru, P. Norcott, M. Fekete, G. G. R. Green, L. A. R. Highton, R. E. Mewis, S. B. Duckett, Proc. Natl. Acad. Sci. U. S. A. 2017, 114, E3188.
- [20] S. S. Roy, P. Norcott, P. J. Rayner, G. G. R. Green, S. B. Duckett, Angew. Chem. Int. Ed. 2016, 128, 15871.
- [21] a] S. S. Roy, P. Norcott, P. J. Rayner, G. G. R. Green, S. B. Duckett, Chem. - Eur. J. 2017, 23, 10496. b] S. S. Roy, P. J. Rayner, M. J. Burns, S. B. Duckett, J. Chem. Phys. 2020, 152, 014201. c] J. F. P. Colell, A. W. J. Logan, Z. Zhou, R. V. Shchepin, D. A. Barskiy, G. X. Ortiz, Q. Wang, S. J. Malcolmson, E. Y. Chekmenev, W. S. Warren, T. Theis, J. Phys. Chem. C. 2017, 121, 6626.

- [22] R. V. Shchepin, M. L. Truong, T. Theis, A. M. Coffey, F. Shi, K. W. Waddell, W. S. Warren, B. M. Goodson, E. Y. Chekmenev, J. Phys. Chem. Lett. 2015, 6, 1961.
- [23] J. Colell, A. W. J. Logan, Z. Zhou, J. R. Lindale, R. Laasner, R. Shchepin, E. Chekmenev, V. Blum, W. S. Warren, S. J. Malcolmson, T. Theis, Chem. Commun. 2020, 56, 9336.
- [24] R. E. Mewis, R. A. Green, M. C. R. Cockett, M. J. Cowley, S. B. Duckett, G. G. R. Green, R. O. John, P. J. Rayner, D. C. Williamson, J. Phys, Chem. B 2015, 119, 1416.
- [25] W. Iali, P. J. Rayner, A. Alshehri, A. J. Holmes, A. J. Ruddlesden, S. B. Duckett, Chem. Sci. 2018, 9, 3677.
- [26] a] D. K. Heldmann, J. Sauer, Tetrahedron Lett. 1997, 38, 5791. b] J. Cioslowski, J. Sauer, J. Hetzenegger, T. Karcher, T. Hierstetter, J. Am. Chem. Soc. 1993, 115, 1353. c] M. D. Helm, J. E. Moore, A. Plant, J. P. A. Harrity, Angew. Chem., Int. Ed. 2005, 44, 3889.
- [27] A. M. Olaru, A. Burt, P. J. Rayner, S. J. Hart, A. C. Whitwood, G. G. R. Green, S. B. Duckett, Chem. Commun. 2016, 52, 14482.
- [28] D. A. Barskiy, A. N. Pravdivtsev, K. L. Ivanov, K. V. Kovtunov, I. V. Koptyug, Phys. Chem. Chem. Phys. 2016, 18, 89.
- [29] a] B. J. A. van Weerdenburg, S. Gloggler, N. Eshuis, A. H. J. Engwerda, J. M. M. Smits, R. de Gelder, S. Appelt, S. S. Wymenga, M. Tessari, M. C. Feiters, B. Blumich, F. P. J. T. Rutjes, Chem. Commun. 2013, 49, 7388. b] L. S. Lloyd, A. Asghar, M. J. Burns, A. Charlton, S. Coombes, M. J. Cowley, G. J. Dear, S. B. Duckett, G. R. Genov, G. G. R. Green, L. A. R. Highton, A. J. J. Hooper, M. Khan, I. G. Khazal, R. J. Lewis, R. E. Mewis, A. D. Roberts, A. J. Ruddlesden, Catal. Sci. Technol. 2014, 4, 3544.
- [30] a] W. Iali, G. G. R. Green, S. J. Hart, A. C. Whitwood, S. B. Duckett, Inorg. Chem. 2016, 55, 11639. b] A. J. Holmes, P. J. Rayner, M. J. Cowley, G. G. R. Green, A. C. Whitwood, S. B. Duckett, Dalton Trans. 2015, 44, 1077.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Rayner PJ, Burns MJ, Fear EJ, Duckett SB. Steric and electronic effects on the ¹H hyperpolarisation of substituted pyridazines by signal amplification by reversible exchange. Magn Reson Chem. 2021;1-12. https:// doi.org/10.1002/mrc.5152