

This is a repository copy of *The Syndemic Challenge of Tuberculosis and Tobacco Use*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/172527/>

Version: Published Version

Article:

Siddiqi, Kamran orcid.org/0000-0003-1529-7778 and Novotny, Thomas E (2021) The Syndemic Challenge of Tuberculosis and Tobacco Use. Tobacco Induced Diseases. 133575. ISSN: 1617-9625

<https://doi.org/10.18332/tid/133575>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

The syndemic challenge of tuberculosis and tobacco use

Kamran Siddiqi¹, Thomas E. Novotny²

The term ‘syndemic’ describes linked health problems involving two or more conditions that act synergistically, causing an excess burden of disease in the population. These interacting conditions often cluster similarly by person, place, and time. Most importantly, interventions against them must address the underlying conditions that bind the problems together¹. The global incidence of tuberculosis (TB) has declined over the last few decades, albeit far slower than envisioned by the World Health Organization (WHO). For 2018–2019, TB incidence only declined by 2.3%, suggesting that WHO objectives for 2030 will not be met². Ten million persons were infected in 2019, but the burden of TB disease is highest in low- and middle-income countries (LMICs), where 95% of TB deaths occur. The high TB burden countries include India, Indonesia, China, Philippines, and Bangladesh, and these nations also rank among the top ten countries for daily smoking prevalence³. Poverty, poor nutrition, and lack of comprehensive healthcare systems in these nations play into the TB and smoking syndemic.

Smoking and secondhand smoke exposure are two of several conditions that exacerbate adverse TB outcomes such as recurrent disease, excess mortality, and treatment failure. These conditions include diabetes, poor nutrition, alcohol use, and HIV infection. Because there are over one billion smokers globally, it is not surprising that 17.6% (95% CI: 8.4–21.4) of new cases and 15.2% (95% CI: 1.8–31.9) of TB deaths are attributable to smoking in high burden countries, regardless of other risk factors⁴.

Given the syndemicity of TB and tobacco use, we might ask what would happen if the smoking prevalence among those infected with TB could be effectively reduced through consistent, integrated treatment for tobacco use. Unfortunately, and despite a higher prevalence of smoking in TB patients compared to the general population⁵, the vast majority of TB patients are neither routinely asked about their smoking status nor advised to quit⁶.

Some progress has been made to meet the syndemic challenge of TB and smoking. There has been increased interest in policies to help TB patients quit smoking⁷ as well as recognition of the need for evidence-based smoking cessation interventions⁸. The WHO and the Union were first to embrace this challenge in the 2007 *Monograph on TB and Tobacco Control*, which called for further research and the application of cessation assistance in TB programs⁹. There are also a few large randomized-controlled trials (RCTs)^{8,10} conducted in high-TB burden countries, which highlight that face-to-face behavioral interventions can achieve high quit rates among TB patients⁸. Those who quit smoking were shown to have better overall clinical outcomes. In these studies, TB staff were able to deliver behavioral interventions for smoking cessation. However, because of several health system barriers¹¹ (e.g. cost, reach, sustainability), no high-TB burden country has so far integrated consistent face-to-face behavioral interventions for smoking cessation within its TB services. Recognizing the challenges of integrating and scaling

AFFILIATION

¹ Department of Health Sciences, University of York, York, United Kingdom

² School of Public Health, San Diego State University, San Diego, United States

CORRESPONDENCE TO

Kamran Siddiqi. Department of Health Sciences, University of York, York, United Kingdom. E-mail: Kamran.siddiqi@york.ac.uk
ORCID ID: <https://orcid.org/0000-0003-1529-7778>

KEYWORDS

syndemic, tuberculosis, tobacco

Received: 15 February 2021

Accepted: 21 February 2021

up face-to-face interventions, WHO has developed a smoking cessation package (mTB-Tobacco) that can be delivered as mobile phone messages to TB patients¹².

The evidence so far points towards a strong syndemic association between adverse TB outcomes and tobacco use. The evidence is also emerging in support of face-to-face behavioral interventions that can be delivered within health services in high-TB burden countries. On the upcoming World TB Day, we want to emphasize the important contribution of tobacco use in sustaining the TB disease burden and to appeal to policymakers and practitioners in high-TB burden countries that they recognize tobacco cessation as an important part of our efforts to end the global TB epidemic.

REFERENCES

1. Merrill S. Introducing Syndemics: A Critical Systems Approach to Public and Community Health. San Francisco: Jossey-Bass. 2009;304.
2. World Health Organization. [Global tuberculosis report 2020: executive summary]. World Health Organization; 2020. Accessed February 15, 2021. <https://apps.who.int/iris/bitstream/handle/10665/337538/9789240016941-ara.pdf>
3. Drope J, Schluger N, Cahn Z, et al. The Tobacco Atlas. American Cancer Society, Vital Strategies; 2018:21-22. Accessed February 15, 2021. https://tobaccoatlas.org/wp-content/uploads/2018/03/TobaccoAtlas_6thEdition_LoRes_Rev0318.pdf
4. Amere GA, Nayak P, Salindri AD, Narayan KMV, Magee MJ. Contribution of Smoking to Tuberculosis Incidence and Mortality in High-Tuberculosis-Burden Countries. *Am J Epidemiol*. 2018;187:1846-1855. doi:10.1093/aje/kwy081
5. Marshall AM, Barua D, Mitchell A, et al. Smoking prevalence among tuberculosis patients: A crosssectional study in Bangladesh and Pakistan. *Tob Induc Dis*. 2020;18:70. doi:10.18332/tid/125452
6. Warsi S, Elsey H, Boeckmann M, et al. Using behaviour change theory to train health workers on tobacco cessation support for tuberculosis patients: a mixed-methods study in Bangladesh, Nepal and Pakistan. *BMC Health Serv Res*. 2019;19(1):71. doi:10.1186/s12913-019-3909-4
7. United Nations Development Programme. Integrating Tobacco Control into Tuberculosis and HIV Responses. September 25, 2018. Accessed September 26, 2019. <https://www.undp.org/content/undp/en/home/librarypage/hiv-aids/integrating-tobacco-control-into-tuberculosis-and-hiv-responses.html>
8. Siddiqi K, Khan A, Ahmad M, et al. Action to Stop Smoking In Suspected Tuberculosis (ASSIST) in Pakistan: a Cluster Randomized, Controlled Trial. *Ann Intern Med*. 2013;158(9):667-675. doi:10.7326/0003-4819-158-9-201305070-00006
9. World Health Organization, International Union Against Tuberculosis and Lung Disease. A WHO/The Union monograph on TB and Tobacco Control: Joining efforts to control two related global epidemics. World Health Organization; 2007. Accessed February 15, 2021. https://apps.who.int/iris/bitstream/handle/10665/43812/9789241596220_eng.pdf?sequence=1&isAllowed=y
10. Dogar OF, Keding A, Gabe R, et al. Cytisine for smoking cessation in patients with tuberculosis: a multicentre, randomised, double-blind, placebo-controlled phase 3 trial. *Lancet Public Health*. 2020;8(11). doi:10.1016/S2214-109X(20)30312-0
11. Dogar O, Elsey H, Khanal S, Siddiqi K. Challenges of Integrating Tobacco Cessation Interventions in TB Programmes: Case Studies from Nepal and Pakistan. *J Smok Cessat*. 2016;11(2):108-115. doi:10.1017/jsc.2015.20
12. World Health Organization, International Telecommunication Union. Be He@lthy, Be Mobile: A handbook on how to implement mTB-Tobacco. World Health Organization, International Telecommunication Union; 2019. Accessed February 15, 2021. <https://apps.who.int/iris/bitstream/handle/10665/325243/9789241515900-eng.pdf>

CONFLICTS OF INTEREST

The authors have each completed and submitted an ICMJE form for disclosure of potential conflicts of interest. The authors declare that they have no competing interests, financial or otherwise, related to the current work. K. Siddiqi reports grants from European Commission, and non-financial support from Aflofarm, during the conduct of the study.

FUNDING

There was no source of funding for this research.

PROVENANCE AND PEER REVIEW

Commissioned; internally peer reviewed.