

This is a repository copy of *Sub-Saharan Africa preparedness and response to the COVID-19 pandemic: A perspective of early career African scientists*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/172749/>

Version: Published Version

Article:

Umviligihozo, Gisele, Mupfumi, Lucy, Sonela, Nelson et al. (10 more authors) (2020) Sub-Saharan Africa preparedness and response to the COVID-19 pandemic: A perspective of early career African scientists. Wellcome Open Research. 163. ISSN 2398-502X

<https://doi.org/10.12688/wellcomeopenres.16070.3>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



OPEN LETTER

REVISED Sub-Saharan Africa preparedness and response to the COVID-19 pandemic: A perspective of early career African scientists [version 3; peer review: 2 approved]

Gisele Umviligihozo¹, Lucy Mupfumi², Nelson Sonela^{3,4}, Delon Naicker⁵, Ekwaro A. Obuku⁶, Catherine Koofhethile⁷, Tuelo Mogashoa⁸, Anne Kapaata⁹, Geoffrey Ombati⁹, Clive M. Michelo¹⁰, Kimani Makobu¹¹, Olamide Todowede^{12,13}, Sheila N. Balinda⁸

¹Faculty of Health Sciences, Simon Fraser University, Burnaby, British Columbia, V5A 1S6, Canada

²Botswana Harvard AIDS Institute Partnership (BHIP), Private Bag BO 320, Bontleng, Gaborone, Botswana

³Chantal Biya International Reference Center for research on the management and prevention of HIV/AIDS (CIRCB), Yaoundé, B.P.: 3077, Cameroon

⁴School of Medicine, Physical and Natural Sciences, University of Rome Tor Vergata, Rome, 1-00133, Italy

⁵HIV Pathogenesis Programme, The Doris Duke Medical Research Institute, Nelson R. Mandela, School of Medicine, University of KwaZulu-Natal, Durban, 4001, South Africa

⁶CoVID Pandemic Rapid Evidence Synthesis Group (CoVPRES), Africa Centre for Systematic Reviews and Knowledge Translation, College of Health Sciences, Makerere University, Kampala, P.O. Box 7072, Uganda

⁷Harvard T.H. Chan School of Public Health, Boston, 651 Huntington Ave, Boston, MA, 02115, USA

⁸International AIDS Vaccine Initiative (IAVI)- Vaccine, Immunology, Science and Technology for Africa (VISTA), Medical Research Council (MRC)/ (Uganda Virus Research Institute) UVRI & London School of Hygiene and Tropical Medicine (LSHTM) Uganda Research Unit, Entebbe, P.O.Box 49, Uganda

⁹Kenya AIDS Vaccine Initiative-Institute of Clinical Research (KAVI-ICR), College of Health Sciences, University of Nairobi, Nairobi, P.O. Box 19676, Kenya

¹⁰Rwanda Zambia HIV Research Group (RZHRG), Kigali Rwanda, Lusaka and Ndola, PostNet 412, P/Bag E891, Zambia

¹¹Kenya Medical Research Institute (KEMRI), Wellcome Trust Research Program (KWTRP), Kilifi, P.O Box 230, Kenya

¹²Center for Health Services Studies, University of Kent, Canterbury, CT2 7NF, UK

¹³School of Nursing and Public Health, College of Health Sciences, University of KwaZulu Natal, Durban, 4001, South Africa

V3 First published: 08 Jul 2020, 5:163
<https://doi.org/10.12688/wellcomeopenres.16070.1>
 Second version: 16 Sep 2020, 5:163
<https://doi.org/10.12688/wellcomeopenres.16070.2>
 Latest published: 10 Dec 2020, 5:163
<https://doi.org/10.12688/wellcomeopenres.16070.3>

Abstract

Emerging highly transmissible viral infections such as SARS-CoV-2 pose a significant global threat to human health and the economy. Since its first appearance in December 2019 in the city of Wuhan, Hubei province, China, SARS-CoV-2 infection has quickly spread across the globe, with the first case reported on the African continent, in Egypt on February 14th, 2020. Although the global number of COVID-19 infections has increased exponentially since the beginning of the pandemic, the number of new infections and deaths recorded in

Open Peer Review

Reviewer Status

Invited Reviewers

1 2

version 3

(revision)

10 Dec 2020

version 2

(revision)



report




report

African countries have been relatively modest, suggesting slower transmission dynamics of the virus on the continent, a lower case fatality rate, or simply a lack of testing or reliable data. Notably, there is no significant increase in unexplained pneumonias or deaths on the continent which could possibly indicate the effectiveness of interventions introduced by several African governments. However, there has not yet been a comprehensive assessment of sub-Saharan Africa's (SSA) preparedness and response to the COVID-19 pandemic that may have contributed to prevent an uncontrolled outbreak so far. As a group of early career scientists and the next generation of African scientific leaders with experience of working in medical and diverse health research fields in both SSA and resource-rich countries, we present a unique perspective on the current public health interventions to fight COVID-19 in Africa. Our perspective is based on extensive review of the available scientific publications, official technical reports and announcements released by governmental and non-governmental health organizations as well as from our personal experiences as workers on the COVID-19 battlefield in SSA. We documented public health interventions implemented in seven SSA countries including Uganda, Kenya, Rwanda, Cameroon, Zambia, South Africa and Botswana, the existing gaps and the important components of disease control that may strengthen SSA response to future outbreaks.

Keywords

Sub-Saharan Africa, SARS-CoV-2, COVID-19, pandemic, preparedness, and response.



This article is included in the [Coronavirus \(COVID-19\)](#) collection.

Invited Reviewers

1

2

16 Sep 2020

version 1


08 Jul 2020


?

report

✓

report

1. Charles Shey Wiysonge , South African Medical Research Council, Cape Town, South Africa

2. Kondwani C. Jambo , Liverpool School of Tropical Medicine (LSTM), Liverpool, UK

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Gisele Umvilighozo (gumvilig@sfu.ca)

Author roles: **Umvilighozo G:** Conceptualization, Funding Acquisition, Investigation, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Mupfumi L:** Funding Acquisition, Investigation, Writing – Review & Editing; **Sonela N:** Funding Acquisition, Investigation, Writing – Review & Editing; **Naicker D:** Funding Acquisition, Investigation, Writing – Review & Editing; **Obuku EA:** Investigation, Writing – Review & Editing; **Koofhethile C:** Funding Acquisition, Investigation, Writing – Review & Editing; **Mogashoa T:** Funding Acquisition, Investigation, Writing – Review & Editing; **Kapaata A:** Funding Acquisition, Investigation, Writing – Review & Editing; **Ombati G:** Funding Acquisition, Investigation, Writing – Review & Editing; **Michelo CM:** Funding Acquisition, Investigation, Writing – Review & Editing; **Makobu K:** Funding Acquisition, Investigation, Writing – Review & Editing; **Todowedo O:** Funding Acquisition, Investigation, Writing – Review & Editing; **Balinda SN:** Investigation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This project was funded by research grants from the Sub-Saharan African Network for TB/HIV Research Excellence (SANTHE; a DELTAS Africa Initiative [grant # DEL-15-006]). The DELTAS Africa Initiative is an independent funding scheme of the African Academy of Sciences (AAS)'s Alliance for Accelerating Excellence in Science in Africa (AESA) and supported by the New Partnership for Africa's Development Planning and Coordinating Agency (NEPAD Agency) with funding from the Wellcome Trust [grant # 107752/Z/15/Z] and the UK government. The views expressed in this publication are those of the author(s) and not necessarily those of AAS, NEPAD Agency, Wellcome Trust or the UK government.

Copyright: © 2020 Umvilighozo G *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Umvilighozo G, Mupfumi L, Sonela N *et al.* **Sub-Saharan Africa preparedness and response to the COVID-19 pandemic: A perspective of early career African scientists [version 3; peer review: 2 approved]** Wellcome Open Research 2020, 5:163 <https://doi.org/10.12688/wellcomeopenres.16070.3>

First published: 08 Jul 2020, 5:163 <https://doi.org/10.12688/wellcomeopenres.16070.1>

REVISED Amendments from Version 2

In response to the reviewer's comments, the following changes were made to this article.

In the introduction and Table 2, "global pandemic" was replaced with "pandemic".

Any further responses from the reviewers can be found at the end of the article

Disclaimer

The views expressed in this article are those of the author(s) and do not reflect those of their employers or institutions. Publication in Wellcome Open Research does not imply endorsement by Wellcome.

Introduction

COVID-19 is caused by a novel beta-coronavirus named severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) that was first reported in December 2019 in the city of Wuhan, Hubei province, China¹. SARS-CoV-2 infection has quickly spread across the globe², with the first case reported on the African continent, in Egypt on February 14th 2020³. COVID-19 was declared a public health emergency of international concern on January 30th 2020 and a pandemic on March 11th 2020 by the World Health Organization (WHO)^{4,5}. Just a few months into the pandemic, COVID-19 has ravaged developed countries, with significant case fatality rates in Europe and the USA⁶. Considering the large number of people that live in poor and crowded informal settings in sub-Saharan Africa (SSA)^{7,8}, the mode of transmission of SARS-CoV-2, the severity of the disease as well as the existing fragile health systems^{9–11} it was hypothesized that Africa may markedly be affected by the COVID-19 pandemic resulting in disastrous consequences with a large number of patients overwhelming hospitals and high case fatality rates^{9,12}.

As of June 6th COVID-19 global infections recorded in 215 countries were 6,663,034 cases and 392,802 deaths [5.9%]. The three countries with the highest burden were the USA with 1,857,872 cases [deaths 5.8%], Brazil 614,941 [5.5%] and Russia 458,689 [1.2%]. Italy and Spain were highly affected in the first quarter of the year but have stabilised their epidemics since the beginning of May 2020 with 234,531 cases [deaths: 14.4%] and 240,978 [11.3%] respectively⁶. At the same time, the African continent accounted for only 2.6% of the global infections with 177,953 cases and 4,936 deaths [2.8%] in 54 countries¹³. Four African countries, South Africa, Egypt, Morocco and Algeria, comprise more than 52% of reported COVID-19 cases in Africa¹³ and only one of these countries is located in the SSA region. Although the global number of SARS-CoV-2 infections increased exponentially since the beginning of the pandemic due to ongoing transmission, the low number of infections and deaths recorded in SSA countries have raised suspicions on whether they represent a slow progression of the pandemic in Africa, a lower case fatality rate, or simply a lack of testing or reliable data. Notably, there is no significant increase in unexplained pneumonias or deaths recorded on the continent which could possibly

indicate the effectiveness of interventions introduced by several African governments, although recent reports from South Africa indicate a surge in weekly deaths from natural causes that may be attributable to COVID-19¹⁴. However, there has not yet been a comprehensive assessment of SSA's preparedness and response to the COVID-19 pandemic that may have contributed to prevent an uncontrolled outbreak so far.

Here, we – a diverse group of early career researchers (graduate students and post-doctoral scientists) and the next generation of African scientific leaders with experience working in various fields of health research including medicine, immunology, molecular biology, microbiology, virology and public health in both SSA countries and in developed countries – conducted an assessment of African preparedness and response to the COVID-19 during the early (first 3 months) of the pandemic on the African continent. The early career researchers involved are fellows of the African Academy of Science's Sub-Saharan African Network for TB/HIV Research Excellence (SANTHE)¹⁵, (part of the DELTAS Africa initiative) and some were at the frontline of the battle against COVID-19 in their respective countries during the time of this assessment.

As of June 6th 2020, the seven countries represented had reported the following caseloads of COVID-19: Cameroon, 7,599 cases [deaths: 2.8%]; Kenya, 2,600 [3.2%]; Zambia, 1,089 [0.6%]; Rwanda, 431 [0.5%]; Uganda, 593 [0%]; Botswana, 40 [2.5%]¹³ and South Africa, which had recorded the highest number of COVID-19 cases in Africa with 45,973 cases [2.1% deaths]¹³. Even though the COVID-19 pandemic unfolded rapidly, and the undertaken public health measures interrupted our studies, our careers and other usual activities, this pandemic has also offered a blueprint on how to deal with epidemics. This analysis presents a unique perspective on the currently developed public health interventions to fight COVID-19 on the African continent. We discuss the challenges and opportunities that exist to improve African capacity to fight future epidemics from our perspective as the next generation of scientists that will oversee these responses in the future.

Our evaluation of sub-Saharan Africa preparedness and response to COVID-19 reviewed country specific preventive measures and critically examined the response to the pandemic in seven African countries. We assessed the public health measures and other crucial interventions that were put in place in the control of COVID-19 in Uganda, Kenya, Rwanda, Cameroon, Zambia, South Africa and Botswana. We argue that these strategies may have helped to prevent a disastrous outcome by reducing rapid transmissions that may happen in clusters and minimizing the number of patients seeking medical assistance at the same time. We also identified and summarized in three categories of biomedical, sociocultural and economic factors; the challenges encountered at different levels of the health system. We presented the gaps existing in the public health intervention programs that may result in delays/failure to halt the spread of the disease as well as the important components of disease control that may strengthen sub-Saharan Africa preparedness and response to future outbreaks.

Assessment and findings

The rapid rise of SARS-CoV-2 infections sent a clear message to the world that quick action was needed to prevent the spread of the disease. WHO warned on February 22nd 2020 that all member states of the African Union should develop an early strong plan of action to tackle the growing threat¹⁶. As emerging African leaders in health research and fellows of the SANthe consortium, we convened a virtual meeting to discuss the threat that the pandemic posed to SSA and agreed that we had a responsibility to critically analyze the response of our governments so far and to offer our own perspective on how this and other similar epidemics should be tackled on the continent. In the meeting, participants discussed the origin of the new SARS-CoV-2 virus, infection preventive measures, diagnostic tests, clinical management of the disease and the development of vaccines and therapeutics by critically reviewing the available scientific literature. We discussed the need for scientific and evidence-based responses that considered Africa's unique health-care, sociocultural and economic challenges. We therefore decided to review articles focusing on SARS-CoV-2 infection dynamics in Africa and the responses that African governments were taking. Specifically, we used the free search engine PubMed to identify articles that discussed the transmission dynamics of the new SARS-CoV-2 virus, or described the sociocultural, economic and the state of health care systems in Africa and their impact on SARS-CoV-2 transmission. Moreover, we searched for popular media reports and performed internet searches at official websites for any documents or articles highlighting interventions against COVID-19 implemented by African governments, local health authorities or nongovernmental health organizations. We also relied on our own personal experiences and perspectives. Considering that the response to the epidemic has been very variable from country to country, we agreed to focus our analysis from reports emanating from Uganda, Kenya, Rwanda, Cameroon, Zambia, South Africa and Botswana, countries that were represented among us. An extensive review of the steps taken in preparedness and response to COVID-19 guided by a representative from each of the seven SSA countries was conducted and a conclusive report encompassing our perspectives was generated.

Sub-Saharan Africa preparedness for COVID-19

Since the beginning of the COVID-19 pandemic, the world's top priority was to contain the spread of SARS-CoV-2, to reduce disease fatalities and to limit the patient burden on health systems. Despite the uncertainty and unanswered questions around the management of the newly emergent SARS-CoV-2 infection, African countries joined the global effort to battle the COVID-19 pandemic as we outline in detail below. It has been reported that the experiences of SSA countries in handling ongoing outbreaks and managing infectious diseases such as Ebola, tuberculosis, malaria and HIV came in handy in the fight of COVID-19^{17–19}. Pre-existing emergency plans on public health interventions, community engagement programs and the work force composed of emergency medical experts and trained health care workers were quickly re-directed to ensure a fast response to COVID-19^{17–21}.

As early as January 2020, prior to the identification of the first case of SARS-CoV-2 infection on the African continent on February 14th 2020³, African countries had already initiated public engagement conversations to inform the population about the new pandemic^{22–28}. Public notices about COVID-19 were issued between January and March in the seven SSA countries that we assessed^{29–35}. In preparation for possible incoming cases the Ministries of Health worked with local health authorities to designate medical teams, testing laboratories and clinical facilities for isolation and care of COVID-19 patients^{25,36–41}. Through media communication, health authorities addressed the new threat, communicating the signs and symptoms of SARS-CoV-2 infection, its mode of transmission as well as the preventive measures and safety guidelines such as hand and respiratory hygiene and social distancing according to WHO guidelines and recommendations^{24,42–48}. Some of the countries that were considered in this analysis such as Rwanda and Kenya quickly instituted widespread hand washing stations or used hand sanitizers in public places such as bus stations and restaurant entrances^{49–51}. Health communication about COVID-19 was made easier by the use of social media platforms such as WhatsApp, Facebook or Twitter, whereas web-based chat or a hotline number that could be contacted for information and inquiries about COVID-19 were made available to the public in all the seven countries^{23,28,52–57}. Furthermore, in an effort guided by Africa CDC to strengthen the emergency response to COVID-19, the readiness of African countries to handle the new disease was assessed and training of health care workers and lab technologists on diagnosis and management of SARS-CoV-2 infection was conducted²².

The majority of African countries lack specialized medical capacity that is critical for handling severe cases of COVID-19, such as intensive care unit (ICU) beds^{11,58} and mechanical ventilators⁵⁹. Therefore, the main priority on the African continent was to contain the infection, initiate immediate testing for suspected cases and to start medical intervention prior to development or progression to severe clinical disease. The countries we assessed focused their efforts on prevention, early identification of new infections and mitigating mass spread of the virus by quickly tracing case contacts based on the available information. All seven countries initiated border screening at ports of entry by March 2020 and a 14-day self-quarantine was recommended for all incoming travellers^{53,60–65}.

Although as a result of few or unavailable laboratory technology, some SSA countries could not test for COVID-19 locally, the early established collaborative model among African countries, coordinated by the Africa CDC, increased testing capacity across the continent^{22,66,67} as an example early samples of suspected COVID-19 cases from the Central African Republic were shipped to Rwanda for testing until the local capacity became available. Nonetheless, testing constraints still remain in most countries⁶⁸, and therefore testing priority was given to most at risk persons such as returning travellers, or the people who have been in contact with confirmed cases, identified through contact tracing by health care workers. All the

seven countries have initiated early testing of suspected cases and established designated facilities for testing and clinical care of COVID-19 patients^{30,36–41}.

Sub-Saharan Africa response to COVID-19

The early implementation of COVID-19 preventive measures delayed the rapid spread of the virus within the African population, but these procedures could not completely halt the spread of the virus in all seven countries. Soon after each country had identified the first case of COVID-19, new infections were reported, with the majority related to returnee travellers or contacts of index COVID-19 cases. To reduce the risk of imported cases, these countries, with the exception of Zambia, swiftly closed borders, shut airports and reduced incoming travellers to essential workers and returning residents^{27,62–65,69–72}. In order to mitigate further spread of the disease, individuals diagnosed with COVID-19 were admitted in designated isolation areas for care and medical assistance while case contact tracing was immediately initiated. Additionally, mass gatherings and non-essential travels were prohibited, government and private business staff were encouraged to work from home and schools were closed^{65,71–77}. With the exception of Zambia, the assessed SSA countries implemented a dusk to dawn curfew and nation-wide lockdown to enforce social distancing measures, limiting movements to essential service providers^{71,78–83}. These measures were mainly put in place to prevent large volumes of new infections that would result in a high demand for hospital services, potentially leading to overwhelming of the fragile medical infrastructure^{84,85}.

Contrary to what was initially expected, the spread of SARS-CoV-2 has been relatively slower in Africa¹², and COVID-19 infections have been generally mild to moderate, leading to more recoveries and lower fatality rates in the seven SSA countries⁸⁶ compared to Western countries⁶. It should also be mentioned that this pandemic started earlier on the other continents, suggesting that it may be too early for SSA to celebrate its relative success as Africa may have not yet faced the highest phase of the disease. However, a comparison of the early phases of the pandemic in some African and European countries has shown a positive impact of early interventions initiated by SSA countries resulting in distinct disease trajectories. For example a comparison of the infection dynamics in the United States, United Kingdom, Italy and Spain vs South Africa and Cameroon has shown a continual exponential peak in non-African countries but slow and gradual increase in both of the SSA countries⁸⁷.

Our analysis suggests that early initiation of preventive measures, a faster response by timely testing of suspected cases and immediate contact tracing done by SSA countries has mitigated a faster and more extensive spread of the virus in the population. Additionally, a contemporary warm climate may have impacted the dynamics of the SARS-CoV-2 transmission in these countries^{88–90}. We posit that the predominantly young demography could be a contributing factor to a mild disease and low case fatality observed in Africa^{20,91}. Furthermore, there are suggestions that cross reactive-immunity resulting

from previous infections that are predominant in the region or the universal BCG vaccine policy⁹², widely recommended for infants vaccination in the assessed countries, may have offered some health benefits such as enhanced lung cells immunity against infections contributing to better clinical outcome of the disease. However, these observations have not yet been confirmed by rigorous evaluations. Altogether, the prevention programs that were put in place and the early response implemented by SSA countries may have mitigated the widespread dissemination of the SARS-CoV-2 virus and fatality due to COVID-19 in SSA countries^{69,93}. The interventions implemented in all seven SSA for prevention and control of the COVID-19 are summarized in [Table 1](#).

Biomedical, sociocultural and economic challenges met in the course of COVID-19 pandemic

The ability to coordinate a rapid response to COVID-19 by African governments, guided by health and scientific experts in the assessed countries, has shown the continent's strengths and the expertise to tackle health threats like the COVID-19. However, the main challenges in SSA countries such as poor infrastructure⁹⁴, clusters of high density populations⁹⁵, highest global burden of infectious diseases⁹⁶, and low GDP per capita^{97,98} have impacted the sustainability of these interventions resulting in early ease of the key measures to prevent other consequences unrelated to the pandemic. Details about the key challenges are summarized in [Table 2](#).

The main challenges that Africa faces in the response to COVID-19 pertain to lack of local biotechnological production and limited research capacity or expertise in speciality fields, thus making African countries unable to conduct sufficient testing and focused research studies related to disease transmissibility, vaccine or cure research relevant to the local context. Many African countries are relying on equipment and reagents imported from outside the continent. Although, most African countries already have the equipment footprint for COVID-19 tests from the key manufacturers Abbott, Cepheid and Roche, testing capacity is hindered by the inability to rapidly expand the technological capacity, limited funds and more recently by the export restrictions imposed on COVID-19-related supplies⁹⁹. It is heartening to note that countries such as Senegal and South Africa have initiated programmes to locally develop reagents and PPE^{100–102}. The shift in focus to production of COVID-19-specific items threatens the pre-existing line products needed for other infections such as HIV and TB. For example procurement of COVID-19 supplies has hampered the existing unstable supply chain for routine medical supplies that are essential for the management of other medical conditions (e.g. antiretroviral therapy, tuberculosis and hepatitis drugs, infant vaccines, anesthesia needed for surgeries and dental treatments, test kits, laboratory reagents and consumables used for other medical/research activities).

The mode of transmission of SARS-CoV-2 has led to enforcement of social distancing measures by restrictions of mass gatherings and a national lockdown in six of the seven countries that we assessed^{62–65,71,72,74–77}. This method of prevention

Table 1. Summary of interventions implemented for prevention and response to COVID-19.

Preparedness and preventive measures (January-March 2020)						
Strengthening medical capacity and testing technology.	<p>In collaboration with WHO, Africa CDC and member states, the following steps were taken:</p> <ul style="list-style-type: none"> January 27th: Africa CDC activated its emergency operations center incident Management system (IMS) for the 2019 n-CoV outbreak. Procurement of SARS-CoV-2 testing kits Establishment of collaborative model among African countries, setting up a specimen referral system. Deployment of health experts, training and technical support of staff from risky areas such as airports. Development of informational materials on the infection. Development/strengthening capacity of local health facilities. Timely communication and weekly updates on high priority areas for coronavirus control. Avail scientific documentation and references on the new coronavirus. 					
	<ul style="list-style-type: none"> Ministries of Health and authorities from all seven countries designated a medical team and indicated health facilities for testing and clinical care of COVID-19 patients. 					
Public engagement and Educational Sessions	<ul style="list-style-type: none"> Health authorities issued public information regarding the new disease, signs and symptoms, health precautions and communicated the WHO/ country specific safety guidelines. Each country issued an official communication independently on different days as shown below: 					
	Kenya	South Africa	Cameroon	Rwanda	Botswana	Uganda
	February 2 nd	March 7 th	January 28 th	January 21 st	March 16 th	March 2 nd
Training health care workers and lab technologists	<ul style="list-style-type: none"> Rwanda and Kenya instituted widespread hand washing stations and used hand sanitizers in public places such as bus stations and restaurant entrances. Mass media use for awareness and health promotion. Social media technology facilitated communication about COVID-19 between health care workers and the public. Each country issued a hotline number specific for COVID-19 information/response. 					
	<ul style="list-style-type: none"> February 6–7th effort guided by Africa CDC for Strengthening the emergence response to COVID-19 by training African laboratories and clinicians on diagnosis and management of SARS-CoV-2 infection was conducted. Upon training completion, documentation about the critical steps of the management of SARS-CoV-2 infection developed by health authorities in each country was made available to health care workers and posted on the official website. 					

Preparedness and preventive measures (January-March 2020)

January 30th, 2020: WHO declares COVID-19 a Public health emergency of International concern

February 14th, 2020: First case of COVID-19 reported on the African continent (Egypt)

March 11th, 2020: WHO declares COVID-19 a pandemic.

Responsive measures (April-May 2020)

All the seven sub-Saharan African countries implemented similar responses to COVID-19 except Zambia that didn't close borders or enforce a national lockdown. The different interventions and their implementation date in each country are presented below:

1. Border closure for non-citizen and non-essential workers. (Except for Zambia)
2. COVID-19 screening at port of entry for all seven countries
3. 14-days self-quarantine recommended for all incoming travellers.
4. Isolation of COVID-19 patients at designated facilities and close medical monitoring
5. Immediate contact tracing and testing.
6. Prohibit mass gathering and non-essential travels inside the country
7. Recommendation to work from home for private and government institutions
8. Schools closure
9. Dusk to dawn curfew and a national lockdown (Except Zambia)

Country	Kenya	South Africa	Cameroon	Rwanda	Botswana	Uganda	Zambia
First positive case	March 12 th	March 5 th	March 6 th	March 14 th	March 30 th	March 21 st	March 18 th
1-3	March 15 th	March 16 th	March 18 th	March 3 rd	March 24 th	March 22 nd	February 21 st
6-8	March 15 th	March 15 th	March 18 th	March 21 st	March 16 th	March 25 th	March 26 th
9	March 28 th April 6 th	March 27 th	March 18 th	March 28 th	April 2 nd	March 30 th	Not done

Note1: Only screening at the port of entry and self-quarantine were implemented in Zambia.

Note2: A curfew was first implemented in Kenya in March 28th followed by a national lockdown in April 6th 2020

Table 2. Biomedical, Sociocultural and economic challenges of outbreak control.

CATEGORY	Type of challenge encountered	Consequences on both preparedness and response to COVID-19 pandemic
Biomedical	Limited capacity for epidemiological techniques such as mathematical modeling to guide the decision making in response to the outbreak, particularly in localised settings.	<ul style="list-style-type: none"> Insufficient scientific references to guide the response to the outbreak.
	Insufficient medical infrastructure including laboratory technology, (Example: RT-PCR testing labs)/medical capacity such as ICU services, mechanical ventilators.	<ul style="list-style-type: none"> Inability/reduced capacity to perform the required tests locally Delays in availability of test results Reduced patient safety that may result in life losses due to unavailability of required medical procedures.
	Shortage of medical supplies and PPE (priority given to COVID-19 medical and research activities)	<ul style="list-style-type: none"> Increased risk of infections among health care workers Interruption/delay of non-COVID-19 related research activities.
	Insufficient testing capacity	<ul style="list-style-type: none"> Inability to attain the testing level needed for adequate disease surveillance and control. Delayed testing that may result in increased disease spread due to late detection of COVID-19 cases.
	Shortage of medical/research and clinical laboratory personnel and space	<ul style="list-style-type: none"> Overworked medical personnel Focused medical attention to COVID-19 delaying non-essential medical services during the pandemic such as the recommended regular medical check-ups and non-life-threatening interventions. Interruption or delay of non-COVID-19 related medical/research activities (Example: minor/elective surgeries).
	Lack of local biotech capacity to conduct advanced biomedical research studies such as transmissibility of SARS-CoV-2 in the African climate conditions, antibody-based therapy, or vaccine and treatment research in the African population.	<ul style="list-style-type: none"> Relying on responses from countries that have the capacity to create solutions. Unavailability of accurate information relevant to the local context that is important for development of adequate preventive measures.
Sociocultural	Interruption of school programs and unavailability of remote education technology	<ul style="list-style-type: none"> Delays in completion of school programs
	Structure of the markets, social aspect of the population and the culture	<ul style="list-style-type: none"> Difficulties to practice social distancing in the communities
	Science is misunderstood, misinformation about the consequences of the safety recommendations (Some interventions being termed harmful or unethical among some communities. Ex: some have suggested that wearing a mask is detrimental to health due to carbon dioxide poisoning ¹⁰³ , misleading myths and numerous faith-related rumors whereby some religious leaders have spread wrong information that they are able to cure COVID-19 ¹⁰⁴ .)	<ul style="list-style-type: none"> Mistrust of health care systems Failure to comply with safety procedures

CATEGORY	Type of challenge encountered	Consequences on both preparedness and response to COVID-19 pandemic
Economic	Insufficient funds	<ul style="list-style-type: none"> • Limited procurement capacity, • Difficulty to expand existing services or scale up to new available technology.
	Borders closure and reduced frequency of international trade	<ul style="list-style-type: none"> • Delay of transport of essential materials that are initially imported (Example shortage of infant vaccines, anesthesia used for minor surgeries or dentistry). • Increased cost for medical supplies and imported food items • Unavailability of needed materials locally
	Interrupted supply chain due to market scarcity/ priority given to non-African countries	<ul style="list-style-type: none"> • Incapacity to obtain suppliers for the African market even when there are available funds. • Shortage of frequently used reagents that need to be imported.
	Poor infrastructure, poverty, informal housing and high population density	<ul style="list-style-type: none"> • Increased risk to get the infection due to unavailability of essential sanitary services • Nearly impossible to comply with social distancing • Hardly able to implement safety measures
	Reduced job security due to lockdown measures	<ul style="list-style-type: none"> • Increased unemployment during COVID-19 pandemic • Loss of income for most of the families who depend on casual labor, informal market that have been severely affected by the lockdown.
	Lockdown resulting in reduced movements between cities, unavailability of public transportation and discontinued non-essential work activities including stopping work for researchers working on non COVID-19 projects	<ul style="list-style-type: none"> • Interruption of pre-existing programs (Ex: HIV prevention programs such as PrEP, ART treatment, TB programs, cancer, maternal health care or non-life-threatening surgeries), • Ironically, the emergence of a new virus has prevented virologists to go to the lab!
	Lack of income due to discontinued earning activities, inability to buy food leading to starvation	<ul style="list-style-type: none"> • Countries unplanned mobilization of emergency fund to feed poor families. • Early ease of the lockdown that may result in new infections • Re-opening work activities to avoid hunger related deaths.

has specifically disrupted the school programmes and created economic crises that resulted in hunger¹⁰⁵ and other hardships for the large SSA population who depend on casual labor and rely on daily income⁷. The fragile health systems coupled with lockdown measures have inadvertently reduced access to health care for non-emergency and other pre-existing medical conditions. For example with a re-prioritisation of human resources, it is estimated that an additional 6.5 million TB cases will occur over the next five years¹⁰⁶. Although the impact on other diseases such as HIV and malaria has not been assessed, it is likely that the COVID-19 pandemic will set back some of the gains made in the countries' responses to these killer diseases.

Recommended solutions to bridge the gaps for improved outbreak preparedness and response

Based on our assessment of the challenges and the gaps that were found in the approach used by the seven SSA to prevent/respond to COVID-19 outbreak, we summarized the potential solutions in Figure 1. We classified these into three interconnected categories of biomedical, sociocultural and economic aspects that we recommended to help improve the preparedness and response to future outbreaks.

In response to the current COVID-19 pandemic, Africa CDC has started planning for the coordination of a centralised procurement system to reach the target of 55 million tests across the continent. Tens of thousands of test kits, PPE and thermometers have already been distributed to countries through a donation from the Jack Ma Foundation⁹⁹. However, sustainability of a strong supply chain requires African governments to mobilize resources and to avail funding for health emergency response and research development on the continent including funded education in speciality fields. This will allow the countries to generate adequate interventions and to maintain a rapid response to outbreaks without overreliance on expertise from non-African countries and urgent importation of supplies. SSA countries should leverage novel medical/research capacity upon the existing structures that were put in place over many decades of fighting other public health threats such as Ebola, HIV, TB and malaria. The establishment of a network of multidisciplinary health care workers competent in various tasks such as community education and testing, in a multi-disease focused approach would allow management of staff shortage rather than having to prioritize the new life-threatening disease over those that were already prevailing on the continent.




Figure 1. Recommended solutions for improved preparedness and response to future outbreaks.

Limitations

Some of the limitations encountered during the development of this assessment were mainly related to the lack of sufficient documentation to address the actual reality in SSA countries, such as the status of health systems or informal housing structure. Documentation on physical capacity on the African ground is needed. Some essential documents lacked the important information such as the date signed and released. Improvement on good record keeping especially for health data of this kind is essential for future references.

Our assessment was not designed to demonstrate with certainty that the implemented interventions were directly linked to the number of infections or COVID-19 deaths in the assessed SSA countries. Clinical and biomedical research studies may be more appropriate to confirm these observations. Further, systematic reviews of effects would be informative. Nevertheless, we believe that our unique perspective on the SSA countries preparedness and response to a great health threat such as the COVID-19 pandemic has provided a valuable contribution to the future interventions.

Conclusion

We assessed the Sub-Saharan Africa preparedness and response to COVID-19. Based on an extensive review of the available scientific publications, the government technical reports and the announcements released by governmental and non-governmental health organizations as well as our personal experiences as workers on the COVID-19 battlefield in SSA countries. This assessment was conducted during the first three month of the COVID-19 pandemic before the virus spread widely in SSA and our conclusions were drawn based on statistical information on the disease presented in this study. We documented the preventive measures and the response put in place to contain the SARS-CoV-2 in seven SSA countries including Uganda, Kenya, Rwanda, Cameroon, Zambia, South Africa and Botswana. We have shown the strengths of early initiated interventions that may have contributed to modest and slower dynamics of COVID-19 in SSA countries. To prepare for the fight against the COVID-19 pandemic, the countries of the sub-Saharan Africa assessed have strengthened their medical capacities by rapidly introducing screening techniques for SARS-CoV-2 and training health care workers in the management of the new disease. Public engagement efforts and information sessions were launched. In response to the COVID-19 pandemic these countries closed their borders and airports, implemented screening for SARS-CoV-2 at port of entry, and introduced a mandatory 14-day quarantine for returning travellers. These SSA countries issued national guidelines on recommended safety measures, initiated

immediate contact tracing, prevented mass gatherings, instituted national lockdown and curfew, closed schools and urged the private sector and the government personnel to work from home. It was noted that Zambia, which did not fully implement the interventions described in this study as well as other countries not included in our review that have struggled to put into action public health interventions^{107,108}, reported no significant increases in deaths compared to countries with more robust responses reported here. There, consistent with the recent study by M. Njenga *et al.*¹⁰⁹ which investigated the causes of low morbidity and low mortality of COVID-19 in African countries, we also suggest that other factors such as warm climate, young population, pre-existing cross reactive immunity may have considerably contributed to the evolution of the COVID-19 pandemic in SSA countries. It should be noted that implementation of the public health interventions for preparedness and response to the COVID-19 pandemic in sub-Saharan Africa faced many challenges. We discussed the need for scientific research and evidence-based responses that considered Africa's unique healthcare, sociocultural and economic challenges. Overall, the assessed countries lacked the local biotechnological capacity for the production of biomedical supplies⁹⁹, they had a limited workforce with the expertise to specifically address the pandemic using evidence¹¹⁰, there was lack of community trust¹⁰³, poor infrastructure¹¹, inability to manage remote education programs¹¹¹, failure to maintain the measures implemented due to the population economic instability^{7,8} and insufficient funds¹⁰. Formal studies of the extent of these challenges and how to address them in the future will be required. While efforts to bridge some of the gaps have been initiated, we recommend that SSA countries develop continued funding streams to support these initiatives as well to increase south to south/north- south collaborations to enhance the capacity of the existing health systems. Therefore, if these problems are addressed in a timely manner, there is no doubt that in the next five years SSA countries will have developed a reliable-strong health system to prevent the newly emerging viral infections to spread at a large scale.

Data availability

No data are associated with this article.

Acknowledgements

We gratefully acknowledge the contributions of health care workers for their helpful discussions. We thank Professor Thumbi Ndung'u for his guidance, Dr Victoria Kaspruwicz and Dr Denis Chopera for their invaluable contribution to this work.

References

1. Lu H, Stratton CW, Tang YW: **Outbreak of pneumonia of unknown etiology in Wuhan, China: The mystery and the miracle.** *J Med Virol.* 2020; **92**(4): 401–402.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Wang C, Horby PW, Hayden FG, *et al.*: **A novel coronavirus outbreak of global health concern.** *Lancet.* 2020; **395**(10223): 470–473.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Egypt Today: **Egypt announces first Coronavirus infection.** 2020.
[Reference Source](#)
4. World Health Organization (WHO): **WHO Director-General's statement on IHR**

- Emergency Committee on Novel Coronavirus (2019-nCoV).** Published 2020. Accessed August 9, 2020.
[Reference Source](#)
5. Cucinotta D, Vanelli M: **WHO declares COVID-19 a pandemic.** *Acta Biomed.* 2020; **91**(1): 157–160.
[PubMed Abstract](#) | [Publisher Full Text](#)
 6. World Health Organization (WHO): **WHO Coronavirus (COVID-19) Dashboard.** 2020.
[Reference Source](#)
 7. Potts D: **The state and the informal in sub-Saharan African urban economies: Revisiting debates on dualism.** *Cris States Res Cent Work Pap.* 2007.
[Reference Source](#)
 8. Potts D: **The urban informal sector in sub-Saharan Africa: From bad to good (and back again?).** *Development Southern Africa.* 2008; 151–167.
[Publisher Full Text](#)
 9. Gilbert M, Pullano G, Pinotti F, *et al.*: **Preparedness and vulnerability of African countries against importations of COVID-19: a modelling study.** *Lancet.* 2020; **395**(10227): 871–877.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 10. Oleribe OO, Momoh J, Uzochukwu BSC, *et al.*: **Identifying key challenges facing healthcare systems in Africa and potential solutions.** *Int J Gen Med.* 2019; **12**: 395–403.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 11. Murthy S, Leligdowicz A, Adhikari NKJ: **Intensive care unit capacity in low-income countries: A systematic review.** *PLoS One.* 2015; **10**(1): e0116949.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 12. Moore M, Gelfeld B, Okunogbe A, *et al.*: **Identifying Future Disease Hot Spots: Infectious Disease Vulnerability Index.** 2016.
[Publisher Full Text](#)
 13. Africa CDC: **Africa CDC dashboard.** 2020; Published 2020. Accessed May 25, 2020.
[Reference Source](#)
 14. South African Medical Research Council: **Report on Weekly Deaths in South Africa.**
[Reference Source](#)
 15. SANTHE: **Sub-saharan African Network for TB/HIV Research Excellence.** 2020.
[Reference Source](#)
 16. The journal.ie: **WHO warns Africa is ill-equipped to deal with coronavirus due to "weaker health systems.** 2020; Published February 22.
[Reference Source](#)
 17. Tasamba J: **Rwanda uses Ebola experience to combat COVID-19.** *Anadolu Agent.* 2020; Published 2020.
[Reference Source](#)
 18. Nordling L: **South Africa hopes its battle with HIV and TB helped prepare it for COVID-19.** *Science (80-).* 2020.
[Reference Source](#)
 19. Africa Renewal: **WHO:How the lessons from Ebola are helping Africa's COVID-19 response.** 2020;
[Reference Source](#)
 20. Ranscombe P: **Rural areas at risk during COVID-19 pandemic.** *Lancet Infect Dis.* 2020; **20**(5): 545.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 21. Geiger D, Harborth L, Mugisha A: **Managing enduring public health emergencies such as COVID-19: lessons from Uganda Red Cross Society' s Ebola virus disease response operation.** 2020; 1–4.
[Publisher Full Text](#) | [Free Full Text](#)
 22. Africa CDC: **Outbreak Brief: Novel Coronavirus (2019-nCoV) Global Epidemic Date of Issue: 04 February 2020.** 2020; 1–3.
[Reference Source](#)
 23. Republic of Kenya Ministry of Health: **General information about COVID-19.** Published. 2020. Accessed May 20, 2020.
[Reference Source](#)
 24. South African Government: **Coronavirus info leaflet.** 2020.
[Reference Source](#)
 25. Ministère de la santé: **République du Camérout. Coronavirus: Mise En Place Progressive Du Système de Prévention.;** 2020.
[Reference Source](#)
 26. Uganda Ministry of Health: **COVID-19 Communication Ressources.** 2020.
[Reference Source](#)
 27. Muzvidziwa O, Muza L: **Botswana – Broad Measures Introduced by Government to Combat COVID-19 Threat.** KPMG.
[Reference Source](#)
 28. Republic of Zambia Ministry of Health: **Coronavirus disease (COVID-19) Protect yourself.** Published. 2020.
[Reference Source](#)
 29. Ministry of Health Republic of Rwanda: **Republic of Rwanda Ministry of Health Public Notice on Novel Coronavirus.** 2020.
[Reference Source](#)
 30. Ministère de la santé: **République du Camérout. Press release: Epidemic caused by a new type of Coronavirus in China.** 2020.
[Reference Source](#)
 31. Kenya Ministry of Health: **Republic of Kenya, Ministry of Health: Update on novel Coronavirus.** Press release Feb 2nd 2020. 2020.
[Publisher Full Text](#)
 32. Uganda Ministry of Health: **Press release: Results for samples of COVID-19 test negative.** 2020.
[Reference Source](#)
 33. Republic of South Africa: **Media Statement- 10th March 2020.** Accessed May 25, 2020.
[Reference Source](#)
 34. Republic of Botswana: **Public notice- Covid19 Press release.pdf.** 2020.
[Reference Source](#)
 35. Republic of Zambia: **Statement by the president of the Republic of Zambia on the COVID-19 Pandemic.** 2020.
[Reference Source](#)
 36. Rwanda Biomedical centre (RBC): **Information on mandatory quarantine on entry into Rwanda.** Accessed May 23, 2020.
[Reference Source](#)
 37. Republic of Zambia: **Zambia: Coronavirus screening measures implemented as of February 21. GARDA WORLD, Alertes de Sécurité.** 2020.
[Reference Source](#)
 38. Republic of Uganda Ministry of Health: **National Guidelines for Management of Covid-19.** Accessed May 20, 2020.
[Reference Source](#)
 39. Botswana COVID-19 Task force: **COVID-19: What are we doing in Botswana.** Accessed May 23, 2020.
[Reference Source](#)
 40. Republic of Kenya Ministry of Health: **Update on Coronavirus disease(COVID-19) as at 13th February 2020.** Accessed May 23, 2020.
[Reference Source](#)
 41. Health Department Republic of South Africa: **Government is identifying accommodation facilities.** Accessed May 23, 2020.
[Reference Source](#)
 42. World Health Organization (WHO): **Coronavirus disease (COVID-19) advice for the public.** Accessed May 8, 2020.
[Reference Source](#)
 43. Uganda Ministry of Health: **Prevention: What must be done.** 2020.
[Reference Source](#)
 44. Ministère de la santé: **République du Camérout. Coronavirus : Protégez Vous et Prévenez La Propagation de l' Infection.** 2020.
 45. Botswana Government Notice 128 of 2020: **Directions for the prevention of the spread of COVID-19.** African Lii. Published 2020.
[Reference Source](#)
 46. Republic of Zambia: **Statement by the Hon. Minister of Finance on further measures aimed at mitigating the impact of Coronavirus (COVID-19) on Zambian economy.** April 2020. 2020; 1–8.
[Reference Source](#)
 47. Kenya Ministry of Health: **How can I protect myself and others.** 2020.
[Reference Source](#)
 48. Rwanda Biomedical centre (RBC): **Now @RwandaHealth Holds a #pressconference to Communicate #Rwanda 's Current Preparedness Status and Guidelines to Prevent Novel #coronavirus.** 2020.
[Reference Source](#)
 49. African News Agency: **Rwanda sets up hand-washing facilities at bus station amid coronavirus fears.** Published 2020.
[Reference Source](#)
 50. Gahutu JB: **Hand washing – an essential protective measure in healthcare settings and in the community.** *Rwanda J Med Heal Sci.* 2020; **3**(1): 4314.
[Publisher Full Text](#)
 51. Al Jazeera English: **Kenya blocks entry for non-residents in virus response.** 2020.
[Reference Source](#)
 52. Ministry of Health Republic of Rwanda: **Cornavirus disease: Minimize your risk of infection.** Published 2020.
[Reference Source](#)
 53. République du Camérout: **Epidémie Causée Par Un Nouveau Type de Coronavirus En Chine.** 2020.
 54. Republic of Uganda Ministry of Health: **Guideline on preventive measures against Coronavirus.** Published 2020.
[Reference Source](#)
 55. Olum R, Bongomin F: **Social Media Platforms for Health Communication and Research in the Face of COVID-19 Pandemic: A Cross Sectional Survey in Uganda.** 2020.
[Publisher Full Text](#)
 56. Government of Botswana: **COVID-19 Contacts.** Published 2020.
[Reference Source](#)
 57. Health Department Republic of South Africa: **Coronavirus: Tool Kits.** Published 2020.
[Reference Source](#)
 58. Atumanya P, Sendagire C, Wabule A, *et al.*: **Assessment of the current capacity of intensive care units in Uganda; A descriptive study.** *J Crit Care.* 2020; **55**: 95–99.
[PubMed Abstract](#) | [Publisher Full Text](#)

59. The times of Africa: **Africa: Shortage of Not Only Ventilators!** Published April 20, 2020.
[Reference Source](#)
60. Republic of Rwanda Ministry of Health: **Instructions on Covid-19 screening at the point of entry.pdf**. 2020.
[Reference Source](#)
61. Republic of Zambia Ministry of Health: **Ministry of Health implements temperature scans at points of entry to detect possible coronavirus (COVID-19) cases as of February 21; individuals arriving from China subject to 14-day quarantine.** *GARDA WORLD, Alertes de Sécurité*. Published 2020.
[Reference Source](#)
62. Government Gazette no. 26 on 24 March 2020: **Public Health (Prevention of Introduction of Covid-19 into Botswana) Order, 2020.** Statutory Instrument 36 of 2020. 2020.
[Reference Source](#)
63. Republic of Uganda Ministry of Health: **Additional guidelines on the COVID-19 (corona virus) preventive measures.** Accessed May 20, 2020.
[Reference Source](#)
64. Health Department Republic of South Africa: **South African closures due to COVID-19.** Accessed May 20, 2020.
[Reference Source](#)
65. Republic of Kenya Ministry of Health: **Press statement on the update of Coronavirus in the country and response measures.** Accessed May 20, 2020.
[Reference Source](#)
66. Nkengasong JN, Mankoula W: **Looming threat of COVID-19 infection in Africa: act collectively, and fast.** *Lancet*. 2020; **395**(10227): 841–842.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
67. Kabatany M: **Coronavirus: Rwanda donates COVID-19 testing device to Central African Republic (CAR).** *Africa newsroom Gr*. 2020.
[Reference Source](#)
68. Our World in data: **Coronavirus (COVID-19) Testing.** Accessed June 5, 2020.
[Reference Source](#)
69. WHO: **COVID-19: situation update for WHO African Region.** *World Heal Organ*. 2020; 1–5.
[Reference Source](#)
70. Development and Cooperation: **Fear, anxiety and panic.** 2020.
[Reference Source](#)
71. République du Camérout Ministère de la Santé Publique: **Déclaration spéciale du Ministre, chef du Gouvernement sure la situation du COVID-19.** Accessed May 20, 2020.
72. Republic of Rwanda Office of the Prime Minister: **Announcement on the enhanced COVID-19 prevention measures.**
[Reference Source](#)
73. Mackenzie A: **COVID-19 - President bans travel and "mass gatherings."** *Southern and East African tourism update*. Published 2020.
74. Republic of Botswana: **Public notice_COVID-19 press release.** Published 2020.
[Reference Source](#)
75. Republic of Uganda Ministry of Health: **COVID-19 Guidelines for safe mass gatherings.** Published 2020.
[Reference Source](#)
76. AfricaNews: **Coronavirus - Zambia: Government Response To The Outbreak Of The Coronavirus 2019 (Covid-19).** *Africanews*. 2020.
[Reference Source](#)
77. Health Department Republic of South Africa: **Social distancing guidelines.** Published 2020.
[Reference Source](#)
78. Brown M: **South Africa Begins Nationwide Coronavirus Lockdown.** *VOA news*. 2020.
[Reference Source](#)
79. Health Department Republic of South Africa: **National lockdown redulations.** Published 2020.
[Reference Source](#)
80. Agence France presse: **Uganda orders two-week lockdown to contain Coronavirus.** 2020.
[Reference Source](#)
81. SABC News: **Kenya starts 21 day lockdown.** 2020.
[Reference Source](#)
82. Republic of Botswana: **Government Gazette Extra Ordinary.** Gaborone, Vol. LVIII, No 32, 2020.
[Reference Source](#)
83. Mugisha IR: **Rwanda begins 14-day complete lockdown.** *The EAST African*. Published 2020.
[Reference Source](#)
84. Mukunya D, Tumwine JK: **Challenges of tackling non COVID-19 emergencies during the unprecedented pandemic.** *Africa Heal Sci*. 2020; **20**(1): 19–20.
85. Chatterjee S: **With simple, fact-informed hygiene measures, Africa can slow COVID-19 spread.** *Africa Renew*. 2020.
[Reference Source](#)
86. WHO Regional office for Africa: **Coronavirus, COVID-19 in the WHO African region.** Published 2020.
[Reference Source](#)
87. Tilford C, Rininsland A, Kao JS, *et al.*: **Coronavirus tracked: has the epidemic peaked near you?** Published 2020.
[Reference Source](#)
88. Beck HE, Zimmermann NE, McVicar TR, *et al.*: **Present and future köppen-geiger climate classification maps at 1-km resolution.** *Sci Data*. 2018; **5**: 1–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
89. Wang J, Tang K, Feng K, *et al.*: **High Temperature and High Humidity Reduce the Transmission of COVID-19.** *SSRN Electron J*. 2020.
[Publisher Full Text](#)
90. Hao K: **Warmer weather could slow the spread of coronavirus—but not by much.** *MIT Technology Review*. Published 2020.
[Reference Source](#)
91. Worldometer: **Africa demographics.** Published 2020.
[Reference Source](#)
92. McGill University: **University of Ottawa. AERAS The BCG world atlas.** Published 2017.
[Reference Source](#)
93. Africa CDC: **Outbreak Brief # 13: Coronavirus Disease 2019 (COVID-19) Pandemic.** Africa CDC: Centres for Disease Control and Prevention.
[Reference Source](#)
94. World Health Organization (WHO): **The state of health in the WHO African Region.**
[Reference Source](#)
95. The World Bank group: **Population density (people per sq. km of land area).** Published 2019.
[Reference Source](#)
96. Breugelmans JG, Makanga MM, Cardoso ALV, *et al.*: **Bibliometric assessment of European and sub-Saharan African research output on poverty-related and neglected infectious diseases from 2003 to 2011.** *PLoS Negl Trop Dis*. 2015; **9**(8): e0003997.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
97. Martinčević I: **Visualizing GDP per Capita by Country.** *howMuch.net*. 2020; Published 2020.
[Reference Source](#)
98. Africa development bank group: **Africa economic outlook 2020.** Published 2020.
[Reference Source](#)
99. Kavanagh MM, Erondou NA, Tomori O, *et al.*: **Access to lifesaving medical resources for African countries: COVID-19 testing and response, ethics, and politics.** *Lancet*. 2020; **395**(10238): 1735–1738.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
100. Yeung P: **Senegal to trial \$1 speedy test for COVID-19.** *New Sci*. 2020; **246**(3282): 13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
101. Awal M: **South Africans develop COVID-19 test kits that give results in an hour.** *Face2Face Africa*. Published April 2020.
[Reference Source](#)
102. The South African Department for Science & Innovation: **South African Department of Science & Innovation COVID-19 research under way.** Science Business. Published March 26, 2020.
[Reference Source](#)
103. Goodman J, Carmichael F: **Coronavirus: "Deadly masks" claims debunked.** BBC News. Published July 2020. Accessed August 9, 2020.
[Reference Source](#)
104. FNR TIGG: **Pastor Who claimed he cured Coronavirus with faith dies from illness.** Complex. Published 2020. Accessed August 9, 2020.
[Reference Source](#)
105. BBC News: **Coronavirus: Most Africans "will go hungry in 14-day lockdown."** *BBC News Africa*. Published May 6, 2020.
[Reference Source](#)
106. STOP TB partnership: **The potential impact of the COVID-19 response on Tuberculosis in high burden countries: A modelling analysis.** 2020; 1–7.
[Reference Source](#)
107. Chibwana MG, Jere KC, Kamn'gona R, *et al.*: **High SARS-CoV-2 seroprevalence in Health Care Workers but relatively low numbers of deaths in urban Malawi.** *COVID-19 SARS-CoV-2 Prepr from medRxiv bioRxiv*. 2020.
[Publisher Full Text](#)
108. Mudge L: **A Perfect Storm Is Brewing in Burundi WHO Experts Expelled, Election Observers Blocked.** Human Rights Watch. Published 2020. Accessed August 25, 2020.
[Reference Source](#)
109. Njenga MK, Dawa J, Nanyingi M, *et al.*: **Why is There Low Morbidity and Mortality of COVID-19 in Africa?** *Am J Trop Med Hyg*. 2020; **103**(2): 564–569.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
110. World Health Organization (WHO), Bulletin of the World Health Organization: **Bulletin of the World Health Organization: Promoting evidence-based health care in Africa.** Published 2017. Accessed September 5, 2020.
[Reference Source](#)
111. Human Rights Watch: **Impact of Covid-19 on Children's Education in Africa.** Published 2020. Accessed September 5, 2020.
[Reference Source](#)

Open Peer Review

Current Peer Review Status:  

Version 2

Reviewer Report 22 September 2020

<https://doi.org/10.21956/wellcomeopenres.17918.r40487>

© 2020 Jambo K. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Kondwani C. Jambo 

Liverpool School of Tropical Medicine (LSTM), Liverpool, UK

The authors have adequately addressed my suggestions. I have no further comments.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Immunology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 17 September 2020

<https://doi.org/10.21956/wellcomeopenres.17918.r40486>

© 2020 Wiysonge C. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Charles Shey Wiysonge 

Cochrane South Africa, South African Medical Research Council, Cape Town, South Africa

Gisele Umvilighozo and colleagues have responded adequately to my comments and suggestions. I have only one discretionary comment at this stage. The authors refer to a "global pandemic" in a few places. Since a pandemic is always global, they should delete the word "global".

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Systematic reviews and meta-analysis, implementation science, evidence-informed policymaking, vaccinology.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 23 Sep 2020

Gisele Umviligihozo, Simon Fraser University, Burnaby, Canada

We thank the reviewer for his positive response, we agree with his comment to replace "global pandemic" with "pandemic"; this correction has been made in the introduction and in Table 2.

Competing Interests: No competing interests were disclosed.

Version 1

Reviewer Report 18 August 2020

<https://doi.org/10.21956/wellcomeopenres.17632.r39736>

© 2020 Jambo K. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Kondwani C. Jambo 

Liverpool School of Tropical Medicine (LSTM), Liverpool, UK

This is an open letter by Umviligihozo *et al.* that aims at documenting the public health interventions against COVID-19 implemented in seven sub-Saharan African countries. Overall, the paper is well written and covers the most pertinent issues on the topic. It highlights the diverse implementation strategies employed by the seven SSA countries but notes the relative similar low mortality experienced in SSA.

However, the article could do with a bit of balancing of viewpoints, especially discussing why despite the relatively different implementation strategies of public health measures the mortality is relatively low but similar among the SSA countries. In countries like Malawi (not discussed in the article), that have struggled to implement public health interventions, the mortality has not been very different from the other SSA countries. Could this observation mean that it is not necessarily the public health interventions that have driven the low mortality in SSA but potentially other factors? The authors have speculated that other factors including population demographics, cross-reactive immunity, and climate could have contributed to the less severe outcomes, but this has been relegated to a short paragraph. In its current state, the article puts a lot of weight on public health interventions but could be strengthened with a bit more discussion on the other factors.

The article would also benefit from a discussion on whether the planned public health interventions in the different countries were successfully implemented. Experiences from other countries show that what was planned on paper is not what happened or is happening on the ground. A discussion on this within the article will aid the reader in the interpretation of the findings and recommendations.

All in all, I commend the authors for a well-thought-out open letter and for their confidence as young scientists in driving opinion in Africa.

Is the rationale for the Open Letter provided in sufficient detail?

Yes

Does the article adequately reference differing views and opinions?

Partly

Are all factual statements correct, and are statements and arguments made adequately supported by citations?

Yes

Is the Open Letter written in accessible language?

Yes

Where applicable, are recommendations and next steps explained clearly for others to follow?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Immunology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 07 Sep 2020

Gisele Umviligihozo, Simon Fraser University, Burnaby, Canada

We thank the reviewer for the constructive comments and important suggestions for improving this article. Our responses are presented below:

- We agree with the reviewer that there are aspects of the COVID-19 presentation in sub-Saharan African that remain unresolved and require additional research. However, our goal here was to review that response in selected African countries and this preliminary analysis suggests a robust and effective response, notwithstanding other factors that may have influenced transmission dynamics and mortality.

In response to the reviewer, we have under the conclusion the following clause that highlighted the importance of other factors driving the observations on infection dynamics of the COVID-19. "It was noted that Zambia, which did not fully implement the interventions described in this study as well as other countries not included in our review that have struggled to put into action public health interventions,^{1,2} reported no significant increases in deaths compared to countries with more robust responses reported here. There, consistent with the recent study by M. Njenga et al³ which investigated the causes of low morbidity and low mortality of COVID-19 in African countries, we also suggest that other factors such as a warm climate, young population, pre-existing cross-reactive immunity may have considerably contributed to the evolution of the COVID-19 pandemic in SSA countries."

- We agree with the reviewer that further studies of the efficiency of implementation of public health interventions is required. This was beyond the scope of the current work.

The following summary discussing the challenges faced in the implementation of the interventions described was added to the text. "It should be noted that implementation of the public health interventions for preparedness and response to the COVID-19 pandemic in sub-Saharan Africa faced many challenges. Overall, the assessed countries lacked the local biotechnological capacity for the production of biomedical supplies,⁴ they had a limited workforce with the expertise to specifically address the pandemic using evidence,⁵ there was lack of community trust⁶, poor infrastructure⁷, inability to manage remote education programs,⁸ failure to maintain the measures implemented due to the population economic instability^{9,10} and insufficient funds.¹¹ Formal studies of the extent of these challenges and how to address them in future will be required."

References

1. Chibwana MG, Jere KC, Kamn'gona R, et al. High SARS-CoV-2 seroprevalence in health care workers but relatively low numbers of 2 deaths in urban Malawi. *COVID-19 SARS-CoV-2 Prepr from medRxiv bioRxiv*. 2020. doi:<https://doi.org/10.1101/2020.07.30.20164970>
2. Mudge L. A Perfect Storm Is Brewing in Burundi WHO Experts Expelled, Election Observers Blocked. Human Rights Watch. <https://www.hrw.org/news/2020/05/14/perfect-storm-brewing-burundi>. Published 2020. Accessed August 25, 2020.
3. Njenga MK, Dawa J, Nanyingi M, et al. Why is There Low Morbidity and Mortality of COVID-19 in Africa? *Am J Trop Med Hyg*. 2020;103(2):564-569. doi:10.4269/ajtmh.20-0474
4. Kavanagh MM, Erondou NA, Tomori O, et al. Access to lifesaving medical resources for African countries : COVID-19 testing and response , ethics , and politics. *Lancet*. 2020;6736(20):19-22. doi:10.1016/S0140-6736(20)31093-X
5. World Health Organization (WHO). Bulletin of the World Health Organization: Promoting evidence-based health care in Africa. <https://www.who.int/bulletin/volumes/95/9/17-030917/en/> . Published 2017. Accessed September 5, 2020.
6. Goodman; J, Carmichael F. Coronavirus: "Deadly masks" claims debunked. BBC News. <https://www.bbc.com/news/53108405>. Published July 2020. Accessed August 9, 2020.
7. Murthy S, Leligdowicz A, Adhikari NKJ. Intensive care unit capacity in low-income countries: A systematic review. *PLoS One*. 2015. doi:10.1371/journal.pone.0116949

8. Human Rights Watch. Impact of Covid-19 on Children's Education in Africa. https://www.hrw.org/news/2020/08/26/impact-covid-19-childrens-education-africa#_edn1 . Published 2020. Accessed September 5, 2020.
9. Potts D. The state and the informal in sub-Saharan African urban economies: Revisiting debates on dualism. *Cris States Res Cent Work Pap*. 2007.
10. Potts D. The urban informal sector in sub-Saharan Africa: From bad to good (and back again?). In: *Development Southern Africa*. ; 2008. doi:10.1080/03768350802090527
11. Oleribe OO, Momoh J, Uzochukwu BSC, et al. Identifying key challenges facing healthcare systems in Africa and potential solutions. *Int J Gen Med*. 2019. doi:10.2147/IJGM.S223882

Competing Interests: No competing interests were disclosed.

Reviewer Report 04 August 2020

<https://doi.org/10.21956/wellcomeopenres.17632.r39461>

© 2020 Wiysonge C. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Charles Shey Wiysonge 

Cochrane South Africa, South African Medical Research Council, Cape Town, South Africa

I read, with keen interest, the article on public health interventions to fight COVID-19 in Africa by Gisele Umviligihozo and her colleagues, all of them early career scientists, who describe themselves as the next generation of African scientific leaders. This portrays a commendable sense of duty.

The article deserves indexing and I have only minor comments and suggestions for improvement.

The authors write that this article is "Based on an extensive review of the available scientific publications, the government technical reports and the announcements released by governmental and nongovernmental health organizations ...". However, the authors do not provide a description of the search and selection of the literature used in this article. I suggest that they should provide this information in the article or as a supplementary document.

The World Health Organization declared Covid-19 a public health emergency of international concern on 30 January 2020 and not in March 2020. Please refer to reference 1¹.

The authors state that "Notably, there is no significant increase in unexplained pneumonias or deaths recorded on the continent which could possibly indicate the effectiveness of interventions introduced by several African governments". While I generally agree with this statement, especially given the time they submitted the article, I would like to alert the authors to reports by the South African Medical Research Council of a recent upsurge in weekly deaths from natural

causes in South Africa. Please refer to reference 2².

In Table 2, what does the sociocultural challenge “suspicion of harmful and unethical interventions” mean? What are the harmful and unethical interventions and why and who offered the harmful and unethical interventions to whom?

What are “faith-related rumors”?

References

1. World Health Organisation: WHO Director-General's statement on IHR Emergency Committee on Novel Coronavirus (2019-nCoV). *Date accessed - 3rd August 2020*. [Reference Source](#)
2. South African Medical Research Council: Report on Weekly Deaths in South Africa. *Date Accessed - 3rd August 2020*. [Reference Source](#)

Is the rationale for the Open Letter provided in sufficient detail?

Yes

Does the article adequately reference differing views and opinions?

Yes

Are all factual statements correct, and are statements and arguments made adequately supported by citations?

Partly

Is the Open Letter written in accessible language?

Yes

Where applicable, are recommendations and next steps explained clearly for others to follow?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Systematic reviews and meta-analysis, implementation science, evidence-informed policymaking, vaccinology.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 07 Sep 2020

Gisele Umvilighozo, Simon Fraser University, Burnaby, Canada

We thank the reviewer for the positive comments and suggestions for improving this article, our responses are as follows:

- To elaborate on the search criteria and to describe the research methodology used in the selection of the literature for this article, the following statement under “Assessment and findings” was updated to reflect these steps. “We therefore decided to review articles focusing on SARS-CoV-2 infection dynamics in Africa and the responses that African governments were taking. Specifically, we used the free search engine PubMed to identify articles that discussed the transmission dynamics of the new SARS-CoV-2 virus, or described the sociocultural, economic and the state of health care systems in Africa and their impact on SARS-CoV-2 transmission. Moreover, we searched for popular media reports and performed internet searches at official websites for any documents or articles highlighting interventions against COVID-19 implemented by African governments, local health authorities or nongovernmental health organizations. We also relied on our own personal experiences and perspectives.”
- We agree with the reviewer's comment on this date, a correction has been made to this statement which now reads “COVID-19 was declared a public health emergency of international concern on January 30th, 2020 and a global pandemic on March 11th, 2020 by the World Health Organization (WHO).¹”
- We agree with the reviewer's comment on the need to draw the reader's attention to the changing dynamics of the COVID-19 pandemic in SSA, to address this concern we updated the introduction to include that recent reports from South Africa indicate a surge in weekly deaths from natural causes that may be attributable to COVID-19.²
We also clarified in conclusion that “This assessment was conducted during the first three month of the COVID-19 pandemic before the virus spread widely in SSA and our conclusions were drawn based on statistical information on the disease presented in this study”
- The sociocultural challenges mentioned in Table 2 refer to the misinformation about the consequences of the safety recommendations. As an example, misinformation has led to some interventions being termed harmful or unethical among some communities. Examples include interventions such as wearing a mask which some have suggested is detrimental to health due to carbon dioxide poisoning,³ numerous faith-related rumors, whereby some religious leaders have spread misleading information that they are able to cure COVID-19⁴. Table 2 was updated to clarify these points.

References

1. World Health Organization (WHO). World Health Organization: WHO Director-General's statement on IHR Emergency Committee on Novel Coronavirus (2019-nCoV). [https://www.who.int/dg/speeches/detail/who-director-general-s-statement-on-ihr-emergency-committee-on-novel-coronavirus-\(2019-ncov\)](https://www.who.int/dg/speeches/detail/who-director-general-s-statement-on-ihr-emergency-committee-on-novel-coronavirus-(2019-ncov)). Published 2020. Accessed August 9, 2020.
2. South African Medical Research Council. Report on Weekly Deaths in South Africa.
3. Goodman; J, Carmichael F. Coronavirus: “Deadly masks” claims debunked. BBC News. <https://www.bbc.com/news/53108405>. Published July 2020. Accessed August 9, 2020.

4. FNR TIGG. Pastor Who claimed he cured Coronavirus with faith dies from illness. Complex. <https://www.complex.com/life/2020/05/pastor-claims-cured-coronavirus-with-faith-dies-from-illness>. Published 2020. Accessed August 9, 2020.

Competing Interests: No competing interests were disclosed.
