

This is a repository copy of *Exposure to second-hand smoke in public places and barriers to the implementation of smoke free regulations in the Gambia : a population-based survey*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/173412/>

Version: Accepted Version

---

**Article:**

Cham, Bai, Mdege, Noreen Dadirai orcid.org/0000-0003-3189-3473, Bauld, Linda et al. (2 more authors) (2021) Exposure to second-hand smoke in public places and barriers to the implementation of smoke free regulations in the Gambia : a population-based survey. *International Journal of Environmental Research and Public Health*. 6263. ISSN 1660-4601

<https://doi.org/10.3390/ijerph18126263>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



1 Article

2 **Exposure to second-hand smoke in public places and**  
3 **barriers to the implementation of smoke free regula-**  
4 **tions in the Gambia: a population-based survey**

5 **Bai Cham**<sup>1,\*</sup>, **Noreen Dadirai Mdege**<sup>2</sup>, **Linda Bauld**<sup>3</sup>, **John Britton**<sup>4</sup> and **Umberto D’Alessandro**<sup>1</sup>

6 <sup>1</sup> Disease Control and Elimination Theme, Medical Research Council Unit The Gambia at  
7 the London School of Hygiene and Tropical Medicine, Atlantic Road, Fajara, P.O.Box  
8 273, Banjul, The Gambia; [bacham@mrc.gm](mailto:bacham@mrc.gm) (BC); [udalessandro@mrc.gm](mailto:udalessandro@mrc.gm) (UDA)  
9 <sup>2</sup> Department of Health Sciences, University of York, United Kingdom;  
10 [noreen.mdege@york.ac.uk](mailto:noreen.mdege@york.ac.uk)  
11 <sup>3</sup> Usher Institute and SPECTRUM Consortium, College of Medicine and Veterinary Med-  
12 icine, University of Edinburgh, United Kingdom; [Linda.Bauld@ed.ac.uk](mailto:Linda.Bauld@ed.ac.uk)  
13 <sup>4</sup> UK Centre for Tobacco and Alcohol Studies, Division of Epidemiology and Public  
14 Health, University of Nottingham, United Kingdom; [j.britton@outlook.com](mailto:j.britton@outlook.com),  
15 \* Correspondence: [bacham@mrc.gm](mailto:bacham@mrc.gm); Tel : +2202329405

16 **Abstract:** Introduction. Second-hand smoke is associated with more than 1.2 mil-  
17 lion deaths per year among non-smokers. Smoking in public places is prohibited  
18 in The Gambia but there is no information on the level of exposure to second-  
19 hand smoke among adolescents and adults 15-64 years. The aim of this study was  
20 to assess the level and predictors of exposure to second-hand smoke in public  
21 places and compliance with smoke free regulations in The Gambia. Methods. A  
22 population-based survey was conducted in an established Health and Demo-  
23 graphic Surveillance System (HDSS). A total of 4547 participants (15-64  
24 years) from households within the Farafenni HDSS were interviewed at their  
25 homes but only 3533 were included in our analysis. Factors associated with expo-  
26 sure to second-hand smoke in public places were assessed by three different mul-  
27 tivariable regression models. Results. Exposure to tobacco smoke in public places  
28 was high (66.1%, ), and higher in men (79.9%) than women (58.7%). Besides being  
29 male, less education, lower household income, urban residence and not aware of  
30 smoke free regulations were strongly associated with exposure to second-hand  
31 smoke. Conclusion. Despite existing smoke-free regulations, reported exposure  
32 to second-hand smoke remains high in public places in The Gambia. The Ministry  
33 of Health should continue to strengthen their advocacy and sensitization pro-  
34 grams to ensure smoke free regulations are fully implemented. Some population  
35 subgroups are at a higher risk of exposure and could be targeted by interventions;  
36 and settings where these subgroups are exposed should be targeted by enforce-  
37 ment efforts.

**Citation:** Cham, B.; Mdege, N.D.;  
Bauld, L.; Britton, J.; Umberto  
D’Alessandr Exposure to second-  
hand smoke in public places and  
barriers to the implementation of  
smoke free regulations in the Gam-  
bia: a population-based survey. *Int.*  
*J. Environ. Res. Public Health* **2021**,  
*x*, <https://doi.org/10.3390/xxxxx>

Received: date  
Accepted: date  
Published: date

**Publisher’s Note:** MDPI stays neu-  
tral with regard to jurisdictional  
claims in published maps and insti-  
tutional affiliations.



**Copyright:** © 2021 by the authors.  
Submitted for possible open access  
publication under the terms and  
conditions of the Creative Commons  
Attribution (CC BY) license  
([http://creativecommons.org/li-  
censes/by/4.0/](http://creativecommons.org/licenses/by/4.0/)).

**Keywords:** Smoking; second-hand smoke; The Gambia; smoke-free regulations

**1. Introduction**

Exposure to second-hand tobacco smoke (hereafter referred to as second-hand smoke) is associated with more than 1.2 million deaths per year worldwide among non-smokers; about 11 million disability-adjusted life years (DALYs) are lost due to second-hand smoke exposure,



45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80

with 61% of them occurring among children[1]. In 2004, 40% of children, 33% of men and 35% of women in 192 countries were exposed to second-hand smoke indoors [1]. There is no safe level of exposure to tobacco smoke [2]. Article 8 of the World Health Organization Framework Convention on Tobacco Control (WHO FCTC) recommends eliminating exposure to tobacco smoke in workplaces and public places, both indoor and outdoor, and this includes public transport [2]. Article 20 (Research, surveillance, and exchange of information) stresses the importance of surveillance for assessing the tobacco epidemic, monitoring tobacco use and the related prevention policies [2].

The Gambia's 2017 Global Youth Tobacco Survey (GYTS) found that 61.8% of students aged 13-15 years were exposed to second-hand smoke in enclosed public places [3]. A recent survey in secondary schools also reported a high prevalence of second-hand smoke exposure among young people aged 12-20 years in The Gambia: 59.2% for indoor public places and 61.4% for outdoor public places [4]. There are several regulations and policies on tobacco control in The Gambia, including the Prohibition of Smoking in Public Places Act [5]. The Tobacco Control Act 2016 was adopted in December 2016 and it came into force in July 2018 [6]. The Tobacco Control Regulations 2019 [7], which supports the enforcement of the Tobacco Control Act 2016 was signed and gazetted in August 2019. Based on the existing international treaties and local regulations, the country has the legal obligation and instruments to protect the public from exposure to tobacco smoke. However, despite the numerous achievements in tobacco control as highlighted above, there are still gaps especially on the implementation of smoke free regulation in The Gambia. Article 20 of the WHO FCTC (Research, surveillance, and exchange of information) stresses the importance of surveillance for assessing the tobacco epidemic, monitoring tobacco use and the related prevention policies including those for the protection of the public and non-smokers, and the level of compliance and the barriers to the implementation of the tobacco control regulations [2]. The aim of this study was to address this critical evidence gap by assessing, for the first time, the level and predictors of exposure to second-hand smoke in indoor and outdoor public places among both adolescents and adults, and compliance with smoke free regulations in The Gambia.

81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95

## 2. Materials and Methods

### 2.1. Study setting and design

The study was conducted in The Gambia from 20<sup>th</sup> January to 26<sup>th</sup> March 2020 as part of the Farafenni Health and Demographic Surveillance System (HDSS) which was established in October 1981 by the Medical Research Council Unit The Gambia (MRCG) and is updated every four months [8]. The HDSS is in the North Bank Region about 170 km from the capital, Banjul; includes the provincial town of Farafenni and 22 peri-urban and 40 rural villages; and covered a population of 59,910 individuals at the time of data collection. Farafenni town and the peri-urban villages were included in July 2002 and designated the *urban demographic surveillance area* [8]. All residents are part of the HDSS except the military staff, students attending boarding schools and staff of the Ministry of Health from other regions but in the North Bank Region on postings. Entry to the HDSS is through initial enumeration, birth, or immigration [8].



---

96 Individuals 15-64 years old were randomly selected from the HDSS data-  
97 base and interviewed at home.

98 *2.2. Study instrument/questionnaire*

99 In addition to the HDSS routine information (age, gender, education,  
100 marital status, ethnicity, and residence), household income, current and  
101 past smoking status, awareness of the tobacco control regulations, expo-  
102 sure to second-hand smoke and several other variables (not included in  
103 this analysis) were also collected. We adapted questions from the tobacco  
104 component of the WHO STEP survey questionnaire [9], as well as the  
105 Global Adults Tobacco Survey (GATS) questionnaire [10]. The question-  
106 naire was translated into Wolof and Mandinka, the two main local lan-  
107 guages and then pilot tested. Interviews were carried out by trained field  
108 workers and data electronically captured using the Research Electronic  
109 Data Capture (REDCap; <https://projectredcap.org/software/>) with tab-  
110 lets and mobile phones programmed with the data collection app.  
111



112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161

### 2.3. Sampling framework and sample size

The target population, i.e. 15-64 years old individuals, represented about half (51.1%) of the total population covered by the Farafenni HDSS. The sample size was estimated assuming the prevalence of smoking was 20%, based on a national prevalence of 15.6% [11], a margin of error of 5%, and a design effect of 1.5 [12]. In addition, since the response rate from the last national STEP survey in The Gambia was 79% and to adjust for non-response and increase the power, the sample size was increased by 20%. This resulted in a sample size of 4,613 which was increased to 6,000 to allow for emigration and non-response.

A multistage sampling technique was applied using the three level of residence within the HDSS (urban, semi urban, and rural). At the first stage, a total of 42 urban, semi and rural villages were randomly selected. Stage two involved sampling of households within the selected villages. There are 3727 compounds and 6037 households within the Farafenni HDSS. The average household size is 10 and on average there are five 15-64 years old individuals per household; a total of 2532 households were randomly selected from the HDSS database. In stage three, one to five people between the ages of 15-64 years were initially randomly selected per household proportional to the size of the household, for a total of 6000. Because of seasonal migration, some selected individuals were unavailable. Therefore, one month after starting the data collection process, 2856 additional individuals were randomly selected, resulting a total sample of 8856 individuals. We were only able to interview 4547 but our analysis was restricted to 3343 participants with complete information.

### 2.4. Exposure and outcome measures

The outcome variable was self-reported exposure to second-hand smoke in public places. For indoor exposure, a participant was defined as exposed if he/she responded *yes* to the question: *Did anyone smoke in your presence, inside any enclosed public place, other than your home (such as school, shops, restaurants, police stations, hospitals, public institutions/offices, shopping malls, video clubs/movie theatres etc) during the past 30 days?* A similar question relating to outdoor public places was asked: *Did anyone smoke in your presence, at any outdoor public place other than your home (such as car park, football field, markets, bantabas, wrestling arenas) during the past 30 days?*

The exposure variables were demographic variables including age, gender, ethnicity, residence, education, marital status, and household income. We also included awareness of regulation on smoking in public places which was assessed by asking the following question: *Are you aware of the law prohibiting smoking in public places in The Gambia?*

### 2.5. Data management and analysis

Immediately after the data collection, the data were cleaned, coded, and analysed using Stata 16. Descriptive statistics was used to describe the socio-demographic characteristics of the respondents, the level of compliance with smoke free regulations and the prevalence of exposure in both indoor and outdoor public places. We estimated the prevalence of exposure to second-hand smoke in both indoor and outdoor public places by our variables of interests as highlighted in section 2.4. Multivariable logistic regression analysis was performed to determine the socio-demographic factors associated with exposure to second-hand smoke in



162  
163  
164  
165  
166  
167  
168

indoor and outdoor public places. We did not conduct multiple imputation for the missing data and hence conducted complete case analysis in all our regression models. Because of inherent difference in policy implications, three separate models for indoor, outdoor, and indoor and/or outdoor public places were run. We used confidence intervals and p-values in both the descriptive and multivariable regression analysis to determine level of significance.

169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180

### 3. Results

Out of the 8856 sampled participants, 4,547 responded to the survey while 16 declined to participate. The rest were either absent at the time of our visits or were not reached due to the early suspension of fieldwork activities because of COVID 19. Our analysis was restricted to 3,343 individuals with complete information on the key variables. The mean age was  $34.6 \pm 14.5$  years; more than two thirds (64.9%) of the respondents were females, 42.0% lived in urban areas, 48.9% in rural areas and 9.1% in semi-urban areas (Table 1). Most of the participants (64.9%) were married and only 10.7% had senior secondary or a higher level of education (Table 1).



**Table 1.** Characteristics of study participants (n=3343).

<b>Variable</b>	<b>N</b>	<b>%</b>
<b>Age group</b>		
15-24	1101	32.9
25-34	679	20.3
35-44	616	18.4
45-54	514	15.4
55-64	433	13.0
<b>Mean age</b>	34.6±14.5	
<b>Sex</b>		
Male	1175	35.2
Female	2168	64.9
<b>Residence</b>		
Rural	1634	48.9
Semi-urban	305	9.1
Urban	1404	42.0
<b>Marital status</b>		
Single (Never married)	989	29.6
Married	2222	66.5
Divorced/Separated	38	1.1
Widowed	94	2.8
<b>Ethnicity</b>		
Wollof	1569	46.9
Fula	716	21.4
Mandinka	920	27.5
Others	138	4.1
<b>Education</b>		
No formal education	284	8.5
Lower Basic	288	8.6
Upper basic/Jun Sec	384	11.5
Senior Sec/College/Uni	358	10.7
Madrassa	655	19.6
Quranic school	1349	40.4
Senegalese	25	0.8
<b>Household income (in Dalasi)</b>		
Under10,000	640	19.1
10,000-19,999	508	15.2
20,000-29,999	383	11.5
30,000-39,999	266	8.0
40,000-50,000	722	21.6





Do not know	824	24.7
<b>Current smoking</b>		
Yes	177	5.2
No	3166	94.7
<b>Awareness of smoking in public regulation</b>		
Aware	2095	62.7
Not aware	1248	37.3

182  
183  
184  
185  
186  
187  
188  
189  
190


The prevalence of smoking was 5.2% and significantly higher among males compared with females (14.7% vs 0.2%,  $p < 0.001$ ). Awareness of the regulation that prohibits smoking in public places was moderate, 62.7% and significantly higher among current smokers than in non-smokers (85.3% vs 61.4%,  $p < 0.001$ ). Half of the participants (50.9%) were not aware of the regulation that protects children from exposure to tobacco, including sale of tobacco to minors. Almost all respondents (98.1%) supported the law that prohibits smoking in public places.

191

### 3.1. The prevalence of exposure to second-hand smoke

192  
193  
194  
195  
196  
197  
198  
199  
200  
201

The prevalence of exposure to second-hand cigarette smoking was higher ) in outdoor (61.3%) than indoor (52.8%) public places. Overall, 66.1% of the participants were exposed, with little variation by age (Figure 1). Levels of exposure were significantly higher in males than in females in all settings (indoor exposure 63.7% in males vs 46.9% in females,  $p < 0.001$ ; outdoor exposure: 75.6% in males vs 53.5% in females,  $p < 0.001$ ). In addition, exposure was higher among urban than rural residents, both indoors (46.1% vs 57.7%,  $p < 0.001$ ) and outdoors (54.1% vs 66.8,  $p < 0.001$ ) (Figure 2).



202

NB: The vertical lines on top of the bars represent the 95% Confidence Intervals


203

204

**Figure 1.** Exposure to second-hand smoke by age.

205





206

207 NB: The vertical lines on top of the bares represent the 95% Confidence Intervals

208

209

**Figure 2.** Exposure to second-hand smoke by residence.

210

211

### 3.2. Factors associated with exposure to second-hand smoke in indoor and outdoor public places

212

213

214

215

216

217

218

219

220

221

222

223

224

Gender, ethnicity, residence, education, income level and level of awareness of the regulation on smoking in public were strongly associated with exposure to second-hand smoke in public places (Table 2). In model I (exposure to second-hand smoking in indoor public places), males were significantly more likely to be exposed than females (AOR=1.78 95% CI: 1.51-2.10). Moreover, urban residents were almost three times more likely to be exposed than rural residents (AOR=2.68, CI: 2.24-3.21). Those with higher education and those from households with higher income were less likely to be exposed in indoor public places (Table 2). The odds of exposure to second-hand smoke were significantly higher among those unaware of smoking regulations (AOR =1.46. CI:1.25-1.71). Models II (outdoor public places) and III (both indoor and outdoor) produced similar results.



225 **Table 2.** Factors associated with exposure to second-hand smoke in public places (n=3343).

	<b>Model I</b>		<b>Model II</b>		<b>Model III</b>	
	<b>Indoor public places</b>		<b>Outdoor public places</b>		<b>Indoor and/or outdoor public places</b>	
	<b>OR</b>	<b>AOR</b>	<b>OR</b>	<b>AOR</b>	<b>OR</b>	<b>AOR</b>
<b>Age group</b>						
15-24	Reference	Reference	Reference	Reference	Reference	Reference
25-34	1.20(1.03-1.41) *	1.05 (0.81-1.35)	1.12(0.95-1.31)	1.04(0.81-1.35)	1.13(0.95-1.34)	1.05 (0.81-1.37)
35-44	1.02(0.86-1.21)	0.78(0.59-1.04)	0.92(0.77-1.10)	0.78(0.59-1.04)	0.92(0.77-1.11)	0.74(0.55-1.00)
45-54	1.18(0.98-1.41)	0.87 (0.64-1.18)	1.06(0.87-1.28)	0.90(0.67-1.23)	1.13(0.93-1.37)	0.88(0.64-1.20)
55-64	1.18(0.97-1.44)	0.95 (0.69-1.31)	0.97(0.79-1.19)	0.90(0.65-1.25)	1.11(0.90-1.37)	0.95(0.68-1.33)
<b>Sex</b>						
Female	Reference	Reference	Reference	Reference	Reference	Reference
Male	2.02(1.78-2.29) ***	1.78(1.51-2.10) ***	2.62(2.29-3.01) ***	2.40(2.02-2.86) ***	2.78(2.40-3.21) ***	2.49 (2.08-2.99) ***
<b>Residence</b>						
Rural	Reference	Reference			Reference	Reference
Semi-urban	1.13(0.92-1.38)	0.95(0.72-1.24)	1.10(0.89-1.35)	1.09(0.82-1.45)	1.02(0.82-1.26)	1.04 (0.78-2.25)
Urban	1.44(1.28-1.63)	2.68 (2.24-3.21) ***	1.55(1.37-1.76) ***	2.34(1.96-2.80) ***	1.23(1.08-1.40) **	1.88 (1.56-2.25) ***
<b>Marital status</b>						
Single	Reference	Reference	Reference	Reference	Reference	Reference
Married	1.41(1.23-1.61)***	1.13(0.89-1.45)	1.27(1.11-1.46) ***	1.05(0.81-1.35)	1.44(1.25-1.66) ***	1.09 (0.84-1.41)
Divorced/Separated	1.76(1.05-2.96)*	1.64(0.79-3.39)	1.50(0.89-2.52)	1.00(0.50-2.02)	1.92(0.91.13-3.23) *	1.48(0.73-2.99)



	Model I Indoor public places		Model II Outdoor public places		Model III Indoor and/or outdoor public places	
	OR	AOR	OR	AOR	OR	AOR
Widowed	2.26(1.55-3.31) ***	1.13(0.67-1.91)	1.98(1.37-2.87) ***	1.28(0.77-2.15)	2.38(1.64-3.46) ***	1.35(0.80-2.27)
<b>Ethnicity</b>						
Wollof	Reference	Reference	Reference	Reference	Reference	Reference
Fula	1.15(0.98-1.34)	1.28(1.06-1.55)**	1.02(0.88-1.20)	1.16(0.96-1.41)	1.02(0.87-1.19)	1.16(0.95-1.41)
Mandinka	0.44(0.38-0.51)***	0.43 (0.36-0.52) ***	0.72(0.63-0.84) **	0.73(0.60-0.89) **	0.56(0.48-0.66) ***	0.62(0.50-0.75) ***
Others	0.49(0.36-0.66) ***	0.53(0.36-0.78) ***	0.56(0.40-0.77) **	0.60(0.40-0.90) **	0.48(0.34-0.68) ***	0.57(0.37-0.88) *
<b>Education</b>						
No formal education	Reference	Reference	Reference	Reference	Reference	Reference
Lower Basic	0.55(0.41-0.75) ***	0.62 (0.43-0.91) *	0.64(0.47-0.88) **	0.67(0.46-0.98) *	0.61(0.44-0.84) **	0.67(0.45-0.99) *
Upper basic/Jun Sec	0.57(0.43-0.75) ***	0.58(0.41-0.83) **	0.70(0.52-0.93) *	0.68(0.48-0.96) *	0.61(0.44-82) ***	0.60(0.42-0.88) **
Senior Sec/College/Uni	0.65(0.49-0.86) ***	0.63(0.44-0.89) **	0.60(0.45-0.81) **	0.57(0.40-0.81) **	0.57(0.42-0.78) ***	0.58(0.40-0.84) **
Madrassa	0.96(0.74-1.24)	1.08(0.79-1.47)	0.91(0.75-1.21)	0.92(0.67-1.28)	1.03(0.79-1.35)	1.01(0.74-1.40)
Quranic school	1.16(0.92-1.47)	1.27(0.96-1.69)	0.95(0.75-1.21)	0.99(0.75-1.31)	1.09(0.85-1.39)	1.10(0.83-1.46)
Senegalese	0.88(0.43-1.82)	0.70(0.30-1.66)	0.62(0.29-1.36)	0.59(0.251.42)	0.70(0.32-1.57)	0.71(0.29-1.72)
<b>Household income</b>						
Under10000	Reference	Reference	Reference	Reference	Reference	Reference
10000-19999	1.17(0.94-1.45)	1.54 (1.20-1.99) ***	1.04(0.84-1.30)	1.36(1.05-1.75)*	1.06(0.85-1.32)	1.24(0.96-1.60)
20000-29999	0.88(0.70-1.13)	1.16(0.89-1.53)	0.84(0.66-1.07)	1.02(0.78-1.35)	0.83(0.65-1.07)	0.96(0.73-1.28)



	Model I Indoor public places		Model II Outdoor public places		Model III Indoor and/or outdoor public places	
	OR	AOR	OR	AOR	OR	AOR
30000-39999	0.90(0.69-1.18)	1.09(0.80-1.49)	0.69(0.52-0.91)**	0.72(0.53-1.00)	0.67(0.50-0.90) **	0.68(0.49-0.95) *
40000-500000	0.86(0.70-1.04)	0.82(0.65-1.04)	0.80(0.66-0.98)*	0.71(0.56-0.90)**	0.73(0.60-0.90) ***	0.70(0.55-0.90) ***
Do not Know	0.84(0.70-1.02)	0.93(0.75-1.17)	0.80(0.66-0.98)	0.82(0.65-1.02)	0.88(0.37-2.15)	0.73(0.58-0.92) **
<b>Awareness of smoking in public regulation</b>						
Aware	Reference	Reference	Reference	Reference	Reference	Reference
Not aware	1.68(1.49-1.90) ***	1.46(1.25-1.71) ***	1.76(1.56-1.99) ***	1.47(1.26-1.73) ***	1.80(1.59-2.04) ***	1.48(1.26-1.73) ***

226 \*p<0.05, \*\*p≤0.01, \*\*\* p≤0.001. OR =Unadjusted odds ratio. AOR= Adjusted odds ratio.

227 Reference= reference category in regression n analysis



228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275

#### 4. Discussion

Exposure to second-hand smoke in public places, despite the existing tobacco control regulations that prohibit smoking in public, remains high in The Gambia. Awareness of such regulations was moderate while it was low for the regulations protecting children. This can serve as a barrier to the implementation of tobacco control regulations. Interestingly, most respondents, including smokers, supported the law against smoking in public places and other smoke free regulations. The factors strongly associated with exposure to second-hand smoke in both indoor and outdoor public places included male gender, lower education, lower household income, urban residence, ethnicity, and lack of awareness of smoke-free regulations.

Exposure to tobacco smoke was high and, although previous surveys focused on different age groups [3, 4], exposure was similarly high in this study. A cross sectional survey in secondary schools in The Gambia found that 59.2% of 12- 20 years old were exposed to second hand-smoke in indoor public places and 61.4% in outdoor public places [4]. Preliminary analysis of the 2017 GYTS revealed that 61.8% of students (13-15 years old) were exposed to second hand tobacco smoke in enclosed public places [3]. In our study, women were much less exposed in both indoor and outdoor public places, a finding that is similar to results from South Africa where females were less likely to be exposed in all locations except their homes [13]. In India, males were also more likely to be exposed in public places, but females were more likely to be exposed at home [14].

In a previous study in The Gambia, higher maternal education and living with parents was significantly associated with lower risks of exposure while higher level of education among fathers was associated with increased odds of exposure in public places [4], suggesting highly educated mothers protected their children from exposure to cigarette smoke. In Farafenni, higher education was also a strong protector of being exposed to second-hand smoke, suggesting increasing public awareness is a good strategy. Our findings also concur with a study in Mongolia and China where among non-smoking women, those with a higher education were less likely to be exposed to second-hand tobacco smoke compared to those with a lower level of education [13].

Individuals with a lower income were more likely to be exposed to tobacco smoke in both indoor and outdoor public places. Lower socioeconomic status was associated with exposure to second-hand smoke in households in the Republic of Korea [15]. A study on exposure to domestic second-hand smoke in 15 low-and-middle-income countries reported that exposure to second-hand smoke, in most countries, was higher among the socioeconomically disadvantaged [16]. These findings on socio-economic status, fairly consistent in several studies, including the current research, reflect the greater risk of second-hand smoke exposure among the less affluent because they are more likely to work or live in settings with smokers and, for public places, poorer enforcement of existing regulations.



276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323

Women were less likely to be exposed to second-hand smoke than men. As suggested in the literature, women have a very important role in protecting themselves, children, and other family members from exposure to cigarette smoke [17]. They may also be less likely to visit public places (such as workplaces) where smokers are present. Women can be powerful advocates for health protection measures and should be involved in tobacco control at local and national level. The Ministry of Health of The Gambia has a long-standing tradition of using organised women groups and traditional communicators (most of them women) in transmitting key health messages to the population. This could be further strengthened and integrated as part of the national campaign to sensitise communities on the dangers of second-hand tobacco smoke and the tobacco control regulations.

Education was a strong protector of being exposed to second-hand smoke which may suggest that measures raising awareness may have an important role to play, although it is also likely that more highly educated individuals are less likely to be living or working with other smokers. The results of the multivariable regression analysis show that non-smokers unaware of the regulations are more likely to be exposed to cigarette smoke in public places. Our findings also suggest that those with a lower level of education are more likely to be exposed to cigarette in public places compared with participants with a high school or college/university education. As people's knowledge of and belief in the adverse health effects of tobacco use increase, the likelihood of using tobacco decreases and their support for protective policies increases [18]. Members of a well-informed society are also more likely to take up and support policies for a tobacco-free environment. Therefore, the Ministry of Health and partners should strengthen their community sensitisation programs to raise awareness of smoke free regulations.

#### *Strengths and limitations*

This is the first study on exposure to second-hand smoke in public places among adults in The Gambia. The study has identified both the population subgroups at a higher risk of exposure and settings where these subgroups are being exposed and thus can be used to guide policy and enforcement efforts. These research findings can be used to inform the development of effective policies and strategies for the control of tobacco and non-communicable diseases in The Gambia, which can be adapted by other low- and middle-income countries. They can also be used as a baseline to monitor over time compliance with the smoke free regulations.

There are some limitations, including the low male response rate, mainly due to high seasonal migration, which may have underestimated overall exposure as men are more likely to smoke and to be exposed to environmental tobacco smoke. The early suspension of field activities due to the COVID 19 pandemic is another limitation as the final sample size was smaller than expected. In addition, the study was based on survey interview data and all the information on exposure to second-hand tobacco smoke was self-reported. An objective measurement of exposure to



324 second-hand smoke, e.g., urine and/or blood analysis to assess cotinine  
325 (nicotine metabolite) was not performed. Participants can therefore un-  
326 derreport or over report their exposure status. Future studies in The Gam-  
327 bia and in other African countries should explore objective measurement  
328 of exposure.

## 329 **5. Conclusion**

330 Our findings illustrate that exposure of both smokers and non-smok-  
331 ers to second-hand smoke is high in The Gambia despite smoke free reg-  
332 ulations, indicating there are still gaps in tobacco control, especially on  
333 the implementation of smoke free laws. People aware of the smoke free  
334 regulations and women with a higher level of education were less likely  
335 to be exposed to second-hand smoke in both indoor and outdoor public  
336 places. The Ministry of Health and its partners, including the members of  
337 the Tobacco Control Committee, should continue to strengthen their ad-  
338 vocacy and sensitisation programs to ensure smoke free regulations are  
339 fully implemented and enforced.

340 **Author Contributions:** All authors contributed to the formulation of the re-  
341 search project; Bai Cham implemented the project, conceptualised the paper, an-  
342 alysed the data and wrote the first draft of the manuscript; Umberto D'Alessandro  
343 supervised all project activities. All the authors reviewed and approved the final  
344 version of the manuscript.

345 **Funding:** Funding for this research comes from the Tobacco Control Capac-  
346 ity Programme (grant MR/P027946/2) supported by UK Research and Innovation  
347 (UKRI) with funding from the Global Challenges Research Fund.

348 **Institutional Review Board Statement:** The study was conducted according  
349 to the guidelines of the Declaration of Helsinki and approved by the Gambia Gov-  
350 ernment/Medical Research Council Unit The Gambia at the London School of Hy-  
351 giene and Tropical Medicine Joint Ethics Committee (Ref 17898, 13/11/2019).

352 **Informed Consent Statement:** Informed consent was obtained from all sub-  
353 jects involved in the study. Participants aged 18 years and above were asked to  
354 provide written informed consent. For minors (15-17 years), parents written as-  
355 sent was obtained in addition to the informed consent.

356 **Data Availability Statement:** Data reported in this study are from an ongo-  
357 ing longitudinal study, but the data is not yet publicly available.

358 **Acknowledgments:** We thank the Farafenni HDSS staff and all the field  
359 workers who took part in the data collection process. Our special thanks and grati-  
360 tude go to the MRCG data management team more especially Lamin Saidy, Si-  
361 mon Donkor and Yusupha Njie for creating the database for the field work and  
362 their support throughout the data collection process. We also thank the Farafenni  
363 HDSS Data Manager, Sawar Golam for his support during the sampling process.

364 **Conflicts of Interest:** The authors declare no conflict of interest.  
365





366

367 **References**

- 368 1. Oberg, M., et al., *Worldwide burden of disease from exposure to second-hand smoke: a*  
369 *retrospective analysis of data from 192 countries*. *Lancet*, 2011. **377**(9760): p. 139-46.
- 370 2. World Health Organization, *WHO framework convention on tobacco control*. 2003: Geneva:  
371 World Health Organization.
- 372 3. Centers for Disease Control and Prevention, *The Gambia 2017 Global Youth Tobacco Survey*  
373 *(GYTS) Factsheet*. 2017.
- 374 4. Jallow, I.K., J. Britton, and T. Langley, *Prevalence and factors associated with exposure to*  
375 *secondhand smoke (SHS) among young people: a cross-sectional study from the Gambia*.  
376 *BMJ Open*, 2018. **8**(3): p. e019524.
- 377 5. The Ministry of Health and Social Welfare, *The Gambia. Prohibition of smoking (Public*  
378 *Places) Act*. 1998.
- 379 6. Campaign for Tobacco-free Kids. *Tobacco control laws: Legislation by country*. 2018; Available  
380 from: <https://www.tobaccocontrol.org/legislation/country/gambia/summary>.
- 381 7. Ministry of Health, *Tobacco Control Regulations 2019*, M.o. Health, Editor. 2019.
- 382 8. Jasseh, M., et al., *Health & demographic surveillance system profile: Farafenni health and*  
383 *demographic surveillance system in The Gambia*. *International journal of epidemiology*,  
384 2015. **44**(3): p. 837-847.
- 385 9. World Health Organization, *World Health Organization STEPS surveillance manual: The WHO*  
386 *STEPwise approach to chronic disease risk factor surveillance*. 2005.
- 387 10. World Health Organization. *Global Adult Tobacco Survey (GATS) Core Questionnaire with*  
388 *Optional Questions. Version 2.0*. 2010; Available from:  
389 [https://www.who.int/tobacco/surveillance/en/tfi\\_gats\\_corequestionnairewithoptionalques](https://www.who.int/tobacco/surveillance/en/tfi_gats_corequestionnairewithoptionalquestions_v2_FINAL_03Nov2010.pdf)  
390 [tions\\_v2\\_FINAL\\_03Nov2010.pdf](https://www.who.int/tobacco/surveillance/en/tfi_gats_corequestionnairewithoptionalques).
- 391 11. Cham, B., et al., *Burden of hypertension in The Gambia: evidence from a national World*  
392 *Health Organization (WHO) STEP survey*. *International journal of epidemiology*, 2018. **47**(3):  
393 p. 860-871.
- 394 12. WHO, *WHO STEPS surveillance manual: The WHO STEPwise approach to chronic disease risk*  
395 *factor surveillance*. 2005.
- 396 13. Ngobese, S.P., et al., *Non-Smoker's Exposure to Second-Hand Smoke in South Africa during*  
397 *2017*. *Int J Environ Res Public Health*, 2020. **17**(21).
- 398 14. Tripathy, J.P., *Secondhand smoke exposure at home and public places among smokers and*  
399 *non-smokers in India: findings from the Global Adult Tobacco Survey 2016-17*. *Environ Sci*  
400 *Pollut Res Int*, 2020. **27**(6): p. 6033-6041.
- 401 15. Park, S., et al., *Socioeconomic disparities in household secondhand smoke exposure among*  
402 *non-smoking adolescents in the Republic of Korea*. *Global Public Health*, 2017. **12**(9): p. 1104-  
403 1121.
- 404 16. Nazar, G.P., et al., *Socioeconomic Inequalities in Secondhand Smoke Exposure at Home and*  
405 *at Work in 15 Low- and Middle-Income Countries*. *Nicotine Tob Res*, 2016. **18**(5): p. 1230-9.
- 406 17. Wipfli, H.L. and J.M. Samet, *Second-hand smoke's worldwide disease toll*. *Lancet*, 2011.  
407 **377**(9760): p. 101-2.
- 408 18. Drope, J., et al., *The Tobacco Atlas*. Atlanta, GA: American Cancer Society. Inc. and Vital  
409 Strategies, 2018.

410