



OPEN ACCESS

Automatic dispensing cabinets and governance of controlled drugs: an exploratory study in an intensive care unit

Valentina Lichtner ,^{1,2} Mirela Prgomet,² Peter Gates,² Bryony Dean Franklin^{3,4}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/ejpharm-2020-002552>).

¹Department of Management, Leeds University Business School, University of Leeds, Leeds, UK

²Centre for Health Systems and Safety Research (CHSSR), Australian Institute of Health Innovation, Macquarie University, Sydney, New South Wales, Australia

³Department of Practice and Policy, UCL School of Pharmacy, University College London, London, UK

⁴Centre for Medication Safety and Service Quality, Pharmacy Department, Imperial College Healthcare NHS Trust, London, UK

Correspondence to

Dr Valentina Lichtner, School of Pharmacy, University College London, London, UK; v.lichtner@leeds.ac.uk

Received 7 October 2020

Revised 22 March 2021

Accepted 23 March 2021

Published Online First

11 May 2021

EAFP Statement 2: Selection, Procurement and Distribution.



© European Association of Hospital Pharmacists 2023. Re-use permitted under CC BY. Published by BMJ.

To cite: Lichtner V, Prgomet M, Gates P, et al. *Eur J Hosp Pharm* 2023;**30**:17–23.

ABSTRACT

Background Governance of controlled drugs (CDs) in hospitals is resource intensive but important for patient safety and policy compliance.

Objectives To explore whether and how storing CDs in an automated dispensing cabinet (ADC) in a children's hospital intensive care unit (ICU) contributes to the effectiveness and efficiency of CD governance.

Methods We conducted a mixed-methods exploratory study, comprising observations, interviews and audits, 3 months after ADC implementation. We observed 54 hours of medications activities in the ICU medication room (with 42 hours of timed data); interviewed nurses (n=19), management (n=1) and pharmacy staff (n=3); reviewed 6 months of ICU incident reports pertaining to CD governance; audited 6 months of CD register data and extracted logs of all ADC transactions for the 3 months following implementation. Data analysis focused on four main CD governance activities: safekeeping/controlling access, documenting use, monitoring, and reporting/investigating.

Results Nurses and pharmacists perceived spending less time on CD governance tasks with the ADC. The ADC supported CD governance through automated documentation of CD transactions; 'blind counts'; automated count discrepancy checks; electronic alerts and reporting functionalities. It changed quality and distribution of governance tasks, such as removing the requirement for 'nurses with keys' to access CDs, and allowing pharmacists to generate reports remotely, rather than reviewing registers on the ward. For CDs in the ADC, auditing and monitoring appeared to be ongoing rather than periodic. Such changes appeared to create positive reinforcing loops. However, the ADC also created challenges for CD governance. Most importantly, it was not suitable for all CDs, leading to workarounds and parallel use of a safe plus paper registers.

Conclusions ADCs can significantly alter CDs governance in clinical areas. Effects of an ADC on efficiency and effectiveness of governance tasks appear to be complex, going beyond simple time savings or more stringent controls.

BACKGROUND

Effective governance of controlled drugs (CDs) in hospitals is necessary for the safety of staff and patients,^{1–3} compliance with the law^{4–6} and CD stewardship.⁷ We define CD governance in hospitals as the structures and processes to organise, secure, monitor and account for, supply and use of CDs. CD governance may involve trade-offs⁸ (such as making CDs more secure but less accessible in

emergencies), and may result in limitations, such as error-prone, time-consuming record keeping via paper-based systems.^{9,10} CD governance may particularly affect clinical areas with high use of CDs, such as intensive care units (ICUs).¹¹

Automated dispensing cabinets (ADCs) have been introduced in hospital wards to make CD governance more effective and efficient.^{10,12,13} However, evidence on the effects of ADCs on CD governance is limited, as most research on ADCs implementations is not focused specifically on CDs or their impact on CD governance.^{14–17} A few studies suggest that ADCs save nurses or pharmacists' time on CDs governance, for example, due to the elimination of (some or all of) the daily counts of CDs, or because of less time spent updating paper registers.^{10,18,19} Reports of initiatives to identify CD diversion in US hospitals^{20,21} show how ADC may contribute to monitoring and investigations. However, the literature also suggests that issues remain with CD governance after ADC implementations.²²

We aimed to explore whether and how an ADC contributes to the effectiveness and efficiency of governance of CDs and through what mechanisms.

METHODS

We carried out an exploratory, mixed-methods study of an ADC in a paediatric ICU, with a focus on CDs.

The setting

The study was conducted in a children's hospital in Sydney, Australia. The hospital implemented an ADC (Omnice²³) in the ICU on 26 March 2019.

Study design

We applied both quantitative methods (structured observations, audits of registers and data extracted from ADC) and qualitative methods (unstructured observations, interviews, reviews of incident reports). The audits were able to capture CD governance data before and after ADC implementation retrospectively, while the other methods were limited to a post-implementation period. We considered medications as CDs if they were treated as such on the ICU concerned for the purpose of governance.

Observations

Observations, structured and unstructured, were carried out by one researcher (VL) in the ICU medication room about 3 months after ADC implementation (June–July 2019). All activities carried

out in the medication room were included. Structured observations were recorded using the Work Observation Method By Activity Timing (WOMBAT 3.0²⁴) tool, which automatically timestamped recorded data. Unstructured observations were captured as field notes. ADC training sessions, where nurses received training as 'super users', were also observed.

Interviews

Qualitative semi-structured interviews were conducted by one researcher (VL) with ICU nurses to gather their perceptions and experiences with using the ADC. All nurses present on the ward on the interview days were invited to participate and could do so individually or as a group. We also sought to interview pharmacy staff, implementation team and hospital management for their views. Sampling was purposive. With consent, interviews were audio-recorded and transcribed; otherwise, the researcher took concurrent field notes.

Incidents

We (VL, PG) carried out a retrospective review of patient safety incidents voluntarily reported through the hospital incident reporting system. All ICU incident reports relating to medications for a period of about 3 months before and 3 months after ADC implementation (January–June 2019) were reviewed.

Audits of controlled drugs registers and ADC transaction logs

Retrospective audits of entries documented in ICU CD paper registers, about 3 months before and 3 months after ADC implementation (January–June 2019) were carried out by two researchers (VL, PG). As a control, we also audited registers from the oncology and orthopaedics wards.

Data extracted from the registers included: date and time; drug name, strength, formulation; amounts (received, given, discarded, balanced); discrepancies (box 1) identified by staff as annotated in registers and number of asterisks to signal amendments to records. We also captured missing amounts. For the denominator, we recorded the number of CD doses administered or discarded.

Two researchers (MP, VL) conducted a retrospective audit of all transactions recorded by the ADC for 3 months following implementation (26 March–30 June 2019). Data extracted included: date and time; drug name, strength, formulation;

transaction types (issue, return, waste, cycle count); amounts (issued, discarded, counted, returned) and discrepancies by type (eg, cycle count discrepancy).

Data analysis

The quantitative data analysis was conducted using descriptive statistics in SPSS (V.23)²⁵ for the WOMBAT data and Excel for the audit data. For the CD register data, we assessed number and frequency of CD discrepancies, calculation errors and other inaccuracies in documentation. The analysis was performed per ward, per type of medication (single doses vs multidose bottles), before and after ADC implementation and as a comparison across wards. For the ADC transaction logs, we calculated the number of ADC-generated discrepancies and time to resolution of discrepancies. For all transactions of CDs removed from the ADC, we matched the corresponding 'partial waste' data and calculated time to waste (the time from removing a medication dose from the ADC and documenting partial waste). For the WOMBAT data, we mapped the corresponding ADC transaction log data to obtain details about the type of medication (CD or non-CD) involved in the recorded activities and calculated frequency and duration of activities for each medication type.

Qualitative data from field notes and interview transcripts were first coded inductively by one researcher (VL) using NVivo (V.12)²⁶ to capture aspects related to systems (eg, ADC, safe and registries), medications (CDs and non-CDs), activities and participants' views as well as the wider context of ICU medication work. Incidents were also analysed thematically in a spreadsheet. Initial analysis was then independently verified by two researchers (MP and PG), with disagreements discussed and resolved via consensus. We then focused the analysis on CD governance. We identified four main activities of CD governance (table 1) in the data and local/national policy⁶ and used this as a framework to analyse mechanisms supporting or hindering CD governance, integrating qualitative and quantitative data within this framework.

RESULTS

Table 2 provides details of the data included in the study. We structure the findings around the main governance activities described in table 1. The focus is on the use of ADC for these activities, compared with a safe plus registers when appropriate. Table 3 provides a summary of the findings.

Safekeeping and controlling access

Prior to ADC implementation, CDs were kept in a locked safe, with CDs in need of refrigeration kept in an unlocked fridge. CD supply and use were documented in paper registers ('CD books').

The ADC was implemented in a new locked medication room created in the ICU, accessible with personal smartcards. Only registered staff members could access the ADC, by fingerprint or individual username and password. CDs were stored in the ADC in drawers which required the fingerprints of two authorised staff for access. However, some CDs could not be stored in the ADC (CDs belonging to patients, those to be discarded and liquid CDs in multidose bottles). Thus, a safe was also placed in the medication room, to store these CDs, leading to parallel use of ADC plus safe for storing CDs. The safe was also needed in case of ADC breakdown.

One CD (clobazam liquid) that needed to be refrigerated was stored in the ADC fridge. The fridge was locked but did not require two fingerprints for opening, thus technically this CD

Box 1 Definition of controlled drug discrepancy

A discrepancy is a difference between a medication balance in the register and the physical balance of that medication in stock in the ward.

Discrepancies with liquid medications in bottles may also be associated with the unaccounted space in syringes ('dead space'). Each syringe would take approximately 0.2–0.4 mL of liquid extra to the intended dose. Over a number of doses taken from the same bottle of medication, this loss from 'dead space' can accumulate to a considerable amount and generate a discrepancy between the balance documented in the register and the actual volume of liquid in the bottle.

Other causes of discrepancies include calculation errors, missing data (eg, a spill not documented), wrong counts of remaining stock and potential diversions of medications.

Discrepancies can be either losses (less medication in stock than the recorded balance) or gains (more in stock than the recorded balance).

Table 1 Activities for the governance of controlled drugs (CDs) in hospital wards

CD governance activity	Description
Safekeeping/Controlling access	<ul style="list-style-type: none"> ▶ CDs (including those owned by patients) stored separately from other medications, in a cabinet or metal safe, kept locked when not in use. ▶ Access to CD cabinets' keys strictly controlled. ▶ Access to CDs restricted to staff members authorised to administer them. ▶ Accesses witnessed by a staff member. ▶ Expired, unusable or unwanted CDs stored in the safe pending destruction.
Documenting transactions	<ul style="list-style-type: none"> ▶ Records kept of CD transactions in a drug register. ▶ Records of transactions signed by two staff members present during the activity (eg, nurse and witness). ▶ Transactions recorded in the CD register included: removal and replacing of CDs from the storage unit; discarding any unused portion of the CD and destruction of CDs.
Monitoring	<ul style="list-style-type: none"> ▶ Audits of CDs in stock performed to confirm records are meeting policy requirements and to detect any possible misappropriation. ▶ The recorded CDs balance checked daily against the physical balance in the cabinet and safe, by a registered nurse/midwife with a witness and documented in the register. ▶ Expired, unusable or unwanted CDs, pending destruction, are included in the routine stock checks.
Reporting and investigating	<ul style="list-style-type: none"> ▶ CDs discrepancies recorded and reported to the hospital management and authorities. ▶ Investigations conducted to explain the discrepancies.

could be accessed by any authorised user accessing the ADC fridge without a witness. Transactions of the CD in the fridge had to be recorded in paper registers. This required a workaround on the ADC that created potential for errors, as shown in a reported incident (Id7). However, this was considered an improvement on previous storage arrangements, as the ADC provided a level of controlled access that was previously absent.

... now is better in the sense that the fridge is locked. [Before the ADC], [...] if [the CD] needed to be in a fridge it would just be in the fridge [...], there was no real, a way to do it, we've got ways to do it now which is exciting. (id4)

Access to the safe required a key, usually held by senior nurses (known locally as 'access nurses'). However, other nurses held the key when the access nurse was away or busy. Nurses reported finding the keys to the safe as time consuming and delaying access to medications for patients. The ADC removed the need for keys and for the presence of specific staff members—any authorised nurse and witness were sufficient to access CDs in the ADC.

However, the ADC only allowed one transaction at a time, while CDs in the safe could be accessed by multiple nurses at the same time. Some nurses reported having to wait to access medications in the ADC, when in use. However, we observed nurses queuing to access the ADC rarely (online supplemental appendix 1). In addition, it was possible (and observed) for a nurse to access the ADC when a previous user had not 'exited' the transaction and therefore to use their login.

Documenting use

Removing (or returning) CDs from (or to) the ADC required entering data in the ADC documenting both transaction and remaining levels of stock. The ADC automatically recorded date, time and staff members involved (through their fingerprints), as well as the patient name, prescriber name and the CD as selected on screen. Additional data entry was required to add any discarded amounts, and remaining levels of stock. In contrast, a CD could be removed from the safe but documentation in the paper register could be delayed and possibly forgotten.

In case of incorrect data entry of remaining stock levels, the ADC alerted the users by inviting a recount, thus contributing to preventing discrepancies due to counting errors.

If you happen to put in like fentanyl 30 and there's 29 ... It actually alerts you. [...] previously you would have just read 30 and three days later someone would have picked it up. [...] this way, I think, it prevents those kinds of incidents (id6-9)

We found that removing a CD from the ADC (also including, when necessary, preparing the medication for administration) took two nurses between ~1 min (66 s) and ~12 min (748 s); the corresponding task performed with the safe and registry took from ~1.5 min (98 s) to ~8 min (472 s) (table 4). However, nurses perceived removing and documenting CDs from the ADC to be less time consuming than with safe and registers.

Table 2 Overview of methods and data type

Methods and data type	Description
Ethnographic observations; field notes	About 54 hours of observations were completed over 14 days in the period 6 June–24 July 2019. Periods of observations were of 2 hours 30 min on average (min 1 hour to max 7 hours). Three one-to-one sessions of nurse ADC super-user training were also observed.
Timed observations with WOMBAT; structured tasks and time stamps in spreadsheet	About 42 hours of observations were timed using the WOMBAT tool. Timed observations took place over 13 days between 6 June and 24 July 2019, average duration of individual observation session: 2 hours 23 min (range 1–47 hours).
Interviews; transcripts of audio recordings and field notes	Two 2-hour 'drop-in sessions' were organised to facilitate interviews with ICU nurses: 19 nurses participated. Interviews took place with ADC back-office pharmacist, ADC trainer, Deputy Director of Pharmacy, Clinical Program Director.
Incidents reports; structured and free text data in spreadsheet	We received a total of 171 incidents that had been reported in ICU in relation to medications in the period January–June 2019, 109 of these pertaining to use/supply/governance of CDs.
Audits of controlled drug registries; structured data in spreadsheet	We audited 6 CDs registers (1231 pages) in ICU, 5 registers (1120 pages) in oncology and 9 registers (1996 pages) in orthopaedics.
ADC transaction logs; structured data in spreadsheet	We received logs of all recorded ICU ADC transactions for the period March–June 2019.
ADC, automated dispensing cabinets; CDs, controlled drugs; ICU, intensive care unit; WOMBAT, Work Observation Method By Activity Timing (a structured data collection tool).	

Table 3 How an automated dispensing cabinet (ADC) may support or challenge governance of controlled drugs (CDs) in hospital wards

CD governance activity	ADC support	ADC challenge
Safekeeping/Controlling access	Access control by fingerprinting/username and password Double lock drawers (two users/fingerprints to access) No requirement for keys	Not all CDs 'storable' in the ADC. Parallel use of ADC and safe and registers required. Access to CD in the fridge only partially controlled, leading to a hybrid system of ADC for storage and paper registers for documentation. Workaround needed to give access to CD in the fridge and alert nurses of the different workflow. Parallel use of ADC by different users, for different transactions, not supported, leading to potential queues to access medications.
Documenting transactions	Documentation synchronous to removal/supply/count of CDs Automatically recorded data for most transactions, through fingerprinting and item selection on screen Automated alerts of potentially incorrect remaining stock balances recorded Users perception of efficiency and completeness of documentation	Reduced flexibility in amending records compared with paper. Returning CDs to the ADC more complex than with safe and registers.
Monitoring	Counts of stock levels blind to documentation of balance Remote monitoring potentially in real-time, through reporting functionalities and remote access to ADC data Users perception of efficiency and greater control	Data recording only partially automated; some data entered by users may not be accurate (eg, reason for discarding a CD), thus affecting monitoring activities.
Reporting and investigating	Easier to identify discrepancies near the time of their occurrence, making it easier to investigate and resolve Investigations involving interviewing staff can be limited to staff that had access to the ADC in the period of interest	Access to CDs in the fridge by unregistered users (while others are still logged in), not detected by the ADC, is possible. Possible risk of illusion of completeness of records.

Compared with paper registers, the ADC restricted flexibility for amending records, such as documenting later discarded amounts, or returning an unused CD to the ADC. A workaround had to be devised for these tasks using the function 'miscellaneous returns'. We observed how some nurses had difficulties completing these tasks on the ADC, and similarly they reported:

... when you're taking [a CD], [...] if you forget to enter the waste, [...] in the book, there will be an empty space and then you might think of it. [...] Even with ADC you have to come back sometimes, but sometimes you tend not to. Like it happened to me once, and then it's a bit tricky [...] I don't know how to do that. (id10-13)

Analysis of the ADC transaction logs showed that for a total of 2632 CDs removed from the ADC, 64% had waste recorded (online supplemental appendix 2—table A) and 81% of these had waste recorded at the time of removing the medication. Comparable data for registers were not available.

Nurses and pharmacy staff commented on how documentation in the ADC, compared with handwritten data in registers, was more legible and complete. In our audits of paper ICU CD registers, we found about 5% of doses annotated with asterisks or with some inaccuracies in documented amounts. Comparable

data patterns were also found in registers on control wards (online supplemental appendix 2—table B).

Monitoring

Monitoring for CD losses was done by: daily counts of available stock against documented balances; periodic audits of registers against stock in the safe and generation of ADC reports.

With the ADC, CD counts were 'blind', that is, the nurses did not know the recorded balance on the system before counting items in drawers. The opposite was the case for counts of CDs in the safe—the level of stock documented was in view while counting and may have influenced the stock check.

[in CDs counts with the ADC] I feel that people actually check more than just writing the book up... ... (id6-9)

Nurses reported daily counts of CDs stock to be faster with the ADC than with the safe and registers: '*ridiculously quicker... So that's really good*'. (id14-20)

... [with the books] you've got to write out every medication and everything. With [the ADC] you can just bang, bang, bang, fingers, and you go through and it's done. It's just so much quicker. (id21)

Table 4 Time on specific tasks with controlled drugs (CDs) in the medication room, using automated dispensing cabinet (ADC) and safe with registers

		Time (s)				
		Number of tasks	Mean	Minimum	Maximum	Median
ADC	Issue	41	360	66	748	331
	Return/Waste	1	108	108	108	108
	Count	1	632	632	632	632
	Other	4	301	195	358	325
Safe	Issue	25	242	98	472	208
	Return/Waste	0
	Count	2	483	469	496	483
	Other	2	225	48	401	225
Both ADC and safe	Issue	1	625	625	625	625

The data exclude tasks related to the CD in the fridge. Times presented in this table must be interpreted with caution. Samples are too small to be considered valid. Tasks performed with ADC are not directly comparable to the corresponding tasks performed with the safe and registries. The medications involved were different (unit doses in the ADC, bottles in the safe) and times of preparation may be significantly different for the different types of medication. The numbers of CDs stored in the ADC and safe were different—potentially affecting CD counts.

The ADC allowed pharmacy staff to monitor CDs in the ADC remotely by generating reports. For example, regular reports were generated on discrepancies with the CDs in the ADC. It also enabled monitoring from a distance that the CD counts had been done in the ward as expected.

... we have more real time reports. So every day at 10:00 am [the ADC] will look at the last 24 hours [...] and sends the ICU Manager a list of discrepancies generated on the ward for her to monitor.
... (id1)

Reporting and investigating

Discrepancies about CDs in the ADC were identified through daily reports and generally resolved within a day, as confirmed by transaction logs analysis (a range of 30 s to 21 hours). Nurses had to document in the ADC a reason for rectifying a discrepancy, but usually entered a generic 'error'.

Discrepancies with CDs in the safe tended to become apparent at the time of pharmacy audits or when the ward unexpectedly ran out of a CD otherwise documented as still available. Explaining these discrepancies involved reviewing all records since the bottle had been opened for any errors, and assessing whether the loss could be explained by dead space in syringes. Such losses could cumulate over large numbers of doses. The process was reported to be very time consuming for pharmacists.

In our audits, we found 54 discrepancies (losses or gains) for CDs in the safe, documented in CD registers in the ICU during a 6-month period (online supplemental appendix 2—table C). As staff told us, and our audits show, discrepancies were more frequent for medications in multidose liquids than for single doses.

The ADC could also be used to generate a list of those who had accessed the specific CD involved in a discrepancy, and thus

limit investigations to these staff instead of questioning everyone on shift.

DISCUSSION

The ADC affected CD governance tasks in the ICU, through a variety of mechanisms. It afforded concurrent removal (or return) and documentation of CDs; it removed the need to control keys to a safe; afforded automated checks (eg, fingerprints) and electronic interaction with users (eg, questioning counts); and allowed real time, remote access to data (individual transactions and aggregate data) through the generation of reports. Nurses' counts were 'blind' and nurses were found to 'check more' when counting remaining level of stock. Pharmacists monitored discrepancies remotely, rather than reviewing registers on the ward (or only doing the latter for CDs in the safe); auditing and monitoring was thus ongoing rather than periodic. These changes, combined, appeared to create positive reinforcing loops (figure 1). However, the ADC only supported governance of CDs in single doses (eg, vials, tablets); a safe and registers were still needed for governance of other CDs. Workarounds were also needed to handle exceptions, such as the single refrigerated drug. Difficulties with the ADCs in relation to multidose liquids (not exclusively CDs) and refrigerated drugs have also been reported elsewhere.¹⁸

Nurses and pharmacists perceived CD governance to be more efficient with the ADC than with safe and registers. This is consistent with nurses' perceptions and time and motion studies in other clinical contexts.^{10 18 19 27} We found the question on whether the ADC saves time on CD governance, compared with safe and registers, requires a nuanced answer. The ADC eliminated some time-consuming tasks




Figure 1 Example of how effects of an automated dispensing cabinet (ADC) on governance of controlled drugs (CDs) may combine. The different effects of using the ADC (such as nurses 'checking more' before recording remaining level of CD stock) are related to (or contribute to) other effects (eg, to making discrepancies easier to monitor and investigate).

(eg, transcribing books) but also created work (eg, entering data on screen to resolve discrepancies) and new roles (eg, ADC trainer and ‘super users’). The ADC changed the quality and distribution of tasks, and created new tasks (eg, pharmacists generating reports), as often happens with technology.^{28 29} Time patterns also changed, with monitoring and fixing discrepancies becoming ongoing. The effects extended beyond the ICU, to pharmacy and management dealing with CDs incidents, affecting their use of time. The ADC eliminated frustrating CD governance inefficiencies and this possibly led to perception of saving time.³⁰

The ADC made CDs safekeeping and access controls more stringent. In another setting, similar ADC technology was associated with fewer misappropriations of medications.¹³ However, the literature shows that diversion occurs also when ADCs are in place.^{3 21} In the setting we studied, greater quantities than prescribed could theoretically be removed from the ADC, not documented, and the refrigerated CD could be removed by any registered user accessing other medications in the fridge without a witness. Discarded CDs are also a recognised risk for diversion,^{3 5} especially in paediatric settings,²¹ which is not directly affected by use of an ADC or safe.

This study is unique in its focus on CD governance. Our analysis however did not include CD governance in prescribing or administration,⁵ as we did not observe these activities. We did not collect demographic data on the nurses who participated in the interviews; however, from the interviews it was clear that some were new to the ward, while others had senior roles. We were not able to compare the ADC to a baseline pre-implementation. We timed and compared tasks with ADC and safe, but our samples were too small to support significance testing.

CONCLUSION

Although creating new challenges, through varied mechanisms, an ADC has the potential to significantly alter CDs governance in clinical areas. But its effects on efficiency and effectiveness seem to be more nuanced and complex beyond

merely producing time savings in governance tasks or more stringent controls.

Twitter Valentina Lichtner @VLichtner

Acknowledgements We thank the hospital ADC implementation team for valuable assistance with research governance, access to wards and participants and extraction of ADC transaction logs. The study could not have been done without their contribution. We thank the nurses of the paediatric intensive care unit and control wards for their participation in this study. We thank Professor Johanna Westbrook, Centre for Health Systems and Safety Research, Australian Institute of Health Innovation, for her insightful comments and advice during the study.

Contributors VL conceived of the study. VL, BDF, MP and PG contributed to study design. MP provided specific input into the observational techniques. VL coordinated data collection and carried out observations and interviews. PG contributed to audits. VL led the analysis of the qualitative data. MP led the quantitative analysis. VL and PG analysed the incident reports. VL drafted the manuscript. All authors contributed to the interpretation of results and provided review and approval of the final manuscript.

Funding The study was funded by a National Health and Medical Research Council Partnership Projects Grant (APP1094878: 2015–2020). VL received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie-Sklodowska Curie Grant Agreement number 740131. This paper represents independent research supported by the NIHR Imperial Patient Safety Translational Research Centre.

Disclaimer The views expressed are those of the authors and not necessarily those of the funders, NHS, the NIHR or the Department of Health and Care.

Competing interests BDF received a grant from Omnicell to conduct a literature review of the evidence base for ADCs in the UK, unrelated to the research described in the present paper. All other authors have no competing interests to declare.

Patient consent for publication Not required.

Ethics approval The hospital research ethics committee approved the study. Interview participants gave written informed consent. ICU staff were informed that observations were taking place and an information poster placed near the medication room. Interviews, observation data and audits were anonymised at the point of data collection. Incidents and ADC log data were deidentified by the hospital prior to transfer to investigators

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. Aggregate data that could be made available have been provided as tables in supplementary files. Our ethics and local approvals preclude making other data, such as interview data, publicly available.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iD

Valentina Lichtner <http://orcid.org/0000-0003-3956-3743>

REFERENCES

- 1 Alroy-Preis S, Daly ER, Adamski C, et al. Large outbreak of hepatitis C virus associated with drug diversion by a healthcare technician. *Clin Infect Dis* 2018;67:845–53.
- 2 Hellinger WC, Bacalis LP, Kay RS, et al. Health care-associated hepatitis C virus infections attributed to narcotic diversion. *Ann Intern Med* 2012;156:477–83.
- 3 Papparella SF. A tale of waste and loss: lessons learned. *J Emerg Nurs* 2016;42:352–4.
- 4 CQC. *The safer management of controlled drugs — annual update 2017: care quality Commission*, 2018.
- 5 Brummond PW, Chen DF, Churchill WW, et al. ASHP guidelines on preventing diversion of controlled substances. *Am J Health Syst Pharm* 2017;74:325–48.

Key messages

What is already known on this subject

- ⇒ Effective governance of controlled drugs (CDs) in hospitals is important for policy compliance and for patient and staff safety.
- ⇒ CD governance through paper-based systems is error-prone and time-consuming.
- ⇒ Automated dispensing cabinets (ADCs) can be used in hospital wards to store and dispense CDs, and for CD governance tasks.

What this study adds

- ⇒ This study describes the effects of implementing a ward-based ADC on four main CD governance areas: safekeeping/controlling access, documenting use, monitoring and reporting/investigating.
- ⇒ It highlights a variety of mechanisms through which the use of an ADC both supports and challenges governance of CDs.
- ⇒ It suggests that an ADC overall improves efficiency of CD governance tasks compared with the use of paper registers, but that these effects are nuanced and complex.

- 6 NSW Health. Policy Directive: Medication Handling in NSW Public Health Facilities [PD2013_043] NSW Government, 2013. Available: https://www1.health.nsw.gov.au/pds/ActivePDSDocuments/PD2013_043.pdf
- 7 Sandbrink F, Uppal R. The time for opioid stewardship is now. *The Joint Commission Journal on Quality and Patient Safety* 2019;45:1–2.
- 8 NICE. *Controlled drugs: safe use and management. NG46*. London: National Institute for Health Care Excellence, 2016.
- 9 Mounsey A, Franklin B, Langfield B. Automated CD management-Its introduction in a UK Hospital pharmacy. *Hospital Pharmacist* 2008;15:424–8.
- 10 Portelli G, Canobbio M, Bitonti R. The impact of an automated dispensing system for supplying narcotics in a surgical unit: the experience of the National cancer Institute Foundation of Milan. *Hospital Pharmacy* 2018.
- 11 Krzyżaniak N, Pawłowska I, Bajorek B. Review of drug utilization patterns in NICUs worldwide. *J Clin Pharm Ther* 2016;41:612–20.
- 12 Epstein RH, Dexter F, Gratch DM, et al. Controlled substance reconciliation accuracy improvement using near real-time drug transaction capture from automated dispensing cabinets. *Anesth Analg* 2016;122:1841–55.
- 13 Hendarto A, Harding AM, Kant JA, et al. Misappropriation of medications and equipment in the emergency department. *Australas Emerg Care* 2019;22:92–6.
- 14 Batson S, Herranz A, Rohrbach N, et al. Automation of in-hospital pharmacy dispensing: a systematic review. *Eur J Hosp Pharm* 2021;28:58–64.
- 15 Tsao NW, Lo C, Babich M, et al. Decentralized automated dispensing devices: systematic review of clinical and economic impacts in hospitals. *Can J Hosp Pharm* 2014;67:138–48.
- 16 Boyd A, Chaffee B. Critical evaluation of pharmacy automation and robotic systems: a call to action. *Hospital Pharmacy* 2018.
- 17 Zheng WY, Lichtner V, Van Dort BA, et al. The impact of introducing automated dispensing cabinets, barcode medication administration, and closed-loop electronic medication management systems on work processes and safety of controlled medications in hospitals: a systematic review. *Res Social Adm Pharm* 2021;17:832–41.
- 18 Rochais E, Atkinson S, Guilbeault M, et al. Nursing perception of the impact of automated dispensing cabinets on patient safety and ergonomics in a teaching health care center. *J Pharm Pract* 2014;27:150–7.
- 19 Roman C, Poole S, Walker C, et al. A 'time and motion' evaluation of automated dispensing machines in the emergency department. *Australas Emerg Nurs J* 2016;19:112–7.
- 20 Epstein RH, Gratch DM, McNulty S, et al. Validation of a system to detect scheduled drug diversion by anesthesia care providers. *Anesth Analg* 2011;113:160–4.
- 21 Brenn BR, Kim MA, Hilmas E. Development of a computerized monitoring program to identify narcotic diversion in a pediatric anesthesia practice. *Am J Health Syst Pharm* 2015;72:1365–72.
- 22 Anyanwu C, Egwim O. The prevalence and determinants of controlled substance discrepancies in a level I trauma Hospital. *Am Health Drug Benefits* 2016;9:128–33.
- 23 Omnicell. Available: <https://www.omnicell.com/>
- 24 Prgomet M, Thomas J, Westbrook J. *Work observation method by activity timing: a guide to the use of WOMBAT 3.0*. Sydney: Centre for Health Systems and Safety Research, Australian Institute of Health Innovation, Macquarie University, 2020.
- 25 IBM. SPSS statistics -. Available: <https://www.ibm.com/products/spss-statistics>
- 26 QSR International. NVivo. Available: <https://www.qsrinternational.com/nvivo-qualitative-data-analysis-software/home>
- 27 Schwarz HO, Brodowy BA. Implementation and evaluation of an automated dispensing system. *Am J Health Syst Pharm* 1995;52:823–8.
- 28 Vikkelso S. Subtle redistribution of work, attention and risks: electronic patient records and organisational consequences. *Scandinavian Journal of Information Systems* 2005;17:10.
- 29 Poissant L, Pereira J, Tamblyn R, et al. The impact of electronic health records on time efficiency of physicians and nurses: a systematic review. *J Am Med Inform Assoc* 2005;12:505–16.
- 30 Flaherty MG. *A watched pot: how we experience time*. NYU Press, 1999.

Appendix 1. WOMBAT data

Summary of data

Data Collection Period - 13 days between 6 June and 24 July

Number of Individual Sessions - 18

Average Duration of Individual Session - 2hours 23mins (Range 1 hour to 4 hours)

Total Hours of Observation - 42 hours 44 mins

Total Tasks Recorded – 686

Inactive Time – 28hours 53mins

Inactive Tasks – 324

Active Time – 15hours 13mins*

Active Tasks - 362

*includes multitasking time (i.e., medication room being used by two or more individuals for separate purposes)

Demand on medication room

There were 56 instances of ‘room multitasking’ (e.g. multiple nurses working on different medications’ tasks).

There were 18 instances where someone was observed to “look in the room and leave” (as if looking for someone or deciding to come back later because the ADC was in use).

There were 9 instances where someone was observed “waiting inside” the medication room (to access the ADC).

Overall time on tasks in medication room

	Time (Seconds)				
	Number of Tasks	Mean	Minimum	Maximum	Median
ADC	207	165	12	1526	78
Safe	29	258	48	496	210
Both ADC & Safe	5	571	177	1282	523
Other*	121	85	5	553	48

*Other includes for example open shelves or EMM.

Time on drug specific tasks in medication room

Times presented in the tables below must be interpreted with caution. Samples are too small to be considered valid. Tasks performed with ADC are not directly comparable to the corresponding tasks performed with the safe and registries. The medications involved were different (unit doses in the ADC, bottles in the safe) and times of preparation may be significantly different for the different types

of medication. The numbers of CDs stored in the ADC and safe were different – potentially affecting CD counts. CD counts always include both ADC and safe, and a ‘hybrid’ fridge item (stored in the fridge but counted with registries).

Regular drugs

		Regular Drugs				
		Number of Tasks	Time (Seconds)			
			Mean	Minimum	Maximum	Median
ADC	Issue	111	80	17	366	61
	Return/Waste	28	52	12	284	42
	Count	0				
	Other	14	335	47	1526	91

Controlled drugs (overall, excluding item in the Fridge)

		Controlled Drug				
		Number of Tasks	Time (Seconds)			
			Mean	Minimum	Maximum	Median
ADC	Issue	41	360	66	748	331
	Return/Waste	1	108	108	108	108
	Count	1	632	632	632	632
	Other	4	301	195	358	325
Safe	Issue	25	242	98	472	208
	Return/Waste	0
	Count	2	483	469	496	483
	Other	2	225	48	401	225
Both ADC & Safe	Issue	1	625	625	625	625

Tasks with both a regular and a controlled drug

		Both Regular and Controlled Drugs				
		Number of Tasks	Time (Seconds)			
			Mean	Minimum	Maximum	Median
ADC	Issue	7	349	72	636	300
	Return/Waste	0				
	Count	0				
	Other	0				
Safe	Issue	0				
	Return/Waste	0				
	Count	0				
	Other	0				
Both ADC & Safe	Issue	4	557	177	1282	385

Controlled drugs – one drug vs multiple drugs

Where the task involves removal of one drug (with no other transactions performed):

		One Drug & Single Transaction				
		Number of Tasks	Time (Seconds)			
			Mean	Minimum	Maximum	Median
ADC	Issue	2	148	111	184	148
Safe	Issue	6	190	98	244	202

Where the task involves removal of one drug, with additional tasks also performed (e.g. recording waste or preparing the drug):

		One Drug & Multiple Transactions				
		Number of Tasks	Time (Seconds)			
			Mean	Minimum	Maximum	Median
ADC	Issue	33	348	66	748	301
Safe	Issue	16	227	119	472	201

When the task involves removal of two or more drugs, with additional tasks also performed:

		Two or More Drugs & Multiple Transactions				
		Number of Tasks	Time (Seconds)			
			Mean	Minimum	Maximum	Median
ADC	Issue	6	496	341	679	486
Safe	Issue	3	431	381	461	451

Appendix 2 – Additional tables

*Analysis of ADC transaction logs***Table A – Controlled Drugs – Time from removing a dose from ADC ('issue') to documenting waste (ADC transaction logs, 26 March – 30 June)**

Drug (Controlled)	Issue Transactions	Waste Transactions		Waste Concurrent with Issue		Waste by Separate Transaction to Issue								
	n	n	% of Issued	n	% of Total Waste	n	% of Total Waste	Average Time	Range Min	Range Max	<10m	10m - 1hr	1hr - 24hr	>24 hr
Bupivacaine + Fentanyl	9	3	33.3%	3	100.0%	0	n/a	n/a	n/a	n/a				
Clobazam	150	52	34.7%	46	88.5%	6	11.5%	0:29:06	0:00:30	2:48:54	5		1	
Clonazepam	2	2	100.0%	1	50.0%	1	50.0%	0:01:57	0:01:57	0:01:57	1			
Diazepam	156	57	36.5%	48	84.2%	9	15.8%	0:22:46	0:00:20	3:18:01	8		1	
Fentanyl	640	390	60.9%	326	83.6%	64	16.4%	2:37:09	0:00:17	84:12:10	44	4	15	1
Ketamine	229	194	84.7%	110	56.7%	84	43.3%	4:17:27	0:00:23	84:12:55	10	10	63	1
Midazolam	765	593	77.5%	495	83.5%	98	16.5%	0:55:14	0:00:15	9:25:36	56	16	26	
Morphine Hydrochloride	75	49	65.3%	44	89.8%	5	10.2%	0:00:36	0:00:14	0:00:54	5			
Morphine Sulfate	325	237	72.9%	195	82.3%	42	17.7%	4:28:15	0:00:19	172:00:21	37	1	3	1
Morphine Sulfate MR	57	49	86.0%	43	87.8%	6	12.2%	0:01:30	0:00:18	0:06:27	6			
Oxycodone	32	25	78.1%	24	96.0%	1	4.0%	0:00:31	0:00:31	0:00:31	1			
Phenobarbital	31	17	54.8%	15	88.2%	2	11.8%	0:02:33	0:01:01	0:04:05	2			
Phenobarbitone	1	0	0.0%	0	n/a	n/a	n/a	n/a	n/a	n/a				
Propofol	157	12	7.6%	2	16.7%	10	83.3%	6:25:49	0:55:18	16:03:49		2	8	
Thiopentone	3	1	33.3%	1	100.0%	0	n/a	n/a	n/a	n/a				
Total *	2632	1681	63.9%	1353	80.5%	328	19.5%	2:40:26	0:00:14	172:00:21	175	33	117	3

* Not including the controlled drug in the fridge – accessed by generic 'fridge item' in the ADC and documented in registries

*Analysis of data from audits of registries of controlled drugs***Table B. Frequency of documentation inaccuracies in controlled drugs registries in the ICU and control wards (January – June 2019)**

Type of documentation inaccuracy	ICU n (%)	Oncology n (%)	Orthopaedics n (%)
	(doses*: 5783)	(doses*:2930)	(doses*:3750)
Asterisks representing corrections to the register	305 (5.3%)	132 (4.5%)	282 (7.5%)
Calculation errors identified by nurses or pharmacists	9 (0.2%)	3 (0.1%)	48 (1.3%)
Other (e.g. balance changed incorrectly)	3 (0.1%)	1 (0.0%)	9 (0.2%)
Missing amounts (given, discarded, balance)	47 (0.8%)	26 (0.9%)	20 (0.5%)

% are calculated on number of doses. * The number of doses is the number of entries documented in registries – i.e. number of times medications have been recorded as given (or not) to patients. The table does not include data about discrepancies, reported in a separate table.

Table C. CD discrepancies (losses and gains) in the ICU and control wards in the study period (January – June 2019)

	ICU (*) [pre/post ADC] n (%)	Oncology [no ADC in use] n (%)	Orthopaedics [no ADC in use] n (%)
	(doses*: 5783)	(doses*:2930)	(doses*:3750)
Discrepancies identified by nurses/pharmacists	54 (0.9%)	11 (0.4%)	37 (1.0%)

(*) Limited to discrepancies identified in registries (excluding discrepancies in ADC). % are calculated on number of doses. * The number of doses is the number of entries documented in registries – i.e. number of times medications have been recorded as given (or not) to patients.

*Comparative analysis of data from audits of controlled drugs registries and ADC transaction logs***Table D. CD discrepancies (losses and gains) in the ICU and control wards in the three months before and after ADC implementation (as recorded in registries and ADC transactions logs)**

	Jan-March 2019 (pre-ADC) *			April – June 2019 (post-ADC)		
	n of doses	Losses N (%)	Gains N (%)	n of doses	Losses N (%)	Gains N (%)
ICU						
Medications in bottles	1597	15 (0.94%)	1 (0.06%)	2273	30 (1.32%)	6 (0.26%)
Medications in unit doses	1792	2 (0.40%)	0	121 [in registries]	0	0
				7366 (*) [in ADC]	15	14
Oncology [no ADC in use]						
Medications in bottles	336	4 (1.19%)	0	409	5 (1.22%)	2 (0.49%)
Medications in unit doses	1045	0	0	1140	0	0
Orthopaedics [no ADC in use]						
Medications in bottles	1123	19 (1.42%)	1 (0.09%)	1031	14 (1.36%)	2 (0.19%)
Medications in unit doses	889	0	0	707	0	0

