



UNIVERSITY OF LEEDS

This is a repository copy of *Scaling behaviour of small-scale dynamos driven by Rayleigh–Bénard convection*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/174148/>

Version: Accepted Version

Article:

Yan, M, Tobias, SM and Calkins, MA (2021) Scaling behaviour of small-scale dynamos driven by Rayleigh–Bénard convection. *Journal of Fluid Mechanics*, 915. A15. ISSN 0022-1120

<https://doi.org/10.1017/jfm.2021.61>

© The Author(s), 2021. Published by Cambridge University Press. This article has been published in a revised form in *Journal of Fluid Mechanics*, <http://doi.org/10.1017/jfm.2021.61>. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Scaling behaviour of small-scale dynamos driven by Rayleigh-Bénard convection

M. Yan^{1*}, S.M. Tobias², and M.A. Calkins¹

¹Department of Physics, University of Colorado, Boulder, CO 80301, USA

²Department of Applied Mathematics, University of Leeds, UK

(Received xx; revised xx; accepted xx)

A numerical investigation of convection-driven dynamos is carried out in the plane layer geometry. Dynamos with different magnetic Prandtl numbers Pm are simulated over a broad range of the Rayleigh number Ra . The heat transport, as characterized by the Nusselt number Nu , shows an initial departure from the heat transport scaling of non-magnetic Rayleigh-Bénard convection (RBC) as the magnetic field grows in magnitude; as Ra is increased further, the data suggests that Nu grows approximately as $Ra^{2/7}$, but with a smaller prefactor in comparison with RBC. Viscous (ϵ_u) and ohmic (ϵ_B) dissipation contribute approximately equally to Nu at the highest Ra investigated; both ohmic and viscous dissipation approach a Reynolds number dependent scaling of the form Re^a scaling, where $a \approx 2.8$. The ratio of magnetic to kinetic energy approaches a Pm -dependent constant as Ra is increased, with the constant value increasing with Pm . The ohmic dissipation length scale depends on Ra in such a way that it is always smaller, and decreases more rapidly with increasing Ra , than the viscous dissipation length scale for all investigated values of Pm .

1. Introduction

Planetary and stellar magnetic fields are ubiquitous throughout the observable universe. These magnetic fields are thought to be actively generated by the convection-driven motion of electrically conducting fluid (Ossendrijver 2003; Jones 2011). Rayleigh-Bénard convection (RBC), consisting of a fluid layer contained between plane parallel boundaries, is a common system in which to study convection due to its simplicity, whilst retaining the primary physical features expected to be important in many natural systems (Meneguzzi & Pouquet 1989; Cattaneo 1999). Although RBC has been investigated in great detail with regard to electrically-insulating fluids, the influence of dynamo action on heat and momentum transport is less well understood. This study reports on numerical results of a broad parameter survey of RBC-driven dynamos.

Natural dynamos can be distinguished by the characteristic length scale of the self-generated magnetic field, relative to that of the forcing length scale. Large-scale dynamos generate magnetic fields that are both system-scale and forcing scale, whereas small-scale dynamos generate magnetic fields with typical length scales that are comparable with, or less than, the typical velocity length scale (Meneguzzi *et al.* 1981; Tobias 2019; Rincon 2019). Breaking the reflectional symmetry of the flow field via the Coriolis force, for instance, is known to be conducive to large-scale dynamo action (Parker 1955; Steenbeck *et al.* 1966; Moffatt 1970; Childress & Soward 1972; Calkins *et al.* 2015); such effects are likely important for the generation of the global-scale components of planetary and

* Email address for correspondence: ming.yan@colorado.edu

stellar magnetic fields. However, small-scale dynamos are also relevant, especially with regard to the Sun’s outer convective layer where intense small-scale field is generated.

Two global diagnostic quantities of central interest in convection are the rate of heat transport through the layer and the typical flow speed, as measured by the non-dimensional Nusselt number, Nu , and the Reynolds number, Re , respectively. For a fixed value of the thermal Prandtl number, $Pr = \nu/\kappa$ (where ν is the kinematic viscosity and κ is the thermal diffusivity), convective flow regimes depend on the non-dimensional Rayleigh number, Ra ; functional relationships of the form $Nu = f(Ra)$ and $Re = g(Ra)$ (where f and g denote generic functions) are sought. For heat transport with $Pr = O(1)$ in non-rotating systems, theory has suggested both a $Nu \sim Ra^{1/3}$ scaling from marginal stability analysis of the thermal boundary layer (Malkus 1954) and a $Nu \sim (Ra/Pr)^{1/2}$ scaling which assumes an ultimate regime in which the entire fluid layer becomes turbulent (Kraichnan 1962; Spiegel 1965); the former is independent of the fluid layer depth, whereas the latter is independent of diffusion coefficients (ν , κ). The convective ‘free-fall’ scaling of $Re \sim (Ra/Pr)^{1/2}$, thought to arise from a balance between nonlinear advection and the buoyancy force, and expected to be valid when $Re \gg 1$, is consistent with the $Nu \sim (Ra/Pr)^{1/2}$ heat transport scaling (e.g. Ahlers *et al.* 2009). Laboratory experiments and numerical simulations observe a $Nu \sim Ra^{2/7}$ scaling over a significant range in Ra (e.g. Castaing *et al.* 1989; Shraiman & Siggia 1990; Cioni *et al.* 1997), and a transition to a $Nu \sim Ra^{1/3}$ scaling at the largest values of Ra (e.g. Cheng *et al.* 2015). Two-dimensional numerical simulations find another transition to a still steeper scaling near $Ra \sim 10^{13}$, where $Nu \sim Ra^{0.35}$ is observed (Zhu *et al.* 2018). Scaling behaviour close to $Re \sim Ra^{1/2}$ has been observed in both low Prandtl number fluids (Vogt *et al.* 2018) and $Pr = O(1)$ fluids (Qiu & Tong 2001). Numerical simulations in a triply periodic geometry show both the $Nu \sim Ra^{1/2}$ and $Re \sim Ra^{1/2}$ scaling (Lohse & Toschi 2003), providing evidence that these ‘ultimate’ scalings are indeed relevant for RBC, and that the presence (or absence) of thermal and kinetic boundary layers dictates the observed scaling exponents.

The work of Meneguzzi & Pouquet (1989) showed that RBC acts as an efficient source of energy for dynamo action, provided the flow is driven sufficiently. Subsequent numerical investigations, both Boussinesq and compressible, have shown that magnetic field tends to be localized to the upwelling and downwelling regions (Cattaneo *et al.* 2003; Bushby & Favier 2014). A common belief is that small-scale non-rotating dynamos equilibrate when both the kinetic and magnetic energy are comparable to each other, a hypothesis that seems to be supported by numerical studies (Cattaneo *et al.* 2003; Haugen *et al.* 2004). However, the heat and momentum transport in RBC-driven dynamos remains largely unexplored; it is currently unknown what influence dynamo action has on the scaling behaviour of both Nu and Re with varying Ra .

Viscous dissipation plays a fundamental role in heat transport in non-magnetic RBC. In a statistically stationary state, the viscous dissipation ϵ_u is directly related to Nu (e.g. Chandrasekhar 1961). The scaling behaviour of Nu with Ra is therefore intimately connected with the scaling of ϵ_u , and therefore also with the scaling of Re . The non-dimensional Taylor microscale λ_u is often used to characterize the length scale at which viscous dissipation becomes dominant; the scaling behaviour of λ_u with Re is therefore thought to control the observed $Nu - Ra$ scaling (e.g. Grossmann & Lohse 2000). In RBC-driven dynamos, both viscous and ohmic dissipation are present. Clearly, the presence of ohmic dissipation provides an additional degree of freedom when determining heat transfer scaling laws (e.g. Zürner *et al.* 2016); understanding heat and momentum transport in RBC-driven dynamos therefore requires an understanding of both λ_u , and an analogous ohmic dissipation scale λ_B .

Two additional parameters that are important in dynamos are the magnetic Prandtl number, $Pm = \nu/\eta$ (where η is the magnetic diffusivity), and the magnetic Reynolds number $Rm = PmRe$. The magnetic Reynolds number characterizes the relative size of magnetic induction to magnetic diffusion. Planetary interiors (French *et al.* 2012; Pozzo *et al.* 2013) and liquid metal experiments (e.g. Cioni *et al.* 2000; Aurnou & Olson 2001) are characterized by $Pm \sim O(10^{-5})$, and typical values in the Sun range from $Pm \sim O(10^{-6})$ to $Pm \sim O(10^{-3})$ (Ossendrijver 2003). These physical values lead one to conclude that $Rm \ll Re$ in planets and stars. On the other hand, Pm can be as large as $O(10^{22})$ in protogalactic plasmas, in which case the opposite limit $Rm \gg Re$ occurs (Schekochihin *et al.* 2002a). Although natural dynamo systems have a widespread range of Pm , direct numerical simulation (DNS) studies are limited to relatively modest values of $10^{-2} \lesssim Pm \lesssim 10$, due primarily to limitations in accessing large values of Re and/or Rm (Sheyko *et al.* 2016; Schaeffer *et al.* 2017; Rincon 2019).

Simulations of isothermal, mechanically forced dynamos in triply periodic domains have yielded insight into the behaviour of (λ_u, λ_B) with varying Re and Rm , and the scaling behaviour of the magnetic and kinetic energy. Brummell *et al.* (2001) found a scaling of the form $\lambda_B \sim Rm^{-1/2}$ that arises when a balance between magnetic field generation and diffusion is present. A scaling law for magnetic energy (at small Rm), $E_{mag} \sim Re^{-1} Rm^{1/2}$, was also derived based on a balance between Lorentz force and the part of the viscous force that is induced by the magnetic field. Haugen *et al.* (2004) showed that, for a fixed value of Pm , the ratio of magnetic energy to kinetic energy (E_{mag}/E_{kin}) converges to a constant value as the Reynolds number increases. It was also found that the ratio of ohmic dissipation to viscous dissipation (ϵ_u/ϵ_B) converges for large Re , while the converged value is weakly influenced by Pm .

A phenomenological model has been proposed to address the saturation of the ratio of magnetic energy to kinetic energy (E_{mag}/E_{kin}) for large Pm (Schekochihin *et al.* 2002b; Tobias *et al.* 2013). The saturation process begins when the magnetic energy becomes comparable with the kinetic energy at the viscous scale. The magnetic field and velocity are modified scale-by-scale until, eventually, an equipartition between the two energies is reached at the integral scale. When Pm is large but $Pm < Re^{1/2}$, only a fraction of the equipartition is reached, and a saturated level of $E_{mag}/E_{kin} \sim Pm/Re^{1/2}$ is derived. For small Pm where $Pm \ll 1$, it is expected that the saturated level of the energy ratio becomes independent of Pm , and the energy ratio E_{mag}/E_{kin} approaches a constant value (Fauve & Pétrélis 2007).

The primary goal of the present study is to investigate the scaling behaviour of heat and momentum transport in RBC-driven dynamos, and the associated balances in the momentum and induction equations. The scaling of viscous and ohmic dissipation and their contribution to heat transport are analyzed. The influence of the magnetic Prandtl number on the length scales associated with the velocity and magnetic fields, as well as the ratio of magnetic energy to kinetic energy will also be discussed. In section 2 the governing equations and numerical methods are discussed; in section 3 the results of the numerical simulations are presented; and concluding remarks are given in section 4.

2. Governing equations and methods

We consider a fluid layer of depth H that is confined between plane parallel boundaries with temperature difference $\Delta T = T_{bot} - T_{top} > 0$, where T_{bot} and T_{top} are the temperatures of the bottom and top surfaces, respectively. The gravitational acceleration has constant magnitude g , and points perpendicular to the bottom boundary. The fluid has density ρ , kinematic viscosity ν , thermal expansion coefficient α , thermal diffusivity

κ , magnetic permeability μ and magnetic diffusivity η . The governing equations are non-dimensionalized using the layer depth H , the large-scale magnetic diffusion timescale H^2/η , and magnetic field scale $\mathcal{B} = \sqrt{\rho\mu\nu\eta}/H$. The equations are then given by

$$(\partial_t - Pm\nabla^2) \mathbf{u} = \mathbf{u} \times (\nabla \times \mathbf{u}) + Pm (\nabla \times \mathbf{B}) \times \mathbf{B} + \frac{RaPm^2}{Pr} \theta \hat{\mathbf{z}} - \nabla p, \quad (2.1)$$

$$(\partial_t - \nabla^2) \mathbf{B} = \nabla \times (\mathbf{u} \times \mathbf{B}), \quad (2.2)$$

$$\left(\partial_t - \frac{Pm}{Pr} \nabla^2 \right) \theta = -\mathbf{u} \cdot \nabla \theta, \quad (2.3)$$

$$\nabla \cdot \mathbf{u} = 0, \quad (2.4)$$

$$\nabla \cdot \mathbf{B} = 0, \quad (2.5)$$

where $\mathbf{u} = (u, v, w)$ is the velocity field, $\mathbf{B} = (B_x, B_y, B_z)$ is the induced magnetic field, θ is the temperature, p is the pressure and the Cartesian coordinate system is denoted by (x, y, z) .

The Rayleigh number (Ra), thermal Prandtl number (Pr) and magnetic Prandtl number (Pm) are defined as

$$Ra = \frac{g\alpha\Delta TH^3}{\nu\kappa}, \quad Pr = \frac{\nu}{\kappa}, \quad Pm = \frac{\nu}{\eta}. \quad (2.6)$$

The particular values used for the fluid properties, as specified by Pr and Pm , are determined by computational restrictions, and an interest in accessing dynamical regimes that are applicable to geophysical and astrophysical systems. Planetary interiors are characterized by $Pr \gg Pm$, with a ratio $Pr/Pm \approx 10^5$. In contrast, the Sun and other stars are composed of plasmas that typically have $Pr \leq Pm$. Both the $Pr/Pm < 1$ and $Pr/Pm > 1$ regimes are therefore of physical interest, though both are also computationally demanding. Extreme spatial resolutions are required to reproduce the wide separation of magnetic and velocity scales (Tobias *et al.* 2013). In this study, Pm is varied from 0.8 to 7, while for the majority of our cases Pr is fixed to unity; a set of simulations with $Pr = 0.05$ and $Pm = 1$ is also presented.

The mechanical boundary conditions are impenetrable and stress-free such that

$$w = \frac{\partial u}{\partial z} = \frac{\partial v}{\partial z} = 0 \quad \text{at} \quad z = 0, 1. \quad (2.7)$$

The thermal boundary conditions are isothermal,

$$\theta = 1 \quad \text{at} \quad z = 0, \quad \text{and} \quad \theta = 0 \quad \text{at} \quad z = 1. \quad (2.8)$$

The magnetic field is enforced to be vertical at the boundaries,

$$B_x = B_y = 0 \quad \text{at} \quad z = 0, 1. \quad (2.9)$$

Since the magnetic field is solenoidal, the above boundary conditions automatically imply that

$$\frac{\partial B_z}{\partial z} = 0 \quad \text{at} \quad z = 0, 1. \quad (2.10)$$

Note that although the boundary conditions on the magnetic field allow for the development of a non-zero horizontally averaged (mean) magnetic field, no appreciable mean field has been observed in the simulations reported here. As the system is non-rotating this is to be expected.

2.1. Energy relations

If we dot the momentum equation (2.1) with \mathbf{u} and volumetrically average the result we obtain

$$\partial_t \left[\frac{1}{2} \mathbf{u}^2 \right] = Pm [\mathbf{u} \cdot \mathbf{J} \times \mathbf{B}] + \frac{Ra Pm^2}{Pr} [w\theta] - Pm [\zeta^2], \quad (2.11)$$

where the square brackets $[\cdot]$ denote a volumetric average only (no time average) and the vorticity vector and the current density vector are denoted by $\zeta = \nabla \times \mathbf{u}$ and $\mathbf{J} = \nabla \times \mathbf{B}$, respectively. Similarly, by dotting the induction equation (2.2) with \mathbf{B} we obtain

$$\partial_t \left[\frac{1}{2} \mathbf{B}^2 \right] = - [\mathbf{u} \cdot \mathbf{J} \times \mathbf{B}] - [\mathbf{J}^2]. \quad (2.12)$$

Multiplying the kinetic energy equation (2.11) by $(1/Pm^2)$ and the magnetic energy equation (2.12) by $(1/Pm)$, and adding the results gives

$$\partial_t [e_{kin} + e_{mag}] = \frac{Ra}{Pr} [w\theta] - \frac{1}{Pm} [\zeta^2] - \frac{1}{Pm} [\mathbf{J}^2], \quad (2.13)$$

where we define the kinetic energy density and the magnetic energy density as, respectively,

$$e_{kin} = \frac{1}{2Pm^2} \mathbf{u}^2, \quad (2.14)$$

$$e_{mag} = \frac{1}{2Pm} \mathbf{B}^2. \quad (2.15)$$

The only requirement for the scalings of the energy densities is that their ratio differs by a non-dimensional factor of Pm . The scalings used above are essentially in viscous diffusion timescale units, and therefore facilitate comparison with RBC simulations (including those reported here). If we now time average equation (2.13) we obtain

$$0 = \frac{Ra}{Pr} \langle w\theta \rangle - \frac{1}{Pm} \langle \zeta^2 \rangle - \frac{1}{Pm} \langle \mathbf{J}^2 \rangle, \quad (2.16)$$

which simply states that the work done by the buoyancy force is exactly balanced by the combined effects of ohmic and viscous dissipation.

The energy balance relationship (2.16) can be put into a slightly more useful form by introducing the Nusselt number, Nu , which is defined as the ratio of total heat transfer (convective and conductive) to conductive heat transfer. In our non-dimensional units this becomes

$$Nu = 1 + \frac{Pr}{Pm} \langle w\theta \rangle. \quad (2.17)$$

Therefore, the energy balance becomes

$$\frac{Ra}{Pr^2} (Nu - 1) = \epsilon_B + \epsilon_u, \quad (2.18)$$

where we define the ohmic and viscous dissipation as

$$\epsilon_B = \frac{1}{Pm^2} \langle \mathbf{J}^2 \rangle, \quad \epsilon_u = \frac{1}{Pm^2} \langle \zeta^2 \rangle. \quad (2.19)$$

We note that, given our isothermal boundary conditions, an equivalent definition of the Nusselt number is given by

$$Nu = - \left. \frac{\partial \bar{\theta}}{\partial z} \right|_{z=0}. \quad (2.20)$$

where $\bar{\theta}$ is the horizontally and time averaged (mean) temperature. Multiplying the heat equation by θ and space-time averaging the resulting equation then gives another equivalent definition of the Nusselt number

$$Nu = \frac{Pm}{Pr} \langle |\nabla \theta|^2 \rangle, \quad (2.21)$$

where the quantity on the righthand side is often referred to as the thermal dissipation.

In reporting our numerical results we shall only make use of the volume and time-averaged kinetic and magnetic densities, which we denote by, respectively

$$E_{kin} \equiv \langle e_{kin} \rangle, \quad (2.22)$$

$$E_{mag} \equiv \langle e_{mag} \rangle. \quad (2.23)$$

Similarly, the Reynolds number is computed as

$$Re = \sqrt{2E_{kin}}. \quad (2.24)$$

2.2. Simulation details

The equations are solved using a standard toroidal-poloidal decomposition of the velocity and magnetic field such that the solenoidal conditions are satisfied exactly (e.g. Jones & Roberts 2000). A fully spectral code (Marti *et al.* 2016) is used for simulating the above equations with Fourier series in the horizontal dimensions and Chebyshev polynomials in the vertical dimension. The non-linear terms are de-aliased with the standard 2/3-rule. The equations are discretized in time with a third-order implicit-explicit Runge-Kutta scheme (Spalart *et al.* 1991). The code was benchmarked with the studies of Meneguzzi & Pouquet (1989) and Cattaneo *et al.* (2003).

While the most extreme three-dimensional RBC simulations have reached Rayleigh numbers as large as $Ra \approx 10^{12}$ (e.g. Stevens *et al.* 2011), the accessible range of Rayleigh numbers in dynamo simulations is restricted to significantly smaller values of Ra . As shown below, the ohmic dissipation scale is always smaller than the viscous dissipation scale for the cases studied here, implying that much higher spatial resolution is required to simulate dynamos in comparison to RBC. For example, for a $Pm = 5$ dynamo, the required resolutions in both the horizontal and the vertical directions are up to approximately two times the resolution needed for an equivalent Rayleigh number for RBC. Moreover, the existence of Alfvén waves in dynamos requires a significantly smaller numerical timestep in comparison to RBC. As Pm (or Rm) is increased the spatiotemporal resolution requirements become increasingly severe.

The aspect ratio of the computational domain is defined as

$$\Gamma = \frac{L}{H}, \quad (2.25)$$

where L is the periodicity length in the x and y dimensions (only domains of square cross-section are considered here). The horizontal dimensions are scaled by integer multiples (n) of the critical horizontal wavelength $\lambda_c = 2\pi/k_c$, where k_c is the critical horizontal wavenumber. For the impenetrable, stress-free, isothermal boundary conditions used in the present work, the critical Rayleigh number and critical wavenumber for the onset of hydrodynamic convection are $Ra_c = 27\pi^4/4 \approx 657.5$ and $k_c = \pi/\sqrt{2} \approx 2.22$, respectively. Thus, the aspect ratio is given by

$$\Gamma = \frac{2\pi n}{k_c} \approx 2.83 n. \quad (2.26)$$




Figure 1: Estimated values of the critical magnetic Reynolds number Rm_d and the critical Rayleigh number Ra_d for the onset of dynamo action: (a) exponential growth rate (γ) of magnetic energy versus magnetic Reynolds number (Rm); (b) exponential growth rate (γ) of magnetic energy versus Rayleigh number (Ra). The interpolated critical magnetic Reynolds number (Rm_d) and the critical Rayleigh number for dynamo action (Ra_d) for each magnetic Prandtl number (Pm) are shown by a ‘x’ symbol; (c) Rm_d versus Pm ; (d) Ra_d versus Pm .

While large aspect ratios are generally preferred, they are obviously more computationally demanding due to the larger resolution requirements. The aspect ratio is known to have an influence on many computed quantities, though it is expected that simulation statistics will converge as Γ is increased. Three-dimensional RBC simulations using aspect ratios up to $\Gamma = 128$ show that whereas bulk quantities such as Nu and Re asymptote to nearly constant values near $\Gamma \approx 4$ for $Ra \geq 2 \times 10^7$, other statistical quantities such as integral scales require significantly larger values of Γ to observe convergence (Stevens *et al.* 2018). Nevertheless, there is a trade-off between reaching larger aspect ratios and reaching larger Rayleigh numbers. In the present work we strive to reach large Rayleigh numbers while achieving convergence in Nu and Re ; the smallest aspect ratio used here is $\Gamma \approx 5.7$. All simulation parameters are listed in the Appendix.

3. Results

3.1. Onset of dynamo action

Dynamo simulations were carried out with five different values of the magnetic Prandtl number, $Pm = (0.8, 1, 3, 5, 7)$. Approximate values for the critical magnetic Reynolds

number for the onset of dynamo action, Rm_d , were determined for each value of Pm by iterating the Rayleigh number Ra . Each simulation was started from an initial state with small random noise in the magnetic field. In this regime the influence of the Lorentz force is negligible in comparison to other forces, and the magnetic energy would undergo an exponential growth (on average) if $Rm(Ra) > Rm_d$ and exponential decay (on average) when $Rm(Ra) < Rm_d$. The growth or decay rate of magnetic energy is denoted by γ , and was computed by a least-squares fit of the form $\ln(E_{mag}) = \gamma t + b$, where b is a constant coefficient for individual cases. We note that γ is twice the dynamo growth rate.

As shown in figure 1(a, c), the critical magnetic Reynolds number is estimated by a linear interpolation between the cases close to the onset of dynamo action. Some additional cases not shown on the plot were also simulated, however, these cases were found to be so close to Rm_d that the magnetic energy oscillated over a wide range and no clear exponential growth or decay was observed, confirming that our estimated Rm_d values are very close to the exact values. The critical magnetic Reynolds number shown in figure 1(a) suggests that Rm_d decreases with increasing Pm ; this result is expected and in agreement with previous studies (e.g. Schekochihin *et al.* 2007; Käpylä *et al.* 2018).

Bushby *et al.* (2012) and Käpylä *et al.* (2018) showed that dynamo action can be excited at a smaller value of Rm_d with the use of a larger aspect ratio. Käpylä *et al.* (2018) found that, when $Pm = Pr = 1$, an aspect ratio $\Gamma \gtrsim 3$ is needed for the growth rate to saturate. In all of our simulations (including those cases with $Pr < 1$), an aspect ratio of at least $\Gamma \gtrsim 8.5$ is used for the determination of Rm_d , thus the simulation domain should be sufficiently large to avoid the issue arising from the use of small values of Γ .

While a small subset of simulations were performed with $Pr = 0.01$ and $Pr = 0.05$, we did not systematically test the role of the aspect ratio for these cases. A single set of tests for $Pr = 0.05$ (where we increased the aspect ratio Γ from ≈ 8.5 to ≈ 14.1) showed that increasing the aspect ratio did decrease the growth rate. However, the estimated values of Rm_d and Ra_d were influenced only slightly.

The critical Rayleigh numbers for dynamo action (Ra_d) are estimated using the same procedure that is used for computing Rm_d . As shown in figure 1(b, d), Ra_d decreases as Pm is increased. For $Pm = (0.8, 1, 3, 5, 7)$ the estimated Rayleigh numbers for the onset of dynamo action are $Ra_d = (4.9 \times 10^5, 2.2 \times 10^5, 1.6 \times 10^4, 5.2 \times 10^3, 3.1 \times 10^3)$. As we will show in the following sections, these computed values of Ra_d can be useful for collapsing specific data.

A selection of $Pr = (0.01, 0.05)$ cases with $Pm = 1$ was also carried out to understand how Pr influences dynamo action. Figure 1(a, c) shows that a smaller value of Pr yields a lower value of Rm_d when Pm is held constant. For $Pr = (0.01, 0.05, 1)$ with $Pm = 1$ we find critical magnetic Reynolds numbers of $Rm_d = (135, 138, 167)$. This effect might occur because cases with lower Pr (at the same Rm and Pm) tend to have a more coherent flow structure (e.g. Goluskin & Spiegel 2012; Vogt *et al.* 2018; Pandey *et al.* 2018), which might be more beneficial to dynamo action. At the same Rm , dynamos with coherent flow are found to have a larger growth rate than dynamos without coherent structures (Tobias *et al.* 2013). However, a previous study of rotating spherical dynamos suggested that higher values of Rm are required for dynamo action if Pr becomes too small, though this effect is due to the influence of rotation (Simitev & Busse 2005). Nevertheless, the Prandtl number appears to play an important role for the onset of small-scale dynamo action; a more systematic investigation, beyond the scope of the present work, is needed to understand this effect in detail.




Figure 2: Heat transport for all cases: (a) Nusselt number (Nu) versus Rayleigh number (Ra); (b) compensated Nusselt number, $Nu/Ra^{2/7}$, versus Rayleigh number.

3.2. Heat transport

In all results presented hereafter we focus solely on the nonlinear regime of small-scale dynamos. Figure 2(a) shows the Nusselt number (Nu) versus Ra for all the cases investigated; the $Nu \sim Ra^{2/7}$ scaling is shown for reference. The compensated Nusselt number ($Nu/Ra^{2/7}$) is also plotted in figure 2(b). RBC cases without magnetic fields, shown as the black circles, are also plotted for comparison. When the dynamos are activated, the (Nu, Ra) curves depart from the RBC scaling. For each value of Pm , a scaling slope slightly smaller than $2/7$ (typically found in RBC at these parameter values) appears. However, as Ra is increased further, these slopes appear to approach a $2/7$ scaling again, suggesting that the influence of Pm on the scaling of Nu is weak. For a fixed value of Ra , the heat transfer is reduced as Pm is increased, or, equivalently, as the strength of magnetic field is increased. Though not shown, the dynamos exhibit similar mean temperature profiles in comparison to RBC, as suggested by the similar heat transport scaling.

In figure 3 both the viscous dissipation, ϵ_u , and the ohmic dissipation, ϵ_B , are plotted versus the Reynolds number. As suggested in figure 3(a) and the corresponding compensated plot shown in the inset, the viscous dissipation strongly depends on the Reynolds number; a numerical fit of $\epsilon_u = 1.14Re^{2.78}$ is found and shown. We find that the influence of Pm on ϵ_u is negligible. A scaling of $\epsilon_u \sim Re^3$ has been derived for the viscous dissipation in the bulk of the turbulent thermal convection (outside of the boundary layers), while $\epsilon_u \sim Re^{5/2}$ has been derived for the boundary layer (Grossmann & Lohse 2000; Scheel & Schumacher 2017). We find that our computed scalings are intermediate between these predicted scalings. Although ohmic dissipation cannot be purely determined by Re , we observe that in figure 3(b), ohmic dissipation is approaching the viscous dissipation scaling line as Re increases. The compensated plot in figure 3(b) shows the asymptotic scaling behaviour of $\epsilon_B \sim Re^{2.78}$ when Re is large enough for each individual Pm . This result suggests that the hydrodynamics properties of the fluid might be controlling both viscous and ohmic dissipation, and inertia appears to play a more important role in the energy cascade than the Lorentz force.

A quantity that provides a useful comparison between viscous and ohmic dissipation is the fraction of ohmic dissipation defined by

$$f_{ohm} \equiv \frac{\epsilon_B}{\epsilon_u + \epsilon_B}. \quad (3.1)$$




Figure 3: Dissipation for all cases: (a) viscous dissipation, ϵ_u , versus Reynolds number Re ; (b) ohmic dissipation ϵ_B versus Re ; (c) fraction of ohmic dissipation f_{ohm} versus Ra ; (d) f_{ohm} versus Ra/Ra_d , where Ra_d is the critical Rayleigh number for dynamo action. A compensated plot is also shown as an inset in (a) and (b).

Figure 3(c) shows f_{ohm} versus Ra for the dynamo cases. As expected, the flow is dominated by viscous dissipation near the onset of dynamo action. For each value of Pm , f_{ohm} initially increases rapidly with increasing Ra , but appears to flatten and approaches $f_{ohm} \rightarrow 0.5$ as Ra is increased. For our most extreme case of $Pm = 5$ and $Ra = 1 \times 10^7$ (our largest value of Rm), a value of $f_{ohm} \approx 0.5$ is reached, suggesting that in the regime of large Ra both ohmic dissipation and viscous dissipation are contributing equally to heat transport.

As shown in figure 3(d), f_{ohm} (for a given Pr) collapses when plotted versus the rescaled Rayleigh number Ra/Ra_d , where Ra_d is the critical Rayleigh number for dynamo action estimated in the previous section. This result suggests that f_{ohm} only has a weak dependence on Pm , while the degree of supercriticality of the Rayleigh number Ra/Ra_d is playing the dominant role. Similar convergent behaviour of f_{ohm} that is weakly dependent on Pm was also observed in the hydromagnetic study of Haugen *et al.* (2004), although their converged fraction of ohmic dissipation is $f_{ohm} \approx 0.7$. We notice that our $Pr = 0.05$ cases suggest that Pr appears to not have a strong influence on the saturated level of f_{ohm} , though the convergence rates appear to be affected.

Vertical profiles of the horizontally and time averaged local dissipation $\epsilon_u(z)$ and $\epsilon_B(z)$ are shown in figure 4(a) for a typical dynamo with $Pm = 5$ and $Ra = 6 \times 10^5$ ($Rm =$

1122). Note that here

$$\epsilon_u(z) = \frac{1}{Pm^2} \overline{(\nabla \times \mathbf{u})^2}, \quad (3.2)$$

$$\epsilon_B(z) = \frac{1}{Pm^2} \overline{(\nabla \times \mathbf{B})^2}. \quad (3.3)$$

The ohmic dissipation is dominant near the boundary, while viscous dissipation is dominant in the bulk when Rm is not too large. We also observe that ohmic dissipation has a markedly thinner boundary layer in comparison to that of the viscous dissipation. Moreover, both profiles show similar structure within the interior, suggesting that both dissipation mechanisms are dynamically linked. The vertical profile of viscous dissipation for the equivalent RBC case is also shown in figure 4(a) for comparison. The viscous dissipation structure of the dynamo remains very similar to that of the RBC case (although their magnitudes are different), suggesting that viscous dissipation has a direct influence on ohmic dissipation in the bulk. The dissipation profiles for our most extreme (largest value of Rm) dynamo case corresponding to $Pm = 5$ and $Ra = 1 \times 10^7$ are plotted in figure 4(b). Here the boundary layers become much thinner and the dissipation is dominated by the contribution in the bulk. Again, we find that both profiles show similar structure within the bulk, while the magnitudes are approaching similar values as the Rayleigh number is increased.

The behaviour of the dissipation near the boundaries is likely influenced by the choice of boundary conditions. Although we did not perform simulations with no-slip mechanical boundary conditions (as opposed to the stress-free conditions used here), three additional simulations with $Pm = 5$ and $Ra = (1 \times 10^4, 1 \times 10^5, 6 \times 10^5)$ were performed in which electrically insulating electromagnetic boundary conditions were used. With these insulating boundary conditions we found that the depth-dependence of the dissipation profiles remained essentially unchanged relative to the vertical magnetic field boundary conditions. Though we found differences in magnitudes of the total dissipation, no systematic variation was investigated. We note that although the ohmic dissipation tends to be largest near the boundaries, the integrated contribution of this boundary layer region to the total dissipation is relatively small due to the thinness of the layer.

3.3. Flow speeds and energy

Figure 5(a,b) shows the Reynolds number Re and compensated Reynolds number $Re/Ra^{1/2}$ versus Ra for all cases. The convective free-fall scaling ($Re \sim Ra^{1/2}$) is shown for reference. Curve fits to the data yield $Re \sim (Ra^{0.45}, Ra^{0.43}, Ra^{0.43}, Ra^{0.44}, Ra^{0.44})$ for $Pr = 1$, $Pm = (0.8, 1, 3, 5, 7)$, respectively. We observe that for a given value of Pr and Ra , the dynamos tend to have smaller flow speeds in comparison to the RBC data since the magnetic energy comes at the cost of kinetic energy. As Ra is increased, the dynamos show a departure from the RBC scaling. The compensated Reynolds number shown in figure 5(b) shows that this departure is very slight, though there is a trend of increased departure with increasing Pm . The $Pr = 0.05$ cases show the most rapid growth of Re with increasing Ra , though there is insufficient data to suggest any significant difference in scaling behaviour between the different Prandtl numbers. Despite the fact that approximately half of the dissipation is ohmic (i.e. $f_{ohm} \approx 0.4$) for the $Pr = 0.05$ cases, there is very little difference in flow speeds between the dynamos and RBC. Of course, the Nusselt numbers for these $Pr = 0.05$ cases are rather small: $Nu \lesssim 4$.

The efficiency of the dynamos can be measured by the ratio of the magnetic energy to the kinetic energy (E_{mag}/E_{kin}). As shown in figure 6(a), cases with different values of Pm show similar behaviour: (E_{mag}/E_{kin}) increases as Ra is increased, and it appears that




Figure 4: Vertical profiles of horizontally and time averaged viscous dissipation $\epsilon_u(z)$ and ohmic dissipation $\epsilon_B(z)$. (a) $Pm = 5$, $Ra = 6 \times 10^5$ ($Rm \approx 1100$). The corresponding non-magnetic case (RBC) is also plotted for comparison. (b) $Pm = 5$, $Ra = 1 \times 10^7$ ($Rm \approx 3900$). The total dissipation ϵ_u and ϵ_B is calculated by depth averaging the profile.




Figure 5: Flow speeds for all cases: (a) Reynolds number versus Rayleigh number; (b) Compensated Reynolds number ($Re/Ra^{1/2}$) versus Rayleigh number.

the ratio approaches constant values at large Rayleigh number. The convergence of the energy ratio was also observed in the mechanically forced dynamo simulations of Haugen *et al.* (2004), however, the dependence on magnetic Prandtl number was not discussed. The data can be reasonably collapsed by rescaling the energy ratio with $Pm^{2/3}$, and rescaling the Rayleigh number with Ra_d , as shown in figure 6(b). Since the growth of Re depends on Pr (e.g. figure 5(a,b)), we do not expect the $Pr = 0.05$ data to follow the same trend as the $Pr = 1$ data. When Ra/Ra_d is large enough, dynamos with larger Pm can transfer kinetic energy to magnetic energy more efficiently. We note that for $Ra = 5 \times 10^5$ and $Pm = 5$ the energy ratio $E_{mag}/E_{kin} \approx 0.2$ agrees with the result of Cattaneo *et al.* (2003). Our results suggest that this value represents the approximate asymptote for the energy ratio when $Pm = 5$, and that the asymptote is Pm dependent.

The magnetic energy is plotted versus Rayleigh number in figure 7(a). The $E_{mag} \sim Ra$




Figure 6: (a) The ratio of magnetic energy to kinetic energy (E_{mag}/E_{kin}) versus the Rayleigh number Ra ; (b) rescaled energy ratio $E_{mag}/(E_{kin} Pm^{2/3})$ versus the rescaled Rayleigh number (Ra/Ra_d) where Ra_d is the critical Rayleigh number for dynamo action.




Figure 7: Magnetic energy for all cases: (a) Magnetic energy versus Rayleigh number; (b) Compensated magnetic energy ($E_{mag}/(Ra Pm^{2/3})$) versus rescaled Rayleigh number Ra/Ra_d .

scaling is shown for reference. The magnetic energy increases relatively rapidly with Ra beyond the onset of dynamo action, but appears to flatten and approaches $E_{mag} \sim Ra$ at large values of Ra/Ra_d . Note that since the energy ratio (E_{mag}/E_{kin}) saturates at large Ra/Ra_d , we expect that the magnitude of the magnetic energy must grow with Ra/Ra_d at the same rate as the flow speed squared. The compensated magnetic energy ($E_{mag}/(Ra Pm^{2/3})$) versus the rescaled Rayleigh number Ra/Ra_d is shown in figure 7(b). The $Pm^{2/3}$ dependence is purely an ad-hoc fit to the data and is only meant to provide a rough scaling with Pm in the asymptotic regime. The curves become relatively flat at large Rayleigh numbers, suggesting that the $E_{mag} \sim Ra$ scaling might be an asymptotic result for high Rm RBC dynamos.

3.4. Length scales

The characteristic length scales are computed for all simulations. Two length scales are computed: (1) the Taylor microscale; and (2) the integral length scale. The Taylor

microscale is the length scale at which the influence of viscous or ohmic dissipation becomes important, and can thus provide an estimate for the dissipation length scales. In contrast, the integral scale is the correlation length scale for the corresponding field. The magnetic Taylor microscale λ_B and the velocity Taylor microscale λ_u are defined by, respectively,

$$\lambda_B = \sqrt{\frac{\langle \mathbf{B}^2 \rangle}{\langle (\nabla \times \mathbf{B})^2 \rangle}}, \quad (3.4)$$

and

$$\lambda_u = \sqrt{\frac{\langle \mathbf{u}^2 \rangle}{\langle (\nabla \times \mathbf{u})^2 \rangle}}. \quad (3.5)$$

Note that these length scales are computed over the entire fluid layer, including the boundary layers. Some tests were done in which the boundary layers were excluded from the calculation, and showed their influence was negligible. We therefore only present calculations that included the entire fluid layer.

The velocity and magnetic Taylor microscales are plotted versus the Rayleigh number in figure 8(a), which shows in detail how these length scales are modified by Pm . The velocity Taylor microscale λ_u shows very little change with increasing Pm . Some of the data points show a small increase in λ_u with increasing Pm ; this effect can be understood by the fact that the dynamo converts kinetic energy into magnetic energy and therefore results in a slight decrease of the Reynolds number for a given value of Ra . However, there is no appreciable difference in the scaling behaviour of λ_u with Ra for the various values of Pm used here: curve fits to the data yield $\lambda_u \sim (Ra^{-0.153}, Ra^{-0.171}, Ra^{-0.170}, Ra^{-0.167}, Ra^{-0.169})$ for $Pm = (0.8, 1, 3, 5, 7)$, respectively. The scaling of λ_B is noticeably steeper than the scaling for λ_u ; the corresponding curve fits are $\lambda_B \sim (Ra^{-0.235}, Ra^{-0.242}, Ra^{-0.245}, Ra^{-0.251}, Ra^{-0.245})$, for $Pm = (0.8, 1, 3, 5, 7)$, respectively. We emphasize that for all of our simulations use $Pm = O(1)$.

Figure 8(b,c) shows the velocity and magnetic Taylor microscales versus Reynolds number and magnetic Reynolds number. We observe that Pm only has an influence on the magnitude of λ_u and λ_B , however, the scaling behaviour of these length scales remains basically the same for all Pm . The scaling $\lambda_B \sim Rm^{-1/2}$ is also plotted in figure 8(c) for reference. Previous studies of mechanically forced isothermal dynamos suggested that λ_B scales as $Rm^{-1/2}$ for fixed Re at moderate Rm (Brummell *et al.* 2001). Both results have a $Rm^{-1/2}$ dependence, however, we note that they are in a relatively low Re and Rm regime, and they assume λ_B does not depend on Re and the magnetic length scale is determined by the balance between advection and diffusion in the induction equation. Our result suggests that λ_B cannot be purely determined by Rm . As we will show in a later section, there does exist a subdominant balance in the induction equation between induction and diffusion for our cases. Of course, given the very different forcing and boundary conditions between our investigation and that of Brummell *et al.* (2001), one might expect differences in the scaling behaviour.

As shown in figure 8(d), the magnetic Taylor microscale collapses when rescaled with Pm , a best-fit yields $\lambda_B Pm^{0.30} = 1.29 Ra^{-0.25}$. Since Re scales with Ra , we also plot the magnetic Taylor microscale as a function of Re in figure 8(e), where a scaling law $\lambda_B Pm^{0.35} = 1.1 Re^{-0.56}$ is found. Figure 8(e) shows the velocity Taylor microscale collapses very well when plotted against Re , which suggests that the influence of Pm on the velocity length scale is very weak, and the magnetic field has a small effect. Curve fits for all dynamo cases as well as RBC cases yields $\lambda_u = 0.97 Re^{-0.39}$. For Kolmogorov turbulence, a scaling law of $\lambda_u \sim Re^{-1/2}$ is obtained (Pope 2000). However, dynamos




Figure 8: Scaling behaviour of the magnetic Taylor microscale (λ_B) (solid marks) and velocity Taylor microscale (λ_u) (empty marks). (a) Taylor microscale plotted against Rayleigh number; (b) Taylor microscale plotted against Reynolds number; (c) Taylor microscale plotted against magnetic Reynolds number; (d) The rescaled magnetic Taylor microscale ($\lambda_B Pm^{0.30}$) versus Rayleigh number; (e) The rescaled magnetic Taylor microscale ($\lambda_B Pm^{0.35}$) (solid marks) versus the Reynolds number and the velocity Taylor microscale (λ_u) (empty marks) versus the Reynolds number.

in the Rayleigh-Bénard convection geometry cannot be characterized by homogeneous isotropic turbulence, so it is reasonable to expect a modified scaling here. The absence of Pm in this scaling law is also expected, as the viscous dissipation appears to be independent of Pm within our accessible parameter range.

It is commonly hypothesized that the ohmic dissipation length scale will occur within




Figure 9: Depth-averaged length scales for $Pm = 5$. The velocity Taylor microscale (λ_u) calculated from energy and dissipation, velocity Taylor microscale (λ'_u) calculated from energy spectra, and velocity integral scale (ℓ_u) plotted versus Rayleigh number. The corresponding magnetic length scales are also shown.

the inertial range of the turbulence when $Pm \ll 1$ ($Re \gg Rm$) (e.g. Rincon 2019; Tobias 2019). Such a regime requires, at a minimum, that $\lambda_B \gg \lambda_u$. Though the simulations in the present study are obviously limited with respect to the accessible parameter range, especially in the restriction to $Pm = O(1)$, we observe $\lambda_u > \lambda_B$ in all cases. Moreover, the results (especially the $Pm = 0.8$ cases) shown in figure 8 indicate that the scaling slope (as a function of Ra , Re , Rm) for the ohmic dissipation length scale is always steeper than the corresponding slope for the viscous dissipation length scale, and that this slope remains mostly independent of Pm . The case $Pm = 0.8$ was chosen to allow $Pm < 1$, while retaining a sufficiently large Rm ; while this particular case is certainly not a small magnetic Prandtl number, we do not observe a change in the scaling slope with this case relative to $Pm \geq 1$. Thus, at least for the limited parameter range of the present study, there is no indication that the slopes change drastically either with increasing Ra or decreasing Pm . However, the Pm we explored here are $Pm \sim O(1)$, it remains unclear whether there would be a change in the scaling slope for ohmic dissipation scale when $Pm \ll 1$. In that case, we might expect a larger ohmic dissipation scale than the viscous scale.

The depth-dependent integral scales for the magnetic field and velocity field are defined by, respectively (e.g. Meneguzzi & Pouquet 1989),

$$\ell_B = \frac{\int k^{-1} \hat{E}_{mag}(k) dk}{\int \hat{E}_{mag}(k) dk}, \quad \ell_u = \frac{\int k^{-1} \hat{E}_{kin}(k) dk}{\int \hat{E}_{kin}(k) dk}, \quad (3.6)$$

where $\hat{E}_{mag}(k)$ and $\hat{E}_{kin}(k)$ are the magnetic and kinetic energy spectra, $\mathbf{k} = (k_x, k_y)$ is the horizontal wavenumber vector and $k = \sqrt{k_x^2 + k_y^2}$. The corresponding depth-dependent Taylor microscales can be computed with the definitions

$$\lambda'_B = \left(\frac{\int \hat{E}_{mag}(k) dk}{\int k^2 \hat{E}_{mag}(k) dk} \right)^{1/2}, \quad \lambda'_u = \left(\frac{\int \hat{E}_{kin}(k) dk}{\int k^2 \hat{E}_{kin}(k) dk} \right)^{1/2}. \quad (3.7)$$

We observe a weak dependence on the depth for all of the computed length scales defined above, and therefore only report the depth-averaged values.

The time averaged values of the spectra-based length scales are calculated for many




Figure 10: Energy spectra. (a) Kinetic energy spectra for $Pm = 5$ dynamos; (b) Magnetic energy spectra for $Pm = 5$ dynamos; (c) Energy spectra at $Ra = 1 \times 10^7$ for $Pm = 0.8$ ($Rm = 742$) and $Pm = 5$ ($Rm = 3948$); (d) Kinetic energy spectra for RBC. The spectral slopes of $k^{1/3}$ and $k^{-5/3}$ are plotted for reference only.

of the $Pm = 5$ cases. As shown in figure 9, we find nearly identical scaling behaviour between both λ'_B and λ_B , and λ'_u and λ_u . As expected, a divergence between the velocity Taylor microscale and the velocity integral scale occurs as the Reynolds number (Rayleigh number) increases. However, figure 9 also shows that the scale separation for the magnetic field is relatively weak since the scaling slopes for the two magnetic length scales are close to one another. The integral scale for the velocity appears to be sensitive to the change of aspect ratio (e.g. Stevens *et al.* 2018). As a result, some velocity integral scale data does appear to follow a clear scaling trend. We should note that the magnetic integral scale appears to have a slope very close to the velocity Taylor microscale, which indicates that there is correlation between the viscous force and the Lorentz force (or magnetic induction).

3.5. Energy spectra

Figure 10(a, b) show how the kinetic and saturated magnetic energy spectra change with Rayleigh number for $Pm = 5$ dynamos. The $\hat{E}_{kin}(k) \sim k^{-5/3}$ scaling (typically

found in the Kolmogorov inertial subrange) is also plotted for reference in figure 10(a), which shows the development of an inertial subrange. The magnetic energy spectra appear to flatten over a large range of wavenumber as Ra is increased. For a fixed Rayleigh number, kinetic energy dominates over magnetic energy at large scales. However, for all our ($Pm \sim O(1)$) cases, magnetic energy spectra exceed kinetic energy spectra for large wavenumbers. As shown in figure 10(c), the magnetic energy becomes slightly larger than kinetic energy at small scales, even for the $Pm = 0.8$ cases. A $k^{1/3}$ scaling behaviour for the magnetic energy spectra at large scales has been observed in the previous hydromagnetic turbulence study of Haugen *et al.* (2004); they also found a $k^{-5/3}$ subrange after the peak of the magnetic energy spectra. These two slopes are plotted in figure 10(b) only for reference, since we do not expect the scalings to be the same for the two different systems. We note that the change in magnetic energy spectra occurs at the wave number where the magnitudes of magnetic energy and kinetic energy are comparable. The kinetic energy spectra for RBC (and the $k^{-5/3}$ slope) are also plotted in figure 10(d) for reference. We observe that when compared with the RBC cases, dynamo action reduces the amplitude of the peak of kinetic energy spectra and makes the spectra tail less steep. The reduction in the peak amplitude is the result of transferring kinetic energy to magnetic energy by dynamo actions, while the slight increase in the spectra at high wavenumber is due to the Lorentz force driving motion at small scales as $Pm > 1$.

3.6. Forces

In this section we analyze the force balance in the momentum equation to understand the role of the Lorentz force. For the vertical component of the momentum equation we remove the hydrostatic part by decomposing the flow variables into a horizontal average and a fluctuation. For instance, $\mathbf{u} = \bar{\mathbf{U}} + \mathbf{u}'$, $\mathbf{b} = \bar{\mathbf{B}} + \mathbf{b}'$, $\Theta = \bar{\Theta} + \theta'$, where $\bar{\mathbf{U}}$, $\bar{\mathbf{B}}$ and $\bar{\Theta}$ are the horizontal mean components, \mathbf{u}' , \mathbf{b}' and θ' are the fluctuating components, and p' is the reduced pressure. Under the present circumstances we find the mean components $\bar{\mathbf{U}}$, $\bar{\mathbf{B}}$ are much smaller than the corresponding fluctuating components \mathbf{u}' , \mathbf{b}' . The fluctuating vertical component of the momentum equation is found by subtracting the mean (hydrostatic) balance to give

$$\underbrace{\partial_t w'}_{F_t} = \underbrace{Pm \nabla^2 w'}_{F_v} + \underbrace{(-\mathbf{u}' \cdot \nabla w' + \partial_z \overline{w'^2})}_{F_a} + \underbrace{Pm(\mathbf{b}' \cdot \nabla b'_z - \partial_z \overline{b'^2_z})}_{F_l} + \underbrace{\frac{Ra Pm^2}{Pr} \theta'}_{F_b} - \underbrace{\partial_z p'}_{F_p}. \quad (3.8)$$

We use the same symbols for denoting the various terms in the horizontal components of the momentum equation.

The time averaged rms values of the different terms in the momentum equation are calculated at the midplane ($z = 0.5$) and shown in figure 11(a,b). The vertical dependence of the forces was also analyzed using vertical profiles as in Yan *et al.* (2019), and it was found that the data shown in figure 11(a,b) was representative of all depths. The Lorentz force initially grows rapidly with increasing Ra , then slows once it becomes comparable in magnitude with the viscous force. We note that even for the largest value of Ra , the Lorentz and viscous forces are approximately the same magnitude. We observe that the time derivative term (inertia) grows at the fastest rate with increasing Ra among all terms in the horizontal components of the momentum equation. The corresponding forces for RBC (without magnetic field) are plotted in figure 11(c,d) for comparison. We observe that the scaling of the buoyancy force is basically the same for both the dynamo and RBC, whereas the scaling slopes for all other forces in the dynamos are reduced relative to RBC.




Figure 11: Time-averaged rms values of forces in the momentum equation at the midplane ($z = 0.5$) for (a,b) dynamos with $Pm = 5$; and (c,d) RBC. (a,c) Forces in the horizontal direction versus Rayleigh number; (b,d) forces in the vertical direction versus Rayleigh number. Advection and the Lorentz, viscous, buoyancy, pressure gradient and inertia forces are denoted by F_a , F_l , F_v , F_b , F_p and F_t , respectively.

Horizontal spectra are computed for all forces in the momentum equation at the midplane, and are shown in figure 12 for (a) $Pm = 5$ dynamo and (b) RBC at $Ra = 6 \times 10^5$. A convective free-fall balance in which the pressure gradient force, buoyancy force and advection are comparable, is present at the largest spatial scales (smallest wavenumbers) for both cases. For the dynamo the Lorentz force is subdominant at large spatial scales ($k \lesssim 70$), and is in a near perfect balance with the viscous force at small spatial scales ($k \gtrsim 40$). The spectra of the sum of the Lorentz force and the viscous force are calculated and plotted as the dashed line. At small scales, the sum of the two forces is one order of magnitude smaller than both Lorentz force and viscous force, suggesting that there is an almost exact balance between Lorentz force and viscous force locally. Part of the velocity driven by the Lorentz force is dissipated by viscosity locally and almost instantaneously. This result suggests that the viscous length scale might be controlling the magnetic field length scales, which explains the similar scaling slopes for the velocity Taylor microscale and the magnetic integral scale. A similar argument for the balance between the viscous




Figure 12: Horizontal spectra of the various forces in (a) dynamo with $Pm = 5$ and (b) RBC. For both cases $Ra = 6 \times 10^5$. (c) Lorentz force and viscous force spectra for $Pm = 5$ dynamos.

force and the Lorentz force is also discussed in previous studies (Brummell *et al.* 2001; Schekochihin *et al.* 2002c), and our spectral result serves as direct evidence for this local balance. Note that the spectra we show here are for $Pm = 5$ cases. For small Pm , it is possible that the Lorentz force might behave quite differently at small scale, though in all of our cases we find that the Lorentz force is driving the flow at small scale. When compared with the RBC case of figure 12(b), broader spectra are observed for the dynamo case; advection, viscous and inertia forces are slightly reduced at large scales and larger in magnitude at small scales, which suggests that the Lorentz force is consuming kinetic energy at large scales while driving the flow at small scales.

Figure 12(c) shows the Lorentz force and viscous force spectra for $Pm = 5$ dynamos at $Ra = (4 \times 10^4, 2 \times 10^5, 6 \times 10^5, 1 \times 10^6)$, respectively. Initially ($Ra \sim 4 \times 10^4$), the Lorentz force is weaker than the viscous force at large scales, and is in balance with the viscous force at small scales. When Ra increases, the Lorentz force exceeds the viscous force at large scales, this result is also reflected in the relatively rapid initial growth of F_l in figure 11(a,b). However, as Ra is increased further ($Ra > 2 \times 10^5$), the growth of the Lorentz force at large scales appears to slow down; both the Lorentz force and the viscous force spectra appear to shift to larger wavenumber, and the length scale where the two forces are balanced becomes smaller. We note that the spectra of the Lorentz force appears to flatten and saturate at small wavenumber as Ra is increased. This result

suggests that the Lorentz force might be limited by the viscous force (at least at small length scales), and the growth rate of the Lorentz force appears to saturate.

3.7. Induction equation balances

Analogous to the force balance analysis of the momentum equation in the previous section, here the relative sizes of the different terms in the magnetic induction equation are analyzed in both physical and spectral space. Figure 13(a,b,c,d) shows vertical profiles of the instantaneous horizontal rms of each term in the induction equations, for two typical cases both with $Ra = 10^6$, and $Pm = 1$ and $Pm = 5$, respectively; these cases correspond to magnetic Reynolds numbers of $Rm \approx 310$ and $Rm \approx 1340$. Though not shown, cases with different values of Pm and Ra show similar behaviour. Not surprisingly, given that $Rm \gg 1$, the primary balance indicates that advection of magnetic field dominates,

$$\partial_t \mathbf{B} + \mathbf{u} \cdot \nabla \mathbf{B} \approx 0. \quad (3.9)$$

In addition, a sub-dominant balance is observed between magnetic induction and magnetic diffusion,

$$\mathbf{B} \cdot \nabla \mathbf{u} \sim \nabla^2 \mathbf{B}. \quad (3.10)$$

We also find that the magnitude of $\partial_t \mathbf{B} + \mathbf{u} \cdot \nabla \mathbf{B}$ is of the same order as induction and diffusion.

The depth-dependence of the different terms is similar for the two different cases shown. Interestingly, even though Rm is approximately an order of magnitude larger for the $Pm = 5$ case, the relative difference in magnitude between the leading order balance and the subdominant balance changes only slightly; for instance $|\mathbf{u} \cdot \nabla \mathbf{B}|/|\nabla^2 \mathbf{B}| \approx 3$ and $|\mathbf{u} \cdot \nabla \mathbf{B}|/|\nabla^2 \mathbf{B}| \approx 4$ for $Pm = 1$ and $Pm = 5$, respectively. This result could be due to the small-scale magnetic Reynolds number not changing appreciably.

The spectral space representation of the different terms in the induction equation is given in figure 13(e,f) for the same $Ra = 10^6$, $Pm = 1$ and $Pm = 5$ cases, respectively. Again, the data is computed at one instant in time. Whereas $\partial_t \mathbf{B}$ and $\mathbf{u} \cdot \nabla \mathbf{B}$ are balanced at all length scales, the stretching ($\mathbf{B} \cdot \nabla \mathbf{u}$) and diffusion ($\nabla^2 \mathbf{B}$) terms are balanced at different length scales. This scale-dependent balance can be understood by a simple scale analysis of the sub-dominant balance. For instance, equation (3.10) yields

$$\frac{Rm}{L_u} \sim \frac{1}{L_B^2}, \quad (3.11)$$

where the velocity has the non-dimensional units of magnetic Reynolds number, L_u is a characteristic length scale for the velocity and L_B is a characteristic length scale for the magnetic field. Since the magnetic length scale is in the diffusion term ($\nabla^2 \mathbf{B}$), it is natural to assume that this length scale can be captured by the magnetic Taylor microscale $L_B = \lambda_B$. Thus we have

$$Rm \sim \frac{L_u}{\lambda_B^2}. \quad (3.12)$$

However, it is not obvious what L_u represents. We plug in the magnetic Taylor microscale (λ_B) and test the velocity Taylor microscale (λ_v) as well as the velocity integral scale (ℓ_u), the results are shown in figure 14. We find that when using the velocity Taylor microscale (λ_v), the curve for the compensated magnetic Reynolds number $Rm \lambda_B^2 / \lambda_u$ is relatively flat, however we can still see a systematic increase of $Rm \lambda_B^2 / \lambda_u$ with increasing Ra . When we apply the velocity integral scale ℓ_u , the compensated magnetic Reynolds number $Rm \lambda_B^2 / \ell_u$ stays nearly constant over a large range of Ra , suggesting that the




Figure 13: Instantaneous balances in the induction equation. Vertical profiles of the horizontal rms of each term present in the vertical component [(a),(b)] and in the horizontal component [(c),(d)] of the induction equation and the corresponding power spectra (of all three components) at the mid-plane [(e),(f)] are shown. Case $Pm = 1, Ra = 1 \times 10^6$ is shown in [(a),(c),(e)]. Case $Pm = 5, Ra = 1 \times 10^6$ is shown in [(b),(d),(f)]. Magnetic induction, diffusion, advection, and time derivative are denoted by $\mathbf{B} \cdot \nabla \mathbf{u}$, $-\mathbf{u} \cdot \nabla \mathbf{B}$, $\nabla^2 \mathbf{B}$, and $\partial_t \mathbf{B}$. w and B_z represent the vertical component of \mathbf{u} and \mathbf{B} , while u_h and B_h represent the vertical component.




Figure 14: Rescaled magnetic Reynolds number versus Rayleigh number for $Pm = 5$ cases. Both $Rm\lambda_B^2/\lambda_u$ and $Rm\lambda_B^2/\ell_u$ results are shown.

velocity integral scale ℓ_u can better characterize the length scale in the induction terms than the Taylor microscale. This result implies that the magnetic diffusion length scale is controlled by Rm and the velocity integral scale:

$$Rm \sim \frac{\ell_u}{\lambda_B^2}. \quad (3.13)$$

4. Discussion

RBC-driven dynamos have been investigated numerically over a range of Rayleigh numbers and magnetic Prandtl numbers. Heat transport in dynamos is reduced relative to RBC, primarily because of the associated reduction in kinetic energy required for the generation of magnetic energy. Simulations at the largest values of Ra suggest a heat transport scaling with Ra that is similar to RBC, i.e. $Nu \sim Ra^{2/7}$ within our investigated range of Ra . The scaling behaviour of the flow speeds with increasing Rayleigh number is also similar to RBC ($Re \sim Ra^{1/2}$). An asymptotic scaling for the magnetic energy $E_{mag} \sim Ra$ is also observed at large Ra/Ra_d . Given a large enough value of Ra/Ra_d , the ratio of magnetic energy to kinetic energy E_{mag}/E_{kin} appears to saturate for individual Pm , while the saturated level depends on Pm ; we find reasonable collapse of the data using $E_{mag}/E_{kin} \sim Pm^{2/3}$. This result suggests that Pm is controlling the efficiency of dynamo action.

Despite the similar scaling behaviour in Nu for both RBC and RBC-driven dynamos, for a fixed value of Pm , the simulations show that ohmic dissipation becomes equally important to viscous dissipation as the Rayleigh number increases, as characterized by an ohmic dissipation fraction $f_{ohm} \rightarrow 0.5$. The scaling of the viscous dissipation in the dynamos is observed as $\epsilon_u \sim Re^{2.8}$, which is nearly identical to equivalent RBC simulations (Grossmann & Lohse 2000). Moreover, the ohmic dissipation is observed to approach the scaling $\epsilon_B \sim Re^{2.8}$ as Ra is increased. The findings suggest that ohmic dissipation is controlled by viscous dissipation.

The Taylor microscale is computed for both the velocity field and magnetic field, thus providing a measure of the viscous and ohmic dissipation scales, respectively. The ohmic dissipation scale is observed to be smaller than the viscous dissipation scale for all simulations, even for the $Pm = 0.8$ cases. Furthermore, the ohmic dissipation length scale decreases more rapidly with increasing Rayleigh number (Reynolds number) in comparison to the viscous dissipation length scale. Curve fits yield scalings of $\lambda_B \sim Re^{-0.56} Pm^{-0.35}$ and $\lambda_u \sim Re^{-0.39}$ for velocity field and magnetic field Taylor microscale,

respectively. Unless a change in scaling occurs at higher Rayleigh numbers beyond those accessible in the present study, this result suggests that the ohmic dissipation scale is always less than the viscous dissipation scale, regardless of the value of the magnetic Prandtl number for the values of Pm considered in this paper.

A numerical analysis of the forces in both physical and spectral space shows that the viscous and Lorentz forces are in balance at small spatial scales. The viscous force in dynamos is enhanced at these same scales relative to RBC. At large spatial scales the Lorentz force dominates the viscous force, but it remains subdominant relative to buoyancy and inertia. A two-dimensional (in the horizontal plane) spectral energy analysis suggests that the Lorentz force is primarily dissipative at large spatial scales, but acts as a source of kinetic energy at small spatial scales; this result might explain why the viscous force at small spatial scales is enhanced in dynamos relative to RBC.

To leading order, the induction equation is characterized by a conservation of magnetic flux. Stretching and diffusion act at higher order and are dominant at large and small spatial scales, respectively. According to this balance, the magnetic diffusion length scale is found to be related to the magnetic Reynolds number and velocity integral scale via: $\lambda_B \sim (\ell_u/Rm)^{1/2}$.

In light of the limited range in Rayleigh numbers accessible to DNS studies of dynamos, several questions remain. Since the buoyancy force has a relatively smaller growth rate with increasing Ra when compared to that of the Lorentz force, it remains unclear whether the growth of the Lorentz force will slow down and approach the scaling of the buoyancy force when Ra becomes large enough, or if the Lorentz force will eventually dominate over the buoyancy force and change the dynamics fundamentally. Simulations with higher Ra are also needed to verify the asymptotic convergence behaviour of flow speeds, magnetic energy and dissipation. For planetary interiors and stars the thermal Prandtl number is less than unity, the dependence of the magnetic to kinetic energy ratio and the ohmic dissipation fraction on Pr remains unclear and might also need to be explored in more detail.

We reiterate that all of the simulations presented here use $Pm = O(1)$. In the small Pm regime in which $Pm \ll 1$, one might expect the stretching of the eddies to be balanced by magnetic diffusion on the ohmic dissipation length scale (Tobias 2019; Rincon 2019). Although the low Pm regime is currently beyond the reach of direct numerical simulation, our results (especially the $Pm = 0.8$ and $Pm = 1$ cases) suggest that instead of a sudden change at $Pm = 1$, the transition to the low Pm regime might be gradual, or occur sharply at much lower Pm . This observation is consistent with calculations (Tobias *et al.* 2012) that suggest numerical resolutions of $(10^4)^3$ modes are required to reach this large- Re regime. Further investigation with smaller Pm will therefore be needed to explore the mechanisms that drive the changes between the scalings.

Acknowledgements

This work utilized the RMACC Summit supercomputer, which is supported by the National Science Foundation (awards ACI-1532235 and ACI-1532236), the University of Colorado Boulder, and Colorado State University. The Summit supercomputer is a joint effort of the University of Colorado Boulder and Colorado State University. The authors acknowledge the Texas Advanced Computing Center (TACC) at The University of Texas at Austin for providing high-performance computing resources that have contributed to the research results reported within this paper.

Funding

This work was supported by the National Science Foundation under grant EAR #1620649 (MY and MAC). SMT would like to acknowledge support from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (Agreement No. 328D5S-DLV-786780).

Declaration of Interests

The authors report no conflict of interest.

REFERENCES

- AHLERS, G., GROSSMANN, S. & LOHSE, D. 2009 Heat transfer and large scale dynamics in turbulent Rayleigh-Bénard convection. *Rev. Mod. Phys.* **81** (2), 503.
- AURNOU, J. M. & OLSON, P. 2001 Experiments on Rayleigh-Bénard convection, magnetoconvection, and rotating magnetoconvection in liquid gallium. *J. Fluid Mech.* **430**, 283–307.
- BRUMMELL, N. H., CATTANEO, F. & TOBIAS, S. M. 2001 Linear and nonlinear dynamo properties of time-dependent ABC flows. *Fluid Dyn. Res.* **28** (4), 237.
- BUSHBY, P. J. & FAVIER, B. 2014 Mesogranulation and small-scale dynamo action in the quiet Sun. *Astronomy & Astrophysics* **562**, A72.
- BUSHBY, P. J., FAVIER, B., PROCTOR, M. R. E. & WEISS, N. O. 2012 Convectively driven dynamo action in the quiet sun. *Geophys. Astrophys. Fluid Dyn.* **106** (4-5), 508–523.
- CALKINS, M. A., JULIEN, K., TOBIAS, S. M. & AURNOU, J. M. 2015 A multiscale dynamo model driven by quasi-geostrophic convection. *J. Fluid Mech.* **780**, 143–166.
- CASTAING, B., GUNARATNE, G., HESLOT, F., KADANOFF, L., LIBCHABER, A., THOMAE, S., WU, X., ZALESKI, S. & ZANETTI, G. 1989 Scaling of hard thermal turbulence in Rayleigh-Bénard convection. *J. Fluid Mech.* **204**, 1–30.
- CATTANEO, F. 1999 On the origin of magnetic fields in the quiet photosphere. *Astrophys. J. Lett.* **515** (1), L39.
- CATTANEO, F., EMONET, T. & WEISS, N. 2003 On the interaction between convection and magnetic fields. *Astrophys. J.* **588** (2), 1183.
- CHANDRASEKHAR, S. 1961 *Hydrodynamic and Hydromagnetic Stability*. U.K.: Oxford University Press.
- CHENG, J. S., STELLMACH, S., RIBEIRO, A., GRANNAN, A., KING, E. M. & AURNOU, J. M. 2015 Laboratory-numerical models of rapidly rotating convection in planetary cores. *Geophys. J. Int.* **201**, 1–17.
- CHILDRESS, S. & SOWARD, A. M. 1972 Convection-driven hydromagnetic dynamo. *Phys. Rev. Lett.* **29** (13), 837–839.
- CIONI, S., CHAUMAT, S. & SOMMERIA, J. 2000 Effect of a vertical magnetic field on turbulent Rayleigh-Bénard convection. *Phys. Rev. E* **62** (4).
- CIONI, S., CILIBERTO, S. & SOMMERIA, J. 1997 Strongly turbulent rayleigh-bénard convection in mercury: comparison with results at moderate Prandtl number. *J. Fluid Mech.* **335**, 111–140.
- FAUVE, S. & PÉTRÉLIS, F. 2007 Scaling laws of turbulent dynamos. *C R Phys* **8** (1), 87–92.
- FRENCH, M., BECKER, A., LORENZEN, W., NETTELMANN, N., BETHKENHAGEN, M., WICHT, J. & REDMER, R. 2012 Ab initio simulations for material properties along the Jupiter adiabat. *Astrophys. J. Supp. Ser.* **202** (1), 5.
- GOLUSKIN, D. & SPIEGEL, E. A. 2012 Convection driven by internal heating. *Phys. Lett. A* **377** (1-2), 83–92.
- GROSSMANN, S. & LOHSE, D. 2000 Scaling in thermal convection: a unifying theory. *J. Fluid Mech.* **407**, 27–56.
- HAUGEN, N. E. L., BRANDENBURG, A. & DOBLER, W. 2004 Simulations of nonhelical hydromagnetic turbulence. *Phys. Rev. E* **70** (1), 016308.

- JONES, C. A. 2011 Planetary magnetic fields and fluid dynamos. *Annu. Rev. Fluid Mech.* **43**, 583–614.
- JONES, C. A. & ROBERTS, P. H. 2000 Convection-driven dynamos in a rotating plane layer. *J. Fluid Mech.* **404**, 311–343.
- KÄPYLÄ, P. J., KÄPYLÄ, M. J. & BRANDENBURG, A. 2018 Small-scale dynamos in simulations of stratified turbulent convection. *Astron. Nachr.* **339** (2-3), 127–133.
- KRAICHNAN, R. H. 1962 Turbulent thermal convection at arbitrary Prandtl number. *Phys. Fluids* **5** (11), 1374–1389.
- LOHSE, D. & TOSCHI, F. 2003 Ultimate state of thermal convection. *Phys. Rev. Lett.* **90** (3), 034502.
- MALKUS, W. V. R. 1954 The heat transport and spectrum of thermal turbulence. *Proc. R. Soc. Lond. A* **225** (1161), 196–212.
- MARTI, P., CALKINS, M. A. & JULIEN, K. 2016 A computationally efficient spectral method for modeling core dynamics. *Geochem. Geophys. Geosys.* **17** (8), 3031–3053.
- MENEGUZZI, M., FRISCH, U. & POUQUET, A. 1981 Helical and nonhelical turbulent dynamos. *Phys. Rev. Lett.* **47** (15), 1060.
- MENEGUZZI, M. & POUQUET, A. 1989 Turbulent dynamos driven by convection. *J. Fluid Mech.* **205**, 297–318.
- MOFFATT, H. K. 1970 Dynamo action associated with random inertial waves in a rotating conducting fluid. *J. Fluid Mech.* **44**, 705–719.
- OSSENDRIJVER, M. 2003 The solar dynamo. *Astron. Astrophys. Rev.* **11** (4), 287–367.
- PANDEY, A., SCHEEL, J. D. & SCHUMACHER, J. 2018 Turbulent superstructures in Rayleigh-Bénard convection. *Nature Comm.* **9** (1), 1–11.
- PARKER, E. N. 1955 Hydromagnetic dynamo models. *Astrophys. J.* **122**, 293–314.
- POPE, S. B. 2000 *Turbulent Flows*. Cambridge: Cambridge University Press.
- POZZO, M., DAVIES, C. J., GUBBINS, D. & ALFÉ, D. 2013 Transport properties for liquid silicon-oxygen-iron mixtures at Earth’s core conditions. *Phys. Rev. B* **87**, 014110.
- QIU, X. L. & TONG, P. 2001 Large-scale velocity structures in turbulent thermal convection. *Phys. Rev. E* **64** (3), 036304.
- RINCON, F. 2019 Dynamo theories. *J. Plasma Phys.* **85** (4).
- SCHAEFFER, N., JAULT, D., NATAF, H.-C. & FOURNIER, A. 2017 Turbulent geodynamo simulations: a leap towards earth’s core. *Geophys. J. Int.* **211** (1), 1–29.
- SCHEEL, J. D. & SCHUMACHER, J. 2017 Predicting transition ranges to fully turbulent viscous boundary layers in low prandtl number convection flows. *Phys. Rev. Fluids* **2**, 123501.
- SCHEKOCIHIN, A. A., BOLDYREV, S. A. & KULSRUD, R. M. 2002a Spectra and growth rates of fluctuating magnetic fields in the kinematic dynamo theory with large magnetic Prandtl numbers. *Astrophys. J.* **567** (2), 828.
- SCHEKOCIHIN, A. A., COWLEY, S. C., HAMMETT, G. W., MARON, J. L. & MCWILLIAMS, J. C. 2002b A model of nonlinear evolution and saturation of the turbulent MHD dynamo. *New J. Phys.* **4** (1), 84.
- SCHEKOCIHIN, A. A., ISKAKOV, A. B., COWLEY, S. C., MCWILLIAMS, J. C., PROCTOR, M. R. E. & YOUSEF, T. A. 2007 Fluctuation dynamo and turbulent induction at low magnetic Prandtl numbers. *New J. Phys.* **9** (8), 300.
- SCHEKOCIHIN, A. A., MARON, J. L., COWLEY, S. C. & MCWILLIAMS, J. C. 2002c The small-scale structure of magnetohydrodynamic turbulence with large magnetic Prandtl numbers. *Astrophys. J.* **576** (2), 806.
- SHEYKO, A., FINLAY, C. C. & JACKSON, A. 2016 Magnetic reversals from planetary dynamo waves. *Nature* **551**, 551–554.
- SHRAIMAN, B. I. & SIGGIA, E. D. 1990 Heat transport in high-Rayleigh-number convection. *Phys. Rev. A* **42** (6), 3650.
- SIMITEV, R. & BUSSE, F. H. 2005 Prandtl-number dependence of convection-driven dynamos in rotating spherical fluid shells. *J. Fluid Mech.* **532**, 365–388.
- SPALART, P. R., MOSER, R. D. & ROGERS, M. M. 1991 Spectral methods for the Navier-Stokes equations with one infinite and two periodic directions. *J. Comp. Phys.* **96**, 297–324.
- SPIEGEL, E. A. 1965 Convective instability in a compressible atmosphere. I. *Astrophys. J.* **141**, 1068–1090.
- STEENBECK, M., KRAUSE, F. & RÄDLER, K.-H. 1966 A calculation of the mean electromotive

- force in an electrically conducting fluid in turbulent motion, under the influence of coriolis forces. *Z. Naturforsch.* **21a**, 369–376.
- STEVENS, R. J., BLASS, A., ZHU, X., VERZICCO, R. & LOHSE, D. 2018 Turbulent thermal superstructures in Rayleigh-Bénard convection. *Phys. Rev. Fluids* **3**, 041501.
- STEVENS, R. J., LOHSE, D. & VERZICCO, R. 2011 Prandtl and Rayleigh number dependence of heat transport in high Rayleigh number thermal convection. *J. Fluid Mech.* **688**, 31–43.
- TOBIAS, S. M. 2019 The turbulent dynamo. *arXiv preprint arXiv:1907.03685* .
- TOBIAS, S. M., CATTANEO, F. & BOLDYREV, S.B. 2012 MHD Dynamos and Turbulence. In *Ten Chapters in Turbulence* (ed. P. A. Davidson, Y. Kaneda & K.R. Sreenivasan), p. 351–404.
- TOBIAS, S. M., CATTANEO, F., BOLDYREV, S., DAVIDSON, P. A., KANEDA, Y. & SREENIVASAN, K. R. 2013 Ten chapters in turbulence. *Cambridge University Press, Cambridge* **1** (1.4), 1–1.
- VOGT, T., HORN, S., GRANNAN, A. M. & AURNOU, J. M. 2018 Jump rope vortex in liquid metal convection. *Proc. Natl. Acad. Sci.* **115** (50), 12674–12679.
- YAN, M., CALKINS, M. A., MAFFEI, S., JULIEN, K., TOBIAS, S. M. & MARTI, P. 2019 Heat transfer and flow regimes in quasi-static magnetoconvection with a vertical magnetic field. *J. Fluid Mech.* **877**, 1186–1206.
- ZHU, X., MATHAI, V., STEVENS, R. J., VERZICCO, R. & LOHSE, D. 2018 Transition to the ultimate regime in two-dimensional Rayleigh-Bénard convection. *Phys. Rev. Lett.* **120** (14), 144502.
- ZÜRNER, T., LIU, W., KRASNOV, D. & SCHUMACHER, J. 2016 Heat and momentum transfer for magnetoconvection in a vertical external magnetic field. *Phys. Rev. E* **94** (4), 043108.

Appendix

The details of our numerical simulations are listed in this section.

Ra	Pr	Pm	$N_x \times N_y \times N_z$	Δt	n	Nu	Re	Rm	E_{mag}
1×10^6	1	0.8	$432 \times 432 \times 192$	4×10^{-6}	2	13.38 ± 0.32	329.9 ± 6.8	263.9 ± 5.4	100 ± 20
2×10^6	1	0.8	$576 \times 576 \times 216$	1×10^{-6}	2	15.91 ± 0.35	447.7 ± 11.1	358.1 ± 8.9	798 ± 47
4×10^6	1	0.8	$576 \times 576 \times 288$	4×10^{-7}	2	19.49 ± 0.49	599.9 ± 13.0	479.9 ± 10.4	$(5.25 \pm 0.62) \times 10^3$
1×10^7	1	0.8	$768 \times 769 \times 288$	2×10^{-7}	2	25.54 ± 0.65	927.5 ± 22.5	742.0 ± 18.0	$(1.51 \pm 0.11) \times 10^4$
1×10^4	1	1	$144 \times 144 \times 64$	1×10^{-4}	5	3.66 ± 0.08	37.6 ± 0.6	37.6 ± 0.6	
2×10^4	1	1	$216 \times 216 \times 64$	1×10^{-4}	5	4.41 ± 0.08	53.9 ± 0.9	53.9 ± 0.9	
4×10^4	1	1	$384 \times 384 \times 72$	1×10^{-4}	5	5.34 ± 0.09	76.4 ± 1.1	76.4 ± 1.1	
1×10^5	1	1	$384 \times 384 \times 96$	6×10^{-5}	5	6.88 ± 0.12	115.9 ± 1.5	115.9 ± 1.5	
2×10^5	1	1	$384 \times 384 \times 96$	4×10^{-5}	5	8.44 ± 0.13	161.2 ± 1.6	161.2 ± 1.6	
3×10^5	1	1	$576 \times 576 \times 108$	1×10^{-5}	4	9.48 ± 0.19	192.3 ± 2.2	192.3 ± 2.2	30 ± 5
4×10^5	1	1	$648 \times 648 \times 108$	5×10^{-6}	4	10.05 ± 0.14	212.2 ± 2.5	212.2 ± 2.5	233 ± 44
6×10^5	1	1	$648 \times 648 \times 144$	2×10^{-6}	4	11.40 ± 0.16	257.9 ± 2.6	257.9 ± 2.6	507 ± 65
1×10^6	1	1	$648 \times 648 \times 192$	1×10^{-6}	3	13.20 ± 0.22	314.6 ± 4.9	314.6 ± 4.9	$(1.52 \pm 0.19) \times 10^3$
2×10^6	1	1	$576 \times 576 \times 216$	6×10^{-7}	2	15.68 ± 0.45	421.4 ± 13.6	421.4 ± 13.6	$(3.96 \pm 0.66) \times 10^3$
4×10^6	1	1	$576 \times 576 \times 288$	2×10^{-7}	2	18.85 ± 0.51	568.4 ± 16.9	568.4 ± 16.9	$(9.33 \pm 0.73) \times 10^3$
1×10^7	1	1	$864 \times 864 \times 432$	1×10^{-7}	2	24.46 ± 0.54	865.4 ± 23.9	865.4 ± 23.9	$(2.02 \pm 0.16) \times 10^4$
1×10^4	1	3	$192 \times 192 \times 72$	2×10^{-5}	5	3.65 ± 0.06	37.3 ± 0.4	111.8 ± 1.3	
1.5×10^4	1	3	$192 \times 192 \times 72$	2×10^{-5}	3	4.14 ± 0.14	46.7 ± 1.5	140.0 ± 4.4	0.2 ± 0.1
2×10^4	1	3	$384 \times 384 \times 72$	1×10^{-5}	3	4.44 ± 0.12	53.4 ± 0.8	160.1 ± 2.4	22 ± 8
3×10^4	1	3	$384 \times 384 \times 96$	1×10^{-5}	3	4.97 ± 0.16	64.8 ± 1.8	194.4 ± 5.3	57 ± 20
4×10^4	1	3	$384 \times 384 \times 96$	1×10^{-5}	3	5.32 ± 0.17	72.7 ± 2.0	218.0 ± 6.1	97 ± 33
6×10^4	1	3	$384 \times 384 \times 96$	4×10^{-6}	3	5.92 ± 0.12	85.8 ± 1.4	257.4 ± 4.2	231 ± 39
1×10^5	1	3	$432 \times 432 \times 108$	2×10^{-6}	3	6.77 ± 0.20	108 ± 2.5	325 ± 7.4	513 ± 79
2×10^5	1	3	$576 \times 576 \times 144$	1×10^{-6}	3	8.15 ± 0.20	145.1 ± 3.3	435 ± 10	967 ± 68
4×10^5	1	3	$648 \times 648 \times 192$	6×10^{-7}	3	9.85 ± 0.23	194.1 ± 3.3	582 ± 10	$(2.05 \pm 0.14) \times 10^3$
1×10^6	1	3	$768 \times 768 \times 288$	2×10^{-7}	3	12.33 ± 0.25	290.2 ± 6.5	871 ± 20	$(5.47 \pm 0.38) \times 10^3$

Table 1: **Details of the numerical simulations for $Pr = 1$, $Pm = (0.8, 1, 3)$ cases.** The non-dimensional parameters are the Rayleigh number Ra , the Prandtl number Pr , the magnetic Prandtl number Pm , the aspect ratio of the domain $\Gamma = \frac{2\pi n}{k_c}$, the Nusselt number Nu , the Reynolds number Re , the magnetic Reynolds number Rm and the magnetic energy E_{mag} . The spatial resolution is quoted in terms of the de-aliased physical space grid points $N_x \times N_y \times N_z$, where (N_x, N_y) is the horizontal resolution and N_z is the vertical resolution. The numerical timestep size is denoted by Δt . The estimated critical Rayleigh numbers for dynamo action (Ra_d) are $(4.9 \times 10^5, 2.2 \times 10^5, 1.6 \times 10^4)$ for $Pm = (0.8, 1, 3)$, respectively.

Ra	Pr	Pm	$N_x \times N_y \times N_z$	Δt	n	Nu	Re	Rm	E_{mag}
2×10^3	1	5	$96 \times 96 \times 48$	5×10^{-4}	5	2.13 ± 0.05	13.4 ± 0.3	67.1 ± 1.3	
4×10^3	1	5	$144 \times 144 \times 48$	2×10^{-4}	5	2.76 ± 0.06	22.1 ± 0.4	110.5 ± 1.8	
5×10^3	1	5	$384 \times 384 \times 96$	2×10^{-4}	5	2.96 ± 0.06	25.4 ± 0.4	126.8 ± 1.9	
6×10^3	1	5	$384 \times 384 \times 96$	1×10^{-5}	5	3.11 ± 0.05	27.8 ± 0.4	139.1 ± 1.9	5 ± 2
7×10^3	1	5	$432 \times 432 \times 96$	1×10^{-5}	5	3.26 ± 0.06	30.5 ± 0.5	152.5 ± 2.7	9 ± 4
8×10^3	1	5	$432 \times 432 \times 96$	1×10^{-5}	5	3.39 ± 0.06	32.8 ± 0.6	163 ± 2.8	15 ± 6
1×10^4	1	5	$432 \times 432 \times 108$	8×10^{-6}	5	3.62 ± 0.07	36.6 ± 0.6	183.0 ± 3.0	23 ± 6
2×10^4	1	5	$432 \times 432 \times 108$	2×10^{-6}	5	4.31 ± 0.06	51.1 ± 0.6	255.5 ± 3.0	97 ± 8
4×10^4	1	5	$576 \times 576 \times 144$	1×10^{-6}	5	5.16 ± 0.07	70.1 ± 0.9	350.4 ± 4.5	262 ± 13
1×10^5	1	5	$576 \times 576 \times 216$	5×10^{-7}	3	6.61 ± 0.21	103.3 ± 2.2	516.8 ± 11.0	744 ± 49
2×10^5	1	5	$768 \times 768 \times 216$	3×10^{-7}	3	8.13 ± 0.18	137.8 ± 3.4	689.3 ± 16.9	$(1.72 \pm 0.08) \times 10^3$
4×10^5	1	5	$648 \times 648 \times 288$	1×10^{-7}	2	9.77 ± 0.40	181.1 ± 3.5	905.7 ± 17.6	$(3.04 \pm 0.28) \times 10^3$
6×10^5	1	5	$648 \times 648 \times 288$	1×10^{-7}	2	10.75 ± 0.42	224.2 ± 4.8	1122 ± 24	$(4.86 \pm 0.24) \times 10^3$
1×10^6	1	5	$864 \times 864 \times 384$	6×10^{-8}	2	12.03 ± 0.42	267 ± 6.1	1338 ± 30	$(6.51 \pm 0.25) \times 10^3$
2×10^6	1	5	$1152 \times 1152 \times 486$	2×10^{-8}	2	14.45 ± 0.47	373 ± 10.3	1869 ± 51	$(1.34 \pm 0.09) \times 10^4$
1×10^7	1	5	$1536 \times 1536 \times 648$	6×10^{-9}	2	22.68 ± 0.67	790 ± 18.9	3948 ± 95	$(5.35 \pm 0.34) \times 10^4$
2×10^3	1	7	$144 \times 144 \times 72$	2×10^{-4}	5	2.13 ± 0.05	13.4 ± 0.2	94.0 ± 1.7	
3×10^3	1	7	$144 \times 144 \times 72$	2×10^{-4}	5	2.50 ± 0.06	18.3 ± 0.3	128.2 ± 2.4	
4×10^3	1	7	$384 \times 384 \times 72$	2×10^{-5}	5	2.74 ± 0.05	21.9 ± 0.4	153.4 ± 3.54	3 ± 1
6×10^3	1	7	$648 \times 648 \times 96$	1×10^{-5}	5	3.09 ± 0.05	27.4 ± 0.4	191.6 ± 3.1	15 ± 3
1×10^4	1	7	$192 \times 192 \times 72$	1×10^{-5}	5	3.57 ± 0.06	36.1 ± 0.7	252.5 ± 4.6	53 ± 6
4×10^4	1	7	$576 \times 576 \times 144$	2×10^{-6}	3	5.09 ± 0.17	66.4 ± 1.8	464.7 ± 12.5	350 ± 43
1×10^5	1	7	$648 \times 648 \times 216$	6×10^{-7}	3	6.52 ± 0.22	98.6 ± 3.1	690.3 ± 21.6	968 ± 86
2×10^5	1	7	$648 \times 648 \times 288$	1×10^{-7}	3	7.66 ± 0.20	132.9 ± 3.6	930.4 ± 25.5	$(1.87 \pm 0.09) \times 10^3$
4×10^5	1	7	$768 \times 768 \times 384$	6×10^{-8}	3	9.27 ± 0.23	180.2 ± 5.1	1261.1 ± 36.0	$(3.75 \pm 0.16) \times 10^3$

Table 2: **Details of the numerical simulations for $Pr = 1$, $Pm = (5, 7)$ cases.** The estimated critical Rayleigh numbers for dynamo action (Ra_d) are $(5.2 \times 10^3, 3.1 \times 10^3)$ for $Pm = (5, 7)$, respectively.

Ra	Pr	Pm	$N_x \times N_y \times N_z$	Δt	n	Nu	Re	Rm	E_{mag}
1×10^3	0.05	1	$96 \times 96 \times 48$	6×10^{-4}	3	1.02 ± 0.01	23.4 ± 1.5	23.4 ± 1.5	
2×10^3	0.05	1	$96 \times 96 \times 48$	2×10^{-4}	3	1.16 ± 0.03	72.0 ± 5.0	72.0 ± 5.0	
4×10^3	0.05	1	$192 \times 192 \times 72$	6×10^{-5}	3	1.40 ± 0.04	133.8 ± 5.5	133.8 ± 5.5	
6×10^3	0.05	1	$192 \times 192 \times 96$	1×10^{-5}	3	1.59 ± 0.05	176.0 ± 7.2	176.0 ± 7.2	171 ± 50
1×10^4	0.05	1	$288 \times 288 \times 144$	4×10^{-6}	3	1.84 ± 0.06	$236.6.0 \pm 7.7$	$236.6.0 \pm 7.7$	735 ± 134
2×10^4	0.05	1	$288 \times 288 \times 192$	1×10^{-6}	2	2.28 ± 0.13	339.96 ± 22.3	339.96 ± 22.3	$(2.91 \pm 0.86) \times 10^3$
4×10^4	0.05	1	$384 \times 384 \times 192$	6×10^{-7}	2	2.83 ± 0.10	500.5 ± 15.5	500.5 ± 15.5	$(6.46 \pm 0.40) \times 10^3$
1×10^5	0.05	1	$768 \times 768 \times 288$	4×10^{-7}	2	3.74 ± 0.13	750.2 ± 17.2	750.2 ± 17.2	$(2.32 \pm 0.24) \times 10^4$

Table 3: **Details of the numerical simulations for $Pr = 0.05$, $Pm = 1$ cases.** The estimated critical Rayleigh numbers for dynamo action (Ra_d) are $2.9 \times 10^3, 4.1 \times 10^3$ for $Pm = 1$, $Pr = (0.01, 0.05)$, respectively.

Ra	Pr	$N_x \times N_y \times N_z$	Δt	n	Nu	Re
2×10^3	1	$96 \times 96 \times 48$	4×10^{-3}	5	2.13 ± 0.05	13.4 ± 0.2
7×10^3	1	$192 \times 192 \times 72$	1×10^{-3}	5	3.30 ± 0.05	31.0 ± 0.4
1×10^4	1	$192 \times 192 \times 72$	5×10^{-4}	5	3.62 ± 0.06	37.5 ± 0.7
4×10^4	1	$384 \times 384 \times 96$	5×10^{-5}	5	5.33 ± 0.07	75.6 ± 1.0
1×10^5	1	$384 \times 384 \times 96$	5×10^{-5}	3	7.01 ± 0.17	119.2 ± 2.3
6×10^5	1	$384 \times 384 \times 144$	1×10^{-5}	2	11.7 ± 0.36	261.0 ± 6.2
1×10^6	1	$432 \times 432 \times 144$	6×10^{-6}	2	13.4 ± 0.34	324.4 ± 9.4
7×10^2	0.05	$96 \times 96 \times 48$	5×10^{-3}	5	1.001 ± 0.000	5.0 ± 0.5
1×10^3	0.05	$96 \times 96 \times 48$	1×10^{-3}	3	1.02 ± 0.00	23.4 ± 1.7
2×10^3	0.05	$96 \times 96 \times 48$	2×10^{-4}	3	1.15 ± 0.03	70.6 ± 5.0
4×10^3	0.05	$144 \times 144 \times 72$	5×10^{-5}	3	1.38 ± 0.04	131.3 ± 6.8
1×10^4	0.05	$288 \times 288 \times 72$	2×10^{-5}	3	1.89 ± 0.07	247.5 ± 8.8
2×10^4	0.05	$432 \times 432 \times 108$	1×10^{-5}	3	2.31 ± 0.09	360.3 ± 13.6
4×10^4	0.05	$432 \times 432 \times 108$	3×10^{-6}	3	2.90 ± 0.12	509.4 ± 13.5

Table 4: Details of the RBC cases.

Ra	Pr	Pm	$N_x \times N_y \times N_z$	Δt	n	Rm	γ
4×10^5	1	0.8	$384 \times 384 \times 96$	3×10^{-5}	4	176	-4.4
6×10^5	1	0.8	$384 \times 384 \times 96$	2×10^{-5}	4	210	5.5
2×10^5	1	1	$384 \times 384 \times 96$	4×10^{-5}	4	161	-1.1
2.5×10^5	1	1	$384 \times 384 \times 96$	2×10^{-5}	4	177	1.7
1×10^4	1	3	$192 \times 192 \times 72$	2×10^{-5}	3	112	-4.3
1.5×10^4	1	3	$192 \times 192 \times 72$	2×10^{-5}	3	141	0.6
2×10^4	1	3	$384 \times 384 \times 72$	1×10^{-5}	3	160	2.4
4×10^3	1	5	$144 \times 144 \times 48$	2×10^{-4}	5	110	-2.2
6×10^3	1	5	$384 \times 384 \times 96$	1×10^{-5}	5	141	1.4
2×10^3	1	7	$144 \times 144 \times 72$	2×10^{-4}	5	94	-3.4
2.5×10^3	1	7	$144 \times 144 \times 72$	2×10^{-4}	5	112	-2.6
3.5×10^3	1	7	$384 \times 384 \times 72$	2×10^{-5}	5	142	1.2
4×10^3	1	7	$384 \times 384 \times 72$	2×10^{-5}	5	152	3.5
3.5×10^3	0.05	1	$192 \times 192 \times 72$	4×10^{-5}	3	122	-1.9
4.5×10^3	0.05	1	$192 \times 192 \times 72$	4×10^{-5}	3	147	1.0
2×10^3	0.01	1	$192 \times 192 \times 72$	2×10^{-5}	3	85	-5.8
2.7×10^3	0.01	1	$192 \times 192 \times 72$	2×10^{-5}	3	125	-1.9
3×10^3	0.01	1	$192 \times 192 \times 72$	2×10^{-5}	3	137	0.9
4×10^3	0.01	1	$192 \times 192 \times 72$	2×10^{-5}	3	188	6.4

Table 5: **Details of the numerical simulations used for the kinematic dynamo regime.** The growth (or decay) rate of magnetic energy is denoted by γ .