



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/174267/>

Version: Accepted Version

Article:

Brimkov, B., Duna, K., Hogben, L. et al. (2020) Graphs that are cospectral for the distance Laplacian. *The Electronic Journal of Linear Algebra*, 36. pp. 334-351.

<https://doi.org/10.13001/ela.2020.4941>

© 2018 The Authors. For re-use permissions, please contact the authors.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

GRAPHS THAT ARE COSPECTRAL FOR THE DISTANCE LAPLACIAN

BORIS BRIMKOV*, KEN DUNA†, LESLIE HOGBEN‡, KATE LORENZEN§,
CAROLYN REINHART§, SUNG-YELL SONG§, AND MARK YARROW¶

Abstract. The distance matrix $\mathcal{D}(G)$ of a graph G is the matrix containing the pairwise distances between vertices, and the distance Laplacian matrix is $\mathcal{D}^L(G) = T(G) - \mathcal{D}(G)$, where $T(G)$ is the diagonal matrix of row sums of $\mathcal{D}(G)$. We establish several general methods for producing \mathcal{D}^L -cospectral graphs that can be used to construct infinite families. We provide examples showing that various properties are not preserved by \mathcal{D}^L -cospectrality, including examples of \mathcal{D}^L -cospectral strongly regular and circulant graphs. We establish that the absolute values of coefficients of the distance Laplacian characteristic polynomial are decreasing, i.e., $|\delta_1^L| \geq \dots \geq |\delta_n^L|$ where δ_k^L is the coefficient of x^k .

Key words. Distance Laplacian matrix, cospectrality, unimodality

AMS subject classifications. 05C50, 05C12, 15A18, 15B57

1. Introduction.

Spectral graph theory is the study of graphs via the eigenvalues of a matrix defined from the graph. Often information about the graph can be recovered from the spectrum of an associated matrix $M(G)$, but the existence of M -cospectral graphs implies there is information that cannot be recovered from the spectrum. Nonisomorphic graphs G and H are said to be M -cospectral if $\text{spec}(M(G)) = \text{spec}(M(H))$. Constructions that switch certain edges to produce adjacency cospectral graphs and distance cospectral graphs have been introduced by Godsil and McKay [13] and Heysse [17]. Aouchiche and Hansen give a thorough discussion of the history of M -cospectrality for various types of matrices M defined from a graph in [3, Section 2].

This paper focuses on the distance Laplacian matrix of a graph, which was introduced in [3] and is defined from the distance matrix in analogy with the way the Laplacian matrix is defined from the adjacency matrix (precise definitions of graph theory terms are provided below). The *distance matrix* $\mathcal{D}(G)$ of a connected graph G is the matrix indexed by the vertices $\{v_1, \dots, v_n\}$ of G having i, j -entry equal to the distance between the vertices v_i and v_j . For $v \in V(G)$, the *transmission index* of v is $t(v) = \sum_{w \in V(G)} d(v, w)$. The *distance Laplacian matrix* is $\mathcal{D}^L(G) = T(G) - \mathcal{D}(G)$ where $T(G) = \text{diag}(t(v_1), \dots, t(v_n))$. Distance matrices were introduced [16] for analysis of the loop switching problem in telecommunications, which involves finding appropriate addresses so that a message can move efficiently through a series of loops from its origin to its destination, choosing the best route at each switching point. Recently there has been renewed interest in the loop switching problem and extensive work on distance spectra; see [4] for a survey. The spectrum of $\mathcal{D}^L(G)$ has been related to various graph parameters such as the connectivity of the complement of G [3].

*Department of Computational and Applied Mathematics, Rice University, Houston, TX, 77005, USA (boris.brimkov@rice.edu)

†Department of Mathematics, University of Kansas, Lawrence, KS, 66045, USA (kduna@ku.edu)

‡Department of Mathematics, Iowa State University, Ames, IA 50011, USA and American Institute of Mathematics, 600 E. Brokaw Road, San Jose, CA 95112, USA (hogben@aimath.org).

§Department of Mathematics, Iowa State University, Ames, IA 50011, USA (lorenkj@iastate.edu, reinh196@iastate.edu, sysong@iastate.edu)

¶School of Mathematics and Statistics, University of Sheffield, Sheffield, S3 7RH, United Kingdom (M-Yarrow1@sheffield.ac.uk)

In Section 2 we provide examples to show that various graph properties are not preserved by \mathcal{D}^L -cospectrality. We establish some general methods for producing \mathcal{D}^L -cospectral graphs and apply one to construct an infinite family of \mathcal{D}^L -cospectral bipartite pairs in Section 3. These constructions do not generally produce cospectral distance matrices because they rely on the fact that $\mathcal{D}^L(G)$ is positive semidefinite with all row sums equal to zero. It is clear that if $T(G)$ is a scalar matrix, then there are formulas that determine the eigenvalues of $\mathcal{D}^L(G)$ from the eigenvalues of $\mathcal{D}(G)$ (and vice versa). Thus, the study of the distance spectrum and the study of the distance Laplacian spectrum are the same for transmission regular graphs; we apply this to \mathcal{D}^L -cospectral strongly regular and circulant graphs in Section 4. In Section 5 we establish that the absolute values of coefficients of the distance Laplacian characteristic polynomial are log-concave, unimodal, and in fact decreasing, i.e., $|\delta_1^L| \geq \dots \geq |\delta_n^L|$ where $|\delta_k^L|$ is the coefficient of x^k . The remainder of this introduction contains additional definitions and terminology.

All graphs in this paper are simple (no loops or multiedges) and undirected. For a graph G , the set of vertices is denoted by $V(G)$ and the set of edges (2-element subsets of vertices) is denoted by $E(G)$; an edge $\{u, v\}$ is also denoted by uv . The order of G is the number $n = |V(G)|$ of vertices. The edge $e = uv$ is *incident* to vertices u and v , and u and v are said to be *adjacent*. Adjacent vertices are called *neighbors*, and the *neighborhood* of vertex v , $N(v)$, is the set of all neighbors of v . The *degree* of a vertex v is $\deg v = |N(v)|$. If there exists a k such that $k = \deg(v)$ for all $v \in V(G)$, then G is said to be *k-regular* or *regular*. A *u, v-path* in G is a list of vertices that start at u and end at v such that any two consecutive vertices in the list form an edge in $E(G)$ and no vertex is repeated. A graph is *connected* if for every pair of vertices u, v there exists a *u, v-path*. The length of a path is one less than the number of vertices (i.e., is the number of edges), and the *distance* between two vertices $d(u, v)$ is the length of the shortest *u, v-path*. A graph must be connected in order for the distance or distance Laplacian matrix to be defined. An *isomorphism* from graph G to graph H is a bijection $f : V(G) \rightarrow V(H)$ such that $uv \in E(G)$ if and only if $f(u)f(v) \in E(H)$ for all $u, v \in V(G)$.

A *subgraph* of G is a graph H such that $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$. If G is a graph, $u, v \in V(G)$, and $uv \notin E(G)$, then $G + uv$ is the graph obtained from G by adding edge uv , i.e., $V(G + uv) = V(G)$ and $E(G + uv) = E(G) \cup \{uv\}$; G is a subgraph of $G + uv$. Given $S \subseteq V(G)$, the *induced subgraph* $G[S]$ is the subgraph of G with vertex set S and edge set consisting of all edges of G that have both endpoints in S . A graph G is *bipartite* if $V(G) = A \cup B$, $A \cap B = \emptyset$, and for every $e \in E$, exactly one vertex of e is in A and exactly one vertex is in B ; in this case A and B are called *parts*. A graph is *planar* if it can be drawn in the plane without any edges crossing. The *diameter* of G , denoted by $\text{diam}(G)$, is the greatest distance between any two vertices in G . A *cycle* is a list of vertices in G such that any two consecutive vertices in the list form an edge in $E(G)$ and no vertex is repeated except that the last vertex equals the first vertex. The *girth* of G is the length of the shortest cycle in G . A *dominating vertex* is a vertex adjacent to every other vertex. A *cut-vertex* in a connected graph is a vertex whose deletion disconnects the graph (to *delete* a vertex from a graph means to delete the vertex and all edges incident with it).

The *adjacency matrix* of a graph G is the matrix indexed by the vertices $\{v_1, \dots, v_n\}$ of G having i, j -entry equal to one if $\{i, j\}$ is an edge and zero otherwise, and is denoted by $A(G)$. The *Laplacian matrix* of G is $L(G) = \text{diag}(\deg v_1, \dots, \deg v_n) - A(G)$. The eigenvalues of $\mathcal{D}(G)$ are called *distance eigenvalues* and are denoted by $\partial_1 \geq \partial_2 \geq \dots \geq \partial_n$. The *distance spectrum* of G is the multiset of distance eigenvalues of G and is denoted by $\text{spec}_{\mathcal{D}}(G)$. The *distance Laplacian eigenvalues* $\partial_1^L = 0 \leq \partial_2^L \leq \dots \leq \partial_n^L = \rho(\mathcal{D}^L(G))$ are eigenvalues of $\mathcal{D}^L(G)$, where ρ denotes the spectral radius (observe that the ordering of the eigenvalues is reversed). The *distance Laplacian spectrum* of G is the multiset of distance Laplacian eigenvalues of G and is denoted by $\text{spec}_{\mathcal{D}^L}(G)$. For a graph G of order n , the *average transmission* of G is $t(G) = \frac{1}{n} \sum_{i=1}^n t(v_i)$.

EXAMPLE 2.2. The graphs G_1 and G_2 shown in Figure 2.2 are \mathcal{D}^L -cospectral because the characteristic polynomials are the same: $p_{\mathcal{D}^L(G_1)}(x) = p_{\mathcal{D}^L(G_2)}(x) = x^9 - 130x^8 + 7369x^7 - 237912x^6 + 4785415x^5 - 61411718x^4 + 491069091x^3 - 2237203064x^2 + 4446151812x$. Observe that G_1 has 13 edges, girth 3, and 5 pairs of vertices at distance 3 from each other, whereas G_2 has 12 edges, girth 4, and 6 pairs of vertices at distance 3. Thus, the number of edges, the girth, and the multiset of distances are not preserved by \mathcal{D}^L -cospectrality.

Figure 2.3 presents a pair of \mathcal{D}^L -cospectral graphs showing that \mathcal{D}^L -cospectrality does not preserve the diameter. Figures 2.4, 2.5, 2.6, and 2.7 present pairs of \mathcal{D}^L -cospectral graphs that establish \mathcal{D}^L -cospectrality does not preserve the following properties: having a leaf, having a dominating vertex, having a cut-vertex, and having a nontrivial automorphism.




Figure 2.3: Observe that $\text{diam}(G_1) = 3$, $\text{diam}(G_2) = 2$, and $p_{\mathcal{D}^L(G_1)}(x) = p_{\mathcal{D}^L(G_2)}(x) = x^{10} - 132x^9 + 7720x^8 - 262558x^7 + 5722578x^6 - 82891102x^5 + 797942816x^4 - 4922528058x^3 + 17658758885x^2 - 28066657350x$.




Figure 2.4: Observe that $\text{spec}_{\mathcal{D}^L}(G_1) = \text{spec}_{\mathcal{D}^L}(G_2) = \{0, 8, 10, 12, 12, 13, 14, 15\}$ and G_1 has a leaf whereas G_2 does not.




Figure 2.5: Observe that $\text{spec}_{\mathcal{D}^L}(G_1) = \text{spec}_{\mathcal{D}^L}(G_2) = \{0, 7, 9, 10, 10, 12, 12\}$ and G_1 has a dominating vertex whereas G_2 does not.

A *clique* is a set $S \subseteq V(G)$ such that $x, y \in S$ implies x and y are adjacent, and an *independent set* is a




Figure 2.6: Observe that $p_{\mathcal{D}^L(G_1)}(x) = p_{\mathcal{D}^L(G_2)}(x) = x^8 - 84x^7 + 3000x^6 - 59072x^5 + 692816x^4 - 4841152x^3 + 18666240x^2 - 30643200x$ and G_1 has a cut vertex whereas G_2 does not.




Figure 2.7: Observe that $p_{\mathcal{D}^L(G_1)}(x) = p_{\mathcal{D}^L(G_2)}(x) = x^8 - 84x^7 + 3007x^6 - 59448x^5 + 700756x^4 - 4923264x^3 + 19080000x^2 - 31449600x$ and G_1 has symmetry, i.e., it has a nontrivial automorphism, whereas G_2 does not, i.e., it only has the trivial automorphism.

set $S \subseteq V(G)$ such that $x, y \in S$ implies x and y are not adjacent. The *clique number* $\omega(G)$ is the maximum cardinality of a clique of G , and *independence number* $\alpha(G)$ is the maximum cardinality of an independent set of G . Example 4.5 below shows that the clique number and independence number are not preserved by \mathcal{D}^L -cospectrality. Example 4.8 shows that the property of being a circulant graph is not preserved (see Section 4.2 for the definition of a circulant graph). The bipartite cospectral pair on eight vertices shown in Figure 2.8 establishes that planarity is not preserved, because $B_1 + v_1v_3$ is planar but $B_1 + v_2v_4$ is not planar. Their \mathcal{D}^L -cospectrality is proved in Theorem 3.14 below.




Figure 2.8: Observe that $p_{\mathcal{D}^L(B_1+v_1v_3)}(x) = p_{\mathcal{D}^L(B_1+v_2v_4)}(x) = x^8 - 98x^7 + 4087x^6 - 94020x^5 + 1288463x^4 - 10517842x^3 + 47349497x^2 - 90671880x$ and $B_1 + v_1v_3$ is planar whereas $B_1 + v_2v_4$ is not.

3. Constructions of \mathcal{D}^L -cospectral graphs. In this section we provide constructions of \mathcal{D}^L -cospectral graphs that use sets of twin vertices, or sets of vertex pairs with a relaxation of the twin structure. One of these constructions explains the only pair of cospectral bipartite graphs on eight vertices and provides a way to build large bipartite \mathcal{D}^L -cospectral graphs on an even number of vertices.

Let v_1, v_2 be vertices of graph G such that $N(v_1) = N(v_2)$. Such vertices are called *twins* (technically, these are *independent* or *non-adjacent* twins, but we use the term *twins* to mean *independent twins*). Note

that twins have the same transmission.

DEFINITION 3.1. If v_1 and v_2 are twins, v_3 and v_4 are twins, and $t(v_1) = t(v_3)$, then we say that $\{\{v_1, v_2\}, \{v_3, v_4\}\}$ is a set of *co-transmission twins*.

In [22], the authors prove that if v_1 and v_2 are twin vertices in graph G where $\text{spec}_{\mathcal{D}^L}(G)$ and $\text{spec}_{\mathcal{D}^L}(G + v_1v_2)$ differ only by one eigenvalue of G decreasing by 2 while the others remain the same, then the changed eigenvalue of G is $t(v_1) + 2$. We establish a stronger statement that we use to prove our first construction.

PROPOSITION 3.2. *Suppose v_1 and v_2 are twins. Then there is an orthogonal basis of eigenvectors of $\mathcal{D}^L(G)$ that includes $[1, -1, 0, \dots, 0]^T$, which is associated with eigenvalue $t_G(v_1) + 2$, and all other eigenvectors in the basis are of the form $[a, a, y_3, \dots, y_n]^T$ for some $a, y_i \in \mathbb{R}$. Furthermore, this basis of eigenvectors for $\mathcal{D}^L(G)$ is also a basis of eigenvectors for $\mathcal{D}^L(G + v_1v_2)$, the eigenvalue of $\mathcal{D}^L(G + v_1v_2)$ for eigenvector $[1, -1, 0, \dots, 0]^T$ is $t_G(v_1)$, and the eigenvalues of $\mathcal{D}^L(G + v_1v_2)$ for other eigenvectors are unchanged.*

Proof. Since v_1 and v_2 are twins, it follows that $(\mathcal{D}^L(G))_{j1} = (\mathcal{D}^L(G))_{j2}$ for all $j = 3, \dots, n$ and $t(v_1) = t(v_2)$. In addition, $v_1v_2 \notin E(G)$ and there exists a path of length two between v_1 and v_2 because they have the same neighborhood. Therefore the 2×2 principal submatrix associated with v_1 and v_2 is of the form $\begin{bmatrix} t_G(v_1) & -2 \\ -2 & t_G(v_2) \end{bmatrix}$. For $\mathbf{x} = [1, -1, 0, \dots, 0]^T$, $\mathcal{D}^L(G)\mathbf{x} = [t_G(v_1) + 2, -t_G(v_2) - 2, 0, \dots, 0]^T$ because after the first two rows, all the entries of the first two columns are equal. Since $t_G(v_1) = t_G(v_2)$, it follows that \mathbf{x} is an eigenvector with eigenvalue $t_G(v_1) + 2$. Since $\mathcal{D}^L(G)$ is a real symmetric matrix, it has a basis of orthogonal eigenvectors. Thus there is a basis of eigenvectors in which every eigenvector $\mathbf{y} \neq \mathbf{x}$ is of the form $\mathbf{y} = [a, a, y_3, \dots, y_n]^T$ for some $a \in \mathbb{R}$.

Since v_1, v_2 share a neighborhood, it follows that adding the edge v_1v_2 to G changes only the distance between v_1 and v_2 . Therefore

$$\mathcal{D}^L(G + v_1v_2) = \mathcal{D}^L(G) + \left(\begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \oplus O_{n-2, n-2} \right).$$

For $\mathbf{x} = [1, -1, 0, \dots, 0]^T$,

$$\mathcal{D}^L(G + v_1v_2)\mathbf{x} = [t_G(v_1) + 2, -t_G(v_1) - 2, 0, \dots, 0]^T + [-2, 2, 0, \dots, 0]^T = [t_G(v_1), -t_G(v_1), 0, \dots, 0]^T.$$

Thus \mathbf{x} is an eigenvector of $G + v_1v_2$ with eigenvalue $t_G(v_1)$. Let $\mathbf{y} = [a, a, y_3, \dots, y_n]^T$ be an eigenvector of $\mathcal{D}^L(G)$. Since $\mathcal{D}^L(G + v_1v_2) = \mathcal{D}^L(G) + \left(\begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \oplus O_{n-2, n-2} \right)$, $\mathcal{D}^L(G + v_1v_2)\mathbf{y} = \mathcal{D}^L(G)\mathbf{y}$. Therefore the eigenvalues of $\mathcal{D}^L(G + v_1v_2)$ associated with other eigenvectors in the orthogonal basis of eigenvectors of $\mathcal{D}^L(G)$ are unchanged. \square

COROLLARY 3.3. *Let G and H be \mathcal{D}^L -cospectral, and let $v_1, v_2 \in V(G)$, $v_3, v_4 \in V(H)$ be such that v_1, v_2 are twins, v_3, v_4 are twins and $t(v_1) = t(v_3)$. Then $G + v_1v_2$ and $H + v_3v_4$ are \mathcal{D}^L -cospectral. In particular, if G is a graph with a set of co-transmission twins $\{\{v_1, v_2\}, \{v_3, v_4\}\}$, then $G + v_1v_2$ and $G + v_3v_4$ are \mathcal{D}^L -cospectral.*

EXAMPLE 3.4. The construction of \mathcal{D}^L -cospectral graphs starting with a single graph and adding edges as described in Corollary 3.3 is illustrated in Figure 3.1. With G , $G_1 = G + v_1v_2$, and $G_2 = G + v_3v_4$ as shown, v_1 and v_2 are twins, v_3 and v_4 are twins, and $t(v_1) = 15 = t(v_3)$. Furthermore, $p_{\mathcal{D}^L(G_1)}(x) = p_{\mathcal{D}^L(G_2)}(x) = x^8 - 104x^7 + 4601x^6 - 112224x^5 + 1629571x^4 - 14083840x^3 + 67065731x^2 - 135702840x$.

Next we show how to create a graph G that has a set of co-transmission twins by *twinning* a vertex v in G , which adds a new vertex v' to G such that $N(v') = N(v)$.




Figure 3.1: The graphs $G_1 = G + v_1v_2$ and $G_2 = G + v_3v_4$ produced as in Corollary 3.3

PROPOSITION 3.5. *Let G be a graph with distinct vertices u and v such that $t(u) = t(v)$. Let G' be the graph obtained from G by twinning each of u and v to obtain u' and v' , respectively. Then $\{\{u, u'\}, \{v, v'\}\}$ is a set of co-transmission twins of G' .*

Proof. Let $d = d_G(u, v)$. We know that u' is distance two away from u and d away from v , and v' is distance d away from u and two away from v . Therefore, we have increased the transmissions of u and v from G to G' by $d + 2$. By construction, $t(u') = t(u)$ and $t(v') = t(v)$. Therefore, $\{\{v, v'\}, \{u, u'\}\}$ are co-transmission twins. \square

Next we extend the construction of \mathcal{D}^L -cospectral graphs from a set of co-transmission twins to sets of vertices called cousins that satisfy somewhat more relaxed conditions.

DEFINITION 3.6. Let G be a graph of order at least five with $v_1, v_2, v_3, v_4 \in V(G)$. Let $C = \{\{v_1, v_2\}, \{v_3, v_4\}\}$ and $U(C) = V(G) \setminus \{v_1, v_2, v_3, v_4\}$. Then C is a set of cousins in G if the following conditions are satisfied:

1. For all $u \in U(C)$, $d_G(u, v_1) = d_G(u, v_2)$ and $d_G(u, v_3) = d_G(u, v_4)$.
2. $\sum_{u \in U(C)} d_G(u, v_1) = \sum_{u \in U(C)} d_G(u, v_3)$.

Note that no structure is assumed between v_1, v_2, v_3, v_4 . In the case when $v_1v_2, v_3v_4 \notin E(G)$, it follows directly from the definition that $N(v_1) \setminus \{v_3, v_4\} = N(v_2) \setminus \{v_3, v_4\}$ and $N(v_3) \setminus \{v_1, v_2\} = N(v_4) \setminus \{v_1, v_2\}$.

PROPOSITION 3.7. *Let G be a graph with a set of cousins $C = \{\{v_1, v_2\}, \{v_3, v_4\}\}$. Then $N(v_1) \cap N(v_2) \neq \emptyset$ and $N(v_3) \cap N(v_4) \neq \emptyset$.*

Proof. Since $|V(G)| \geq 5$, $U(C) \neq \emptyset$. Since G is connected, there must be some vertex $u \in U(C)$ adjacent to at least one of v_1, v_2, v_3, v_4 . Without loss of generality, suppose $u \in N(v_1)$. Since u must also be the same distance from v_2 , u is also adjacent to v_2 , and thus $N(v_1) \cap N(v_2) \neq \emptyset$. Suppose to the contrary that $N(v_3) \cap N(v_4) = \emptyset$, which implies $N(v_3) \cap U(C) = \emptyset$ and $N(v_4) \cap U(C) = \emptyset$. Since G is connected, it follows without loss of generality that v_3 is adjacent to v_1 . Then for all $u \in U(C)$, the shortest path between u and v_3 uses the edge v_1v_3 . Thus $d(u, v_3) = d(u, v_1) + 1$, and $\sum_{u \in U(C)} d(u, v_3) = (n - 4) + \sum_{u \in U(C)} d(u, v_1)$, which is a contradiction. \square

LEMMA 3.8. *Let G be a graph with set of cousins $C = \{\{v_1, v_2\}, \{v_3, v_4\}\}$ such that $v_1v_2, v_3v_4 \notin E(G)$. Then $d_{G+v_1v_2}(u, v) = d_G(u, v) = d_{G+v_3v_4}(u, v)$ for all $u \in U(C)$ and $v \in V(G)$.*

Proof. Let $u \in U(C)$ and $v \in V(G)$. If a shortest path between u and v in $G + v_1v_2$ does not contain edge v_1v_2 and a shortest path between u and v in $G + v_3v_4$ does not contain edge v_3v_4 , then the distance between u and v is the same in G , $G + v_1v_2$, and $G + v_3v_4$. We show that if a shortest path \mathcal{P} from u to v in $G + v_1v_2$ contains v_1v_2 , then there is a shorter path between u and v in $G + v_1v_2$, contradicting the choice of \mathcal{P} . The argument for $G + v_3v_4$ is similar.

Suppose a shortest path \mathcal{P} from u to v in $G + v_1v_2$ contains the edge v_1v_2 ; without loss of generality, v_1 precedes v_2 in the path order starting from u . Since $u \in U(C)$, there is a predecessor w of v_1 in \mathcal{P} , i.e. \mathcal{P} contains the subpath (w, v_1, v_2) . If $w \in U(C)$, then $w \in N(v_1) \cap N(v_2)$ and so replacing (w, v_1, v_2) by (w, v_2) produces a shorter subpath. So w is one of v_3 or v_4 , without loss of generality let $w = v_3$. Since $v_3 \notin U(C)$, \mathcal{P} contains the subpath (u', v_3, v_1, v_2) for some $u' \in V(G)$. Since $v_3v_4 \notin E(G + v_1v_2)$, $u' \neq v_4$. Thus $u' \in U(C)$. If $u'v_1 \in E(G)$ there would be a shorter path between u and v obtained by replacing (u', v_3, v_1) by (u', v_1) in \mathcal{P} . Since this would contradict the choice of \mathcal{P} , $u'v_1 \notin E(G)$. Thus $d_G(u', v_1) = 2$, which implies $d_G(u', v_2) = 2$ by the definition of a set of cousins. So there exists $x \in N(u') \cap N(v_2)$. Then there is a shorter path between u and v obtained by replacing (u', v_3, v_1, v_2) by (u', x, v_2) . \square

LEMMA 3.9. *Let H be a graph with $V(H) = \{u_1, u_2, u_3, u_4\}$ such that $u_1u_2, u_3u_4 \notin E(H)$ and suppose $H + u_1u_2$ is isomorphic to $H + u_3u_4$. Then at least one of the permutations $\sigma_1 = (u_1u_4)(u_2u_3)$ or $\sigma_2 = (u_1u_3)(u_2u_4) = (u_1u_2)\sigma_1(u_1u_2)$ in the symmetric group on $V(H)$ is an isomorphism of $H + u_1u_2$ and $H + u_3u_4$.*

Proof. This lemma was verified by exhaustively checking every labeling of every graph on four vertices using Sage [10]. \square

THEOREM 3.10. *Let G be a graph with a set of cousins $C = \{\{v_1, v_2\}, \{v_3, v_4\}\}$ satisfying the following conditions:*

- *Vertices v_1, v_2 are not adjacent and v_3, v_4 are not adjacent.*
- *The subgraph of $G + v_1v_2$ induced by $\{v_1, v_2, v_3, v_4\}$ is isomorphic to the subgraph of $G + v_3v_4$ induced by $\{v_1, v_2, v_3, v_4\}$.*

If $G_1 = G + v_1v_2$ and $G_2 = G + v_3v_4$ are not isomorphic, then they are \mathcal{D}^L -cospectral.

Proof. By Lemma 3.8, we know that the entries of $\mathcal{D}^L(G_1)$, $\mathcal{D}^L(G_2)$, and $\mathcal{D}^L(G)$ are equal everywhere except the 4×4 principal submatrix corresponding to v_1, v_2, v_3, v_4 . Let $M_1 = \mathcal{D}^L(G_1)[\{1, 2, 3, 4\}]$ and $M_2 = \mathcal{D}^L(G_2)[\{1, 2, 3, 4\}]$. Then we can partition $\mathcal{D}^L(G_1)$ and $\mathcal{D}^L(G_2)$ as block matrices:

$$(3.1) \quad \mathcal{D}^L(G_1) = \begin{bmatrix} M_1 & Q \\ Q^T & B \end{bmatrix} \text{ and } \mathcal{D}^L(G_2) = \begin{bmatrix} M_2 & Q \\ Q^T & B \end{bmatrix}$$

where Q is a $4 \times (n - 4)$ matrix and B is an $(n - 4) \times (n - 4)$ matrix.

From the definition of a set of cousins, $\sum_{u \in U(C)} d(u, v_1) = \sum_{u \in U(C)} d(u, v_2) = \sum_{u \in U(C)} d(u, v_3) = \sum_{u \in U(C)} d(u, v_4)$. Thus the row sums of Q are constant. Since every row sum of a distance Laplacian matrix is equal to zero, it follows that the row sums of M_1 are constant, the row sums of M_2 are constant, and these constants are equal; denote the constant row sum of $M_i, i = 1, 2$ by a . Thus M_1 has the form

$$M_1 = \begin{bmatrix} a + 1 + c + d & -1 & -c & -d \\ -1 & a + 1 + f + h & -f & -h \\ -c & -f & a + c + f + 2 & -2 \\ -d & -h & -2 & a + d + 2 + h \end{bmatrix}$$

for some $c, d, f, h \in \mathbb{Z}$. The induced subgraph $G_1[\{v_1, v_2, v_3, v_4\}]$ is isomorphic to the induced subgraph of $G_2[\{v_1, v_2, v_3, v_4\}]$ by hypothesis. By Lemma 3.9, we can choose the isomorphism to be either σ_1 or σ_2 . So

M_2 has one of the forms

$$M_2^{\sigma_1} = \begin{bmatrix} a+d+h+2 & -2 & -h & -d \\ -2 & a+c+f+2 & -f & -c \\ -h & -f & a+1+f+h & -1 \\ -d & -c & -1 & a+1+c+d \end{bmatrix}$$

or

$$M_2^{\sigma_2} = \begin{bmatrix} a+d+h+2 & -2 & -c & -f \\ -2 & a+c+f+2 & -d & -h \\ -c & -d & a+1+f+h & -1 \\ -f & -h & -1 & a+1+c+d \end{bmatrix}$$

corresponding to σ_1 or σ_2 , respectively.

Define

$$S_1 = \begin{bmatrix} 1/2 & 1/2 & 1/2 & -1/2 \\ 1/2 & 1/2 & -1/2 & 1/2 \\ 1/2 & -1/2 & 1/2 & 1/2 \\ -1/2 & 1/2 & 1/2 & 1/2 \end{bmatrix}, \quad S_2 = \begin{bmatrix} 1/2 & 1/2 & -1/2 & 1/2 \\ 1/2 & 1/2 & 1/2 & -1/2 \\ -1/2 & 1/2 & 1/2 & 1/2 \\ 1/2 & -1/2 & 1/2 & 1/2 \end{bmatrix},$$

and $\mathcal{S}_i = S_i \oplus I_{n-4}$ for $i = 1, 2$. Observe that each of S_1, S_2, \mathcal{S}_1 , and \mathcal{S}_2 is its own inverse and transpose (S_1 and S_2 are scalar multiples of Hadamard matrices). Suppose σ_1 maps $G_1[\{v_1, v_2, v_3, v_4\}]$ to $G_2[\{v_1, v_2, v_3, v_4\}]$ and consider

$$\mathcal{S}_1 \mathcal{D}^L(G_1) \mathcal{S}_1 = \begin{bmatrix} S_1 M_1 S_1 & S_1 Q \\ (S_1 Q)^T & B \end{bmatrix}.$$

By direct computation [10], we can verify that $S_1 M_1 S_1 = M_2^{\sigma_1}$. Every column of Q is of the form $[-p, -p, -q, -q]^T$, and $S_1[-p, -p, -q, -q]^T = [-p, -p, -q, -q]^T$. Therefore $S_1 Q = Q$. We have shown that $\mathcal{S}_1 \mathcal{D}^L(G_1) \mathcal{S}_1 = \mathcal{D}^L(G_2)$ and thus the graphs are cospectral. The proof that similarity by \mathcal{S}_2 transforms $\mathcal{D}^L(G_1)$ to $\mathcal{D}^L(G_2)$ when σ_2 is the isomorphism is analogous. \square

Using a set of co-transmission twins to construct \mathcal{D}^L -cospectral graphs (as in Corollary 3.3) is a special case of Theorem 3.10: A straightforward computation shows that set $\{\{v_1, v_2\}, \{v_3, v_4\}\}$ of co-transmission twins is a set of cousins. The twins relationships require that the subgraph $G[\{v_1, v_2, v_3, v_4\}]$ is four isolated vertices or the complete bipartite graph with parts $\{v_1, v_2\}$ and $\{v_3, v_4\}$, which implies the subgraph of $(G + v_1 v_2)[\{v_1, v_2, v_3, v_4\}]$ is isomorphic to $(G + v_3 v_4)[\{v_1, v_2, v_3, v_4\}]$.

Our next main result uses a set of cousins $C = \{\{v_1, v_2\}, \{v_3, v_4\}\}$ but toggles edges $v_1 v_3$ and $v_2 v_4$ rather than $v_1 v_2$ and $v_3 v_4$. We need a preliminary lemma.

LEMMA 3.11. *Let G be a graph with a set of cousins $C = \{\{v_1, v_2\}, \{v_3, v_4\}\}$ such that $v_1 v_3, v_2 v_4 \notin E(G)$, for every $x \in N(v_1) \cap N(v_2)$ there exists $y \in N(v_3) \cap N(v_4)$ such that $xy \in E(G)$, and for every $y \in N(v_3) \cap N(v_4)$ there exists $x \in N(v_1) \cap N(v_2)$ such that $xy \in E(G)$. Then $d_{G+v_1 v_3}(u, v) = d_G(u, v) = d_{G+v_2 v_4}(u, v)$ for all $u \in U(C)$ and $v \in V(G)$.*

Proof. Let $u \in U(C)$ and $v \in V(G)$. If a shortest path between u and v in $G + v_1 v_3$ does not contain edge $v_1 v_3$ and a shortest path between u and v in $G + v_2 v_4$ does not contain edge $v_2 v_4$, then the distance between u and v is the same in $G, G + v_1 v_3$, and $G + v_2 v_4$. We show that there is a shortest path that does not include $v_1 v_3$ between u and v in $G + v_1 v_3$; analogously, there is a shortest path in $G + v_2 v_4$ that does not contain edge $v_2 v_4$.

Suppose a shortest path \mathcal{P} from u to v in $G + v_1v_3$ contains the edge v_1v_3 ; without loss of generality, v_1 precedes v_3 in the path order starting from u . Since $u \in U(C)$, there is a predecessor x of v_1 in \mathcal{P} , i.e. \mathcal{P} contains the subpath (x, v_1, v_3) . We first show that $x \in U(C)$. Since \mathcal{P} is a path, $x \neq v_1, v_3$. Since (x, v_1, v_3) is a subpath of a shortest path between u and v_3 , it cannot be replaced by a shorter subpath. For $u' \in U(C)$, a subpath of the form $(u', v_2, \dots, v_1, v_3)$ can be replaced by (u', v_1, v_3) , and a subpath of the form $(u', v_4, \dots, v_1, v_3)$ can be replaced by (u', v_3) . Thus $x \neq v_2, v_4$, so $x \in U(C)$. Because $x \in U(C)$, there exists $y \in N(v_3) \cap N(v_4)$ such that $xy \in E(G)$, the subpath (x, v_1, v_3) can be replaced by (x, y, v_3) , creating a path of equal length that avoids edge v_1v_3 . \square

THEOREM 3.12. *Let G be a graph with a set of cousins $C = \{\{v_1, v_2\}, \{v_3, v_4\}\}$ satisfying the following conditions:*

- *Vertices v_1, v_3 are not adjacent and v_2, v_4 are not adjacent.*
- *The subgraph of $G + v_1v_3$ induced by $\{v_1, v_2, v_3, v_4\}$ is isomorphic to the subgraph of $G + v_2v_4$ induced by $\{v_1, v_2, v_3, v_4\}$ via the permutation $\sigma_1 = (14)(23)$.*
- *For every $x \in N(v_1) \cap N(v_2)$ there exists $y \in N(v_3) \cap N(v_4)$ such that $xy \in E(G)$, and for every $y \in N(v_3) \cap N(v_4)$ there exists $x \in N(v_1) \cap N(v_2)$ such that $xy \in E(G)$.*

If $G_1 = G + v_1v_3$ and $G_2 = G + v_2v_4$ are not isomorphic, then they are \mathcal{D}^L -cospectral.

Proof. By Lemma 3.11, the entries of $\mathcal{D}^L(G_1)$ are equal to $\mathcal{D}^L(G_2)$ everywhere except the 4×4 principal submatrix corresponding to v_1, v_2, v_3, v_4 . The 4×4 principal submatrices have structure analogous to M_1 and to M_2 with σ_1 , and the remainder of the proof follows the same method. \square

The reason we assume that the isomorphism of $G_1[\{v_1, v_2, v_3, v_4\}]$ and $G_2[\{v_1, v_2, v_3, v_4\}]$ is $\sigma_1 = (14)(23)$ is because isomorphic graphs are not cospectral by definition. Applying Lemma 3.9 with $v_2 = u_3$ and $v_3 = u_2$ produces $\sigma_1 = (v_1 v_4)(v_2 v_3)$ and $\sigma_2 = (v_1 v_2)(v_3 v_4)$. Suppose $G_1[\{v_1, v_2, v_3, v_4\}]$ is isomorphic to $G_2[\{v_1, v_2, v_3, v_4\}]$ via σ_2 . Let P_{σ_2} denote the 4×4 permutation matrix representing $\sigma_2 = (12)(34)$ and $P_2 = P_{\sigma_2} \oplus I_{n-4}$. With the partition of $\mathcal{D}^L(G_1)$ as in (3.1), $P_{\sigma_2}Q = Q$. Thus $P_2\mathcal{D}^L(G_1)P_2 = \mathcal{D}^L(G_2)$ and G_1 is isomorphic to G_2 .

We now discuss an example of Theorem 3.12 on a family of cospectral bipartite graphs. This family contains the only bipartite pair of cospectral graphs on eight vertices.

DEFINITION 3.13. *For $k \in \mathbb{Z}_+$, B_k is the graph defined as follows (and shown in Figure 3.2 not including the dashed edges): Five vertices are denoted by v_0, v_1, v_2, v_3 , and v_4 , and the set of edges of the induced subgraph $B_k[v_0, v_1, v_2, v_3, v_4]$ is $\{v_0v_1, v_0v_2, v_2v_3\}$. There are k twin leaf vertices adjacent to v_0 ; the set of these vertices is denoted by L . There are $k + 1$ twin vertices of degree three each adjacent to v_0, v_3 , and v_4 ; the set of these vertices is denoted by R .*

THEOREM 3.14. *The graphs $B_k + v_1v_3$ and $B_k + v_2v_4$ are \mathcal{D}^L -cospectral, and both are bipartite.*

Proof. By construction, B_k is a bipartite graph parts $\{v_0, v_3, v_4\}$ and $V(B_k) \setminus \{v_0, v_3, v_4\}$. The edges v_1v_3 or v_2v_4 each have endpoints in different parts, so adding these edges keeps the graph bipartite.

We show that B_k satisfies Theorem 3.12. First we show that $\{\{v_1, v_2\}, \{v_3, v_4\}\}$ are a set of cousins. By inspection, we see that for all $u \in L$, $d(u, v_1) = 2 = d(u, v_2)$ and $d(u, v_3) = 3 = d(u, v_4)$. For all $u \in R$, $d(u, v_1) = 2 = d(u, v_2)$ and $d(u, v_3) = 1 = d(u, v_4)$. Additionally $d(v_0, v_1) = 1 = d(v_0, v_2)$ and




Figure 3.2: A bipartite family B_k where the dashed edges are toggled to create \mathcal{D}^L -cospectral graphs. The set L contains k twin vertices and the set R contains $(k+1)$ twin vertices.

$d(v_0, v_3) = 2 = d(v_0, v_4)$. Finally,

$$\sum_{u \in U(C)} d(u, v_1) = \sum_{u \in L} d(u, v_1) + d(v_0, v_1) + \sum_{u \in R} d(u, v_1) = 2k + 1 + 2(k+1) = 4k + 3.$$

$$\sum_{u \in U(C)} d(u, v_3) = \sum_{u \in L} d(u, v_3) + d(v_0, v_3) + \sum_{u \in R} d(u, v_3) = 3k + 2 + (k+1) = 4k + 3.$$

Therefore, $C = \{\{v_1, v_2\}, \{v_3, v_4\}\}$ is a set of cousins. In B_k , we have that v_1 is not adjacent to v_3 and v_2 is not adjacent to v_4 . The subgraph of $B_k + v_1v_3$ induced by $\{v_1, v_2, v_3, v_4\}$ is isomorphic to the subgraph of $B_k + v_2v_4$ induced by $\{v_1, v_2, v_3, v_4\}$. Additionally, the only vertex in $N(v_1) \cap N(v_2) = \{v_0\}$ is adjacent to every vertex in $R = N(v_3) \cap N(v_4)$. Thus, $B_k + v_1v_3$ and $B_k + v_2v_4$ are \mathcal{D}^L -cospectral by Theorem 3.12. \square

The family B_k grows by increasing k , or twining two vertices (one in L and one in R) at the same time to preserve the cousin property. This growth is not unique to this cospectral family: We can change the set of edges between the vertices $\{v_1, v_2, v_3, v_4\}$ provided the cousin property and the induced graph isomorphism are preserved. Furthermore, L and R can have arbitrary structure since it does not affect the cousin property nor the induced graph isomorphism; however adding edges within L or R will result in the graphs that are not bipartite.

4. Cospectral transmission regular graphs. If two graphs are transmission regular and \mathcal{D} -cospectral, then they are \mathcal{D}^L -cospectral (see Remark 4.3 for more detail). Thus, results for \mathcal{D} -cospectrality can be combined with characterizations of transmission regular graphs in order to obtain results about \mathcal{D}^L -cospectrality. In this section we discuss transmission regular \mathcal{D}^L -cospectral graphs, including strongly regular graphs, distance-regular graphs, and circulant graphs. We begin our discussion of transmission regular graphs by generalizing a well-known example of a transmission regular graph that is not regular (the graph in Figure 4.1) to an infinite family of graphs with this property.




Figure 4.1: The graph H_3 , which is transmission regular but not regular.

DEFINITION 4.1. Let H_n be the graph with vertex set $V = \{v_1, \dots, v_{2n}, u_1, \dots, u_n\}$ and edge set $E = \{v_i v_{i+1} : 1 \leq i \leq 2n-1\} \cup \{v_{2n} v_1\} \cup \{u_i v_{2i-1}, u_i v_{2i}, u_i v_{2i+1} : 1 \leq i \leq n-1\} \cup \{u_n v_{2n-1}, u_n v_{2n}, u_n v_1\}$.

PROPOSITION 4.2. For each $n \geq 2$, H_n is not regular. For each odd $n \geq 3$, H_n is transmission regular and $t(H_n) = \frac{3n^2+1}{2}$.

Proof. Since H_n has vertices of degrees three and four, H_n is not regular. By the symmetry of H_n , all vertices u_i and v_{2i} in the definition of H_n have the same transmission for $1 \leq i \leq n$; likewise, all vertices v_{2i-1} have the same transmission as v_1 . Thus, to show that H_n is transmission regular, it suffices to show that $t(u_1) = t(v_1)$. Using the identities $1 + 3 + \dots + k = \left(\frac{k+1}{2}\right)^2$ and $2 + 4 + \dots + k = \frac{k}{2} \left(\frac{k}{2} + 1\right)$, we have

$$\begin{aligned} t(u_1) &= 3(1) + 4(2) + 2(3) + 4(4) + 2(5) + \dots + 2(n-2) + 4(n-1) + 1(n) \\ &= 4\left(\frac{n-1}{2}\left(\frac{n-1}{2} + 1\right)\right) + 2\left(\left(\frac{n-2+1}{2}\right)^2\right) + 1 + n \\ &= \frac{3n^2+1}{2} \\ &= 4\left(\frac{n-2+1}{2}\right)^2 + 2\left(\frac{n-1}{2}\left(\frac{n-1}{2} + 1\right)\right) + 2n \\ &= 4(1) + 2(2) + 4(3) + 2(4) + 4(5) + \dots + 4(n-2) + 2(n-1) + 2(n) \\ &= t(v_1), \end{aligned}$$

where the first and last equalities can be easily verified by inspection by counting the distances from u_1 and v_1 to all other vertices. \square

REMARK 4.3. Suppose G is a transmission regular graph of order n . The spectral radius of the distance matrix $\mathcal{D}(G)$ is the transmission of G and the trace of the distance Laplacian is $\text{tr}(\mathcal{D}^L(G)) = nt(G)$. Observe that $\text{spec}_{\mathcal{D}^L}(G) = \{\partial_1^L = 0, \partial_2^L, \dots, \partial_n^L\}$ where $\partial_k^L = t(G) - \partial_k$ for $k = 1, \dots, n$ (recall that distance eigenvalues are ordered largest to smallest, whereas distance Laplacian eigenvalues are ordered smallest to largest). Thus, two transmission regular graphs are \mathcal{D} -cospectral if and only if they are \mathcal{D}^L -cospectral.

4.1. Strongly regular and distance-regular graphs. Strongly regular, and more generally distance-regular graphs, provide important examples of transmission regular \mathcal{D}^L -cospectral graphs. A k -regular graph G of order n is *strongly regular* with parameters (n, k, λ, μ) if every pair of adjacent vertices has λ common neighbors and every pair of nonadjacent distinct vertices has μ common neighbors. A connected strongly regular graph G has $\text{diam}(G) \leq 2$. There are well known formulas for computing the adjacency eigenvalues of G and their multiplicities from the strongly regular graph parameters (see [14, Section 10.2]). Formulas for the distance eigenvalues and their multiplicities have also been determined [5, p. 262] and [1]. Thus the distance Laplacian eigenvalues can be readily determined, as done in the next remark.

REMARK 4.4. Let G be a strongly regular graph with parameters (n, k, λ, μ) . Then any vertex v has k neighbors and $n - k - 1$ vertices at distance two, so G is $(2n - k - 2)$ -transmission regular. Thus the distance Laplacian eigenvalues of G are

$$\begin{aligned} &0 \text{ of multiplicity } 1 \\ &2n - k + \frac{1}{2} \left(\lambda - \mu + \sqrt{(\lambda - \mu)^2 + 4(k - \mu)} \right) \text{ of multiplicity } \frac{1}{2} \left(n - 1 - \frac{2k + (n-1)(\lambda - \mu)}{\sqrt{(\lambda - \mu)^2 + 4(k - \mu)}} \right) \\ &2n - k + \frac{1}{2} \left(\lambda - \mu - \sqrt{(\lambda - \mu)^2 + 4(k - \mu)} \right) \text{ of multiplicity } \frac{1}{2} \left(n - 1 + \frac{2k + (n-1)(\lambda - \mu)}{\sqrt{(\lambda - \mu)^2 + 4(k - \mu)}} \right). \end{aligned}$$

Furthermore, two strongly regular graphs with the same set of parameters are \mathcal{D}^L -cospectral.

There are examples, such as those presented next, of non-isomorphic strongly regular graphs with the same parameters (n, k, λ, μ) ; such graphs are therefore \mathcal{D}^L -cospectral and transmission regular (so neither strong regularity nor transmission regularity implies a graph is spectrally determined). The notation $\lambda^{(m)}$ means that the multiplicity of eigenvalue λ is m .

EXAMPLE 4.5. The Shrikhande graph, shown in Figure 4.2, is the graph $S = (V, E)$ where $V = \{0, 1, 2, 3\} \times \{0, 1, 2, 3\}$ and $E = \{\{(a, b), (c, d)\} : (a, b) - (c, d) \in \{\pm(0, 1), \pm(1, 0), \pm(1, 1)\}\}$. Both S and the 2,4-Hamming graph $H(2, 4) := K_4 \square K_4$ are strongly regular with parameters $(16, 6, 2, 2)$. They are 24-transmission regular with distance spectrum $\text{spec}_{\mathcal{D}}(S) = \{24, 0^{(9)}, (-4)^{(6)}\}$ and distance Laplacian spectrum $\text{spec}_{\mathcal{D}^L}(S) = \{0, 24^{(9)}, 28^{(6)}\}$. Since S and $H(2, 4)$ are cospectral and strongly regular, so are their complements \bar{S} and $\bar{H}(2, 4)$. Observe that $\omega(S) = 3 = \alpha(\bar{S})$ and $\omega(H(2, 4)) = 4 = \alpha(\bar{H}(2, 4))$.




Figure 4.2: The Shrikhande graph S

EXAMPLE 4.6. Let $i, j, k \geq 0$ be integers. The graph $G = (V, E)$ is *distance-regular* if for any choice of $u, v \in V$ with $d(u, v) = k$, the number of vertices $w \in V$ such that $d(u, w) = i$ and $d(v, w) = j$ is independent of the choice of u and v . Distance spectra of several families of distance-regular graphs were determined in [5] and the distance spectra of all distance-regular graphs having exactly one positive distance eigenvalue are listed in [1]. A *Doob graph* $D(m, n)$ of diameter $2m+n$ is the Cartesian product of m copies of the Shrikhande graph with one copy of the Hamming graph $H(n, 4)$. It is known [1] that the Doob graph $D(m, n)$ and the Cartesian product of m copies of $H(2, 4)$ with $H(n, 4)$ are \mathcal{D} -cospectral and thus \mathcal{D}^L -cospectral.

EXAMPLE 4.7. The triangular graph T_8 (which is the line graph of K_8) and the three Chang graphs are all strongly regular with the parameters $(28, 12, 6, 4)$ (see [20] for definitions of the Chang graphs). Thus all four have the same distance spectrum $\{42, 0^{(20)}, (-6)^{(7)}\}$ and the same \mathcal{D}^L -spectrum $\{0, 42^{(20)}, 48^{(7)}\}$.

4.2. Cospectral circulant graphs. The ring of integers mod n is denoted by \mathbb{Z}_n , with elements $0, \dots, n-1$, and we view an element of \mathbb{Z}_n as also having a meaningful interpretation as an integer, where we can use the standard ordering of elements, although arithmetic is performed modulo n . For an integer $n \geq 3$ and a set $S \subseteq \mathbb{Z}_n$ that does not contain zero, the *circulant graph* defined by n and S is the graph with vertex set \mathbb{Z}_n and edges of the form ij where $i-j$ or $j-i$ is in S ; this graph is denoted by $\text{Circ}_n(S)$. The set S is called the *connection set* of $\text{Circ}_n(S)$. By convention we assume that $1 \leq x \leq \lfloor \frac{n}{2} \rfloor$ for every $x \in S$. If q is relatively prime to n then $\text{Circ}_n(S)$ is isomorphic to $\text{Circ}_n(qS)$ where $qS = \{qx : x \in S\}$. Thus it is common to require that $1 \in S$ if S contains any element relatively prime to n , and we make this assumption also. Note that $\text{Circ}_n(S)$ is connected if and only if the greatest common divisor of the elements of S together with n is one. A circulant graph is both regular and transmission regular. Circulants are an interesting family of graphs that play an important role in some applications of graph theory such as computing and

quantum spin networks [6, 19]. They also provide examples of distance Laplacian (and distance) cospectral mates.

Table 4.1 lists the orders and connection sets for all sets of \mathcal{D}^L -cospectral circulant graphs of order at most 20; the data in the table were produced by computations in *Sage* [11]. The smallest order is 16, and the \mathcal{D}^L -cospectral graphs $\text{Circ}_{16}(\{1, 2, 8\})$ and $\text{Circ}_{16}(\{1, 6, 8\})$ are shown in Figure 4.3.

n	S
16	$\{1, 2, 8\}, \{1, 6, 8\}$
18	$\{1, 2, 9\}, \{1, 4, 9\}$
18	$\{1, 6, 8\}, \{2, 3, 6\}$
20	$\{1, 2, 3, 8\}, \{1, 2, 7, 8\}$
20	$\{1, 2, 4, 9\}, \{1, 6, 8, 9\}$
20	$\{1, 2, 4, 10\}, \{1, 6, 8, 10\}$
20	$\{1, 2, 4, 9, 10\}, \{1, 6, 8, 9, 10\}$
20	$\{1, 2, 3, 8, 10\}, \{1, 2, 7, 8, 10\}$
20	$\{1, 2, 4, 5, 9\}, \{1, 5, 6, 8, 9\}$
20	$\{1, 2, 3, 5, 8\}, \{1, 2, 5, 7, 8\}$
20	$\{1, 2, 4, 5, 9, 10\}, \{1, 5, 6, 8, 9, 10\}$
20	$\{1, 2, 3, 5, 8, 10\}, \{1, 2, 5, 7, 8, 10\}$

Table 4.1: Order n and connection sets S for small circulant \mathcal{D}^L -cospectral pairs




Figure 4.3: The circulant graphs $\text{Circ}_{16}(\{1, 2, 8\})$ and $\text{Circ}_{16}(\{1, 6, 8\})$

Although Table 4.1 contains only cospectral pairs of even order, by using *Sage* to perform a search we found that the graphs $\text{Circ}_{27}(\{1, 2, 9\})$ and $\text{Circ}_{27}(\{1, 4, 9\})$ on 27 vertices are \mathcal{D}^L -cospectral.

The *Sage* search that produced Table 4.1 was restricted to circulant graphs, so it would not detect a pair of \mathcal{D}^L -cospectral graphs where one is a circulant and the other is not. Recall a result from [8, 18, 21] that links strongly regular graphs and circulant graphs: If a connected circulant graph $\text{Circ}_n(S)$ is strongly regular, then n must be a prime p satisfying $p \equiv 1 \pmod{4}$ and $\text{Circ}_p(S)$ must be isomorphic to a Paley graph on the field of order p (see [14] for the definition of a Paley graph).

EXAMPLE 4.8. The graph $\text{Circ}_{29}(\{1, 4, 5, 6, 7, 9, 13\})$, which is shown in Figure 4.4, is a Paley graph on 29 vertices that is strongly regular with parameters $(29, 14, 6, 7)$. There are 40 additional nonisomorphic strongly regular graphs on 29 vertices with the same strongly regular graph parameters [23] that are necessarily \mathcal{D}^L -cospectral. The \mathcal{D}^L -spectrum of $\text{Circ}_{29}(\{1, 4, 5, 6, 7, 9, 13\})$ is $\left\{0^{(1)}, \frac{87-\sqrt{29}}{2}^{(14)}, \frac{87+\sqrt{29}}{2}^{(14)}\right\}$; the \mathcal{D}^L -

spectrum can be computed from the formulas for the \mathcal{D}^L -spectrum of a strongly regular graph (see Remark 4.4 and [10]).




Figure 4.4: The circulant graph $\text{Circ}_{29}(\{1, 4, 5, 6, 7, 9, 13\})$

A *circulant matrix* is a square matrix in which each row is formed from the row above by a right cyclic shift of plus one. A trivial example of a circulant matrix is that of the identity matrix I_n where this shift pattern can be seen along the lead diagonal. The adjacency, distance, and distance Laplacian matrices of a circulant graph are circulant matrices.

For $j = 1, \dots, n - 1$, define $d_j^{(r)}$ as the distance between vertex 0 and vertex j in $\text{Circ}_n([r])$. Thus, $d_j^{(r)} = \lceil \frac{j}{r} \rceil$ for $1 \leq j \leq \lfloor \frac{n}{2} \rfloor$ and due to the symmetry of the circulant graphs $d_{n-j}^{(r)} = d_j^{(r)}$. Define $d_0^{(r)} = t(\text{Circ}_n([r])) = \sum_{j=1}^{n-1} d_j^{(r)}$. Then the first row of the circulant matrix $\mathcal{D}^L(\text{Circ}_n([r]))$ is given by

$$\left(d_0^{(r)}, -d_1^{(r)}, \dots, -d_{n-1}^{(r)} \right)$$

The remaining entries of the matrix $\mathcal{D}^L(\text{Circ}_n([r]))$ are defined by using the circulant matrix property.

PROPOSITION 4.9. *The transmission of any consecutive circulant graph $\text{Circ}_n([r])$ is*

$$t(\text{Circ}_n([r])) = \left(\left\lfloor \frac{n-1}{2r} \right\rfloor + 1 \right) \left((n-1) - r \left\lfloor \frac{n-1}{2r} \right\rfloor \right).$$

Proof. For any vertex v of $\text{Circ}_n([r])$, there are $2r$ vertices at distance one from v , $2r$ vertices at distance two, and so on, ending with $2r$ vertices at distance $\lfloor \frac{n-1}{2r} \rfloor$. The remaining $n - 1 - 2r \lfloor \frac{n-1}{2r} \rfloor$ vertices are at distance $\lfloor \frac{n-1}{2r} \rfloor + 1$. Thus the transmission is

$$\begin{aligned} t(G) = t(v) &= 2r \left(\sum_{i=0}^{\lfloor \frac{n-1}{2r} \rfloor} i \right) + \left((n-1) - 2r \left\lfloor \frac{n-1}{2r} \right\rfloor \right) \left(\left\lfloor \frac{n-1}{2r} \right\rfloor + 1 \right) \\ &= \left(\left\lfloor \frac{n-1}{2r} \right\rfloor + 1 \right) \left((n-1) - r \left\lfloor \frac{n-1}{2r} \right\rfloor \right). \quad \square \end{aligned}$$

We can use Proposition 4.9 to easily calculate the Wiener index from the transmission.

COROLLARY 4.10. *The Wiener index of a consecutive circulant is*

$$W(\text{Circ}_n([r])) = \frac{n}{2}t(G) = \frac{n}{2} \left(\left\lfloor \frac{n-1}{2r} \right\rfloor + 1 \right) \left((n-1) - r \left\lfloor \frac{n-1}{2r} \right\rfloor \right).$$

Since the pair (n, r) determines both the number of eigenvalues and the transmission (by Proposition 4.9), and the transmission equals the distance spectral radius, the next result is immediate.

COROLLARY 4.11. *For $r_j \leq \lfloor \frac{n_j}{2} \rfloor, j = 1, 2$, the consecutive circulants $\text{Circ}_{n_1}([r_1])$ and $\text{Circ}_{n_2}([r_2])$ are distance cospectral (and hence distance Laplacian cospectral) if and only if $n_1 = n_2$ and $r_1 = r_2$.*

5. Unimodality of the coefficients of the distance Laplacian characteristic polynomial. In their study of distance spectra of trees, Graham and Lovász [15] showed that for a tree T and $\delta_k(T)$ denoting the coefficient of x^k in $\det(\mathcal{D}(T) - xI) = (-1)^n p_{\mathcal{D}(T)}(x)$, the quantity $d_k(T) = (-1)^{n-1} \delta_k(T) / 2^{n-k-2}$ is determined as a fixed linear combination of the numbers of certain subtrees in T ; the values $d_k(T)$ are called the *normalized coefficients* of T . A sequence a_0, a_1, \dots, a_n of real numbers is *unimodal* if there is a k such that $a_{i-1} \leq a_i$ for $i \leq k$ and $a_i \geq a_{i+1}$ for $i \geq k$. Graham and Lovász [15] conjectured that the sequence $d_0(T), \dots, d_{n-2}(T)$ of normalized coefficients is unimodal with the maximum value occurring at $\lfloor \frac{n}{2} \rfloor$ for a tree T of order n . Collins [9] showed that the location of the maximum for a path on n vertices is approximately $n \left(1 - \frac{1}{\sqrt{5}}\right)$ and with modern computers it is straightforward to verify that this location of the peak is strictly greater than $\lfloor \frac{n}{2} \rfloor$ for $n \geq 18$ vertices (and for odd $n \geq 9$). Aalipour et al. [2] established the unimodality of the normalized coefficients and also of the absolute values of the coefficients of the distance characteristic polynomial. These ideas naturally suggest exploring the behavior of the coefficients of distance Laplacian characteristic polynomial.

The unimodality of the sequence of absolute values of the coefficients of the distance characteristic polynomial is equivalent to that of the normalized coefficients [2, 15], and the proof used log-concavity and its connection to polynomials with real roots. A sequence a_1, a_2, \dots, a_n of real numbers is *log-concave* if $a_j^2 \geq a_{j-1}a_{j+1}$ for all $j = 2, \dots, n-1$. A polynomial p is *real-rooted* if all roots of p are real (by convention, constant polynomials are considered real-rooted). The next result describes the connections between real-rootedness, log-concavity, and unimodality, and was the key to the proof.

LEMMA 5.1. [2, 7]

- (a) *If $p(x) = a_n x^n + \dots + a_1 x + a_0$ is a real-rooted polynomial, then the coefficient sequence a_i of $p(x)$ is log-concave.*
- (b) *If a sequence $a_0, a_1, a_2, \dots, a_n$ is positive and log-concave, then $a_0, a_1, a_2, \dots, a_n$ is unimodal.*

COROLLARY 5.2. *Let A be a real symmetric matrix. The coefficient sequence $a_0, a_1, \dots, a_{n-1}, a_n = 1$ of $p_A(x)$ is log-concave. If a subsequence a_s, \dots, a_t of $a_0, a_1, \dots, a_{n-1}, a_n$ is positive then this subsequence is also unimodal. If all entries of a subsequence of $a_0, a_1, \dots, a_{n-1}, a_n$ alternate in sign or all are negative, then the subsequence of absolute values $|a_s|, \dots, |a_t|$ is unimodal.*

Proof. Since all the eigenvalues of A are real, the log-concavity of a_0, \dots, a_{n-1}, a_n follows immediately from Lemma 5.1, and any positive subsequence is unimodal. If the signs of the subsequence a_s, \dots, a_t alternate, or if all values are negative, then $|a_j^2| = a_j^2 \geq a_{j-1}a_{j+1} = |a_{j-1}||a_{j+1}|$ for $s+1 \leq j \leq t-1$, so the positive subsequence $|a_s|, \dots, |a_t|$ is also log-concave, and therefore unimodal. \square

The ℓ th symmetric function of $\alpha_1, \dots, \alpha_n$, denoted by $S_\ell(\alpha_1, \dots, \alpha_n)$ is the sum over all products of

ℓ distinct elements of $\{\alpha_1, \dots, \alpha_n\}$. Observe that if $0 = \alpha_1 = \dots = \alpha_c < \alpha_{c+1} < \dots < \alpha_n$, then $S_\ell(\alpha_1, \dots, \alpha_n) = S_\ell(\alpha_{c+1}, \dots, \alpha_n)$ for $\ell \leq n - c$ and $S_\ell(\alpha_1, \dots, \alpha_n) = 0$ for $\ell > n - c$.

THEOREM 5.3. *Let G be a connected graph, and let $p_{\mathcal{D}^L(G)}(x) = x^n + \delta_{n-1}^L x^{n-1} + \dots + \delta_1^L x$. Then the sequence $\delta_1^L, \dots, \delta_{n-1}^L, \delta_n^L = 1$ is log-concave and $|\delta_1^L|, \dots, |\delta_n^L|$ is unimodal. In fact, $|\delta_1^L| \geq \dots \geq |\delta_n^L|$.*

Proof. It is well known that the distance Laplacian is positive semidefinite (this follows immediately from Gershgorin's disk theorem), and that zero is a simple eigenvalue (see, for example, [3]). Since $\delta_k^L = (-1)^{n-k} S_{n-k}(\partial_1^L, \partial_2^L, \dots, \partial_n^L) = (-1)^{n-k} S_{n-k}(\partial_2^L, \dots, \partial_n^L)$ and $\partial_2^L, \dots, \partial_n^L$ are positive, all the nonzero coefficients alternate in sign. By Corollary 5.2 the sequence $\{|\delta_k^L|\}_{k=1}^n$ is unimodal.

To show that $|\delta_1^L| \geq \dots \geq |\delta_n^L|$, by unimodality it is sufficient to show that $|\delta_1^L| > |\delta_2^L|$. Since $\partial_2^L \geq n$ [3, Corollary 3.6],

$$|\delta_2^L| = \sum_{i=2}^n \prod_{j \neq i} \partial_j^L \leq (n-1) \prod_{j=3}^n \partial_j^L < \prod_{j=2}^n \partial_j^L = |\delta_1^L|. \quad \square$$

6. Concluding remarks. In this paper, we studied graph cospectrality with respect to the distance Laplacian matrix. We showed that several global properties of a graph, such as degree sequence, transmission sequence, diameter, and number of nontrivial automorphisms, and also several local properties, such as the presence of a leaf, a dominating vertex, and a cut vertex, are not preserved by \mathcal{D}^L -cospectrality. It would be interesting to determine whether or not other properties, such as those listed in the next question, are preserved by \mathcal{D}^L -cospectrality.

QUESTION 6.1. *Are the following properties preserved by \mathcal{D}^L -cospectrality?*

1. *Acyclicity (being a tree).*
2. *Bipartiteness.*
3. *Transmission regularity.*
4. *Regularity.*
5. *Strong regularity.*

It is not known whether trees are \mathcal{D}^L -spectrally determined [3]; if so, then the property of being a tree would necessarily be preserved by \mathcal{D}^L -cospectrality. For each of the other properties in Question 6.1, examples of a \mathcal{D}^L -cospectral pair with both graphs having the property and a \mathcal{D}^L -cospectral pair with neither graph having the property have been presented (in Sections 2 and 4). However, we do not have examples of \mathcal{D}^L -cospectral pairs where one member has the property and the other does not.

We also established several general methods for producing \mathcal{D}^L -cospectral graphs based on sets of vertices sharing common neighborhoods, transmissions, and partial transmissions. We used these constructions to produce infinite families of \mathcal{D}^L -cospectral graphs. Future work could include finding additional methods for constructing \mathcal{D}^L -cospectral graphs. Conversely, it would also be interesting to find certain local structures or general properties of a graph that guarantee the graph has no \mathcal{D}^L -cospectral mates; one conjectured such property is acyclicity.

Since two transmission regular graphs are \mathcal{D} -cospectral if and only if they are \mathcal{D}^L -cospectral, results about \mathcal{D} -cospectrality of transmission regular graphs lead to results about \mathcal{D}^L -cospectrality. To this end, we studied various families of strongly regular, distance-regular, and circulant graphs, and showed that many of them have \mathcal{D}^L -cospectral mates. We also constructed an infinite family of graphs which are transmission

regular but not regular; it is an open question to determine whether this family of graphs is \mathcal{D}^L -spectrally determined.

Finally, we established that the absolute values of coefficients of the distance Laplacian characteristic polynomial are unimodal, and in fact decreasing. It would be interesting to establish other structural results about the coefficients of the distance Laplacian characteristic polynomial and their relation to \mathcal{D}^L -cospectrality.

Acknowledgements. We gratefully acknowledge financial support for this research from the following grants and organizations: NSF grants #1603823 and #1604458, NSA grant #H98230-18-1-0017, and the Combinatorics Foundation.

REFERENCES

- [1] G. Aalipour, A. Abiad, Z. Berikkyzy, J. Cummings, J. De Silva, W. Gao, K. Heysse, L. Hogben, F.H.J. Kenter, J.C.-H. Lin, M. Tait. On the distance spectra of graphs. *Linear Algebra Appl.* 497 (2016), 66–87.
- [2] G. Aalipour, A. Abiad, Z. Berikkyzy, L. Hogben, F.H.J. Kenter, J.C.-H. Lin, M. Tait. Proof of a conjecture of Graham and Lovasz concerning unimodality of coefficients of the distance characteristic polynomial of a tree. *Electron. J. Linear Algebra* 34 (2018), 373–380.
- [3] M. Aouchiche, P. Hansen. Two Laplacians for the distance matrix of a graph. *Linear Algebra Appl.* 439 (2013), 21–33.
- [4] M. Aouchiche, P. Hansen. Distance spectra of graphs: A survey. *Linear Algebra Appl.* 458 (2014), 301–386.
- [5] F. Atik, P. Panigrahi. On the distance spectrum of distance regular graphs. *Linear Algebra Appl.* 478 (2015), 256–273.
- [6] M. Basić, M. Petković, D. Stevanović. Perfect state transfer in integral circulant graphs. *Applied. Math. Let.* 22 (2009), 1117–1121.
- [7] P. Brändén. Unimodality, log-concavity, real-rootedness, and beyond. In *Handbook of Enumerative Combinatorics*, M. Bona, Editor, CRC Press, Boca Raton, 2015.
- [8] W.G. Bridge, R.A. Mena. Rational circulant with rational spectra and cyclic strongly regular graphs. *Ars Combin.* 8 (1979), 143–161,
- [9] K.L. Collins. On a conjecture of Graham and Lovász about distance matrices. *Discrete Appl. Math.* 25 (1989), 27–35.
- [10] K. Duna, L. Hogben, K. Lorenzen. Sage verifications of examples and computational lemmas. PDF available at <https://orion.math.iastate.edu/lhogben/BDGLRSY18-Distance.Laplacian--Sage.pdf>. Sage worksheet available at <https://sage.math.iastate.edu/home/pub/90/>.
- [11] K. Duna, K. Lorenzen. Sage code for finding \mathcal{D}^L -copsectral graphs. PDF available at https://orion.math.iastate.edu/lhogben/Sage_Code.BDHLSY18--Sage.pdf. Sage worksheet available at <https://sage.math.iastate.edu/home/pub/91/>.
- [12] K. Duna, K. Lorenzen. List of all \mathcal{D}^L -cospectral graphs of order at most 10 in *graph6* format. PDF available at https://orion.math.iastate.edu/lhogben/Data_orders78910.BDHLSY18.pdf.
- [13] C.D. Godsil, B.D. McKay. Constructing cospectral graphs. *Aequationes Math.* 25 (1982), 257–268.
- [14] C. Godsil, G. Royle. *Algebraic Graph Theory*. Springer-Verlag, New York, 2001.
- [15] R.L. Graham, L. Lovász. Distance matrix polynomials of trees. *Adv. Math.* 29 (1978), 60–88.
- [16] R.L. Graham, H.O. Pollak. On the addressing problem for loop switching. *Bell Syst. Tech. J.* 50 (1971), 2495–2519.
- [17] K. Heysse. A construction of distance cospectral graphs. *Linear Algebra Appl.* 535 (2017), 195–212.
- [18] D.R. Hughes, J.V. van Lint, R.M. Wilson. Announcement at the Seventh British Combinatorial Conference, Cambridge, 1979 (unpublished).
- [19] A. Ilić. Distance spectra and distance energy of integral circulant graphs. *Linear Algebra Application.* 433 (2010), 1005–1014.
- [20] J.H. Koolen, S.V. Shpectorov. Distance-regular Graphs the Distance Matrix of which has Only One Positive Eigenvalue. *European J. Combin.* 15 (1994), 269–275.
- [21] S.L. Ma. Partial difference sets. *Discrete Math.* 52 (1984), 75–89.
- [22] M. Nath, S. Paul. On the distance Laplacian spectra of graphs. *Linear Algebra Appl.* 460 (2014), 97-110.
- [23] E. Spence. Strongly Regular Graphs on at most 64 vertices. Available at <http://www.maths.gla.ac.uk/~es/srgraphs.php>.