



UNIVERSITY OF LEEDS

This is a repository copy of *Meat consumption is associated with higher dementia prevalence: a cross-sectional analysis of UK Biobank*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/174928/>

Version: Accepted Version

Conference or Workshop Item:

Zhang, H, Hardie, LJ, Greenwood, DC orcid.org/0000-0001-7035-3096 et al. (1 more author) (2021) Meat consumption is associated with higher dementia prevalence: a cross-sectional analysis of UK Biobank. In: Nutrition Society Winter Conference Live 2020, 08-09 Dec 2020, Online.

<https://doi.org/10.1017/S0029665121000100>

This article has been published in a revised form in Proceedings of the Nutrition Society [<https://doi.org/10.1017/S0029665121000100>]. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © The Authors 2021

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Meat consumption is associated with higher dementia prevalence: a cross-sectional analysis of UK Biobank

H. Zhang, L.J. Hardie, D.C. Greenwood and J.E. Cade

Abstract

Dementia represents an increasing burden of disease in aging societies worldwide; modifiable lifestyle factors, especially diet, have been implicated and represent potential targets for primary prevention(1). Although the role of processed and red meat consumption in relation to certain cancers has been established(2), its involvement in dementia development currently remains debatable, while there is an increasing trend of meat intakes globally.

The intake frequencies per week of total meat comprising red meat, processed meat, and poultry were assessed via a 47- item food frequency questionnaire in 502493 UK Biobank participants mean age 56.5 years (SD: 8.1) at recruitment between 2006–2010. Prevalent dementia cases were self-reported or hospital- and mortality- registered. Odds ratios (OR) for each meat type in relation to dementia risk were estimated in logistic regression models adjusting for age, sex, ethnicity, Townsend deprivation index, education level, smoking status, alcohol status, body mass index, physical activity level, sleeping hours, stroke history, family history of dementia.

There were 564 prevalent dementia cases mean age 59.8 years (SD: 7.5). Our results show that increased odds of dementia were associated with total-meat consumption by 5% (adjusted OR: 1.05, 95% CI: 1.01 to 1.10, P = 0.015) for each additional portion of meat/week, processed meat by 14% (adjusted OR: 1.14, 95% CI: 1.07 to 1.22, P <0.001), and red meat by 9% (adjusted OR: 1.09, 95% CI: 1.02 to 1.16, P = 0.017), but not poultry (adjusted OR: 0.94, 95% CI: 0.87 to 1.03, P = 0.195). However, sex-specific associations between dementia prevalence and consumption of red meat or total meat were pronounced in men (adjusted OR: 1.11, 95%CI: 1.04 to 1.19, P = 0.002 for red meat; adjusted OR: 1.07, 95% CI: 1.03 to 1.12, P = 0.002 for total meat), but not in women (adjusted OR: 1.02, 95% CI: 0.86 to 1.21, P = 0.783 for red meat; adjusted OR: 1.02, 95% CI: 0.93 to 1.12, P = 0.730 for total meat).

Our findings suggest that higher frequency of meat consumption, especially processed meat, is associated with greater dementia prevalence, especially among men in the UK. However, due to the cross-sectional nature of this study the direction of effect is uncertain; these associations need to be examined in longitudinal studies further.

References

- 1 Kivipelto, M, Mangialasche, F & Ngandu, T (2018) *Nat Rev Neurol* 14, 653–666.
- 2 Bouvard, V, Loomis, D, Guyton, KZ, et al. (2015) *Lancet Oncol* 16, 1599–1600.