



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/175122/>

Version: Accepted Version

Book Section:

Johnston, J (2021) Pharmacology of A-Type K⁺ Channels. In: Gamper, N and Wang, K, (eds.) Pharmacology of Potassium Channels. Handbook of Experimental Pharmacology, 267. Springer, Cham, Switzerland, pp. 167-183. ISBN: 978-3-030-84051-8. ISSN: 0171-2004.

https://doi.org/10.1007/164_2021_456

© The Author(s), under exclusive license to Springer Nature Switzerland AG 2021. This is an author produced version of a book chapter published in Handbook of Experimental Pharmacology. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Pharmacology of A-type K⁺ channels

Jamie Johnston

Faculty of Biological Sciences, University of Leeds, Leeds, UK

Abstract

Transient outward potassium currents were first described nearly 60 years ago, since then major strides have been made in understanding their molecular basis and physiological roles. From the large family of voltage-gated potassium channels members of 3 subfamilies can produce such fast-inactivating A-type potassium currents. Each subfamily gives rise to currents with distinct biophysical properties and pharmacological profiles and a simple workflow is provided to aid the identification of channels mediating A-type currents in native cells. Their unique properties and regulation enable A-type K⁺ channels to perform varied roles in excitable cells including repolarisation of the cardiac action potential, controlling spike and synaptic timing, regulating dendritic integration and long-term potentiation as well as being a locus of neural plasticity.

Keywords: A-type K⁺ channel, Voltage-clamp, potassium channel, Kv₄, Kv_{1.4}, Kv_{3.4}

1. Introduction

A-type refers to neuronal K⁺ currents generating a transient outward current in response to depolarising voltage-clamp steps. This stereotypical phenotype was first described in neurons from sea slug *Onchidium* (Hagiwara et al., 1961). These early experiments described the now stereotypical hallmarks of A-type currents. They identified a transient outward voltage-gated K⁺ current activated by depolarising steps from a hyperpolarised potential (Fig. 1 C & D). In current clamp when hyperpolarising steps were used to remove A-current inactivation, they observed an increase in the latency to the first spike in response to depolarising current steps (Fig. 1 A & B). Such transient outward currents were later designated as I_A, a term which has persisted (Connor and Stevens, 1971). Work over the subsequent decades has described A-type channels in the heart (the corresponding current fraction is called I_{to} in the heart; see Chapter 4 for more details) where they contribute to the early repolarisation phase (Dixon et al., 1996; Yeola and Snyders, 1997) and in many different types of neurons where they contribute to such varied functions as controlling spiking, neurotransmitter release, dendritic integration and long term potentiation (Molineux et al., 2005; Imai et al., 2019; Watanabe et al., 2002; Frick et al., 2004; Chen et al., 2006; Shevchenko et al., 2004; Kim et al., 2005; Kuo et al., 2017).

The transient nature of A-type currents is due to rapid voltage-dependent inactivation, with inactivation decay time constants typically in the 10s of milliseconds. The subunit composition and presence of accessory subunits or other modulatory factors determine the exact biophysics and pharmacological profile of these currents. Alpha subunits from three subfamilies of voltage-gated K⁺ channel generate rapid inactivating K⁺ currents when expressed as homomers: Kv1.4, Kv3.3, Kv3.4 and all three Kv4 members, Kv4.1, Kv4.2 & Kv4.3 (Coetzee et al., 1999). The presence of the β subunit Kvβ1 can also confer A-type kinetics to other, normally delayed rectifier subunits from the Kv1 family (Rettig et al., 1994). These different subfamilies have distinct biophysical properties, enabling the subfamily mediating a native A-type current to be narrowed down with simple voltage-clamp protocols and confirmed with their pharmacological profiles.

Figure 1 near here

2. Identifying native A-type channels

2.1 Biophysics

2.1.1 Activation range. Canonical A-type K⁺ currents have sub-threshold activation ranges, i.e. they begin to activate at voltages more negative than that required to generate an action potential (Fig. 1). A hall mark of the Kv3 subfamily is their supra-threshold activation range, they typically do not begin to activate until voltages greater than -30 mV with half activation voltages around +10 mV (Rudy and McBain, 2001). This

property restricts their physiological activation window to that of the action potential, in which Kv₃ channels act to rapidly repolarise the cell to the resting state (Rudy and McBain, 2001; Johnston et al., 2010). A-type K⁺ currents mediated by channels containing Kv_{3.3} and/or Kv_{3.4} can therefore be distinguished from other A-type subunits by their positively shifted activation curve (Fig. 2 B).

2.1.2 Inactivation properties. The molecular basis of rapid K⁺ channel inactivation has been extensively studied with Kv_{1.4} used as the prototype. Two forms of inactivation have been demonstrated; fast N-type or "ball and chain" inactivation involves the N-terminal inactivation peptide of the alpha subunit binding within the central cavity to occlude the inner pore (Hoshi et al., 1990; Zhou et al., 2001). The rate of this type of inactivation is directly related to the number of N-terminal peptides, e.g. Kv_{1.4} homomers, possessing 4 N-terminal peptides, show faster inactivation than Kv_{1.4}/Kv_{1.1} heteromers, which have 3 or less depending on the stoichiometry (Ruppertsberg et al., 1990; MacKinnon et al., 1993). The presence of Kvβ₁ subunits confer a similar N-terminal inactivation peptide (one per subunit), which further accelerates the inactivation rate of Kv_{1.4} or can endow Kv_{1.1} with fast A-type inactivation (Rettig et al., 1994). Kv_{3.3} and Kv_{3.4} are thought to inactivate through a similar mechanism showing a similar dose dependency of N-terminal inactivating peptide on inactivation rate (Rudy et al., 1999; Weiser et al., 1994). In addition to the described N-type inactivation, Kv_{1.4} channels also inactivate by a slower C-type inactivation attributed to partial collapse of the selectivity filter / outer pore. N and C-type inactivation in Kv_{1.4} are allosterically coupled (Hoshi et al., 1990; Hoshi et al., 1991; Bett and Rasmusson, 2004) and the rate of recovery from inactivation is controlled by the slower C-type inactivation (Rasmusson et al., 1995). As a result Kv_{1.4} channels recover from inactivation with a time constant of seconds (Bett and Rasmusson, 2004; Roeper et al., 1997).

In contrast, the molecular details concerning inactivation of Kv₄ channels have not been determined with such precision. Kv₄ subunits use neither the N nor C-type inactivation processes identified for Kv_{1.4}, instead inactivation seems to involve the inner vestibule of the pore and readily occurs through the closed state (Jerng et al., 1999; Bähring et al., 2001; Beck and Covarrubias, 2001; Shahidullah and Covarrubias, 2003). Kv₄ channels possess a N-terminal inactivating domain but rather than contributing to inactivation this region is the binding site of the β-subunits of Kv₄ channels known as K⁺ channel interacting proteins (KChIPs). These accessory proteins appear to be obligatory subunits for native channels (Rhodes et al., 2004). KChIPs and DPPX, another group of accessory proteins, dramatically increase the surface expression of Kv₄ subunits and alter the kinetics of and recovery from inactivation of Kv₄ subunits (Beck et al., 2002; Jerng et al., 2004).

The kinetics of inactivation cannot reliably distinguish between Kv_{1.4} and Kv₄ subunits as their decay time constants can overlap, largely depending on the composition of accessory subunits. However, the recovery from inactivation of Kv₄ channels is usually much more rapid, with time constants around 10s of milliseconds (Kim et al., 2005; Beck et al., 2002; Johnston et al., 2008; Amadi et al., 2007), compared to Kv_{1.4} with recovery rates of 100s of milliseconds to seconds (Ruppertsberg et al., 1990; Rasmusson et al., 1995; Roeper

et al., 1997).

Two simple voltage protocols can therefore help to determine the subfamily responsible for native A-type currents. First the activation curve should be attained with a simple I/V protocol with test steps from ~ -100 mV, to ensure removal of inactivation (Fig. 2 A). The peak current soon after the test step (red arrow in Fig. 2 A) should be corrected for the non-linear driving force before being normalised and fit with a Boltzmann function (Clay, 2000). With the protocol shown in Fig. 2 A the steady-state inactivation curve can also be constructed by plotting the current amplitudes measured at the yellow arrow vs the voltage of the preceding test step. The activation curve will distinguish between Kv3 mediated A-type currents with positive shifted activation ranges and Kv4 or Kv1 mediated A-type currents which have more negative sub-threshold activation ranges (Fig. 2 B). If necessary, the A-type current can be isolated from other endogenous currents with a subtraction protocol as shown in (Johnston et al., 2008). If the A-current has a sub-threshold activation range, the rate of recovery from inactivation can be determined with a protocol similar to that shown in Fig. 2 B right; rapid recovery from inactivation, a $\tau < \sim 100$ ms, is indicative of Kv4 subunits whereas currents displaying time constants of several hundreds of milliseconds or longer are likely to be mediated by Kv1.4 subunits.

2.2 Pharmacology

Pharmacology of ion channels can serve multiple purposes as illustrated by the holy grail of pharmacology, a selective antagonist, a molecule that will antagonise one kind of channel and no other. Such a molecule can be used to unambiguously determine the presence of a channel, its contribution to total current and the role of this channel in normal physiology. There are some striking examples of this with biologically derived toxins, e.g. Dendrotoxin-K is selective for any channel containing Kv1.1 subunits and blocks such channels in the 10 nM range (Wang et al., 1999; Robertson et al., 1996; Dodson et al., 2002). Unfortunately, such molecules are not always available particularly for the large diversity of Kv channels. However, less selective antagonists can still serve a useful role; when combined with other antagonists or biophysical characteristics they can aid identification of the subfamily mediating an A-type current. With these considerations in mind the following discussion largely focuses on molecules that fit these criteria.

2.2.1 Kv3 A-type currents. All Kv3 channels are blocked by 1 mM tetraethylammonium (TEA) which is readily washed on and off tissue (Johnston et al., 2010). TEA has an IC₅₀ for Kv3.3 and Kv3.4 in the 100-200 μ M range and acts by occluding the pore (De Miera et al., 1992; Rettig et al., 1992). TEA sensitivity in combination with a positive activation range provides a strong indication that a Kv3 subunit mediates an A-type current (Fig. 2 B). Block of an A-type current by 1 mM TEA rules out Kv4 subunits which are insensitive to even 100 mM (Johnston et al., 2008; Johnston et al., 2010), though the presence of DPPX can increase their sensitivity to TEA somewhat. Kv1.4 homomers are also insensitive to TEA (Ruppersberg et al., 1990) however heteromeric Kv1.1/Kv1.4 channels are endowed with a mix of properties from their homomeric

counterparts; they display A-type kinetics with slower inactivation rates and have moderate TEA sensitivity with an IC₅₀ of ~10 mM (Ruppersberg et al., 1990). Although 1 mM TEA can be used to unambiguously identify a high-voltage activated A-type currents, it is of limited use in exploring their functional role as many other non-A-type channels are sensitive to TEA e.g. some Kv7 subunits and BK channels (Johnston et al., 2010). The sea anemone toxins BD-I and BDS-II were thought to be selective antagonists of Kv_{3.4} channels which would provide a powerful tool for elucidating the functional role Kv_{3.4} A-type currents in physiology (Diochot et al., 1998). However, later detailed investigation of the interaction of BDS toxins with Kv₃ subunits revealed them to act as gating modifiers and with little selectivity between Kv₃ subunits (Yeung et al., 2005). BDS-I is also highly potent for Nav1.7 channels, slowing inactivation rates with a consequent broadening of action potentials and increased resurgent currents during repolarisation (Liu et al., 2012). Consequently, BDS toxins have limited potential for revealing the physiological role of Kv₃ A-type currents. This also highlights a pitfall of using poorly characterised pharmacological tools; it took 14 years after their first characterisation as selective Kv_{3.4} blockers to realise a more complete pharmacological profile of BDS toxins. To date there are no sufficiently selective antagonists of Kv_{3.3} or Kv_{3.4} channels suitable for studying their physiological roles.

Figure 2 near here

2.1.2 Kv_{1.4} A-type currents. Kv_{1.4} channels are relatively insensitive to TEA (Ruppersberg et al., 1990) and have a lower sensitivity (IC₅₀ ~12.5 mM) to the broad spectrum K⁺ channel blocker 4-AP (Stühmer et al., 1989). In fact, there is a paucity of pharmacological tools available for Kv_{1.4} channels, they are somewhat unique in their resistance to biological toxins which are very potent against many of the other Kv_{1.x} members. At 1 μM the small molecule CP-339,818 shows comparative selectivity for Kv_{1.4} and Kv_{1.3} over other Kv channels, including Kv₃ and Kv₄ subfamilies (Nguyen et al., 1996). CP-339,818 preferentially binds to the C-type inactivated state and therefore shows a use dependent block. At concentrations greater than 1 μM off-target effects become a problem with many other channels being inhibited e.g. Kv_{2.1}, HCN₁, Kv_{3.2} (Lee et al., 2008; Nguyen et al., 1996; Sforza et al., 2015). Another small molecule inhibitor, UK-78,282, has a very similar pharmacological profile to CP-339,818 (Hanson et al., 1999). To date there are no well-established selective antagonists of Kv_{1.4} channels suitable for studying its physiological roles, though if a contribution from Kv_{1.3} can be ruled out, CP-339,818 may serve this purpose.

2.1.3 Kv₄ A-type currents. The anti-arrhythmic drug, flecainide, at 10 μM inhibits Kv_{4.2} & Kv_{4.3} leaving Kv_{1.4} unaffected and has been used to determine the contribution of these channels to the cardiac I_{to} current (Dixon et al., 1996; Yeola and Snyders, 1997). Although flecainide can be used to distinguish between A-type currents mediated by Kv_{1.4} and Kv₄ subunits it also inhibits a number of other channels including the

cardiac Nav1.7 channel (Ramos and O'leary, 2004), ether-a-go-go related gene (ERG) K⁺ channels (Paul et al., 2002) and ryanodine receptors (Mehra et al., 2014). The spider toxins, phrixotoxin 1 & 2, are potent and selective inhibitors of Kv4.2 and Kv4.3 with IC₅₀s between 5 - 70 nM (Diochot et al., 1999), these toxins are gating modifiers stabilising the closed state (Chagot et al., 2004) which shifts the activation curve to more positive potentials (Diochot et al., 1999).

Kv4.2 and Kv4.3 show strong expression in many areas of the brain and heart (Serodio et al., 1996) and consequently have received the most attention. The first member of this family to be cloned, Kv4.1, shows much weaker expression in both heart and brain (Serodio et al., 1996) and as a result the pharmacological profiles of many molecules have mostly omitted this subunit. However, Kv4.1 is expressed in the suprachiasmatic nucleus (Hermansteyne et al., 2017), striatum (Song et al., 1998), basolateral amygdala (Dabrowska and Rainnie, 2010) and nociceptive neurons in the dorsal root ganglion (Phuket and Covarrubias, 2009). The spider toxin *Heteroscodra maculata* 1 (HmTx-1) is reported to inhibit all 3 Kv4 members but also inhibits Kv2.1 (Escoubas et al., 2002). Phrixotoxin-1, although only initially tested on Kv4.2 and Kv4.3, also inhibits Kv4.1 but with 3 times lower affinity; this block seems to be voltage-independent (Yunoki et al., 2014). At present phrixotoxin-1 provides a useful tool for identifying the presence of Kv4 currents and exploring their physiological roles, without being able to distinguish between different Kv4 subunits.

3. Physiological roles of A currents

3.1 Neural excitability and spike properties

A-type channels are rapidly activated upon depolarisation and therefore contribute to action potential repolarisation (Kim et al., 2005; Carrasquillo et al., 2012; Rudy and McBain, 2001), including in the heart (Dixon et al., 1996; Yeola and Snyders, 1997). Kv4 subunits are predominantly located in the somato-dendritic compartments of neurons (Trimmer, 2015), which regulates back-propagation of action potentials into the dendritic tree. The sub-threshold activation range of Kv4 and Kv1.4 channels enables them to regulate intrinsic firing/bursting of neurons in many areas including the: suprachiasmatic nucleus (Hermansteyne et al., 2017), hypothalamus (Mendonça et al., 2018; Imai et al., 2019), dorsal root ganglion (Zemel et al., 2018), hippocampus (Bourdeau et al., 2007), substantia nigra (Liss et al., 2001) and cortex (Carrasquillo et al., 2012). A key factor determining the ability of A-type channels to exert influence is the availability of channels in their non-inactivated state, for example fast spiking will result in more inactivated A-type channels and consequent spike broadening (Kim et al., 2005). The steady-state inactivation curve (Fig. 2A yellow) describes the proportion of channels available at resting membrane potentials. This property is influenced by the presence of accessory subunits (Jerng et al., 1999) and can be dynamically regulated by phosphorylation (Rosenkranz et al., 2009) and Ca²⁺ (Anderson et al., 2010a). Cav3 T-type

channels which are also activated at sub-threshold potentials form complexes with Kv4 channels and provide a source of Ca²⁺ that shifts the inactivation curve of Kv4 channels to more positive potentials (Anderson et al., 2010a; Anderson et al., 2010b), increasing their availability. The presence of Cav3 channels can therefore alter the contribution that A-type channels make to neural excitability (Heath et al., 2014).

Figure 3 near here

Due to the lack of specific inhibitors of Kv1.4, Kv3.3 and Kv3.4 there are few concrete demonstrations of their physiological roles. Throughout the brain Kv1.4 is often found in axons and presynaptic terminals (Veh et al., 1995; Cooper et al., 1998) and may therefore play a role in regulating transmitter release. Kv1.4's slow recovery from inactivation means that with repetitive firing the repolarisation rate of presynaptic action potentials will reduce as Kv1.4 accumulates in its inactivated state. Slower repolarisation of the action potential causes larger presynaptic Ca²⁺ influx and an increase in release probability (Yang and Wang, 2006), suggesting that Kv1.4 may play a role in short term synaptic plasticity. Consistent with this idea, knockdown of Kv1.4 in the hippocampus reduced paired pulse facilitation (Meiri et al., 1998), a form of presynaptic short-term plasticity. Ca²⁺ can also regulate the rate of recovery from inactivation of Kv1.4 in a CamKinase II dependent manner (Roeper et al., 1997), providing scope for regulation of such short-term plasticity. Kv3.3 and Kv3.4 are also located in axons and synaptic terminals (Laube et al., 1996; Ishikawa et al., 2003; Trimmer, 2015) and could have similar roles.

3.2 Dendritic integration and LTP

In hippocampal pyramidal neurons the density of Kv4 channels increases from the soma towards the distal dendrites (Fig. 3A), this gradient depends on expression of the accessory subunit DPP6 (Sun et al., 2011). This gradient of Kv4 channels controls the ability of action potentials to back-propagate and invade the dendritic tree (Hoffman et al., 1997; Johnston et al., 2000) (Fig. 3B), a phenomenon required for induction of synaptic plasticity in these cells (Magee and Johnston, 1997; Buchanan and Mellor, 2007). Consistent with this idea, action potentials more readily back-propagate into the dendritic tree when Kv4.2 channels are knocked down (Fig. 3 B) and consequently the threshold for LTP induction is lowered (Chen et al., 2006). Kv4 channels are also a locus of plasticity in CA1 dendrites being targeted by mitogen-activated protein kinase (Rosenkranz et al., 2009). Induction of LTP results in a shift in the steady-state inactivation of Kv4 channels to more negative-potentials (Fig. 3C) (Frick et al., 2004) and internalisation of channels (Fig. 3D) (Kim et al., 2007), effects which are localised to the site of synaptic input. Both of these changes reduce the available A-type current and consequently increase dendritic excitability and therefore the likelihood of further plasticity.

A similar attenuation of back propagating action potentials by A-type currents is observed in the lateral dendrites of mitral cells of the olfactory bulb (Margrie et al., 2001; Christie and Westbrook, 2003), this influences the extent of lateral dendrodendritic interactions with inhibitory granule cells. A-type K⁺ channels also influence the inhibitory side of the reciprocal synapse between granule and mitral cell dendrites. Granule cells receive excitation from mitral cell dendrites via AMPA and NMDA receptors and provide GABAergic feedback inhibition. The A-type current in granule cells attenuates excitation for the fast AMPA component; spiking and feedback inhibition then occur with a delay, following the time course of the NMDA component (Schoppa and Westbrook, 1999).

The varied biophysics, sub-cellular distributions and dynamic modulation of A-type K⁺ channels enable them to perform a multitude of roles in regulating cellular excitability. Large strides have been made in the 60 years since their discovery but much still remains to be elucidated.

References

- Amadi, C. C., Brust, R. D., Skerritt, M. R., and Campbell, D. L. (2007). Regulation of Kv4.3 closed state inactivation and recovery by extracellular potassium and intracellular KChIP2b. *Channels (Austin)* *1*, 305-314.
- Anderson, D., Mehaffey, W. H., Iftinca, M., Rehak, R., Engbers, J. D., Hameed, S., Zamponi, G. W., and Turner, R. W. (2010a). Regulation of neuronal activity by Cav3-Kv4 channel signaling complexes. *Nat Neurosci* *13*, 333-337.
- Anderson, D., Rehak, R., Hameed, S., Mehaffey, W. H., Zamponi, G. W., and Turner, R. W. (2010b). Regulation of the KV4.2 complex by CaV3.1 calcium channels. *Channels (Austin)* *4*, 163-167.
- Bähring, R., Boland, L. M., Varghese, A., Gebauer, M., and Pongs, O. (2001). Kinetic analysis of open- and closed-state inactivation transitions in human Kv4.2 A-type potassium channels. *The Journal of Physiology* *535*, 65.
- Beck, E. J., Bowlby, M., An, W. F., Rhodes, K. J., and Covarrubias, M. (2002). Remodelling inactivation gating of Kv4 channels by KChIP1, a small-molecular-weight calcium-binding protein. *The Journal of physiology* *538*, 691-706.
- Beck, E. J., and Covarrubias, M. (2001). Kv4 channels exhibit modulation of closed-state inactivation in inside-out patches. *Biophysical Journal* *81*, 867-883.
- Bett, G. C. L., and Rasmusson, R. L. (2004). Inactivation and recovery in Kv1.4 K⁺ channels: lipophilic interactions at the intracellular mouth of the pore. *J Physiol (Lond)* *556*, 109-120.
- Bourdeau, M. L., Morin, F., Laurent, C. E., Azzi, M., and Lacaille, J. C. (2007). Kv4.3-mediated A-type K⁺ currents underlie rhythmic activity in hippocampal interneurons. *J Neurosci* *27*, 1942-1953.

- Buchanan, K. A., and Mellor, J. R. (2007). The development of synaptic plasticity induction rules and the requirement for postsynaptic spikes in rat hippocampal CA1 pyramidal neurones. *The Journal of Physiology* 585, 429-445.
- Carrasquillo, Y., Burkhalter, A., and Nerbonne, J. M. (2012). A-type K⁺ channels encoded by Kv4.2, Kv4.3 and Kv1.4 differentially regulate intrinsic excitability of cortical pyramidal neurons. *J Physiol* 590, 3877-3890.
- Chagot, B., Escoubas, P., Villegas, E., Bernard, C., Ferrat, G., Corzo, G., Lazdunski, M., and Darbon, H. (2004). Solution structure of Phrixotoxin 1, a specific peptide inhibitor of Kv4 potassium channels from the venom of the theraphosid spider Phrixotrichus auratus. *Protein Sci* 13, 1197-1208.
- Chen, X., Yuan, L.-L., Zhao, C., Birnbaum, S. G., Frick, A., Jung, W. E., Schwarz, T. L., Sweatt, J. D., and Johnston, D. (2006). Deletion of Kv4.2 gene eliminates dendritic A-type K⁺ current and enhances induction of long-term potentiation in hippocampal CA1 pyramidal neurons. *J Neurosci* 26, 12143-12151.
- Christie, J. M., and Westbrook, G. L. (2003). Regulation of backpropagating action potentials in mitral cell lateral dendrites by A-type potassium currents. *J Neurophysiol* 89, 2466-2472.
- Clay, J. R. (2000). Determining K⁺ channel activation curves from K⁺ channel currents. *European Biophysics Journal* 29, 555-557.
- Coetzee, W. A., Amarillo, Y., CU, J. O. A. N. N. A., CHOW, A. L. A. N., LAU, D. A. V. I. D., McCORMACK, T. O. M., MORENA, H. E. R. M. A. N., NADAL, M. A. R. C. E. L. A. S., OZAITA, A. N. D. E. R., POUNTNEY, D. A. V. I. D., SAGANICH, M. I. C. H. A. E. L., DE MIERA, E. L. E. A. Z. A. R. V. E. G. A. -S. A. E. N. Z., and RUDY, B. E. R. N. A. R. D. O. (1999). Molecular Diversity of K⁺ Channels. *Ann NY Acad Sci* 868, 233-255.
- Colinas, O., Pérez-Carretero, F. D., López-López, J. R., and Pérez-García, M. T. (2008). A role for DPPX modulating external TEA sensitivity of Kv4 channels. *J Gen Physiol* 131, 455-471.
- Connor, J. A., and Stevens, C. F. (1971). Voltage clamp studies of a transient outward membrane current in gastropod neural somata. *The Journal of physiology* 213, 21.
- Cooper, E. C., Milroy, A., Jan, Y. N., and Jan..., L. Y. (1998). Presynaptic localization of Kv1. 4-containing A-type potassium channels near excitatory synapses in the hippocampus. *Journal of ...*
- Dabrowska, J., and Rainnie, D. G. (2010). Expression and distribution of Kv4 potassium channel subunits and potassium channel interacting proteins in subpopulations of interneurons in the basolateral amygdala. *Neuroscience* 171, 721-733.
- De Miera, E. V.-S., Moreno, H., Fruhling, D., Kentros, C., and Rudy, B. (1992). Cloning of ShIII (Shaw-like) cDNAs encoding a novel high-voltage-activating, TEA-sensitive, type-A K⁺ channel. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 248, 9-18.
- Diochot, S., Schweitz, H., Béress, L., and Lazdunski, M. (1998). Sea anemone peptides with a specific

blocking activity against the fast inactivating potassium channel Kv3.4. *J Biol Chem* 273, 6744-6749.

Diochot, S., Drici, M., Moinier, D., Fink, M., and Lazdunski, M. (1999). Effects of phrixotoxins on the Kv4 family of potassium channels and implications for the role of Ito1 in cardiac electrogenesis. *British journal of pharmacology* 126, 251-263.

Dixon, J. E., Shi, W., Wang, H. S., McDonald, C., Yu, H., Wymore, R. S., Cohen, I. S., and McKinnon, D. (1996). Role of the Kv4.3 K⁺ channel in ventricular muscle. A molecular correlate for the transient outward current. *Circ Res* 79, 659-668.

Dodson, P. D., Barker, M. C., and Forsythe, I. D. (2002). Two Heteromeric Kv1 Potassium Channels Differentially Regulate Action Potential Firing. *J. Neurosci.* 22, 6953-6961.

Escoubas, P., Diochot, S., Celerier, M.-L., Nakajima, T., and Lazdunski, M. (2002). Novel Tarantula Toxins for Subtypes of Voltage-Dependent Potassium Channels in the Kv2 and Kv4 Subfamilies. *Mol Pharmacol* 62, 48-57.

Frick, A., Magee, J., and Johnston, D. (2004). LTP is accompanied by an enhanced local excitability of pyramidal neuron dendrites. *Nat Neurosci* 7, 126-135.

Hagiwara, S., Kusano, K., and Saito, N. (1961). Membrane changes of Onchidium nerve cell in potassium-rich media. *The Journal of Physiology* 155, 470.

Hanson, D. C., Nguyen, A., Mather, R. J., Rauer, H., Koch, K., Burgess, L. E., Rizzi, J. P., Donovan, C. B., Bruns, M. J., and Canniff, P. C. (1999). UK-78,282, a novel piperidine compound that potently blocks the Kv1.3 voltage-gated potassium channel and inhibits human T cell activation. *British journal of pharmacology* 126, 1707-1716.

Heath, N. C., Rizwan, A. P., Engbers, J. D., Anderson, D., Zamponi, G. W., and Turner, R. W. (2014). The expression pattern of a Cav3-Kv4 complex differentially regulates spike output in cerebellar granule cells. *J Neurosci* 34, 8800-8812.

Hermanstynne, T. O., Granados-Fuentes, D., Mellor, R. L., Herzog, E. D., and Nerbonne, J. M. (2017). Acute Knockdown of Kv4.1 Regulates Repetitive Firing Rates and Clock Gene Expression in the Suprachiasmatic Nucleus and Daily Rhythms in Locomotor Behavior. *eNeuro* 4,

Hoffman, D. A., Magee, J. C., Colbert, C. M., and Johnston, D. (1997). K⁺ channel regulation of signal propagation in dendrites of hippocampal pyramidal neurons. *Nature* 387, 869-875.

Hoshi, T., Zagotta, W. N., and Aldrich, R. W. (1990). Biophysical and molecular mechanisms of Shaker potassium channel inactivation. *Science* 250, 533-538.

Hoshi, T., Zagotta, W. N., and Aldrich, R. W. (1991). Two types of inactivation in Shaker K⁺ channels: effects of alterations in the carboxy-terminal region. *Neuron* 7, 547-556.

- Imai, R., Yokota, S., Horita, S., Ueta, Y., Maejima, Y., and Shimomura, K. (2019). Excitability of oxytocin neurons in paraventricular nucleus is regulated by voltage-gated potassium channels Kv4.2 and Kv4.3. *Bioscience, Biotechnology, and Biochemistry* 83, 202-211.
- Ishikawa, T., Nakamura, Y., Saitoh, N., Li, W.-B., Iwasaki, S., and Takahashi, T. (2003). Distinct Roles of Kv1 and Kv3 Potassium Channels at the Calyx of Held Presynaptic Terminal. *J. Neurosci.* 23, 10445-10453.
- Jerng, H. H., Shahidullah, M., and Covarrubias, M. (1999). Inactivation gating of Kv4 potassium channels: molecular interactions involving the inner vestibule of the pore. *The Journal of general physiology* 113, 641-660.
- Jerng, H. H., Pfaffinger, P. J., and Covarrubias, M. (2004). Molecular physiology and modulation of somatodendritic A-type potassium channels. *Mol Cell Neurosci* 27, 343-369.
- Johnston, D., Hoffman, D. A., Magee, J. C., Poolos, N. P., Watanabe, S., Colbert, C. M., and Migliore, M. (2000). Dendritic potassium channels in hippocampal pyramidal neurons. *J Physiol* 525 Pt 1, 75-81.
- Johnston, J., Forsythe, I. D., and Kopp-Scheinpflug, C. (2010). Going native: voltage-gated potassium channels controlling neuronal excitability. *J Physiol* 588, 3187-3200.
- Johnston, J., Griffin, S. J., Baker, C., and Forsythe, I. D. (2008). Kv4 (A-type) potassium currents in the mouse medial nucleus of the trapezoid body. *Eur J Neurosci* 27, 1391-1399.
- Kim, J., Jung, S. C., Clemens, A. M., Petralia, R. S., and Hoffman, D. A. (2007). Regulation of dendritic excitability by activity-dependent trafficking of the A-type K⁺ channel subunit Kv4.2 in hippocampal neurons. *Neuron* 54, 933-947.
- Kim, J., Wei, D.-S., and Hoffman, D. A. (2005). Kv4 potassium channel subunits control action potential repolarization and frequency-dependent broadening in rat hippocampal CA1 pyramidal neurones. *J Physiol* 569, 41-57.
- Kuo, Y.-L., Cheng, J.-K., Hou, W.-H., Chang, Y.-C., Du, P.-H., Jian, J.-J., Rau, R.-H., Yang, J.-H., Lien, C.-C., and Tsaur, M.-L. (2017). K⁺ channel modulatory subunits KChIP and DPP participate in Kv4-mediated mechanical pain control. *Journal of Neuroscience* 37, 4391-4404.
- Laube, G., Röper, J., Pitt, J. C., Sewing, S., Kistner, U., Garner, C. C., Pongs, O., and Veh, R. W. (1996). Ultrastructural localization of Shaker-related potassium channel subunits and synapse-associated protein 90 to septate-like junctions in rat cerebellar Pinceaux. *Brain Res Mol Brain Res* 42, 51-61.
- Lee, Y. T., Vasilyev, D. V., Shan, Q. J., Dunlop, J., Mayer, S., and Bowlby, M. R. (2008). Novel pharmacological activity of loperamide and CP-339,818 on human HCN channels characterized with an automated electrophysiology assay. *Eur J Pharmacol* 581, 97-104.
- Liss, B., Franz, O., Sewing, S., Bruns, R., Neuhoff, H., and Roeper, J. (2001). Tuning pacemaker frequency of

individual dopaminergic neurons by Kv4.3L and KChip3.1 transcription. *EMBO J* 20, 5715-5724.

Liu, P., Jo, S., and Bean, B. P. (2012). Modulation of neuronal sodium channels by the sea anemone peptide BDS-I. *J Neurophysiol* 107, 3155-3167.

MacKinnon, R., Aldrich, R. W., and Lee, A. W. (1993). Functional stoichiometry of Shaker potassium channel inactivation. *Science* 262, 757-759.

Magee, J. C., and Johnston, D. (1997). A synaptically controlled, associative signal for Hebbian plasticity in hippocampal neurons. *Science* 275, 209-213.

Margrie, T. W., Sakmann, B., and Urban, N. N. (2001). Action potential propagation in mitral cell lateral dendrites is decremental and controls recurrent and lateral inhibition in the mammalian olfactory bulb. *Proc Natl Acad Sci U S A* 98, 319-324.

Mehra, D., Imtiaz, M. S., van Helden, D. F., Knollmann, B. C., and Laver, D. R. (2014). Multiple modes of ryanodine receptor 2 inhibition by flecainide. *Mol Pharmacol* 86, 696-706.

Meiri, N., Sun, M. K., Segal, Z., and Alkon, D. L. (1998). Memory and long-term potentiation (LTP) dissociated: normal spatial memory despite CA1 LTP elimination with Kv1.4 antisense. *Proc Natl Acad Sci U S A* 95, 15037-15042.

Mendonça, P. R. F., Kyle, V., Yeo, S. H., Colledge, W. H., and Robinson, H. P. C. (2018). Kv4.2 channel activity controls intrinsic firing dynamics of arcuate kisspeptin neurons. *J Physiol* 596, 885-899.

Molineux, M. L., Fernandez, F. R., Mehaffey, W. H., and Turner, R. W. (2005). A-type and T-type currents interact to produce a novel spike latency-voltage relationship in cerebellar stellate cells. *J Neurosci* 25, 10863-10873.

Nguyen, A., Kath, J. C., Hanson, D. C., Biggers, M. S., Canniff, P. C., Donovan, C. B., Mather, R. J., Bruns, M. J., Rauer, H., Aiyar, J., Lepple-Wienhues, A., Gutman, G. A., Grissmer, S., Cahalan, M. D., and Chandy, K. G. (1996). Novel nonpeptide agents potently block the C-type inactivated conformation of Kv1.3 and suppress T cell activation. *Mol Pharmacol* 50, 1672-1679.

Paul, A. A., Witchel, H. J., and Hancox, J. C. (2002). Inhibition of the current of heterologously expressed HERG potassium channels by flecainide and comparison with quinidine, propafenone and lignocaine. *British journal of pharmacology* 136, 717-729.

Phuket, T. R., and Covarrubias, M. (2009). Kv4 Channels Underlie the Subthreshold-Operating A-type K-current in Nociceptive Dorsal Root Ganglion Neurons. *Front Mol Neurosci* 2, 3.

Ramos, E., and O'leary, M. E. (2004). State-dependent trapping of flecainide in the cardiac sodium channel. *J Physiol* 560, 37-49.

Rasmusson, R. L., Morales, M. J., Castellino, R. C., Zhang, Y., Campbell, D. L., and Strauss, H. C. (1995). C-

type inactivation controls recovery in a fast inactivating cardiac K⁺ channel (Kv1.4) expressed in *Xenopus* oocytes. *The Journal of Physiology* 489, 709-721.

Rettig, J., Heinemann, S. H., Wunder, F., Lorra, C., Parcej, D. N., Dolly, J. O., and Pongs, O. (1994). Inactivation properties of voltage-gated K⁺ channels altered by presence of β -subunit. *Nature* 369, 289-294.

Rettig, J., Wunder, F., Stocker, M., Lichtinghagen, R., Mastiaux, F., Beckh, S., Kues, W., Pedarzani, P., Schröter, K. H., and Ruppersberg, J. P. (1992). Characterization of a Shaw-related potassium channel family in rat brain. *The EMBO Journal* 11, 2473-2486.

Rhodes, K. J., Carroll, K. I., Sung, M. A., Doliveira, L. C., Monaghan, M. M., Burke, S. L., Strassle, B. W., Buchwalder, L., Menegola, M., Cao, J., An, W. F., and Trimmer, J. S. (2004). KChIPs and Kv4 alpha subunits as integral components of A-type potassium channels in mammalian brain. *J Neurosci* 24, 7903-7915.

Robertson, B., Owen, D., Stow, J., Butler, C., and Newland, C. (1996). Novel effects of dendrotoxin homologues on subtypes of mammalian Kv1 potassium channels expressed in *Xenopus* oocytes. *FEBS Letters* 383, 26-30.

Roeper, J., Lorra, C., and Pongs, O. (1997). Frequency-dependent inactivation of mammalian A-type K⁺ channel KV1.4 regulated by Ca²⁺/calmodulin-dependent protein kinase. *Journal of Neuroscience* 17, 3379-3391.

Rosenkranz, J. A., Frick, A., and Johnston, D. (2009). Kinase-dependent modification of dendritic excitability after long-term potentiation. *J Physiol* 587, 115-125.

Rudy, B., Chow, A., Lau, D., Amarillo, Y., Ozaita, A., Saganich, M., Moreno, H., Nadal, M. S., HERNANDEZ-PINEDA, R. I. C. A. R. D. O., and HERNANDEZ-CRUZ, A. R. T. U. R. O. (1999). Contributions of Kv3 channels to neuronal excitability. *Annals of the New York Academy of Sciences* 868, 304-343.

Rudy, B., and McBain, C. J. (2001). Kv3 channels: voltage-gated K⁺ channels designed for high-frequency repetitive firing. *Trends in Neurosciences* 24, 517-526.

Ruppersberg, J. P., Schröter, K. H., Sakmann, B., Stocker, M., Sewing, S., and Pongs, O. (1990). Heteromultimeric channels formed by rat brain potassium-channel proteins. *Nature* 345, 535-537.

Schoppa, N. E., and Westbrook, G. L. (1999). Regulation of synaptic timing in the olfactory bulb by an A-type potassium current. *Nat Neurosci* 2, 1106-1113.

Serodio, P., Vega-Saenz de Miera, E., and Rudy, B. (1996). Cloning of a novel component of A-type K⁺ channels operating at subthreshold potentials with unique expression in heart and brain. *Journal of neurophysiology* 75, 2174-2179.

Sforna, L., D'Adamo, M. C., Servettini, I., Guglielmi, L., Pessia, M., Franciolini, F., and Catacuzzeno, L. (2015). Expression and function of a CP339,818-sensitive K⁺ current in a subpopulation of putative nociceptive

neurons from adult mouse trigeminal ganglia. *Journal of Neurophysiology* 113, 2653-2665.

Shahidullah, M., and Covarrubias, M. (2003). The link between ion permeation and inactivation gating of Kv4 potassium channels. *Biophysical journal* 84, 928-941.

Shevchenko, T., Teruyama, R., and Armstrong, W. E. (2004). High-Threshold, Kv3-Like Potassium Currents in Magnocellular Neurosecretory Neurons and Their Role in Spike Repolarization. *J Neurophysiol* 92, 3043-3055.

Song, W.-J., Tkatch, T., Baranauskas, G., Ichinohe, N., Kitai, S. T., and Surmeier, D. J. (1998). Somatodendritic depolarization-activated potassium currents in rat neostriatal cholinergic interneurons are predominantly of the A type and attributable to coexpression of Kv4. 2 and Kv4. 1 subunits. *Journal of Neuroscience* 18, 3124-3137.

Stühmer, W., Ruppersberg, J. P., Schröter, K. H., Sakmann, B., Stocker, M., Giese, K. P., Perschke, A., Baumann, A., and Pongs, O. (1989). Molecular basis of functional diversity of voltage-gated potassium channels in mammalian brain. *The EMBO Journal* 8, 3235-3244.

Sun, W., Maffie, J. K., Lin, L., Petralia, R. S., Rudy, B., and Hoffman, D. A. (2011). DPP6 establishes the A-type K(+) current gradient critical for the regulation of dendritic excitability in CA1 hippocampal neurons. *Neuron* 71, 1102-1115.

Trimmer, J. S. (2015). Subcellular localization of K+ channels in mammalian brain neurons: remarkable precision in the midst of extraordinary complexity. *Neuron* 85, 238-256.

Veh, R. W., Lichtinghagen, R., Sewing, S., Wunder, F., Grumbach, I. M., and Pongs, O. (1995). Immunohistochemical localization of five members of the Kv1 channel subunits: contrasting subcellular locations and neuron-specific co-localizations in rat brain. *European Journal of Neuroscience* 7, 2189-2205.

Wang, F. C., Parcej, D. N., and Dolly, J. O. (1999). Subunit compositions of Kv1.1-containing K+channel subtypes fractionated from rat brain using dendrotoxins. *Eur J Biochem* 263, 230-237.

Watanabe, S., Hoffman, D. A., Migliore, M., and Johnston, D. (2002). Dendritic K+ channels contribute to spike-timing dependent long-term potentiation in hippocampal pyramidal neurons. *Proc Natl Acad Sci U S A* 99, 8366-8371.

Weiser, M., De Miera, E. V.-S., Kentros, C., Moreno, H., Franzen, L., Hillman, D., Baker, H., and Rudy, B. (1994). Differential expression of Shaw-related K+ channels in the rat central nervous system. *Journal of Neuroscience* 14, 949-972.

Yang, Y.-M., and Wang, L.-Y. (2006). Amplitude and Kinetics of Action Potential-Evoked Ca₂⁺ Current and Its Efficacy in Triggering Transmitter Release at the Developing Calyx of Held Synapse. *J Neurosci.* 26, 5698-5708.

- Yeola, S. W., and Snyders, D. J. (1997). Electrophysiological and pharmacological correspondence between Kv4.2 current and rat cardiac transient outward current. *Cardiovascular research* *33*, 540-547.
- Yeung, S. Y., Thompson, D., Wang, Z., Fedida, D., and Robertson, B. (2005). Modulation of Kv3 subfamily potassium currents by the sea anemone toxin BDS: significance for CNS and biophysical studies. *J Neurosci* *25*, 8735-8745.
- Yunoki, T., Takimoto, K., Kita, K., Funahashi, Y., Takahashi, R., Matsuyoshi, H., Naito, S., and Yoshimura, N. (2014). Differential contribution of Kv4-containing channels to A-type, voltage-gated potassium currents in somatic and visceral dorsal root ganglion neurons. *J Neurophysiol* *112*, 2492-2504.
- Zemel, B. M., Ritter, D. M., Covarrubias, M., and Muqeem, T. (2018). A-Type K_v Channels in Dorsal Root Ganglion Neurons: Diversity, Function, and Dysfunction. *Front Mol Neurosci* *11*, 253.
- Zhou, M., Morais-Cabral, J. H., Mann, S., and MacKinnon, R. (2001). Potassium channel receptor site for the inactivation gate and quaternary amine inhibitors. *Nature* *411*, 657-661.




Figure 1. Initial observations of A-type currents in *Onchidium*. **A)** Current clamp recording of action potentials evoked from rest with short spike latencies. **B)** A preceding hyperpolarising step removes steady-state inactivation of the A-type current and results in a lag to the first spike as the A-current inactivates (yellow arrow). **C)** Voltage-clamp recording showing a fast inactivating outward current activated at voltage around resting membrane potentials after hyperpolarising steps, yellow arrow. **D)** The A-current is present at more positive potentials only with a hyperpolarising pre-step, yellow arrow. The delayed rectifier can be isolated by inactivating the A-current. Adapted from Hagiwara et al., (1961) reproduced with permission.




Figure 2. Biophysical and pharmacological identification of channels mediating native A-type K⁺ currents. **A)** *Left:* Voltage protocol to determine the activation and inactivation curves for A-type channels. The activation curve can be constructed by measuring the peak current soon after the test step (red arrow) and correcting it for the non-linear driving force (see text). The steady-state inactivation curve can be obtained by measuring the current at the yellow arrow as a function of the preceding test step. Note: the duration of the initial hyperpolarising pre-pulse and the test step should be adjusted depending on the kinetics of that current being measured, e.g. the duration pre-pulse should ensure complete recovery from inactivation. *Right:* The rate of recovery from inactivation can be determined with a pair of test steps of

sufficient duration to allow complete inactivation of the A-current and separated by a variable delay (Δt). The time constant of recovery can then be determined by fitting an exponential to the current amplitude of the second step as a function of Δt . Note the current amplitude of the first step should be constant. Adapted from Johnston *et al*, (2008). **B)** A biophysical and pharmacological procedure to aid identification of the channel mediating a native A-current (see text for further description).




Figure 3. Dendritic Kv4 A-type channels regulate and are a locus of plasticity in hippocampal CA1 neurons. **A**) Dendritic recordings of Kv₄ A-type K⁺ currents demonstrate a gradient of current that increases with distance from soma adapted from Hoffman et al., (1997) with permission. **B**) Imaging of fura-2 loaded CA1 neurons reveal attenuation of back-propagating action potential evoked Ca²⁺ transients. This attenuation is dramatically reduced when Kv_{4.2} is absent, adapted from Chen, *et al.* (2006, Copyright 2006, Society for Neuroscience) with permission. **C**) The steady-state inactivation curve of dendritically located Kv₄ channels is shifted to more hyperpolarised potentials after induction of LTP with theta burst stimuli. Note that this decreases the available Kv₄ current at resting membrane potentials by ~50%, adapted from Frick et al., (2004) with permission. **D**) Fluorescently tagged A-type channels (Kv_{4.2g}) are trafficked from the membrane following activation of synaptic receptors, adapted from Kim et al., (2007) with permission.