

This is a repository copy of *Climate change impacts on water security in global drylands*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/175447/>

Version: Accepted Version

Article:

Stringer, Lindsay Carman orcid.org/0000-0003-0017-1654, Mirzabaev, Alisher, Benjaminsen, Tor A. et al. (5 more authors) (2021) Climate change impacts on water security in global drylands. *One Earth*. ISSN 2590-3322

<https://doi.org/10.1016/j.oneear.2021.05.010>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

One Earth

Climate change impacts on water security in global drylands

--Manuscript Draft--

Manuscript Number:	ONE-EARTH-D-20-00360R2
Full Title:	Climate change impacts on water security in global drylands
Article Type:	Review
Keywords:	Adaptation; aridity; integrated water resources management; greening; browning; projections
Corresponding Author:	Lindsay Stringer University of York Environment Department York, United Kingdom UNITED KINGDOM
First Author:	Lindsay Stringer
Order of Authors:	Lindsay Stringer Alisher Mirzabaev Tor A Benjaminsen Rebecca M B Harris Mostafa Jafari Tabea Lissner Nicola Stevens Cristina Tirado-von der Pahlen
Abstract:	<p>Water scarcity affects 1-2 billion people globally, most of whom live in drylands. Under projected climate change, millions more people will be living under conditions of severe water stress in the coming decades. This review examines observed and projected climate change impacts on water security across the world's drylands to the year 2100. We find that efficient water management, technology and infrastructure, and better demand and supply management, can offer more equitable access to water resources. People are already adapting but need to be supported with coherent system-oriented policies and institutions that situate water security at their core, in line with the components of Integrated Water Resources Management. Dryland water governance urgently needs to better account for synergies and trade-offs between water security and other dimensions of sustainable development, to support an equitable approach in which no one gets left behind.</p>
Opposed Reviewers:	
Suggested Reviewers:	

Cover letter

21 May 2021

Dear Erika

Please find our revised article herewith.

Best wishes

Lindsay

21/5/2021

Dear Erika

Editorial feedback has been addressed and we hope all is in order so that this can be published in the June issue as indicated.

Best wishes

Lindsay

Climate change impacts on water security in global drylands

Lindsay C. Stringer^{1*}, Alisher Mirzabaev², Tor A. Benjaminsen³, Rebecca M. B. Harris⁴, Mostafa Jafari,⁵ Tabea K. Lissner⁶, Nicola Stevens⁷, Cristina Tirado-von der Pahlen^{8,9}

¹Department of Environment and Geography, University of York, York, United Kingdom

²Center for Development Research (ZEF), University of Bonn, Bonn, Germany

³Department of International Environment and Development Studies, Faculty of Landscape and Society, Norwegian University of Life Sciences, Norway

⁴School of Geography, Planning and Spatial Science, University of Tasmania, Hobart, Australia

⁵Research Institute of Forests and Rangelands (RIFR), Agricultural Research Education and Extension Organisation (AREEO), Tehran, Iran

⁶Climate Analytics gGmbH, Berlin, Germany

⁷Environmental Change Institute, School of Geography and the Environment, University of Oxford, Oxford, United Kingdom

⁸Loyola Marymount University, Department of International Relations and Political Science, Los Angeles, USA

⁹University of California Los Angeles, Institute of Environment and Sustainability, Los Angeles, United States of America

* Correspondence: lindsay.stringer@york.ac.uk

Summary

Water scarcity affects 1-2 billion people globally, most of whom live in drylands. Under projected climate change, millions more people will be living under conditions of severe water stress in the coming decades. This review examines observed and projected climate change impacts on water security across the world's drylands to the year 2100. We find that efficient water management, technology and infrastructure, and better demand and supply management, can offer more equitable access to water resources. People are already adapting but need to be supported with coherent system-oriented policies and institutions that situate water security at their core, in line with the components of Integrated Water Resources Management. Dryland water governance urgently needs to better account for synergies and trade-offs between water security and other dimensions of sustainable development, to support an equitable approach in which no one gets left behind.

Introduction

Drylands are the hyper-arid, arid, semi-arid and dry sub-humid parts of the Earth, found on all continents. Grasslands, savannas and woodlands in these environments are rich in biodiversity¹, and store substantial amounts of the world's terrestrial carbon in their soils and biomass. Drylands possess a varied and rich geological, cultural and historical heritage^{2,3} and are home to approximately 40 percent of the world's human population⁴ (Figure 1; based on data from: Centre for International Earth Science Information Network (CIESIN), Columbia University (2018)⁵. Dryland extent is based on Millennium Ecosystem Assessment delineation⁶). Major land uses include agriculture and pastoralism, with the majority of livelihoods directly reliant upon natural resources. A number of megacities including New Delhi, Beijing, Los Angeles, Cairo, Tehran and Mexico City, all of which have complex and diversified economies, are also located in these water-limited environments⁷. Life can be very difficult for people living in dryland areas, particularly in developing regions, where around 70% of the world's drylands are found⁸. In many of these areas, people already face stark challenges related to poverty, food insecurity and malnourishment, poor access to healthcare, poor governance, economic hardship and marginalisation⁹. These difficulties are often exacerbated by land degradation, flooding, drought, and climate change. Drylands in hot, tropical areas have already experienced temperature rises that are higher than the global average, and temperatures are projected to increase by 2-4°C by

2100 under higher emissions scenarios (Representative Concentration Pathway (RCP) 4.5 and 8.5)¹⁰. Understanding what these changes mean for water security in drylands is therefore vital.

Several different measures are relevant to the assessment of observed and projected future water security in drylands, and because they consider slightly different aspects of the system, they can highlight different trends. Climatological indices measure the physical components of water security, and include the Aridity Index (AI), drought indices such as the Palmer Drought Severity Index (PDSI) and the Standardized Precipitation Evapotranspiration Index (SPEI), the Ecohydrological index (EI), and Soil Moisture and Terrestrial water storage (TWS). There are also measures that indicate biological responses to physical or climatological variables, such as changes to dryland vegetation indicated by the Normalized Difference Vegetation Index (NDVI). Box 1 explains these terms for each of the indices referred to in the main text in relation to observed and projected impacts of climate change on water security.

Box 1: The Aridity Paradox: Defining and delineating the drylands

'Dryland extent' describes both the physical boundaries of dry areas (a climatological definition, commonly measured using the Aridity Index), and the extent of dryland vegetation (an ecological definition). Climatological and ecological definitions do not always delineate the same geographical areas when projecting future changes to dryland extent.

The **Aridity Index (AI)** is the ratio of annual precipitation to potential evapotranspiration. It has a long use history, defining drylands as areas with $AI < 0.65$ ¹¹. Sub-categories include: (i) dry sub-humid ($0.5 \leq AI < 0.65$), (ii) semi-arid ($0.2 \leq AI < 0.5$), (iii) arid ($0.05 \leq AI < 0.2$) and (iv) hyper-arid ($AI < 0.05$) areas (Figure 1). Cold (polar) drylands (not considered here), are where potential evapotranspiration is < 400 mm/year¹². The AI usually projects increasing aridity under climate change, leading to projections of widespread dryland expansion^{13,14}. However, while the AI has been decreasing over the last 50 years¹⁵, dryland vegetation has been increasing globally¹⁶⁻¹⁹. Hence, correspondence between changes in AI and changes in dryland vegetation over recent decades is limited¹⁵. The AI overestimates the role of potential evapotranspiration compared with rainfall¹⁵, and neglects CO₂ impacts on evapotranspiration and seasonality in rainfall and evapotranspiration. Increased annual potential evapotranspiration due to higher temperatures may have little impact if temperature and actual evapotranspiration are not increasing during the wet season when there is vegetation growth. Given the AI's limitations, indices that incorporate the influence of plant physiology on evapotranspiration, such as precipitation minus actual evapotranspiration, soil moisture, runoff and land water storage, may be more suitable for future projections. The **Ecohydrological index (EI)** is directly based on observations of land surface ecohydrological properties using Coupled Model Intercomparison Project Phase 5 (CMIP5) models¹⁵. The EI aims to capture the role of vegetation responses under higher CO₂ levels. Results show that EI decreases in some regions (reflecting increased aridity) and increases in others, better capturing observed dryland changes than the AI.

The **Palmer Drought Severity Index (PDSI)** is a standardized index, generally spanning -10 (dry) to +10 (wet). Values lower than -3 represent severe to extreme meteorological drought. The PDSI incorporates prior precipitation, moisture supply, runoff and evaporation demand at the surface level to estimate relative dryness²⁰. It is based on temperature data and a physical water balance model, so can capture global warming effects on drought through changes in potential evapotranspiration. However, it does not compare well across regions, and is not amenable to assessing short timescales, making it difficult to correlate with specific water resources²¹. The **Standardized Precipitation Index (SPI)** characterizes meteorological drought on a range of timescales. The SPI can be calculated for periods of 1-36 months, using monthly input data, so can characterize drought at timescales corresponding with the temporal availability of different water resources (such as soil moisture, groundwater, river discharge and reservoir storage). The SPI is more comparable across regions than the PDSI because it is calculated in relation to climatological norms for the location and season. However, the SPI does not consider evapotranspiration, so does not capture the effect of increased temperatures associated with climate change on moisture demand and availability.

The **Standardized Precipitation Evapotranspiration Index (SPEI)** is a drought index, calculated as the difference between precipitation and potential evapotranspiration (PET). By incorporating evaporation, it captures the main impact of increased temperatures on water demand. It can be calculated at different timescales (e.g. monthly or weekly). The SPEI is used to measure drought severity in terms of intensity and duration and can identify the onset and end of drought episodes. **Potential evapotranspiration (PET)** (or Potential Evaporation) describes the amount of evaporation that would occur if unlimited water were available. It is influenced by surface and air temperatures, radiation and wind, vegetation characteristics, such as ground cover and plant density, and soil type. By definition, annual potential evaporation exceeds annual precipitation in drylands. PET is estimated using various methods, such as the Penman–Monteith equation. The **surface water balance**, the difference between precipitation and actual evapotranspiration, describes the availability of surface water on land, while **soil moisture** is the water content stored in a given layer of soil. Global analyses rely on satellite data or model simulations because in situ observations are still unavailable for most of the world.

Terrestrial water storage (TWS) is the sum of continental water stored in vegetation, rivers, lakes and reservoirs, wetlands, soil and groundwater. It is critical in the global water and energy budget, influencing water resource availability and water flux interactions among Earth system components²². While groundwater is an important component of the dryland ecohydrological system, its role in TWS remains poorly quantified²³. The **Normalized Difference Vegetation Index (NDVI)** is based on global satellite data which characterises vegetation growth by assessing absorption and reflection of photosynthetically active radiation over a given time period, relative to the regional norm. The NDVI describes the relative density of vegetation and is used as an indicator of agricultural drought.

By definition, precipitation in drylands balances evaporation from the land and vegetation surfaces. As a result, water is a key limiting factor that shapes the drylands, with these systems being highly sensitive to precipitation and potential evapotranspiration dynamics. Globally, water scarcity already affects between one and two billion people, the vast majority of whom live in drylands, where the gap

between the demand for and supply of water is the highest in the world. This challenge means that the impacts of climate change, combined with water management decisions, will have profound impacts on drylands and their inhabitants into the future. Projected climate changes indicate that in a matter of just a few decades, millions more people (approximately half the world's population in total) will be living under conditions of high water stress²⁴. This will have impacts not just in dryland areas, but also for neighbouring countries and beyond, particularly because water and its impacts do not respect national political and administrative boundaries.

The impacts of climate change on water security in drylands go beyond access to clean water and sanitation; they are highly intertwined with many other dimensions of sustainable development, including eradicating hunger and reducing poverty, peace and security, gender equality, education and health²⁵ (Figure 2). For example, water scarcity can negatively affect women more so than men because of women's key role in household water provisioning in many developing countries²⁶ resulting in more time spent by female household members, including children, in fetching water for domestic consumption. Climate change affects the structure and functioning of multiple ecosystem attributes across the world's drylands²⁷, with important implications for agriculture and vegetation productivity, as well as the livelihoods they support. Increased water scarcity due to climate change will make attainment of the Sustainable Development Goals (SDGs) more difficult in many drylands, especially in the developing world¹⁰. Improved knowledge about new emerging hotspots of climate change-driven water insecurity will be critical for measuring progress toward the achievement of the SDGs²⁸.

Taking into account the magnitude and importance of these emerging challenges, this review focuses on the impacts of climate change on water security in drylands, considering both environmental and human systems. Water security has been defined in various ways depending on the discipline, method and scale of analysis that is adopted²⁹, but the different definitions agree that water security and water scarcity are multi-faceted. Water scarcity is often perceived as determined by the (non-)availability of water resources, also termed physical water scarcity, however scarcity can also result from lack of infrastructure (economic water scarcity), poor water quality (clean water scarcity) or insufficient retention of water resources in ecosystems (environmental water scarcity). Most global freshwater resources are transboundary, adding a political water security dimension: potential disruption to the availability, access and quality of water resources when managed across two or more jurisdictions. These determinants vary in their spatial distribution, where physically water abundant regions can be water scarce due to high pollution levels or lack of infrastructure or uncoordinated transboundary water use (Figure 3, which uses AQUASTAT/FAO values for 2015³⁰, calculations based on Falkenmark et al. 1989³¹; and water quality risk based on Damania et al., 2019³²). Water security therefore encompasses aspects of availability, accessibility, quality, and stability²⁸ (Figure 4). Stability refers to the time dimension of water availability, access and quality. If availability, access and quality of water resources fluctuate substantially, people cannot be considered water secure. Political water security is a major factor that can affect water stability, and climate change and associated increases in rainfall variability are becoming a key source of instability in water security. While water security can be quantitatively measured^{28,33} across physical, economic and quality dimensions, it also involves contested views, particularly regarding what is implied by stability and political water security. The dynamic and subjective nature of these concepts makes it difficult to create maps showing water stability and political water security. Water security may be linked to food security, to energy security, to physical water scarcity generating conflicts, or even viewed as a potential weapon in conflicts³⁴. While physical water scarcity is real in many locations, for instance as a result of dwindling groundwater aquifer supplies or increased salinity, a discourse of scarcity may also be used to justify certain political interests (e.g. building a high dam for hydro-power development)³⁵. Climate change directly affects all four dimensions of water security.

The remainder of this review synthesises: a) observed climate changes in drylands to date and their impacts on water security; b) projected future changes under different climate change projections, considering what may be anticipated in terms of water security challenges in the future; and c) the management of water security challenges in drylands. Effective water management under a changing climate needs to target all dimensions of water (in)security in a holistic way. The review concludes by presenting the major features of such a multidimensional approach through the concept of integrated water resource management (IWRM) system for drylands.

Observed climate changes and their impacts on water security

Observed changes in temperature, rainfall and evaporation over recent decades have already affected dryland extent and water security in many areas of the world¹⁰. In some drylands, rising temperatures have augmented aridity where increases in potential evapotranspiration outpace those of precipitation^{36,37}, and temperature and aridity increases are exacerbated by the sparse vegetation cover and lower soil moisture of dryland ecosystems¹³. However, this is not a global trend as many drylands are experiencing increases in vegetation cover and rainfall.

Both the amount of rainfall and its seasonality have changed in many dryland areas, associated with both natural decadal variability and anthropogenic warming³⁶, affecting both availability and stability dimensions of water security. Annual rainfall has increased in some locations (e.g. the West African Sahel, the Karoo in South Africa, Gobi Desert in China, and central/west Australia) and decreased in others (e.g. east Australia, and parts of East Asia), often with fewer, more intense rainfall events and increased unpredictability¹⁴. Such changes have had major impacts on water security in desert and semi-arid areas. For those people who directly depend on the natural resource base for their livelihoods, such variability and change present a huge challenge.

Changes in surface water availability (the difference between precipitation and actual evapotranspiration) over drylands have been demonstrated globally over the period 1982 to 2011, with significant decreasing trends occurring in south-western North America and west Asia and significant increasing trends in northern Australia and central and southern Africa¹⁴. Over the same period, a weak decrease in soil moisture has occurred in drylands globally, although in some regions, including parts of central western Australia, southern North America, southern South America, west Asia and the Mongolian Plateau, reductions in soil moisture have been significant¹⁴. A strong increasing trend in mean NDVI over the past three decades has occurred, consistent with the widely reported greening over Africa, Australia and south Asia. Greening is associated with precipitation increase in most regions, with the exception of some irrigated regions in western USA, Arab regions and north-eastern parts of China, where precipitation has decreased but NDVI increased¹⁴.

Terrestrial water storage (TWS) in drylands has also declined globally over the period 2002 to 2017, with stronger trends found in hyper-arid and arid regions³⁸. Model experiments suggest that the observed reductions in TWS are due to anthropogenic warming over south-western North America and the Middle East, while water withdrawals have contributed more to the recent declines in TWS over North China³⁸.

When measured using the AI, dryland area has increased by ~4% from 1948 to 2004^{12,39,40}, with the semi-arid zone expanding at the greatest rate (~7%)¹³. However as described in Box 1, the AI approach is widely critiqued and contested. The rapid rate of warming in drylands and the observed increase in climatological extent of dryland climates, as measured by the AI, has led to assertions that changing aridity would result in a global trend of declining dryland vegetation productivity, and the expansion of drylands^{36,41}. However, interactions with land-use and variable changes in rainfall and carbon-fertilization mean that dryland ecosystems have shown mixed trends of decreases and increases in vegetation and biodiversity in different locations^{10,42,43}, both of which can create long-term failure to

meet demands for and supply of ecosystem goods and services⁷. Drought and warming-associated tree cover decline and tree mortality have been recorded in localised areas in North Africa's drylands^{44,45}, parts of the arid diagonal in South America⁴⁶, patches in North America's Mojave, Chihuahuan and Sonoran deserts^{43,47} and parts of the Middle East and southwest Asia^{43,48}, with desert succulent species appearing susceptible to both heat and drought induced mortality^{49,50}. Hot droughts in particular may also reduce the resilience of the system, making plants vulnerable to secondary agents of mortality like disturbance or disease⁵¹ and altering vegetation recovery after disturbances (e.g. fire)⁵².

Desertification or vegetation browning has been observed in the western USA, eastern Brazil, Iraq, Syria, Jordan, Kazakhstan, Uzbekistan, Mongolia, and parts of Australia⁴³ with desertification generally driven by interactions between climate change, land use and land management. Ecosystems experiencing reduced water availability are more sensitive to unsustainable land management practices like sustained heavy grazing and heavy biomass utilisation⁴³. In other regions, widespread dryland greening has been observed^{53,54}, with trends driven by large-scale increases in woody cover and a limited increase in herbaceous production at desert-grassland interfaces^{45,55,56}. Shrub encroachment and increases in woody vegetation have been recorded widely in North and central American drylands⁵⁷, the West African Sahel^{58,59}, Southern African shrublands^{60,61}, Central Asia⁶² and most tropical savannas⁶³, with changes driven by seasonal combinations of changing temperature and rainfall, CO₂ fertilization and changes in land management (e.g. decreases in large browsers, increased sustained heavy grazing and fire suppression^{42,57,64}). Increases in woody plants are sometimes accompanied by decreases in palatable grass species⁶⁵ and significantly reduced grass biomass⁶⁶. These changes have important implications for livestock productivity and livelihoods, especially where herds cannot utilise increasing shrub and tree layers. Research in grasslands and savannas in North and South America identified that for every 1 % increase in tree cover a 0.6-1.6 reduction of reproductive cows per km² occurred⁶⁷.

Changes in climate and ecosystems both directly and indirectly alter the ecosystem water balance, affecting the availability dimension of water security. Extreme droughts, which are expected to become more frequent⁶⁸, directly alter local hydrology and cause reduced plant productivity and increased tree mortality^{69,70} and reduce delivery of water-derived ecosystem services^{71,72}, including e.g. nutrient cycling, hydropower and flood control. Yet simultaneously, woody plant encroachment, invasion and afforestation alter the ecosystem water balance through increased precipitation interception⁷³ and increased evapotranspiration⁷⁴⁻⁷⁶. Increased woody cover, in combination with extreme droughts, will further reduce soil water availability⁷⁷ but can also increase carbon storage in vegetation and soils. Droughts also have considerable impacts on water quality⁷⁸, changes in which can be associated with increased salinity due to lower dilution, enhanced algal production, as well as changes in turbidity levels⁷⁸. While these observed changes demonstrate a range of impacts on different ecosystem components, changes to water regimes also have substantial impacts on people.

Droughts have had the greatest adverse impact on human populations out of any natural hazard during the 20th century⁷⁹, and in the period 2000-2019, represented 11 % of extreme climate events in Africa, but impacted 80 % of affected people (270 million)⁸⁰. The Dry Corridor in Central America (Honduras, El Salvador, Guatemala and Nicaragua) has been affected by the worst droughts in decades in the past ten years, and >1.3 million subsistence farmers and workers, most of whom are considered to be disadvantaged Indigenous People, have lost their livelihoods, experienced severe food insecurity (Integrated Phase Classification 3 or above) and have adapted their migration patterns⁸¹. In the drylands of El Salvador and Honduras, high levels of out-migration have left many women single-handedly managing farms and tending to families⁸¹. The severe droughts in this case were followed by fires, heavy rains, flooding, landslides and serious outbreaks of climate-sensitive vector-borne diseases including dengue, chikungunya and zika⁸¹. Similar disease impacts, including malaria,

diarrhoea and cholera have been experienced following increased dryland irrigation to produce food, for example, in southern Africa⁸².

Climate extremes, including droughts, have exacerbated seasonal dryland food shortages, impacted livelihoods and human resilience, and have been major contributors to food insecurity and malnutrition. This provides a useful example of how instability in water security spills over into instability of food security. Food security challenges affected around 166 million people in 26 countries in Africa and Central America who required urgent humanitarian assistance to safeguard their lives between 2015 and 2019, the majority of whom were located in drylands⁸³⁻⁸⁵. Links between droughts and malnutrition are nevertheless complex⁸⁶. Recent studies in Ethiopia, Senegal and India indicate that drought exposure can contribute to malnutrition, with malnutrition occurring not only during the drought itself, but also for several years after the event⁸⁷⁻⁸⁹. Children affected by under-nutrition during their first 1,000 days of life can experience lifelong impacts^{90,91}, leading to stunted growth, which is linked with impaired cognitive ability and reduced educational and future work performance⁹². Associated costs of stunting in terms of lost economic growth can be of the order of 10 % of GDP per year in Africa⁹³. On top of this, a lack of access to water and sanitation services causes almost 1,000 deaths among children under 5 every day, and increases the risk of numerous diseases through intake of unsafe water.

A lack of water security is not just problematic for human health and wellbeing in rural areas, but also in urban dryland areas. Box 2 provides insights into the importance of appropriate water management in relation to various dimensions of water security challenges linked to observed climate changes in Tehran, one of the world's dryland megacities.

Box 2: Water security challenges linked to observed climate changes in Tehran

Urbanisation is one of many global megatrends, as people move to both existing and emerging cities around the world. Urbanisation is taking place against the backdrop of overall population growth, particularly in sub-Saharan Africa and other tropical and Mediterranean drylands^{94,95}. Many large cities around the world, not just in the drylands, are already experiencing prolonged droughts⁹⁶, with water shortages seen recently in Rome (Italy), Chennai (India), Cape Town (South Africa), Tehran (Iran) and Sao Paulo (Brazil).

Iran is currently facing high water stress in terms of reduced water availability during the current prolonged drought, resulting from a combination of climate variability and change, alongside water mismanagement^{97,98}. This has led to water insecurity in Iran's capital, Tehran, a megacity home to around nine million people. Treated water use per capita in Tehran is 2-3 times greater than the international average, while leakage rates are high due to the use of old, poorly maintained pipes⁹⁹. The situation highlights economic water security challenges in the city, related to insufficient investment in water infrastructure. Water quality is also strongly affected, with knock-on impacts on accessibility to clean water. Pollution linked to poor soil nutrient management, inadequate wastewater treatment, and sewage, is widespread. Around 60% of water withdrawal in Tehran relies on groundwater extraction from wells, springs and qanats (gently sloping interconnected tunnels in hillsides that use gravity to deliver groundwater to lower altitude areas)¹⁰⁰. However, after this water is used, it tends to exit the city as wastewater, while investment in reuse, revision of water allocation, and assessments of water supply chains are limited^{97,99}. Tehran as a major urban centre is not the only part of the country facing water challenges. Even northern and north-western areas that are historically less prone to water insecurity have seen substantial rainfall decreases. Given that the agricultural sector uses 92% of the country's renewable water per year (an amount substantially greater than in other countries¹⁰¹), it is clear that these challenges will only be exacerbated by further climate change and that significant adaptation efforts are needed to enable a water secure future¹⁰².

Projected climate changes and impacts on water security to 2100

The accelerated warming that has been observed over drylands in recent decades is expected to continue in the future, with deserts expected to warm at a faster rate than many other terrestrial areas^{10,103}. Surface warming over drylands is projected to reach ~6.5°C under the high emissions scenario (RCP8.5) and ~3.5°C under low-moderate emissions (RCP4.5) by the end of this century, relative to the historical period (1961-1990)^{13,36}. Associated with increased temperatures, potential evapotranspiration is projected to increase in all regions globally, under all RCPs¹⁰, resulting in drier conditions and lower soil moisture in some locations, even where precipitation is unchanged.

Projected changes in precipitation are more uncertain than temperature projections, because rainfall is naturally highly variable, and the fine temporal and spatial scale of the associated physical processes pose challenges to modelling. Improvements in simulating variability of precipitation, long term trends and extremes have occurred in recent years, however, as resolution has increased parametrization of physical processes has improved¹⁰⁴. Nevertheless, the large variability in regional precipitation results in widely varying projected changes in annual precipitation over global drylands. For example, projections suggest that annual precipitation could increase by more than 40 % over central Asia and the Sahara and Sahel, but decrease by approximately 20 % over southern Africa and north-eastern South America³⁶.

In general, annual precipitation is projected to increase over most of Eurasia, tropical Africa, and the extratropical North America, but to decrease in the subtropical regions, including areas near the Mediterranean Sea, south-western North America, southern Africa, and most of Australia and South America^{37,105-107}. The picture overall though is mixed. Wet regions are generally expected to become wetter and dry regions drier with the intensification of the hydrological cycle under climate change. Drying trends may be most significant in semi-arid and arid regions¹⁰⁸⁻¹¹⁰ but recent research using the EI has shown how variable the impacts are on drylands, highlighting no overall anticipated dryland expansion¹⁵.

The frequency, severity and duration of drought conditions are expected to substantially worsen in many regions of the world¹¹¹. In a 1.5°C warmer world, historical 50-year droughts (based on the SPEI) could double across 58 % of global landmasses, an area that increases to 67 % under 2°C of warming¹¹². Multi-year drought events of magnitudes exceeding historical baselines will increase by 2050 in Australia, Brazil, Spain, Portugal, and the USA¹¹³. Declines in TWS are projected to continue, with future changes driven primarily by climate forcing rather than land and water management activities²².

The magnitude of drought stress in different regions differs depending on the metric used (see Box 1)¹¹⁴. Projections based on the PDSI suggest drought stress will increase by more than 70 % globally, while a substantially lower estimate of 37 % is found when precipitation minus evapotranspiration is used¹¹⁵. However, the two metrics agree on increasing drought stress in regions with more robust decreases in precipitation, such as southern North America, north-eastern South America and southern Europe¹¹⁵, all of which contain substantial dryland areas. An increase in drought hazards in the later quarter of the 21st century compared to the period 1971–2000 in Mediterranean, southern and eastern Africa, and southern Australia's dryland areas is indicated from the literature¹¹⁶. Nevertheless, Coupled Model Intercomparison Project (CMIP) 5 models for all types of droughts exhibit substantial differences, so confidence remains low in relation to drought projections¹¹⁷.

Modelling studies based on the AI suggest that the extent of hyperarid, arid, semiarid and dry-subhumid drylands could expand globally by 7 % by 2100 relative to 1981-2010 under a 4 °C above preindustrial warming scenario¹⁰⁴. In a 4 °C warmer world, 11.2 % of global land area is projected to shift towards drier types and 4.24 % to wetter¹⁰⁴ with the majority of newly expanded dryland areas occurring in developing countries¹³. Expansion of arid regions is likely in southwest North America, the northern fringe of Africa, parts of southern Africa and Australia^{13,104}. In contrast, India, northern China, eastern equatorial Africa and most regions south of the Sahara are projected to have shrinking drylands^{106,118-120}. The global picture nevertheless remains mixed under the EI ¹⁵.

Vegetation models which incorporate CO₂ demonstrate a similar paradox, where with warming, widespread greening with patches of browning are likely^{54,121,122}. These models project notable increases in leaf area index (LAI) and woody cover for arid grasslands, desert margins and tropical

savannas¹²² to the extent that losses of savannas of magnitudes of between 5 % (in Australia) and 55 % (South America) are expected to occur through conversion to closed canopy systems. Browning and aridification have been projected for parts of the Catinga in South America, Northern Morocco, and parts of the Namib desert⁵⁴. Projections that incorporate CO₂ most closely reflect the changes in drylands that have been observed in recent decades.

Where areas are likely to experience increased aridification, the warming and drying can reduce soil organic carbon (SOC) storage¹³. Soil degradation and reduced soil moisture also substantially limit gross primary productivity and affect the rate of photosynthesis, which absorbs CO₂ and stores carbon. In combination with land degradation, which also contributes to greenhouse gas emissions, this is a positive feedback cycle, with the warming and drying reinforcing each other, and dryland soils storing less carbon and emitting more CO₂ into the atmosphere, exacerbating global warming. Nevertheless, dryland areas in which greening is occurring can experience an increase in soil carbon¹²³ and reduced freshwater availability as the new vegetation uses water that would previously have run off¹²⁴.

Projections show that the number of people residing in drylands exposed to water stress, drought intensity and habitat degradation will reach ~974 million, ~1267 million and ~1285 million people, respectively, by 2050, at 1.5 °C, 2 °C and 3 °C of global warming under the Shared Socio-economic Pathway 2 (SSP2; business as usual) scenario²⁴. At the same time, projected climate change impacts are far exceeded by the hydrological effects of past and present water extraction in many semi-arid continental river systems¹²⁵. This suggests there are important opportunities that could emerge from improved water governance in drylands into the future¹²⁶.

Limited access to water, land and livestock, low agriculture productivity and increases in food prices and household food insecurity are among many factors leading to migration and displacement: 'survival migration'⁸⁵. However, the literature remains inconclusive on the direct attribution of migration to water insecurity and how this may develop under future climate change scenarios, given myriad contextual factors linked to socio-economic, institutional, cultural and political aspects²⁵. Projections tend to focus on modelling exposure to risk, rather than quantifying how many people will actually migrate as a response to that exposure^{127,128}. Food and water insecurity, combined with poverty may potentially increase the likelihood and intensity of armed conflict in some dryland contexts¹²⁹, although agricultural output and violent conflict tend to be only weakly and inconsistently connected¹³⁰. For example, in the Sahel, droughts and water insecurity have only played a minor role compared to politics and governance in explaining the conflicts that have emerged in the last few decades^{131,132}.

Ensuring effective and sustainable access to water and sanitation systems is a challenge, particularly because weather extremes such as floods, storms, heatwaves, and droughts are becoming more common in drylands under climate change. Displaced populations, including those in conflict regions, often have limited access to safe drinking water and/or sanitation and are at increased risk of infectious disease outbreaks such as measles or cholera, which frequently cannot be prevented, treated or controlled, due to lack of access to health and/or preventive services such as micronutrient supplementation and immunization⁸³. It is anticipated that without intervention, these aspects will worsen into the future. In addition, while water insecurity and physical human health links are well established, there is little understanding of the relationship between water and mental health¹³³, including in drylands, where economic impacts of droughts have been associated with increases in suicide, particularly among farmers in India¹³⁴ and Australia¹³⁵.

Challenges and opportunities for water security in drylands

Proactive strategies are needed to plan for global change issues, including planning for a water secure future¹³⁶. Drylands offer a long history of adaptation to water constraints and may offer useful insights

and lessons for other locations that will experience water scarcity. Water conservation and redistribution efforts in drylands have attempted to improve the location, timing and quality of water to support water security, either by re-allocating the water itself, e.g. through investments in water infrastructure, or relocating the human and animal populations that depend on it. For example, in Somalia, Ethiopia, and other parts of East Africa, herders have traditionally moved with their livestock to graze different areas depending on water and pasture availability. While this movement still happens today, these kinds of traditional practices are becoming more difficult due to a range of factors including sedentarisation, conflict, rural to urban migration and the breakdown of traditional institutions and property rights¹³⁷. Similar challenges emerge from Botswana's Kalahari where decades of policies supporting sedentarisation have led to increased borehole drilling and widespread pasture degradation^{138,139}. Geopolitics also affect mobility. In dryland Central Asia, pastoralists used to cover extensive distances with their herds, but with the dissolution of the Union of Soviet Socialist Republics (USSR), movement across what then became international borders became more difficult¹⁴⁰. These kinds of challenges most starkly and disproportionately impact societal groups who directly depend upon the natural resource base for their livelihoods.

Alleviating economic water scarcity by increasing access to water through installation of large-scale irrigation systems, can deliver short-term benefits for agricultural productivity in dry areas. However, in the context of climate change, increasing demand, and inefficient management of water and associated infrastructure, can result in complex outcomes over the longer term. For example, irrigation systems once operating in the Aral Sea region have resulted in large-scale landscape degradation and downstream water shortages, threatening livelihoods and human health, as levels of inflows into the Aral sea from the Amu Darya and Syr Darya rivers decreased, and infrastructure became outdated and inefficient¹⁴¹. Irrigation has also been shown to affect local climate conditions. While cooling effects have been observed, and are especially pronounced for parts of south Asia^{142,143}, there is also the potential for irrigation to increase heat stress, despite a cooler surface¹⁴³.

The Global Water Partnership (GWP) recognises the interlinked nature of the various dimensions of water security, and strongly supports the integration of water concerns across sectors through approaches such as Integrated Water Resources Management (IWRM). IWRM is defined as "a process which promotes the coordinated development and management of water, land and related resources in order to maximise economic and social welfare in an equitable manner without compromising the sustainability of vital ecosystems and the environment"¹⁴⁴. In drylands and other areas where water is a key limiting factor, such coherent approaches can be vital in managing water, energy, and food (WEF) security, as these challenges often compete for resources and funding¹⁴⁵. Considering WEF as a nexus provides a framework to analyse these components as an integrated system involving multiple research disciplines¹³⁶, and can be used to support decision-making and the assessment of adaptation options^{145,146}.

Often, policies lack sufficient evaluation of trade-offs: temporally and spatially, as well as across different sectors and societal groups^{9,146}. This is particularly apparent with large-scale engineering megaprojects seeking to increase water access, which reconfigure desert landscapes (e.g. the Great Manmade River (Libya), the South-to-North Water Transfer Scheme (China), the Central Arizona Project (United States) and the Greater Anatolia Project (Turkey)¹⁴⁷). Efforts to solve problems originating in other sectors have important impacts on water security across the landscape too. For example, large-scale initiatives to support dryland greening through afforestation programmes (e.g. in northwest China¹⁴⁸) aimed to reduce land degradation and restore watershed ecosystem services, support climate change mitigation and adaptation as well as address poverty challenges. However, these endeavours, particularly in drylands, can lead to trade-offs where increases in above-ground carbon gain¹⁴⁹ can decrease biodiversity and water availability¹⁵⁰. In arid drylands where afforestation of previously non-forested ecosystems is likely, the potential increase in above-ground carbon is

balanced against increased water interception and evapotranspiration and a decrease in run-off and groundwater recharge, which can result in local and regional water shortages^{75,151,152}.

Interactions between climate change and land use change have resulted in extensive dryland regions experiencing significant woody plant encroachment and alien plant invasion^{63,64,153}. Increased native woody cover also causes higher evapotranspiration and more water to be removed from the soil, especially from deeper horizons¹⁵⁴. These processes significantly reduce water resources and further exacerbate drought effects, even though carbon storage may be increased⁷³. A key adaptation that has been used to improve water security in this context is large-scale selective woody plant clearing in dryland water catchments and riparian zones. The Working For Water programme in South Africa is clearing invasive shrubs to improve water supplies and enhance other ecosystem services (e.g. increase grazer forage availability), while providing employment to local communities¹⁵⁵. Current estimates indicate that targeted clearing of encroaching shrubs and alien invasives has restored ecosystem services valued at US\$8 billion with the largest benefits coming from increased water resources, timber products, woody fuels and improved grazing¹⁵⁶. However, studies valuing encroaching bushes and the ecosystem services they provide, generally remain limited⁶⁵.

Along with large-scale infrastructure and landscape adaptations, a wealth of small-scale water harvesting and soil and water conservation measures are used in drylands in the pursuit of water security¹⁵⁷, including crop diversification, switching to more drought-resistant crops and varieties, and adopting conservation agriculture practices which improve the water holding capacity of the soil by increasing organic matter and soil organic carbon¹⁰. In some areas, locally important methods are based on traditional knowledge, however, many of these methods are becoming less widespread. For example, traditional irrigation in Iran over the past 2,500 years has used qanats, but as western solutions have proliferated, only around half of the estimated 72,000 qanats remain in use¹⁵⁸. Another example comes from the pastoral Borana in southern Ethiopia, where the *tula* well system dates back five centuries and has played a critical role in the sustainable management of pastures, in shaping cultural identity, and the organization of water management¹⁵⁹. Wells are dug down to deep water aquifers where water is brought to the surface manually. Human labour is regularly demanded for maintenance of the wells and the capacity to organize workers for re-excavation and repair is crucial for the sustainability of this pastoral system¹⁵⁹. More recently, hired labour has started to replace clan-based labour and payments for well rehabilitation have changed from cattle to cash. In addition, plastic buckets are replacing leather buckets, and mechanization also replaces human labour. It is uncertain how these transformations will affect the sustainability of the Borana water management system¹⁶⁰.

While traditional efforts to improve water security can deliver useful local benefits, they are often difficult to scale up and out to other locations. Development status matters when it comes to upscaling. In parallel with measures for increasing the supply of available water through re-use of wastewater and sea water desalination, which will become increasingly necessary when faced with growing water scarcity¹⁶¹ an essential adaptation is to increase water use efficiency in irrigated agriculture through wider application of well-established technologies such as drip and sprinkler irrigation¹⁶². In many rainfed areas in drylands of sub-Saharan Africa and Central Asia, agricultural economic water scarcity is a major concern. Whereas successful adaptation to climate change in these areas may require expansion of irrigation, lack of investment into irrigation infrastructure and in some instances, limited institutional capacities to effectively manage expanded irrigation, hinders climate change adaptation opportunities¹⁶³. In more advanced dryland economies (e.g. Middle Eastern countries such as Israel, Saudi Arabia etc; Australia; USA) deployment of technologies such as desalination plants and precision irrigation systems is facilitated by the institutional set-ups, economic incentives (subsidies and/or credit systems), political will and greater levels of state investment than is feasible in low income economies. Water processed at many desalination plants,

e.g. in the Persian Gulf, is largely for industrial use, although opportunities exist for improved agricultural use¹⁰¹. Nevertheless, use of these technologies can increase inequalities both within and between nations, and could lead to increased tensions. The reuse of marginal quality waters can also increase soil salinity and lead to negative impacts on agricultural productivity and human health¹⁶⁴, so proper management of water quality and more effective wastewater treatment becomes critical. A major problem from the water quality dimension of water security is that not only in practice, but also in hydro-economic modelling and planning, differentiated economic valuation of different quality waters is still absent from decision processes.

Overall, governance design and implementation deficiencies need urgent attention to better manage water-related challenges in drylands. Water governance benchmarking to monitor progress is essential, with research from Egypt, Jordan, Morocco, Oman, Turkey and Yemen providing useful insights¹⁶⁵. Climate-sensitive political leadership is also paramount, especially where transboundary water resources are concerned and political water security challenges are present¹⁶⁶. Ensuring dryland water security into the future requires governance structures and processes that recognise interlinkages and which support coherent policies that take into account climate change impacts¹⁶⁷ and the nexus between sectors¹⁶⁸. Such approaches need to carefully balance the short and long term, as well as engage the necessary stakeholders at different levels to enable an equitable approach¹⁶⁹.

Conclusion

The challenges associated with reaching and maintaining water security in the world's drylands are set to become even more difficult, given the plethora of observed climate impacts in drylands to date, and climate projections for the future. Profound ecosystem changes are already being observed and experienced in drylands, alongside impacts on human systems that limit the access of vast numbers of people to sufficient clean and safe water supplies, including those living in rapidly expanding dryland cities. While current adaptation strategies are varied and offer a useful starting point, including those grounded in traditional knowledge and practice, the changes to dryland areas anticipated to 2100 exceed those that have been experienced previously, suggesting new approaches are needed.

IWRM sets out a useful framework that is applicable to many environments, but is urgently required in drylands. Figure 5 establishes the core components of an integrated approach to water resource management in drylands under a changing climate, emphasising the need to manage demand, optimise supply, provide equitable access to water resources, as well as establish improved and integrated policy, regulatory and institutional frameworks, to tackle all the dimensions of water security.

Ostensibly, achieving water security is less of an environmental challenge (availability dimension) and more of a governance issue (access, quality and stability dimensions), requiring political will, capacity, resourcing and leadership in the development of a truly integrated and coherent approach to deliver water-related decisions that also align with the needs of other sectors in drylands. Impacts of a lack of water security on food, health, energy, livelihoods, migration and conflict demonstrate the crucial role of water as a connector. Technological fixes alone will be insufficient and could exacerbate inequalities. Stakeholder engagement is becoming increasingly important, particularly in complex contexts where dryland rivers flow through multiple national boundaries, highlighting the importance of IWRM in shaping more equitable water resource allocation in a transboundary context, as well as in multi-sector SDG contexts.

Adaptations that enhance water security in drylands offer the potential to inform solutions that can be shared with other (dryland and non-dryland) locations, such that climate change impacts in areas that will only just be starting to experience increased water security challenges in the future, can

benefit. Drylands represent useful climate analogues in this sense but require increased research activity to improve decision making on water management and water security. Improved knowledge is needed to better understand integrated risks, and to assess costs, benefits and trade-offs associated with different water-linked adaptations, across different societal groups and time frames, as well as across different aspects of the environmental system. Such efforts offer considerable scope to accelerate progress towards multiple SDGs, both in the lead up to 2030, and beyond.

Declaration of interests

The authors declare no competing interests

Acknowledgement

Thank you to Pierre Hiernaux for useful feedback and to the journal reviewers for their thoughtful comments.

References cited

1. Maestre, F.T., Quero, J.L., Gotelli, N.J., Escudero, A., Ochoa, V., Delgado-Baquerizo, M., García-Gómez, M., Bowker, M.A., Soliveres, S., Escolar, C., et al. (2012). Plant Species Richness and Ecosystem Multifunctionality in Global Drylands. *Science* 335, 214. [10.1126/science.1215442](https://doi.org/10.1126/science.1215442).
2. Teff-Seker, Y., and Orenstein, D.E. (2019). The 'desert experience': Evaluating the cultural ecosystem services of drylands through walking and focusing. *People and Nature* 1, 234-248. <https://doi.org/10.1002/pan3.28>.
3. Richards, J., Mayaud, J., Zhan, H., Wu, F., Bailey, R., and Viles, H. (2020). Modelling the risk of deterioration at earthen heritage sites in drylands. *Earth Surface Processes and Landforms* 45, 2401-2416. <https://doi.org/10.1002/esp.4887>.
4. van der Esch, S., ten Brink, B., Stehfest, E., Bakkenes, M., Sewell, A., Bouwman, A., Meijer, J., Westhoek, H., van den Berg, M., van den Born, G.J., et al. (2017). Exploring Future Changes in Land Use and Land Condition and the Impacts on Food, Water, Climate Change and Biodiversity: Scenarios for the UNCCD Global Land Outlook. Policy Report. PBL Netherlands Environmental Assessment Agency.
5. Center for International Earth Science Information Network (CIESIN), Columbia University. (2018). Documentation for the Gridded Population of the World, Version 4 (GPWv4), Revision 11 Data Sets. Palisades NY: NASA Socioeconomic Data and Applications Center (SEDAC).
6. Millennium Ecosystem Assessment (2005). MA Ecosystems. Palisades, NY: NASA Socioeconomic Data and Applications Center (SEDAC). <http://dx.doi.org/10.7927/H4KW5CZ6>.
7. Cherlet, M., Hutchinson, C., Reynolds, J., Hill, J., Sommer, S., von Maltitz, G (2018). World Atlas of desertification: rethinking land degradation and sustainable land management. In M. Cherlet, C. Hutchinson, J. Reynolds, J. Hill, S. Sommer, G. Von Maltitz, and European Union. Publications, eds. Publications Office of the European Union.
8. Plaza, C., Zaccone, C., Sawicka, K., Méndez, A.M., Tarquis, A., Gascó, G., Heuvelink, G.B.M., Schuur, E.A.G., and Maestre, F.T. (2018). Soil resources and element stocks in drylands to face global issues. *Scientific Reports* 8, 13788. [10.1038/s41598-018-32229-0](https://doi.org/10.1038/s41598-018-32229-0).
9. Stringer, L.C., Reed, M.S., Fleskens, L., Thomas, R.J., Le, Q.B., and Lala-Pritchard, T. (2017). A New Dryland Development Paradigm Grounded in Empirical Analysis of Dryland Systems Science. *Land Degradation & Development* 28, 1952-1961. <https://doi.org/10.1002/ldr.2716>.
10. Mirzabaev, A., Wu, J., Evans, J., F. García-Oliva, I.A.G. Hussein, M.H. Iqbal, J. Kimutai, T. Knowles, F. Meza, D. Nedjraoui, et al. (2019). Desertification. In *Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems*, P.R. Shukla, J. Skea, E. Calvo Buendia, V. Masson-Delmotte, H.-O. Pörtner, D.C. Roberts, P. Zhai, R. Slade, S. Connors, R. van Diemen, et al., eds. (In press).

11. UNEP. (1992). World Atlas of Desertification. United Nations Environment Programme, Nairobi, Kenya.
12. Spinoni, J., Vogt, J., Naumann, G., Carrao, H., and Barbosa, P. (2015). Towards identifying areas at climatological risk of desertification using the Köppen–Geiger classification and FAO aridity index. *International Journal of Climatology* 35, 2210-2222. <https://doi.org/10.1002/joc.4124>.
13. Huang, J., Yu, H., Guan, X., Wang, G., and Guo, R. (2016). Accelerated dryland expansion under climate change. *Nature Climate Change* 6, 166-171. 10.1038/nclimate2837.
14. He, B., Wang, S., Guo, L., and Wu, X. (2019). Aridity change and its correlation with greening over drylands. *Agricultural and Forest Meteorology* 278, 107663. <https://doi.org/10.1016/j.agrformet.2019.107663>.
15. Berg, A., and McColl, K.A. (2021). No projected global drylands expansion under greenhouse warming. *Nature Climate Change*. 10.1038/s41558-021-01007-8.
16. Fensholt, R., Langanke, T., Rasmussen, K., Reenberg, A., Prince, S.D., Tucker, C., Scholes, R.J., Le, Q.B., Bondeau, A., Eastman, R., et al. (2012). Greenness in semi-arid areas across the globe 1981–2007 — an Earth Observing Satellite based analysis of trends and drivers. *Remote Sensing of Environment* 121, 144-158. <https://doi.org/10.1016/j.rse.2012.01.017>.
17. Andela, N., Liu, Y., van Dijk, A.I.J.M., de Jeu, R.A.M., and McVicar, T. (2013). Global changes in dryland vegetation dynamics (1988-2008) assessed by satellite remote sensing: comparing a new passive microwave vegetation density record with reflective greenness data. *Biogeosciences* 10, 6657 - 6676. <https://doi.org/10.5194/bg-10-6657-2013>
18. Yang, Y., Roderick, M.L., Zhang, S., McVicar, T.R., and Donohue, R.J. (2019). Hydrologic implications of vegetation response to elevated CO₂ in climate projections. *Nature Climate Change* 9, 44-48. 10.1038/s41558-018-0361-0.
19. Zhu, Z., Piao, S., Myneni, R.B., Huang, M., Zeng, Z., Canadell, J.G., Ciais, P., Sitch, S., Friedlingstein, P., Arneeth, A., et al. (2016). Greening of the Earth and its drivers. *Nature Climate Change* 6, 791-795. 10.1038/nclimate3004.
20. Palmer, W.C. (1968). Keeping Track of Crop Moisture Conditions, Nationwide: The New Crop Moisture Index. *Weatherwise* 21, 156-161. 10.1080/00431672.1968.9932814.
21. Dai, A. (2011). Characteristics and trends in various forms of the Palmer Drought Severity Index during 1900–2008. *Journal of Geophysical Research: Atmospheres* 116. <https://doi.org/10.1029/2010JD015541>.
22. Pokhrel, Y., Felfelani, F., Satoh, Y., Boulange, J., Burek, P., Gädeke, A., Gerten, D., Gosling, S.N., Grillakis, M., Gudmundsson, L., et al. (2021). Global terrestrial water storage and drought severity under climate change. *Nature Climate Change* 11, 226-233. 10.1038/s41558-020-00972-w.
23. Yao, Y., Tian, Y., Andrews, C., Li, X., Zheng, Y., and Zheng, C. (2018). Role of Groundwater in the Dryland Ecohydrological System: A Case Study of the Heihe River Basin. *Journal of Geophysical Research: Atmospheres* 123, 6760-6776. <https://doi.org/10.1029/2018JD028432>.
24. Byers, E., Gidden, M., Leclère, D., Balkovic, J., Burek, P., Ebi, K., Greve, P., Grey, D., Havlik, P., Hillers, A., et al. (2018). Global exposure and vulnerability to multi-sector development and climate change hotspots. *Environmental Research Letters* 13, 055012. 10.1088/1748-9326/aabf45.
25. Pradhan, P., Costa, L., Rybski, D., Lucht, W., and Kropp, J.P. (2017). A Systematic Study of Sustainable Development Goal (SDG) Interactions. *Earth's Future* 5, 1169-1179. <https://doi.org/10.1002/2017EF000632>.
26. Harris, L., Kleiber, D., Goldin, J., Darkwah, A., and Morinville, C. (2017). Intersections of gender and water: comparative approaches to everyday gendered negotiations of water access in underserved areas of Accra, Ghana and Cape Town, South Africa. *Journal of Gender Studies* 26, 561-582. 10.1080/09589236.2016.1150819.

27. Berdugo, M., Delgado-Baquerizo, M., Soliveres, S., Hernández-Clemente, R., Zhao, Y., Gaitán, J.J., Gross, N., Saiz, H., Maire, V., Lehmann, A., et al. (2020). Global ecosystem thresholds driven by aridity. *Science* 367, 787. [10.1126/science.aay5958](https://doi.org/10.1126/science.aay5958).
28. Gain, A.K., Giupponi, C., and Wada, Y. (2016). Measuring global water security towards sustainable development goals. *Environmental Research Letters* 11, 124015. [10.1088/1748-9326/11/12/124015](https://doi.org/10.1088/1748-9326/11/12/124015).
29. Cook, C., and Bakker, K. (2012). Water security: Debating an emerging paradigm. *Global Environmental Change* 22, 94-102. <https://doi.org/10.1016/j.gloenvcha.2011.10.011>.
30. AQUASTAT/FAO (2020). AQUASTAT/FAO database. In Food and Agriculture Organisation, ed. Rome.
31. Falkenmark, M., Lundqvist, J., and Widstrand, C. (1989). Macro-scale water scarcity requires micro-scale approaches. *Natural Resources Forum* 13, 258-267. <https://doi.org/10.1111/j.1477-8947.1989.tb00348.x>.
32. Damania, R.A., Desbureaux, S., Rodella, A.S., Russ, J., Zaveri, E. (2019). Quality Unknown: The Invisible Water Crisis *10.1596/978-1-4648-1459-4*.
33. Lissner, T.K., Sullivan, C.A., Reusser, D.E., and Kropp, J.P. (2014). Determining regional limits and sectoral constraints for water use. *Hydrol. Earth Syst. Sci.* 18, 4039-4052. [10.5194/hess-18-4039-2014](https://doi.org/10.5194/hess-18-4039-2014).
34. Allouche, J., Nicol, A., and Mehta, L. (2011). Water security: Towards the human securitization of water? *Journal of Diplomacy and International Relations* XII, 153-171.
35. Mehta, L. (2003). Contexts and Constructions of Water Scarcity. *Economic and Political Weekly* 38, 5066-5072.
36. Huang, J., Li, Y., Fu, C., Chen, F., Fu, Q., Dai, A., Shinoda, M., Ma, Z., Guo, W., Li, Z., et al. (2017). Dryland climate change: Recent progress and challenges. *Reviews of Geophysics* 55, 719-778. <https://doi.org/10.1002/2016RG000550>.
37. Zhao, T., and Dai, A. (2015). The Magnitude and Causes of Global Drought Changes in the Twenty-First Century under a Low-Moderate Emissions Scenario. *Journal of Climate* 28, 4490-4512. [10.1175/jcli-d-14-00363.1](https://doi.org/10.1175/jcli-d-14-00363.1).
38. Chang, L.-L., Yuan, R., Gupta, H.V., Winter, C.L., and Niu, G.-Y. (2020). Why Is the Terrestrial Water Storage in Dryland Regions Declining? A Perspective Based on Gravity Recovery and Climate Experiment Satellite Observations and Noah Land Surface Model With Multiparameterization Schemes Model Simulations. *Water Resources Research* 56, e2020WR027102. <https://doi.org/10.1029/2020WR027102>.
39. Ji, M., Huang, J., Xie, Y., and Liu, J. (2015). Comparison of dryland climate change in observations and CMIP5 simulations. *Advances in Atmospheric Sciences* 32, 1565-1574. [10.1007/s00376-015-4267-8](https://doi.org/10.1007/s00376-015-4267-8).
40. Huang, J., Ji, M., Xie, Y., Wang, S., He, Y., and Ran, J. (2016). Global semi-arid climate change over last 60 years. *Climate Dynamics* 46, 1131-1150. [10.1007/s00382-015-2636-8](https://doi.org/10.1007/s00382-015-2636-8).
41. Právělie, R., Bandoc, G., Patriche, C., and Sternberg, T. (2019). Recent changes in global drylands: Evidences from two major aridity databases. *Catena* 178, 209-231. <https://doi.org/10.1016/j.catena.2019.03.016>.
42. Donohue, R.J., Roderick, M.L., McVicar, T.R., and Farquhar, G.D. (2013). Impact of CO2 fertilization on maximum foliage cover across the globe's warm, arid environments. *Geophysical Research Letters* 40, 3031-3035. <https://doi.org/10.1002/grl.50563>.
43. Burrell, A.L., Evans, J.P., and De Kauwe, M.G. (2020). Anthropogenic climate change has driven over 5 million km2 of drylands towards desertification. *Nature Communications* 11, 3853. [10.1038/s41467-020-17710-7](https://doi.org/10.1038/s41467-020-17710-7).
44. le Polain de Waroux, Y., and Lambin, E.F. (2012). Monitoring degradation in arid and semi-arid forests and woodlands: The case of the argan woodlands (Morocco). *Applied Geography* 32, 777-786. <https://doi.org/10.1016/j.apgeog.2011.08.005>.

45. Zhang, W., Brandt, M., Penuelas, J., Guichard, F., Tong, X., Tian, F., and Fensholt, R. (2019). Ecosystem structural changes controlled by altered rainfall climatology in tropical savannas. *Nature Communications* 10, 671. [10.1038/s41467-019-08602-6](https://doi.org/10.1038/s41467-019-08602-6).
46. Barbosa, H.A., Kumar, T.V.L., and Silva, L.R.M. (2015). Recent trends in vegetation dynamics in the South America and their relationship to rainfall. *Natural Hazards* 77, 883-899. [10.1007/s11069-015-1635-8](https://doi.org/10.1007/s11069-015-1635-8).
47. Becerril-Piña, R., Mastachi-Loza, C.A., González-Sosa, E., Díaz-Delgado, C., and Bâ, K.M. (2015). Assessing desertification risk in the semi-arid highlands of central Mexico. *Journal of Arid Environments* 120, 4-13. <https://doi.org/10.1016/j.jaridenv.2015.04.006>.
48. Jiang, L., Bao, A., Jiapaer, G., Guo, H., Zheng, G., Gafforov, K., Kurban, A., and De Maeyer, P. (2019). Monitoring land sensitivity to desertification in Central Asia: Convergence or divergence? *Science of The Total Environment* 658, 669-683. <https://doi.org/10.1016/j.scitotenv.2018.12.152>.
49. Aragón-Gastélum, J.L., Flores, J., Yáñez-Espinosa, L., Badano, E., Ramírez-Tobías, H.M., Rodas-Ortiz, J.P., and González-Salvatierra, C. (2014). Induced climate change impairs photosynthetic performance in *Echinocactus platyacanthus*, an especially protected Mexican cactus species. *Flora - Morphology, Distribution, Functional Ecology of Plants* 209, 499-503. <https://doi.org/10.1016/j.flora.2014.06.002>.
50. Musil, C., Schmiedel, U., and Midgley, G. (2005). Lethal effects of experimental warming approximating a future climate scenario on southern African quartz-field succulents: A pilot study. *The New phytologist* 165, 539-547. [10.1111/j.1469-8137.2004.01243.x](https://doi.org/10.1111/j.1469-8137.2004.01243.x).
51. Allen, C.D., Breshears, D.D., and McDowell, N.G. (2015). On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* 6, art129. <https://doi.org/10.1890/ES15-00203.1>.
52. Slingsby, J.A., Merow, C., Aiello-Lammens, M., Allsopp, N., Hall, S., Kilroy Mollmann, H., Turner, R., Wilson, A.M., and Silander, J.A. (2017). Intensifying postfire weather and biological invasion drive species loss in a Mediterranean-type biodiversity hotspot. *Proceedings of the National Academy of Sciences* 114, 4697. [10.1073/pnas.1619014114](https://doi.org/10.1073/pnas.1619014114).
53. Lu, X., Wang, L., and McCabe, M. (2016). Elevated CO₂ as a driver of global dryland greening. *Scientific Reports* 6, 20716. [10.1038/srep20716](https://doi.org/10.1038/srep20716).
54. Piao, S., Wang, X., Park, T., Chen, C., Lian, X., He, Y., Bjerke, J.W., Chen, A., Ciais, P., Tømmervik, H., et al. (2020). Characteristics, drivers and feedbacks of global greening. *Nature Reviews Earth & Environment* 1, 14-27. [10.1038/s43017-019-0001-x](https://doi.org/10.1038/s43017-019-0001-x).
55. Collins, S.L., and Xia, Y. (2015). Long-Term Dynamics and Hotspots of Change in a Desert Grassland Plant Community. *The American Naturalist* 185, E30-E43. [10.1086/679315](https://doi.org/10.1086/679315).
56. Masubelele, M.L., Hoffman, M.T., Bond, W.J., and Gambiza, J. (2014). A 50 year study shows grass cover has increased in shrublands of semi-arid South Africa. *Journal of Arid Environments* 104, 43-51. <https://doi.org/10.1016/j.jaridenv.2014.01.011>.
57. Archer, S.R., Andersen, E.M., Predick, K.I., Schwinning, S., Steidl, R.J., and Woods, S.R. (2017). Woody Plant Encroachment: Causes and Consequences. In *Rangeland Systems: Processes, Management and Challenges*, D.D. Briske, ed. (Springer International Publishing), pp. 25-84. [10.1007/978-3-319-46709-2_2](https://doi.org/10.1007/978-3-319-46709-2_2).
58. Benjaminsen, T.A., and Hiernaux, P. (2019). From Desiccation to Global Climate Change: A History of the Desertification Narrative in the West African Sahel, 1900-2018. *Global Environment* 12, 206-236. [10.3197/ge.2019.120109](https://doi.org/10.3197/ge.2019.120109).
59. Brandt, M., Hiernaux, P., Rasmussen, K., Tucker, C.J., Wigneron, J.-P., Diouf, A.A., Herrmann, S.M., Zhang, W., Kergoat, L., Mbow, C., et al. (2019). Changes in rainfall distribution promote woody foliage production in the Sahel. *Communications Biology* 2, 133. [10.1038/s42003-019-0383-9](https://doi.org/10.1038/s42003-019-0383-9).
60. Rohde, R.F., Hoffman, M.T., Durbach, I., Venter, Z., and Jack, S. (2019). Vegetation and climate change in the Pro-Namib and Namib Desert based on repeat photography: Insights into

- climate trends. *Journal of Arid Environments* 165, 119-131. <https://doi.org/10.1016/j.jaridenv.2019.01.007>.
61. Timm Hoffman, M., Rohde, R.F., and Gillson, L. (2019). Rethinking catastrophe? Historical trajectories and modelled future vegetation change in southern Africa. *Anthropocene* 25, 100189. <https://doi.org/10.1016/j.ancene.2018.12.003>.
 62. Li, Z., Chen, Y., Li, W., Deng, H., and Fang, G. (2015). Potential impacts of climate change on vegetation dynamics in Central Asia. *Journal of Geophysical Research: Atmospheres* 120, 12345-12356. <https://doi.org/10.1002/2015JD023618>.
 63. Stevens, N., Lehmann, C.E.R., Murphy, B.P., and Durigan, G. (2017). Savanna woody encroachment is widespread across three continents. *Global Change Biology* 23, 235-244. <https://doi.org/10.1111/gcb.13409>.
 64. Venter, Z.S., Cramer, M.D., and Hawkins, H.J. (2018). Drivers of woody plant encroachment over Africa. *Nature Communications* 9, 2272. 10.1038/s41467-018-04616-8.
 65. Reed, M.S., Stringer, L.C., Dougill, A.J., Perkins, J.S., Atlhopheng, J.R., Mulale, K., and Favretto, N. (2015). Reorienting land degradation towards sustainable land management: Linking sustainable livelihoods with ecosystem services in rangeland systems. *Journal of Environmental Management* 151, 472-485. <https://doi.org/10.1016/j.jenvman.2014.11.010>.
 66. Scholes, R.J. (2003). Convex Relationships in Ecosystems Containing Mixtures of Trees and Grass. *Environmental and Resource Economics* 26, 559-574. 10.1023/B:EARE.0000007349.67564.b3.
 67. Anadon, J.D., Sala, O.E., Turner, B.L., and Bennett, E.M. (2014). Effect of woody-plant encroachment on livestock production in North and South America. *Proceedings of the National Academy of Sciences* 111, 12948-12953. 10.1073/pnas.1320585111.
 68. Bradford, J.B., Schlaepfer, D.R., Lauenroth, W.K., and Palmquist, K.A. (2020). Robust ecological drought projections for drylands in the 21st century. *Global Change Biology* 26, 3906-3919. <https://doi.org/10.1111/gcb.15075>.
 69. Bodner, G.S., and Robles, M.D. (2017). Enduring a decade of drought: Patterns and drivers of vegetation change in a semi-arid grassland. *Journal of Arid Environments* 136, 1-14. <https://doi.org/10.1016/j.jaridenv.2016.09.002>.
 70. Choat, B., Brodribb, T.J., Brodersen, C.R., Duursma, R.A., López, R., and Medlyn, B.E. (2018). Triggers of tree mortality under drought. *Nature* 558, 531-539. 10.1038/s41586-018-0240-x.
 71. Banerjee, O., Bark, R., Connor, J., and Crossman, N.D. (2013). An ecosystem services approach to estimating economic losses associated with drought. *Ecological Economics* 91, 19-27. <https://doi.org/10.1016/j.ecolecon.2013.03.022>.
 72. Heitschmidt, R.K., Klement, K.D., and Haferkamp, M.R. (2005). Interactive Effects of Drought and Grazing on Northern Great Plains Rangelands. Society for Range Management.
 73. Honda, E.A., and Durigan, G. (2016). Woody encroachment and its consequences on hydrological processes in the savannah. *Philosophical Transactions of the Royal Society B: Biological Sciences* 371, 20150313. doi:10.1098/rstb.2015.0313.
 74. Nosetto, M.D., Jobbágy, E.G., Brizuela, A.B., and Jackson, R.B. (2012). The hydrologic consequences of land cover change in central Argentina. *Agriculture, Ecosystems & Environment* 154, 2-11. <https://doi.org/10.1016/j.agee.2011.01.008>.
 75. Schwärzel, K., Zhang, L., Montanarella, L., Wang, Y., and Sun, G. (2020). How afforestation affects the water cycle in drylands: A process-based comparative analysis. *Global Change Biology* 26, 944-959. <https://doi.org/10.1111/gcb.14875>.
 76. Villegas, J.C., Dominguez, F., Barron-Gafford, G.A., Adams, H.D., Guardiola-Claramonte, M., Sommer, E.D., Selvey, A.W., Espeleta, J.F., Zou, C.B., Breshears, D.D., and Huxman, T.E. (2015). Sensitivity of regional evapotranspiration partitioning to variation in woody plant cover: insights from experimental dryland tree mosaics. *Global Ecology and Biogeography* 24, 1040-1048. <https://doi.org/10.1111/geb.12349>.

77. Caldeira, M.C., Lecomte, X., David, T.S., Pinto, J.G., Bugalho, M.N., and Werner, C. (2015). Synergy of extreme drought and shrub invasion reduce ecosystem functioning and resilience in water-limited climates. *Scientific reports* 5, 15110-15110. 10.1038/srep15110.
78. Mosley, L.M. (2015). Drought impacts on the water quality of freshwater systems; review and integration. *Earth-Science Reviews* 140, 203-214. <https://doi.org/10.1016/j.earscirev.2014.11.010>.
79. Mishra, A.K., and Singh, V.P. (2010). A review of drought concepts. *Journal of Hydrology* 391, 202-216. <https://doi.org/10.1016/j.jhydrol.2010.07.012>.
80. Yaghmaei, N., and Below, R. (2019). Issue No. 56 CRED Crunch Disasters in Africa: 20 Year Review (2000-2019).
81. OCHA (2019). OCHA Annual Report 2019. United Nations Office for the Coordination of Humanitarian Affairs. <https://www.unocha.org/sites/unocha/files/2019OCHAAnnualreport.pdf>.
82. Mabhaudhi, T.N., L.; Mpandeli, S.; Nhemachena, C.; Senzanje, A.; Sobratee, N.; Chivenge, P.P.; Slotow, R.; Naidoo, D.; Liphadzi, S.; Modi, A.T. (2019). The Water–Energy–Food Nexus as a Tool to Transform Rural Livelihoods and Well-Being in Southern Africa. *Int. J. Environ. Res. Public Health* 16, 2970. 10.3390/ijerph16162970.
83. FSIN, and GNAFC (2020). Global Report on Food Crises 2020
84. FSIN (2017). Global Report on Food Crises 2016. World Food Programme.
85. FAO/IFAD/UNICEF/WFP/WHO (2018). The State of Food Security and Nutrition in the World 2018. Building climate resilience for food security and nutrition.
86. Belesova, K., Agabiirwe, C.N., Zou, M., Phalkey, R., and Wilkinson, P. (2019). Drought exposure as a risk factor for child undernutrition in low- and middle-income countries: A systematic review and assessment of empirical evidence. *Environment International* 131, 104973. <https://doi.org/10.1016/j.envint.2019.104973>.
87. Kumar, S., Molitor, R., and Vollmer, S. (2016). Drought and Early Child Health in Rural India. *Population and Development Review* 42, 53-68. <https://doi.org/10.1111/j.1728-4457.2016.00107.x>.
88. Gari, T., Loha, E., Deressa, W., Solomon, T., Atsbeha, H., Assegid, M., Hailu, A., and Lindtjørn, B. (2017). Anaemia among children in a drought affected community in south-central Ethiopia. *PLOS ONE* 12, e0170898. 10.1371/journal.pone.0170898.
89. Lazzaroni, S., and Wagner, N. (2016). Misfortunes never come singly: Structural change, multiple shocks and child malnutrition in rural Senegal. *Economics & Human Biology* 23, 246-262. <https://doi.org/10.1016/j.ehb.2016.10.006>.
90. Delbiso, T.D., Rodriguez-Llanes, J. M., Donneau, A.-F., Speybroeck, N., Guha-Sapir, D. (2017). Drought, conflict and children's undernutrition in Ethiopia 2000-2013: A meta-analysis. *Bulletin of the World Health Organization* 95, 94–102.
91. IFPRI (2016). 2016 Global Food Policy Report. International Food Policy Research Institute (IFPRI) <https://www.ifpri.org/publication/2016-global-food-policy-report>.
92. UNICEF/WHO/WBG (2019). Levels and trends in child malnutrition: Key findings of the 2019 Edition Joint Child Malnutrition Estimates. World Health Organisation. <https://www.who.int/nutgrowthdb/estimates/en/>.
93. Galasso, E., Wagstaff A., Naudeau S., Shekar M. (2017). The Economic Costs of Stunting and How to Reduce Them. World Bank Group. <http://pubdocs.worldbank.org/en/536661487971403516/PRN05-March2017-Economic-Costs-of-Stunting.pdf>.
94. Tabutin, D., and Schoumaker, B. (2004). La démographie de l'Afrique au sud du Sahara des années 1950 aux années 2000. Synthèse des changements et bilan statistique. *Population*, 519-621.
95. Denis E., and Moriconi-Ebrard F. (2009). La croissance urbaine en Afrique de l'Ouest : De l'explosion à la prolifération.

96. Kookana, R.S., Drechsel, P., Jamwal, P., and Vanderzalm, J. (2020). Urbanisation and emerging economies: Issues and potential solutions for water and food security. *Science of The Total Environment* 732, 139057. <https://doi.org/10.1016/j.scitotenv.2020.139057>.
97. Rezaei Kalvani, S., Sharaai, A., Manaf, L., Hamidian, A. (2019). Assessing ground and surface water scarcity indices using ground and surface water footprints in the Tehran province of Iran. *Applied Ecology and Environmental Research* 17, 4985-4997.
98. Madani, K. (2014). Water management in Iran: what is causing the looming crisis? *Journal of Environmental Studies and Sciences* 4, 315-328. 10.1007/s13412-014-0182-z.
99. Ardalan, A., Khaleghy Rad, M., and Hadi, M. (2019). Urban Water Issues in the Megacity of Tehran. In *Urban Drought: Emerging Water Challenges in Asia*, B. Ray, and R. Shaw, eds. (Springer Singapore), pp. 263-288. 10.1007/978-981-10-8947-3_16.
100. Naghibi, S.A., Pourghasemi, H.R., Pourtaghi, Z.S., and Rezaei, A. (2015). Groundwater qanat potential mapping using frequency ratio and Shannon's entropy models in the Moghan watershed, Iran. *Earth Science Informatics* 8, 171-186. 10.1007/s12145-014-0145-7.
101. Badawi, T. (2018). Iran's water problem. <https://carnegieendowment.org/sada/77935>.
102. Heydari, N., and Morid, S. (2020). Water and agricultural policies in Iranian macro-level documents from the perspective of adaptation to climate change. *Irrigation and Drainage*. 10.1002/ird.2498.
103. Zhou, L., Chen, H., and Dai, Y. (2015). Stronger warming amplification over drier ecoregions observed since 1979. *Environmental Research Letters* 10, 064012. 10.1088/1748-9326/10/6/064012.
104. Koutroulis, A.G. (2019). Dryland changes under different levels of global warming. *Science of The Total Environment* 655, 482-511. <https://doi.org/10.1016/j.scitotenv.2018.11.215>.
105. Feng, S., and Fu, Q. (2013). Expansion of global drylands under a warming climate. *Atmos. Chem. Phys.* 13, 10081-10094. 10.5194/acp-13-10081-2013.
106. Feng, S., Hu, Q., Huang, W., Ho, C.-H., Li, R., and Tang, Z. (2014). Projected climate regime shift under future global warming from multi-model, multi-scenario CMIP5 simulations. *Global and Planetary Change* 112, 41-52. <https://doi.org/10.1016/j.gloplacha.2013.11.002>.
107. Zhao, T., Chen, L., and Ma, Z. (2014). Simulation of historical and projected climate change in arid and semiarid areas by CMIP5 models. *Chinese Science Bulletin* 59, 412-429. 10.1007/s11434-013-0003-x.
108. Chou, C., Neelin, J.D., Chen, C.-A., and Tu, J.-Y. (2009). Evaluating the "Rich-Get-Richer" Mechanism in Tropical Precipitation Change under Global Warming. *Journal of Climate* 22, 1982-2005. 10.1175/2008jcli2471.1.
109. Held, I.M., and Soden, B.J. (2006). Robust Responses of the Hydrological Cycle to Global Warming. *Journal of Climate* 19, 5686-5699. 10.1175/jcli3990.1.
110. Seager, R., Naik, N., and Vecchi, G.A. (2010). Thermodynamic and Dynamic Mechanisms for Large-Scale Changes in the Hydrological Cycle in Response to Global Warming*. *Journal of Climate* 23, 4651-4668. 10.1175/2010jcli3655.1.
111. Liu, F., Zhao, T., Wang, B., Liu, J., and Luo, W. (2018). Different Global Precipitation Responses to Solar, Volcanic, and Greenhouse Gas Forcings. *Journal of Geophysical Research: Atmospheres* 123, 4060-4072. <https://doi.org/10.1029/2017JD027391>.
112. Gu, L., Chen, J., Yin, J., Sullivan, S.C., Wang, H.-M., Guo, S., Zhang, L., and Kim, J.-S. (2020). Projected increases in magnitude and socioeconomic exposure of global droughts in 1.5 and 2 °C warmer climates. *Hydrology and Earth System Sciences* 24, 451-472. 10.5194/hess-24-451-2020.
113. Jenkins, K., and Warren, R. (2015). Quantifying the impact of climate change on drought regimes using the Standardised Precipitation Index. *Theoretical and Applied Climatology* 120, 41-54. 10.1007/s00704-014-1143-x.
114. Vicente-Serrano, S.M., Domínguez-Castro, F., McVicar, T.R., Tomas-Burguera, M., Peña-Gallardo, M., Noguera, I., López-Moreno, J.I., Peña, D., and El Kenawy, A. (2020). Global

- characterization of hydrological and meteorological droughts under future climate change: The importance of timescales, vegetation-CO₂ feedbacks and changes to distribution functions. *International Journal of Climatology* 40, 2557-2567. <https://doi.org/10.1002/joc.6350>.
115. Swann, A.L.S., Hoffman, F.M., Koven, C.D., and Randerson, J.T. (2016). Plant responses to increasing CO₂ reduce estimates of climate impacts on drought severity. *Proceedings of the National Academy of Sciences* 113, 10019. 10.1073/pnas.1604581113.
 116. Carrão, H., Naumann, G., and Barbosa, P. (2018). Global projections of drought hazard in a warming climate: a prime for disaster risk management. *Climate Dynamics* 50, 2137-2155. 10.1007/s00382-017-3740-8.
 117. Ukkola, A.M., Pitman, A.J., De Kauwe, M.G., Abramowitz, G., Herger, N., Evans, J.P., and Decker, M. (2018). Evaluating CMIP5 Model Agreement for Multiple Drought Metrics. *Journal of Hydrometeorology* 19, 969-988. 10.1175/jhm-d-17-0099.1.
 118. Biasutti, M. (2013). Forced Sahel rainfall trends in the CMIP5 archive. *Journal of Geophysical Research: Atmospheres* 118, 1613-1623. <https://doi.org/10.1002/jgrd.50206>.
 119. Biasutti, M., and Giannini, A. (2006). Robust Sahel drying in response to late 20th century forcings. *Geophysical Research Letters* 33. <https://doi.org/10.1029/2006GL026067>.
 120. Rowell, D.P., Senior, C.A., Vellinga, M., and Graham, R.J. (2016). Can climate projection uncertainty be constrained over Africa using metrics of contemporary performance? *Climatic Change* 134, 621-633. 10.1007/s10584-015-1554-4.
 121. Greve, P., Roderick, M.L., Ukkola, A.M., and Wada, Y. (2019). The Aridity Index under global warming. *Environmental Research Letters* 14, 124006. 10.1088/1748-9326/ab5046.
 122. Moncrieff, G.R., Bond, W.J., and Higgins, S.I. (2016). Revising the biome concept for understanding and predicting global change impacts. *Journal of Biogeography* 43, 863-873. <https://doi.org/10.1111/jbi.12701>.
 123. Du, L., Zeng, Y., Ma, L., Qiao, C., Wu, H., Su, Z., and Bao, G. (2021). Effects of anthropogenic revegetation on the water and carbon cycles of a desert steppe ecosystem. *Agricultural and Forest Meteorology* 300, 108339. <https://doi.org/10.1016/j.agrformet.2021.108339>.
 124. Mankin, J.S., Seager, R., Smerdon, J.E., Cook, B.I., and Williams, A.P. (2019). Mid-latitude freshwater availability reduced by projected vegetation responses to climate change. *Nature Geoscience* 12, 983-988. 10.1038/s41561-019-0480-x.
 125. Grafton, R.Q., Pittock, J., Davis, R., Williams, J., Fu, G., Warburton, M., Udall, B., McKenzie, R., Yu, X., Che, N., et al. (2013). Global insights into water resources, climate change and governance. *Nature Climate Change* 3, 315-321. 10.1038/nclimate1746.
 126. Biggs, E.M., Duncan, J.M.A., Atkinson, P.M., and Dash, J. (2013). Plenty of water, not enough strategy: How inadequate accessibility, poor governance and a volatile government can tip the balance against ensuring water security: The case of Nepal. *Environmental Science & Policy* 33, 388-394. <https://doi.org/10.1016/j.envsci.2013.07.004>.
 127. McLeman, R. (2013). Developments in modelling of climate change-related migration. *Climatic Change* 117, 599-611. 10.1007/s10584-012-0578-2.
 128. Gemenne, F. (2011). Why the numbers don't add up: A review of estimates and predictions of people displaced by environmental changes. *Global Environmental Change* 21, S41-S49. <https://doi.org/10.1016/j.gloenvcha.2011.09.005>.
 129. Holleman, C., Jackson, J., Sánchez, M.V., Vos, R. (2017). Sowing the seeds of peace for food security - Disentangling the nexus between conflict, food security and peace
 130. Buhaug, H., Benjaminsen, T.A., Sjaastad, E., and Magnus Theisen, O. (2015). Climate variability, food production shocks, and violent conflict in Sub-Saharan Africa. *Environmental Research Letters* 10, 125015. 10.1088/1748-9326/10/12/125015.
 131. Benjaminsen, T.A. (2016). Does Climate Change Lead to Conflicts in the Sahel? In *The End of Desertification? : Disputing Environmental Change in the Drylands*, R. Behnke, and M. Mortimore, eds. (Springer Berlin Heidelberg), pp. 99-116. 10.1007/978-3-642-16014-1_4.

132. Benjaminsen, T.A., Alinon, K., Buhaug, H., and Buseeth, J.T. (2012). Does climate change drive land-use conflicts in the Sahel? *Journal of Peace Research* 49, 97-111. 10.1177/0022343311427343.
133. Wutich, A., Brewis, A., and Tsai, A. (2020). Water and mental health. *WIREs Water* 7, e1461. <https://doi.org/10.1002/wat2.1461>.
134. Carleton, T.A. (2017). Crop-damaging temperatures increase suicide rates in India. *Proceedings of the National Academy of Sciences* 114, 8746. 10.1073/pnas.1701354114.
135. Edwards, B., Gray, M., and Hunter, B. (2019). The social and economic impacts of drought. *Australian Journal of Social Issues* 54, 22-31. <https://doi.org/10.1002/ajs4.52>.
136. Fu, C. (2017). From climate to global change: Following the footprint of Prof. Duzheng YE's research. *Advances in Atmospheric Sciences* 34, 1159-1168. 10.1007/s00376-017-6300-6.
137. Wario, H.T., Roba, H.G., and Kaufmann, B. (2016). Responding to mobility constraints: Recent shifts in resource use practices and herding strategies in the Borana pastoral system, southern Ethiopia. *Journal of Arid Environments* 127, 222-234. <https://doi.org/10.1016/j.jaridenv.2015.12.005>.
138. Basupi, L.V., Dougill, A.J., and Quinn, C.H. (2019). Institutional challenges in pastoral landscape management: Towards sustainable land management in Ngamiland, Botswana. *Land Degradation & Development* 30, 839-851. <https://doi.org/10.1002/ldr.3271>.
139. Basupi, L.V., Quinn, C.H., and Dougill, A.J. (2019). Adaptation strategies to environmental and policy change in semi-arid pastoral landscapes: Evidence from Ngamiland, Botswana. *Journal of Arid Environments* 166, 17-27. 10.1016/j.jaridenv.2019.01.011.
140. Alimaev, I.I., and Jr, R.H.B. (2008). Ideology, Land Tenure and Livestock Mobility in Kazakhstan. In *Fragmentation in Semi-Arid and Arid Landscapes: Consequences for Human and Natural Systems*, K.A. Galvin, R.S. Reid, R.H.B. Jr, and N.T. Hobbs, eds. (Springer Netherlands), pp. 151-178. 10.1007/978-1-4020-4906-4_7.
141. Bekchanov, M., Ringler, C., Bhaduri, A., and Jeuland, M. (2016). Optimizing irrigation efficiency improvements in the Aral Sea Basin. *Water Resources and Economics* 13, 30-45. <https://doi.org/10.1016/j.wre.2015.08.003>.
142. Thiery, W., Visser, A.J., Fischer, E.M., Hauser, M., Hirsch, A.L., Lawrence, D.M., Lejeune, Q., Davin, E.L., and Seneviratne, S.I. (2020). Warming of hot extremes alleviated by expanding irrigation. *Nature Communications* 11, 290. 10.1038/s41467-019-14075-4.
143. Mishra, V., Ambika, A.K., Asoka, A., Aadhar, S., Buzan, J., Kumar, R., and Huber, M. (2020). Moist heat stress extremes in India enhanced by irrigation. *Nature Geoscience* 13, 722-728. 10.1038/s41561-020-00650-8.
144. Global Water Partnership (2011). What is IWRM? <https://www.gwp.org/en/GWP-CEE/about/why/what-is-iwrm/>
145. Putra, M.P.I.F., Pradhan, P., and Kropp, J.P. (2020). A systematic analysis of Water-Energy-Food security nexus: A South Asian case study. *Science of The Total Environment* 728, 138451. <https://doi.org/10.1016/j.scitotenv.2020.138451>.
146. Stringer, L.C., Quinn, C.H., Le, H.T.V., Msuya, F., Pezzuti, J., Dallimer, M., Afionis, S., Berman, R., Orchard, S.E., and Rijal, M.L. (2018). A New Framework to Enable Equitable Outcomes: Resilience and Nexus Approaches Combined. *Earth's Future* 6, 902-918. <https://doi.org/10.1029/2017EF000694>.
147. Sternberg, T. (2016). Water megaprojects in deserts and drylands. *International Journal of Water Resources Development* 32, 301-320. 10.1080/07900627.2015.1012660.
148. Chen, J., John, R., Sun, G., Fan, P., Henebry, G.M., Fernández-Giménez, M.E., Zhang, Y., Park, H., Tian, L., Groisman, P., et al. (2018). Prospects for the sustainability of social-ecological systems (SES) on the Mongolian plateau: five critical issues. *Environmental Research Letters* 13, 123004. 10.1088/1748-9326/aaf27b.

149. Wang, J., Feng, L., Palmer, P.I., Liu, Y., Fang, S., Bösch, H., O'Dell, C.W., Tang, X., Yang, D., Liu, L., and Xia, C. (2020). Large Chinese land carbon sink estimated from atmospheric carbon dioxide data. *Nature* 586, 720-723. [10.1038/s41586-020-2849-9](https://doi.org/10.1038/s41586-020-2849-9).
150. Abreu, R.C.R., Hoffmann, W.A., Vasconcelos, H.L., Pilon, N.A., Rossatto, D.R., and Durigan, G. (2017). The biodiversity cost of carbon sequestration in tropical savanna. *Science Advances* 3, e1701284. [10.1126/sciadv.1701284](https://doi.org/10.1126/sciadv.1701284).
151. Bryan, B.A., Gao, L., Ye, Y., Sun, X., Connor, J.D., Crossman, N.D., Stafford-Smith, M., Wu, J., He, C., Yu, D., et al. (2018). China's response to a national land-system sustainability emergency. *Nature* 559, 193-204. [10.1038/s41586-018-0280-2](https://doi.org/10.1038/s41586-018-0280-2).
152. Jackson, R.B., Jobbágy, E.G., Avissar, R., Roy, S.B., Barrett, D.J., Cook, C.W., Farley, K.A., le Maitre, D.C., McCarl, B.A., and Murray, B.C. (2005). Trading Water for Carbon with Biological Carbon Sequestration. *Science* 310, 1944. [10.1126/science.1119282](https://doi.org/10.1126/science.1119282).
153. O'Connor, T.G., and van Wilgen, B.W. (2020). The Impact of Invasive Alien Plants on Rangelands in South Africa. In *Biological Invasions in South Africa*, B.W. van Wilgen, J. Measey, D.M. Richardson, J.R. Wilson, and T.A. Zengeya, eds. (Springer International Publishing), pp. 459-487. [10.1007/978-3-030-32394-3_16](https://doi.org/10.1007/978-3-030-32394-3_16).
154. Rolo, V., and Moreno, G. (2019). Shrub encroachment and climate change increase the exposure to drought of Mediterranean wood-pastures. *Science of The Total Environment* 660, 550-558. <https://doi.org/10.1016/j.scitotenv.2019.01.029>.
155. van Wilgen, B.W., Le Maitre, D.C., and Cowling, R.M. (1998). Ecosystem services, efficiency, sustainability and equity: South Africa's Working for Water programme. *Trends in Ecology & Evolution* 13, 378. [10.1016/S0169-5347\(98\)01434-7](https://doi.org/10.1016/S0169-5347(98)01434-7).
156. Stafford, W., Birch, C., Etter, H., Blanchard, R., Mudavanhu, S., Angelstam, P., Blignaut, J., Ferreira, L., and Marais, C. (2017). The economics of landscape restoration: Benefits of controlling bush encroachment and invasive plant species in South Africa and Namibia. *Ecosystem Services* 27, 193-202. <https://doi.org/10.1016/j.ecoser.2016.11.021>.
157. Oweis, T.Y. (2017). Rainwater harvesting for restoring degraded dry agro-pastoral ecosystems: a conceptual review of opportunities and constraints in a changing climate. *Environmental Reviews* 25, 135-149. [10.1139/er-2016-0069](https://doi.org/10.1139/er-2016-0069).
158. N., B. (2014). Iran: Dried out. *Financial Times*, <https://www.ft.com/content/5a5579c6-0205-11e4-ab5b-00144feab7de> (15.02.2019).
159. Tiki, W., Oba, G., and Tvedt, T. (2011). Human stewardship or ruining cultural landscapes of the ancient Tula wells, southern Ethiopia. *Geogr J* 177, 62-78. [10.1111/j.1475-4959.2010.00369.x](https://doi.org/10.1111/j.1475-4959.2010.00369.x).
160. Tiki, W., and Oba, G. (2019). Transforming Labour and Technology of The Ancient Tula Wells for Watering Livestock In Borana, Ethiopia. *Nomadic Peoples* 23, 218-240. [10.3197/np.2019.230204](https://doi.org/10.3197/np.2019.230204).
161. Tal, A. (2016). Rethinking the sustainability of Israel's irrigation practices in the Drylands. *Water Research* 90, 387-394. <https://doi.org/10.1016/j.watres.2015.12.016>.
162. Burney, J., Woltering, L., Burke, M., Naylor, R., and Pasternak, D. (2010). Solar-powered drip irrigation enhances food security in the Sudano-Sahel. *Proceedings of the National Academy of Sciences*. [10.1073/pnas.0909678107](https://doi.org/10.1073/pnas.0909678107).
163. Rosa, L., Chiarelli, D.D., Rulli, M.C., Dell'Angelo, J., and D'Odorico, P. (2020). Global agricultural economic water scarcity. *Science Advances* 6, eaaz6031. [10.1126/sciadv.aaz6031](https://doi.org/10.1126/sciadv.aaz6031).
164. Faour-Klingbeil, D., and Todd, E.C.D. (2018). The Impact of Climate Change on Raw and Untreated Wastewater Use for Agriculture, Especially in Arid Regions: A Review. *Foodborne Pathog Dis* 15, 61-72. [10.1089/fpd.2017.2389](https://doi.org/10.1089/fpd.2017.2389).
165. De Stefano, L., Svendsen, M., Giordano, M., Steel, B.S., Brown, B., and Wolf, A.T. (2014). Water governance benchmarking: concepts and approach framework as applied to Middle East and North Africa countries. *Water Policy* 16, 1121-1139. [10.2166/wp.2014.305](https://doi.org/10.2166/wp.2014.305).

166. Okpara, U.T., Stringer, L.C., and Dougill, A.J. (2018). Integrating climate adaptation, water governance and conflict management policies in lake riparian zones: Insights from African drylands. *Environmental Science & Policy* 79, 36-44. <https://doi.org/10.1016/j.envsci.2017.10.002>.
167. England, M.I., Dougill, A.J., Stringer, L.C., Vincent, K.E., Pardoe, J., Kalaba, F.K., Mkwambisi, D.D., Namaganda, E., and Afionis, S. (2018). Climate change adaptation and cross-sectoral policy coherence in southern Africa. *Regional Environmental Change* 18, 2059-2071. 10.1007/s10113-018-1283-0.
168. Conway, D., van Garderen, E.A., Deryng, D., Dorling, S., Krueger, T., Landman, W., Lankford, B., Lebek, K., Osborn, T., Ringler, C., et al. (2015). Climate and southern Africa's water–energy–food nexus. *Nature Climate Change* 5, 837-846. 10.1038/nclimate2735.
169. Bautista, S., Llovet, J., Ocampo-Melgar, A., Vilagrosa, A., Mayor, Á.G., Murias, C., Vallejo, V.R., and Orr, B.J. (2017). Integrating knowledge exchange and the assessment of dryland management alternatives – A learning-centered participatory approach. *Journal of Environmental Management* 195, 35-45. <https://doi.org/10.1016/j.jenvman.2016.11.050>.

Figure titles and captions


Figure 1: Human population density in drylands. The map shows the population density (number of people/km²) in drylands, as determined by the Aridity Index (see Box 1). High latitude (polar) regions where potential evapotranspiration $\leq 400 \text{ mm y}^{-1}$ constitute cold drylands. Cold drylands are excluded from this review given their sparse human populations compared with hot (tropical) drylands.

Figure 2: Interlinkages between water security and attainment of SDGs in drylands. All SDGs show links to water, underscoring the importance of water security for both environmental and human systems.

Figure 3: Different dimensions of water scarcity in global dryland regions. Different measures highlight the spatial diversity of different dimensions of water scarcity: (a) physical water scarcity (internal renewable water resources per capita); (b) economic water scarcity (total population with access to safe drinking water: and (c) clean water scarcity (water quality risk). Measures are shown for the world's dryland areas, based on the same Millennium Ecosystem Assessment delineation as Figure 1. All variables are normalised to values between 0 and 1 with equal interval classification for comparability.

Figure 4. Conceptualization of water security

Figure 5: Components of an enabling environment for water security in drylands under climate change, drawing on relevant aspects of Integrated Water Resources Management (IWRM)




Goal	Links to water security in drylands	Goal	Links to water security in drylands
 <p>1 NO POVERTY</p>	Water security is vital in drylands to support vast populations dependent on agricultural and natural resource based incomes	 <p>10 REDUCED INEQUALITIES</p>	Inequalities in access to water crosscut multiple aspects of intersectionality
 <p>2 ZERO HUNGER</p>	Rainfed/irrigated food production requires water security to support adequate yields, food security and good nutrition	 <p>11 SUSTAINABLE CITIES AND COMMUNITIES</p>	Drought and floods in urban areas can affect large numbers of people given rapid urbanisation trends in many drylands
 <p>3 GOOD HEALTH AND WELL-BEING</p>	Water-borne diseases and can be reduced with improved access to clean, safe water	 <p>12 RESPONSIBLE CONSUMPTION AND PRODUCTION</p>	Water footprint analysis can provide insights into water use and allocation at different points in the production and consumption process
 <p>4 QUALITY EDUCATION</p>	Child morbidity due to water-borne diseases can be reduced with improved knowledge and education; reduced time spent collecting water increases time for formal education	 <p>13 CLIMATE ACTION</p>	Adaptation is a key response to water scarcity under climate change in drylands
 <p>5 GENDER EQUALITY</p>	Lack of water security has differentiated impacts on women and men / girls and boys	 <p>14 LIFE BELOW WATER</p>	Wastewater flows and eutrophication can affect coastal biodiversity
 <p>6 CLEAN WATER AND SANITATION</p>	Core element with links to all other SDGs	 <p>15 LIFE ON LAND</p>	Irrigation, secondary salinization, ecosystem characteristics
 <p>7 AFFORDABLE AND CLEAN ENERGY</p>	Water security supports hydro-energy generation and plays a key role in energy security	 <p>16 PEACE, JUSTICE AND STRONG INSTITUTIONS</p>	Water can be at the centre of conflicts and competition, as well as being a focus for cooperation
 <p>8 DECENT WORK AND ECONOMIC GROWTH</p>	Water sector jobs and jobs in other sectors rely on water (energy, agriculture etc) and link to youth migration, while earning a decent living can enable people to pay for clean water	 <p>17 PARTNERSHIPS FOR THE GOALS</p>	Transboundary water governance, technology and knowledge exchange, alongside stakeholder participation can help achieve water security

9 INDUSTRY, INNOVATION
AND INFRASTRUCTURE




Industry creates challenges linked to water pollution, water distribution/sewage but can also help by providing flood defences, dams, precision irrigation systems


A - Physical water scarcity



B - Economic water scarcity



C - Clean water scarcity



Risks to water security

