



UNIVERSITY OF LEEDS

This is a repository copy of *Patterns of infection in a native and an invasive crayfish across the UK*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/176521/>

Version: Accepted Version

---

**Article:**

Anderson, LG, Bojko, J, Bateman, KS et al. (3 more authors) (2021) Patterns of infection in a native and an invasive crayfish across the UK. *Journal of Invertebrate Pathology*, 184. 107595. ISSN 0022-2011

<https://doi.org/10.1016/j.jip.2021.107595>

---

© 2021, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Patterns of infection in a native and an invasive crayfish across the UK

Lucy G. Anderson<sup>1</sup>, Jamie Bojko<sup>2,3</sup>, Kelly S. Bateman<sup>4</sup>, Paul D. Stebbing<sup>4,5</sup>, Grant D. Stentiford<sup>4</sup>, Alison M. Dunn<sup>1,\*</sup>

<sup>1</sup>Faculty of Biological Sciences, University of Leeds, Leeds, LS2 9JT, UK. <sup>2</sup>School of Health and Life Sciences, Teesside University, Middlesbrough, TS1 3BA, UK. <sup>3</sup>National Horizons Centre of Excellence in Bioscience Industry, Teesside University, Darlington, DL1 1HG, UK. <sup>4</sup>Centre for environment, fisheries and aquaculture science, Weymouth, Dorset, DT4 8UB, UK. <sup>5</sup>APEM Limited, International House, International Business Park, Southampton SO18 2RZ, UK.

Correspondence: a.dunn@leeds.ac.uk

Keywords: *Nudiviridae*, *Psorospermium*, *Pacifastacus*, *Austropotamobius*, Conservation

## Abstract

Invasive crayfish and the introduction of non-native diseases pose a significant risk for the conservation of endangered, white-clawed crayfish (*Austropotamobius pallipes*). Continued pollution of waterways is also of concern for native species and may be linked with crayfish disease dynamics. We explore whether crayfish species or environmental quality are predictors of infection presence and prevalence in native *A. pallipes* and invasive signal crayfish (*Pacifastacus leniusculus*). We use a seven-year dataset of histology records, and a field survey comparing the presence and prevalence of infectious agents in three isolated *A. pallipes* populations; three isolated *P. leniusculus* populations and three populations where the two species had overlapped in the past. We note a lower diversity of parasites (Simpson's Index) in *P. leniusculus* ('*Pacifastacus leniusculus* Bacilliform Virus' - PIBV) (n= 1 parasite) relative to native *A. pallipes* (n= 4 parasites), which host *Thelohania contejeani*, 'Austropotamobius pallipes bacilliform virus' (ApBV), *Psorospermium haeckeli* and *Branchiobdella astaci*, at the sites studied. The infectious group present in both species was an intranuclear bacilliform virus of the hepatopancreas. The prevalence of *A. astaci* in *A. pallipes* populations was higher in more polluted water bodies, which may reflect an effect of water quality, or may be due to increased chance of transmission from nearby *P. leniusculus*, a species commonly found in poor quality habitats.

## 1. Introduction

Conservationists have long sought to understand the traits that make invasive non-native species (INNS) successful in their introduced ranges (Van Kleunen et al., 2010; Parker et al., 2013). Understanding these attributes can help in assessing comparative risk and formulating preventative management measures. Infectious agents can play an important role in biological invasions, influencing the success of an introduced species and the resilience of native species in the introduced range; ultimately determining the outcome of the invasion (Hatcher et al., 2012; Dunn et al., 2012; Dunn and Hatcher, 2015).

We know little about the diversity and potential for parasite transmission in invasive species, particularly for the Crustacea (Bojko et al. 2020a). Those non-native species that vector infectious agents may result in parasite 'spill over' into native populations, resulting in an emerging disease (Kelly et al., 2009; Tompkins et al., 2011; Okamura and Feist, 2011; Hatcher et al., 2012). Acquisition of non-native diseases by native species depends on the number of infected individuals introduced; host-specificity of the infectious agent; the immunity of native species; and suitable environmental conditions (water chemistry, presence of secondary hosts) for the infectious agent to survive (Okamura and Feist, 2011; Hatcher et al., 2012). If the non-native species is a competent host, it may act as a reservoir for the agent, increasing its prevalence and resulting in 'spill back' into susceptible native populations (Kelly et al., 2009; Poulin et al., 2011; Strauss et al., 2012). In contrast, if the INNS is a less competent host than the native species it may act as a sink for the agent, reducing infection prevalence in native species through a dilution effect (Poulin et al., 2011).

Stochastic and selective pressures during the invasion process may result in non-native species losing their infectious agents, a concept termed 'enemy release' (Keane and Crawley, 2002). Factors surrounding enemy release depend on the size of the invading propagule; selective pressures in the introduced habitat favouring fitter (i.e. uninfected/resistant) hosts; reduced transmission opportunities in the introduced range due to low (founder) population density, or absence of an intermediate host; or, the founder population could be an uninfected life history stage (Torchin et al., 2002; Dunn and Hatcher, 2015).

At the species scale of disease introduction, biogeographical studies have reported that INNS may escape up to 75% of their native infectious agents (Torchin and Mitchell, 2004). Such studies may over-represent the effects of enemy release if they do not compare the invasive population with the specific source population from which it was founded (Colautti et al., 2004; Colautti et al., 2005) as there may be genetic heterogeneity in different native populations, which could influence their resistance to infectious agents as well as spatial heterogeneity in prevalence (MacLeod et al., 2010). Community studies comparing the diversity of infectious

agents in native and invasive conspecifics in the introduced range have often identified similar levels of infectious agents in both species (Colautti et al., 2004).

Under natural conditions, host-parasite interactions are likely to be affected by external factors in the host population's abiotic environment (Sures, 2008; Johnson and Paull, 2011). Freshwater habitats are affected by multiple environmental stressors including pollution, habitat degradation, agricultural run-off and flow modification, as well as INNS (Dudgeon et al., 2006). Environmental factors can affect the competitive ability of native and non-native hosts and their resistance to disease, as well as altering the survival and virulence of the parasite (Keane and Crawley, 2002; Prenter et al., 2004; Sures, 2008; Poulin et al., 2011). Such studies highlight the importance of considering enemy release in the context of other environmental factors (Torchin et al., 2001; Roy et al., 2011).

UK freshwater environments are increasingly imperilled by INNS (Dudgeon et al., 2006; Jackson and Grey, 2012; Gallardo and Aldridge, 2014). One of the impactful freshwater INNS is the American signal crayfish *Pacifastacus leniusculus*, which was first introduced into the UK during the 1970s for the purposes of aquaculture but subsequently escaped and dispersed forming widespread wild populations (Holdich et al., 2014). It commonly out competes native *Austropotamobius pallipes* for food and habitat (Vorburger and Ribi, 1999; Dunn et al., 2008) and is an asymptomatic carrier of the oomycete *Aphanomyces astaci* (causative agent of the disease crayfish plague) (Alderman, 1983). Crayfish plague is associated with sudden and acute mortality events in crayfish (Alderman, 1983; Longshaw, 2011) including subsequent local extinction in *A. pallipes*, and is a major contributor to its listing as an endangered species on the IUCN Red List (Füreder et al., 2010).

Although mortalities are an obvious impact of this invasion, the role that sub-lethal infectious agents play in invasion dynamics have received less attention in crayfish (Longshaw, 2011). Infectious agents in *A. pallipes* show high diversity and prevalence of infection in some cases, while similar data of non-native crayfish imports show that 66.4% of non-native crayfish were free from infection (Longshaw et al., 2012a; Longshaw et al., 2012b). Although these studies indicate that enemy release may play a role in invasion success, others suggest that the fitness of native crayfish, and prevalence of some infections in native crayfish, may be affected by the quality of the local environment (Imhoff, 2010).

We explore the pathogen profile of invasive populations of *P. leniusculus* in the UK and correlate this information with environmental measurements. We also explore the pathogen profile of UK native *A. pallipes*. In both cases we explore specimens from populations that have and have not interacted and compare diversity indices. We discuss these pathological,

geographical, and environmental data in relation to the ongoing crayfish invasion across the UK.

## 2. Methods

### 2.1 Curation of a seven-year dataset on crayfish disease

All histological surveys of *A. pallipes* (25 populations; 210 individuals) and *P. leniusculus* (44 populations; 818 individuals) conducted by the Centre for Environment, Fisheries and Aquaculture Science (Cefas) between 2007 and 2014 were combined into a single seven-year dataset (hereby “Sample A”) (Table 1). Only crayfish populations subjected to full histological examinations were included in the dataset to investigate the influence of co-infections. Other *A. pallipes* populations (25 populations; 123 individuals, Sample B) that exist within a Cefas dataset exploring mortality events [Table 2 were analysed separately to investigate putative predictors of *A. astaci* prevalence, using molecular screening (PCR) instead of histology].

### 2.2 Specimen collection for field studies

To explore any relationships between water quality (chemical and ecological status of the catchment) and the presence/prevalence of parasites, crayfish were collected from three *A. pallipes* populations and six *P. leniusculus* populations under license from Natural England (Licence number: 20122156) and the Environment Agency (FR2 licenses), between June and October 2012 (Sample C). Efforts to locate overlapping mixed-species populations of the two crayfish species were unsuccessful, in part due to a lack of available documentation on their distribution. As a proxy for mixed populations, three single-species *P. leniusculus* populations (Bookill Gill Beck, Cawthorne Dike, and the River Ure; Table 2) where the co-occurrence of *A. pallipes* had been recorded within the previous two years were used (Dunn et al., 2008; Haddaway et al., 2012). The other six populations (three of *A. pallipes* and three of *P. leniusculus*) were selected because they had been isolated from introductions of other crayfish species in the past, according to the Rivers Trusts, local records centres, National Biodiversity Network Gateway, and ecological consultants.

### 2.3 Histological screening

Samples were prepared and histologically analysed in accordance with a standard crustacean disease screening protocol (Longshaw 2011; Longshaw et al. 2012a; Longshaw et al. 2012b). All crayfish were examined for external abnormalities (e.g. missing claws, damage to carapace), sexed and measured (carapace length) before being exposed to an overdose of chloroform vapours to humanely euthanize them prior to sampling for histology.

Following the standard protocol (described in Longshaw et al. 2012a), Juvenile crayfish ( $\leq 10\text{mm}$  carapace length) were euthanized and fixed whole by direct injection of Davidson's freshwater fixative. Larger animals ( $>10\text{mm}$  carapace length) were dissected and samples of the carapace, gill, gonad, gut, hepatopancreas and tail muscle were collected immediately and preserved in Davidson's freshwater fixative for 24h before being transferred to 70% industrial methylated spirits (IMS). If required, the tissue samples were decalcified in a rapid decalcification solution prior to paraffin infiltration. The tissues were processed to wax block using an automatic vacuum infiltration tissue processor (Peloris, Vision Biosystems). Sections were cut at 3 to 5  $\mu\text{m}$  and routinely stained with haematoxylin and eosin (H&E) in an automatic tissue stainer (Tribune Autostainer, Surgipath). The tissues were examined on a light microscope using brightfield illumination. A record was made of any pathologies or infectious agents in organs and tissues including an indication of the infection severity. Images were captured using a LUCIA™ (Nikon, UK) screen measurement system.

#### 2.4 PCR screening for *Aphanomyces astaci*

Biopsies were collected from the cuticle and sub-cutis of all adult crayfish and individually placed in 100% ethanol and stored at  $-20^{\circ}\text{C}$  (sample numbers are located in Table 1 and Table 2). The full methods used to extract DNA from tissues have been previously described in (Oidtmann, 2004). DNA was extracted using a DNeasy tissue kit (Qiagen) following the manufacturer's instructions. Animals were screened for *A. astaci* using the PCR protocol described in (Oidtmann et al., 2006).

#### 2.5 Environmental parameters

The Water Framework Directive requires all European Union (EU) Member States to assess and classify the status of their river catchments according to a unified set of ecological and chemical standards (Water Framework Directive, 2012). The ecological classification ('high' to 'bad' on a five-point scale), scores each water body against a set of biological quality (abundance of fish and rooted plants), general chemical and physio-chemical (temperature and nutrient levels) water quality with respect to specific pollutants (synthetic and non-synthetic) and hydro-morphological (water flow and physical habitat) criteria (Water Framework Directive, 2012). The chemical classification ('good' or 'fail') examines the presence of polluting substances that could adversely affect the ecology of the catchment by checking whether the water meets Environmental Quality Standards (EQSs) for substances listed in Annex IX (Dangerous Substances Directive and associated daughter Directives) and Annex X (WFD Priority List Substances) (Water Framework Directive, 2012). According to the Environment Agency, a status of good means that concentrations of priority

substances and priority hazardous substances do not exceed the environmental quality standards in the EQS Directive (cite EA glossary). To ensure that the environmental parameters were policy-relevant as well as biologically robust, the chemical and ecological status of the sub-catchment of each crayfish population in the Cefas dataset was checked using the Environment Agency's Catchment Data Explorer website ([environment.data.gov.uk/catchment-planning/](http://environment.data.gov.uk/catchment-planning/)).

## 2.5 Statistical analysis

Generalised linear mixed-effects models (GLMMs) were used to determine which variables were predictors of the presence/absence, prevalence, and diversity (Simpson's index) of parasites in *A. pallipes* and *P. leniusculus* populations in the UK based on the Cefas dataset. Explanatory variables included crayfish species; presence of other infectious agents in the population; prevalence of other infectious agents in the population; chemical status of the sub-catchment; and ecological status of the sub-catchment. For single-species *P. leniusculus* sites the former presence of *A. pallipes* crayfish in the catchment was added to the model. Due to the wide temporal and geographical range of the data, site and year were included as random factors in each model. All models were fitted with a binomial error distribution and a logit link function since the response variables consisted of both binary and proportion data (Crawley, 2007).

The models met the assumptions of homogeneity of variance based on visual assessment of the plots of residuals vs. fitted values. The normality of residuals was checked using quantile-quantile plots and histograms. Log-likelihood tests were used to compare simplified models to null models (random effects only) and conditional  $R^2$  values were calculated to describe the proportion of variance explained by both the fixed and random factors (Johnson, 2014).

## 3. Results

### 3.1 Predictors of infection (Cefas dataset)

*Austropotamobius pallipes* were found to have a higher number of parasites than *P. leniusculus* (4:1) based on histopathology results. Histology results revealed that *A. pallipes* populations were infected with four agents: *Thelohania contejeani*, 'Austropotamobius pallipes bacilliform virus' (ApBV), *Psorospermium haeckeli* and *Branchiobdella astaci* (mean Simpson's index = 0.17), while *P. leniusculus* were infected by 'Pacifastacus leniusculus Bacilliform Virus' only (PIBV) (Fig. 1). The crayfish species was the only significant predictor of infectious agent diversity in the minimum adequate model (Simpson's index) (Table 2). Crayfish plague data were collected for *A. pallipes* using PCR and is explored in section 3.2. Individual signal crayfish were screened for crayfish plague, using PCR, by Cefas and

reported for 5 sites over 2011 and 2014 (Table 1). This constituted 34 individual signal crayfish in total, but the overall sample size was too small for reliable statistical comparison between sites.

None of the variables (ecological status, chemical status, or crayfish species) were significant predictors of the presence/absence of bacilliform viruses in crayfish populations; however, crayfish species was a significant predictor of the prevalence of infection (Table 2), with a higher prevalence observed in *A. pallipes* populations (13/25 populations infected, mean prevalence = 0.39) than *P. leniusculus* populations (11/44 populations infected, mean prevalence = 0.13).

### 3.2 Predictors of infection in *A. pallipes*

The presence and prevalence of *T. contejeani*, *P. haeckeli* and *B. astaci* were explored in the *A. pallipes* dataset (Sample B) *Thelohania contejeani* was present in 11/25 *A. pallipes* populations in the long-term dataset and the mean prevalence among infected populations was 27 % (range 9 – 37 %). *Branchiobdella astaci* and *P. haeckeli* were identified histologically (Rosewarne et al., 2012) and present in 2/25 populations each (not the same two populations) and infected populations had a mean prevalence of 19 % and 60 %, respectively. Neither the ecological nor chemical status of the catchment were significant predictors of the presence or prevalence of the three parasites (model P values >0.05). The presence of *T. contejeani* in the population was a significant predictor of ApBV prevalence (Table 2). The mean prevalence of ApBV was 58 % in *A. pallipes* populations with *T. contejeani* and 23 % in populations without *T. contejeani* infection.

Populations of *A. pallipes* that were tested positive for *A. astaci* (Sample B; Table 1) included 13 of the 25 populations in the UK. Infection prevalence in infected populations ranged from 10% to 100%. A significant relationship was found between the chemical status of the water body and the presence of crayfish plague: waterbodies with a “fail” for chemical status appeared more likely to test positive for *A. astaci*; however, none of the variables were significant predictors of the prevalence of *A. astaci*.

### 3.3 Field sampling to investigate enemy release

Nine sites with three different population compositions were assessed to compare enemy release: i) isolated *A. pallipes* populations; ii) isolated *P. leniusculus* populations; iii) *P. leniusculus* with recent *A. pallipes* overlap. *Austropotamobius pallipes* populations at these sites were infected with three parasites (*T. contejeani*, ApBV, *Branchiobdella astaci*), while signal crayfish populations were only infected with PIBV (Fig. 2). Crayfish population composition (single species vs. proxy mixed) was a significant predictor of bacilliform virus



prevalence (Estimate =  $2.62 \pm 0.89$ ,  $t = 2.96$ ,  $p < 0.05$ ; Model  $R^2 = 0.21$ ), with higher prevalence associated with single-species *A. pallipes* populations (72 %) and *P. leniusculus* populations that had recently overlapped with *A. pallipes* (20%) and low prevalence among *P. leniusculus* only populations (<1 %) (Table 2; Fig. 2).

#### 4. Discussion

Our study identified that non-native *P. leniusculus* are hosts to a significantly lower diversity of infectious agents when compared to native *A. pallipes*, in the UK (Table 2). Our results did not provide evidence that the presence and prevalence of sub-lethal crayfish infections are affected by the chemical or environmental status of their habitat; however, this data did suggest that water quality may be linked with a tentatively increased prevalence of crayfish plague. We sub-divide our discussion by parasite group to explore pathology, impacts and conservation but provide a single section to discuss environmental quality and the presence of disease.

##### 4.1 Intranuclear bacilliform virus

Hepatopancreatic bacilliform viruses have been identified from a range of crustacean species including multiple observations among crayfish (Bateman and Stentiford, 2017; Bojko et al. 2017; Bojko and Ovcharenko, 2019). Recent work has identified that these viruses are likely novel members of the *Nudiviridae*, based on genomic and ultrastructural data (Yang et al. 2014; Holt et al. 2019; Allain et al. 2020). Their impact on crustacean hosts is sparse, including data pertaining to biological invasions. Some evidence has been gathered using amphipod models, where infection level is associated with increased activity (Bojko et al. 2018); however, this has not been explored in crayfish invasion, nor has the distribution of the two viruses in UK populations. To date, five 'bacilliform viruses' have been identified in crayfish, but genomic data is necessary to determine their taxonomy (see Table 1 of Bojko et al. 2017).

Bacilliform viruses in both *P. leniusculus* and *A. pallipes* populations in this study were most prevalent in single-species *A. pallipes* populations. A high prevalence of PIBV was also detected in *P. leniusculus* populations that had previously overlapped with *A. pallipes*. Low prevalence, or absence, was detected in *P. leniusculus* populations where no native crayfish were located. Greater genomic data is necessary to determine if these viruses are part of the same species complex or if they are different, species-specific, viruses of each crayfish species.

In previous studies, no gross pathological changes were observed in crayfish infected with bacilliform viruses, suggesting that this virus has a minimal impact on host health, and it may not be a major driver of crayfish mortality, nor a determinant of competitive success (Stentiford

et al., 2004; Longshaw, 2011; Longshaw et al., 2012a; Longshaw et al., 2012c). Interestingly, a higher prevalence of *T. contejeani* in *A. pallipes* populations was a predictor for bacilliform virus presence, suggesting the presence of shared stressors that could increase susceptibility to these two parasites. Alternatively, the virus and microsporidian may share a relationship that is yet to be discovered. The presence of microsporidian-virus coinfection was described by Bojko et al. (2019) at high prevalence in amphipods, and there may be a relationship to untangle between the two during an invasion dynamic.

#### 4.2 *Thelohania contejeani* and other Microsporidia

Crayfish have been associated with several microsporidian lineages. These include members from clades III (*T. contejeani*), IV (AM261754) and V (*Camabaraspora floridanus* and *Ovipleistophora diplostumuri*) (Bojko et al. 2020b; Bojko et al. 2020c). In the UK, aside from the detection of a *Bacillidium*-like microsporidian via PCR, this is dominated by the detection of *T. contejeani* (Dunn et al. 2008). This microsporidian is thought to be a native parasite of *A. pallipes* and results in extensive muscular degeneration, resulting in limited movement, feeding (30% reduction) and predator behaviour (Alderman and Polglase, 1988; Oidtmann et al., 1996; Longshaw et al. 2011). Sub-lethal impacts that facilitate competitive exclusion and are thought to play an important role in competitive interactions between *A. pallipes* and *P. leniusculus* (Haddaway et al., 2012).

*Thelohania contejeani* was common among *A. pallipes* populations (present in 46% of populations from the Cefas dataset) with a maximum prevalence of 37%, consistent with other studies (Cossins and Bowler, 1974; Dieguez-Urbeondo et al. 1997; Mori and Salvidio, 2000; Rodgers et al., 2003; Dunn et al., 2008; Quaglio et al. 2011; Longshaw et al., 2012c). Despite examining 50 populations (966 individuals) of *P. leniusculus* in the field study and Cefas datasets combined, there were not any histologically identifiable microsporidian infections found in *P. leniusculus*. The lack of detection is surprising given that *P. leniusculus* populations had previously tested positive for microsporidian infection and were re-sampled as part of this study (Dunn et al., 2008). This may suggest that although *T. contejeani* is present in *P. leniusculus* using PCR (Dunn et al., 2008; Imhoff et al., 2012), it may be a less competent host, resulting in lower parasite burden and preventing an infection from reaching detectable levels using histopathology alone. Imhoff et al. (2012) found that *P. leniusculus* could become infected with *T. contejeani* by consuming infected tissue from *A. pallipes*, confirming the capability to infect; however, in the wild this may be rare.

The native range of *P. leniusculus* has been little studied for parasitic associations; however, one study isolated microsporidian parasites from this species in California (McGriff and Modin, 1983). Since this data includes morphological comparison but lacks genetic identification, it

remains speculative to suggest this is *T. contejeani*. Such information could however suggest that this parasite was introduced by the signal crayfish invasion. Our data suggest this is unlikely, because multiple white clawed crayfish populations that are not affected by the invasion exhibit this microsporidian infection, suggesting that the parasite found in California could be a different species.

#### 4.3 *Branchiobdella astaci*

Branchiobdellids are ectobionts generally considered to be commensal. In our study we found their presence in two *A. pallipes* populations. Histology images show an association between the presence of *B. astaci* and gill melanisation, a localised immune response that may impair the gill function (Alderman and Polglase, 1988; Rosewarne et al., 2012). *Branchiobdella astaci* has not been associated with crayfish mortality (Longshaw, 2011), nor co-infection, and did not appear to cause any gross pathological signs in *A. pallipes*, suggesting that it would have minimal impact on competitive interactions between native species and INNS.

#### 4.4 Environmental parameters

The environment of the host is an important determinant of disease dynamics (Strayer, 2010; Johnson and Paull, 2011). Here we show that this may also be the case for crayfish and some of the disease groups we explore. A relatively small-scale study of *A. pallipes* in the River Wharfe catchment in Yorkshire reported a positive correlation between the presence of *T. contejeani* and the levels of zinc, lead, and dissolved oxygen (Imhoff, 2010). Similarly, a laboratory experiment showed a trend with penaeid shrimp that revealed the prevalence of a shrimp-specific bacilliform virus increased from 23 % to 75 % after 35 days, when shrimp were exposed to aquatic pollutants (polychlorinated biphenyls) but only increased from 23 % to 46 % in the control group (Couch and Courtney, 1977). In contrast to these previous studies, we did not reveal a relationship between the presence or prevalence of either *T. contejeani* or bacilliform virus and the chemical status of the catchment.

There was a significant relationship between the chemical status of the water body and the prevalence of *A. astaci*, the causative agent of crayfish plague. The prevalence of *A. astaci* was higher in catchments that received 'fail' status for their chemical classification. This may be because the immunity of *A. pallipes* is reduced in more polluted river catchments, or because these catchments provide optimal conditions for oomycete growth. The motility of *A. astaci* is dependent on water temperature, while high magnesium levels and low calcium levels are considered less favourable for spores (Oidtmann, 2000). Alternatively, external factors such as pollution are considered to reduce the resilience of freshwater ecosystems to invasion (Dudgeon et al., 2006; Strayer, 2010). The correlation we observe may have been

the result of polluted catchments being more susceptible to invasion by *P. leniusculus*, an asymptomatic carrier of *A. astaci*, rather than resulting in decreased host immunity. Such a theory requires testing, but the basis is hinted at from our environmental data.

The Water Framework Directive catchment classification takes multiple environmental parameters into account to produce an overall “status”, which may omit the impacts of particularly important stressors that were not measured in this study. Moreover, localised pollution events affecting acute areas of a river may have been missed at the broad spatial scale at which the catchments are assessed, and therefore recommend that more localised studies are conducted in future to explore these findings in higher resolution.

#### 4.5 Study conclusions

The objectives of this study were to determine whether either crayfish species or environment were predictors of infection in UK habitats. This information could be highly informative for conservation efforts, providing the best chances to develop crayfish havens where water quality may avoid further disease outbreaks. We did not find a relationship between environmental quality and the presence or prevalence of sublethal infections in crayfish. Our results did suggest a link between water quality and the presence of *A. astaci*, it is difficult to disentangle the contribution of water quality from that of *P. leniusculus* presence, as the species is an asymptomatic carrier of *A. astaci* and is more tolerant of low-quality waterbodies.

#### Acknowledgements

The authors acknowledge Cefas/Defra (FB002) funding for the development of the dataset and histological processing. Specific thanks to Tracy Bull in the Fish Health Inspectorate (Cefas) for her work on the database. This research was part of LGA's PhD thesis, funded by BBSRC.

#### References

- Alderman DJ (1993) Crayfish plague in Britain, the first twelve years. *Freshwater Crayfish*. 9, 266–272.
- Alderman DJ, Polglase JL, Frayling M, Hogger J (1984) Crayfish plague in Britain. *Journal of Fish Diseases*. 7(5), 401–405.
- Allain, T. W., Stentiford, G. D., Bass, D., Behringer, D. C., & Bojko, J. (2020). A novel nudivirus infecting the invasive demon shrimp *Dikerogammarus haemobaphes* (Amphipoda). *Scientific Reports*, 10(1), 1-13.

390 Bateman KS, Stentiford GD (2017) A taxonomic review of viruses infecting crustaceans with  
 391 an emphasis on wild hosts. *Journal of Invertebrate Pathology*. 147, 86-110.

392 Bojko, J., Burgess, A. L., Baker, A. G., & Orr, C. H. (2020a). Invasive Non-Native Crustacean  
 393 Symbionts: Diversity and Impact. *Journal of Invertebrate Pathology*, In Press.

394 Bojko, J., Behringer, D. C., Moler, P., Stratton, C. E., & Reisinger, L. (2020b). A new lineage  
 395 of crayfish-infecting Microsporidia: The *Cambaraspora floridanus* n. gen. n. sp. (Glugeida:  
 396 Glugeidae) complex from Floridian freshwaters (USA). *Journal of Invertebrate Pathology*, 171,  
 397 107345.

398 Bojko, J., Behringer, D. C., Moler, P., & Reisinger, L. (2020c). *Ovipleistophora diplostomuri*,  
 399 a parasite of fish and their trematodes, also infects the crayfish *Procambarus bivittatus*.  
 400 *Journal of Invertebrate Pathology*, 169, 107306.

401 Bojko J, Ovcharenko M (2019) Pathogens and other symbionts of the Amphipoda: taxonomic  
 402 diversity and pathological significance. *Diseases of Aquatic Organisms*. 136, 3-36.

403 Bojko J, Stentiford GD, Stebbing PD, Hassall C, Deacon A, Cargill B, Pile B, Dunn AM (2019)  
 404 Pathogens of *Dikerogammarus haemobaphes* regulate host activity and survival, but also  
 405 threaten native amphipod populations in the UK. *Diseases of Aquatic Organisms*. 136, 63-78.

406 Bojko J, Bącela-Spychalska K, Stebbing PD, Dunn AM, Grabowski M, Rachalewski M,  
 407 Stentiford GD (2017) Parasites, pathogens and commensals in the “low-impact” non-native  
 408 amphipod host *Gammarus roeselii*. *Parasites & Vectors*. 10(1), 193.

409 Colautti RI, Muirhead JR, Biswas RN, MacIsaac HJ (2005) Realized vs apparent reduction in  
 410 enemies of the European starling. *Biological Invasions*. 7(4), 723–732.

411 Colautti RI, Ricciardi A, Grigorovich IA, MacIsaac HJ (2004) Is invasion success explained by  
 412 the enemy release hypothesis? *Ecology Letters*. 7(8), 721–733.

413 Crawley MJ (2007) *The R book*. Chichester, England; Hoboken, N.J.: Wiley.

414 Dudgeon D, Arthington AH, Gessner MO, Kawabata ZI, Knowler DJ, Lévêque C, Naiman RJ,  
 415 Prieur-Richard AH, Soto D, Stiassny MLJ, Sullivan C (2006) Freshwater biodiversity:  
 416 importance, threats, status and conservation challenges. *Biological reviews of the Cambridge*  
 417 *Philosophical Society*. 81(2), 163–82.

418 Dunn AM, Hatcher MJ (2015) Parasites and biological invasions: parallels, interactions, and  
 419 control. *Trends in parasitology*. 31(5), 189-199.

420 Dunn AM, Torchin ME, Hatcher MJ, Kotanen PM, Blumenthal DM, Byers JE, Coon CAC,  
 421 Frankel VM, Holt RD, Hufbauer RA, Kanarek AR, Schierenbeck KA, Wolfe LM, Perkins SE

422 (2012) Indirect effects of parasites in invasions C. Fox, ed. *Functional Ecology*. 26(6), 1262–  
 423 1274.

424 Dunn JC, McClymont HE, Christmas M, Dunn AM (2008) Competition and parasitism in the  
 425 native White-clawed Crayfish *Austropotamobius pallipes* and the invasive Signal Crayfish  
 426 *Pacifastacus leniusculus* in the UK. *Biological Invasions*. 11(2), 315–324.

427 Füreder L, Gherardi F, Holdich D, Reynolds J, Sibley P, Souty-Grosset C (2010)  
 428 *Austropotamobius pallipes*. The IUCN red list of threatened species. 11, 2010-3.

429 Gallardo B, Aldridge DC (2015) Is Great Britain heading for a Ponto–Caspian invasional  
 430 meltdown?. *Journal of Applied Ecology*. 52(1), 41-49.

431 Haddaway NR, Wilcox RH, Heptonstall REA, Griffiths HM, Mortimer RJG, Christmas M, Dunn  
 432 AM (2012) Predatory functional response and prey choice identify predation differences  
 433 between native/invasive and parasitised/unparasitised crayfish. *PLoS ONE*. 7(2), e32229.

434 Hatcher MJ, Dick JTA, Dunn AM (2012) Disease emergence and invasions S. Perkins, ed.  
 435 *Functional Ecology*. 26(6), 1275–1287.

436 Hatcher MJ, Dunn AM (2011) *Parasites in ecological communities: from interactions to  
 437 ecosystems*. Cambridge, UK; New York: Cambridge University Press.

438 Holdich DM, James J, Jackson C, Peay S (2014) The North American signal crayfish, with  
 439 particular reference to its success as an invasive species in Great Britain. *Ethology Ecology  
 440 & Evolution*. 26(2-3), 232–262.

441 Holt, C. C., Stone, M., Bass, D., Bateman, K. S., van Aerle, R., Daniels, C. L., ... & Stentiford,  
 442 G. D. (2019). The first clawed lobster virus *Homarus gammarus nudivirus* (HgNV n. sp.)  
 443 expands the diversity of the Nudiviridae. *Scientific Reports*, 9(1), 1-15.

444 Imhoff E (2010) The impact of water chemistry and parasitism by *Thelohania contejeani*  
 445 (Microspora) on native (*Austropotamobius pallipes*) and invasive (*Pacifastacus leniusculus*)  
 446 crayfish and their interactions. PhD Thesis. University of Leeds.

447 Jackson MC, Grey J (2013) Accelerating rates of freshwater invasions in the catchment of the  
 448 River Thames. *Biological Invasions*. 15(5), 945-951.

449 Johnson PTJ, Paull SH (2011) The ecology and emergence of diseases in fresh waters.  
 450 *Freshwater Biology*. 56(4), 638–657.

451 Keane RM, Crawley MJ (2002) Exotic plant invasions and the enemy release hypothesis.  
 452 *Trends in Ecology & Evolution*. 17(4), 164–170.

453 Kelly DW, Paterson RA, Townsend CR, Poulin R, Tompkins DM (2009) Parasite spillback: a  
 454 neglected concept in invasion ecology? *Ecology*. 90(8), 2047–2056.

455 Longshaw M (2011) Diseases of crayfish: A review. *Journal of Invertebrate Pathology*. 106(1),  
 456 54–70.

457 Longshaw M, Bateman KS, Stebbing P, Stentiford GD, Hockley FA (2012a) Disease risks  
 458 associated with the importation and release of non-native crayfish species into mainland  
 459 Britain. *Aquatic Biology*. 16(1), 1–15.

460 Longshaw M, Stebbing PD, Bateman KS, Hockley FA (2012b) Histopathological survey of  
 461 pathogens and commensals of white-clawed crayfish (*Austropotamobius pallipes*) in England  
 462 and Wales. *Journal of Invertebrate Pathology*. 110(1), 54–59.

463 MacLeod CJ, Paterson AM, Tompkins DM, Duncan RP (2010) Parasites lost - do invaders  
 464 miss the boat or drown on arrival? *Ecology Letters*. 13(4), 516–527.

465 McGriff, D., & Modin, J. (1983). *Thelohania contejeani* parasitism of the crayfish, *Pacifastacus*  
 466 *leniusculus*, in California. *California Fish and Game*, 69(3), 178-183.

467 Oidtmann B (2000) Disease in freshwater crayfish. In Leeds Crayfish Conference.  
 468 Environment Agency, pp. 9–18.

469 Oidtmann BC, Thrush MA, Denham KL, Peeler EJ (2011) International and national  
 470 biosecurity strategies in aquatic animal health. *Aquaculture*. 320(1-2), 22–33.

471 Oidtmann B, Heitz E, Rogers D, Hoffmann RW (2002) Transmission of crayfish plague.  
 472 *Diseases of Aquatic Organisms*. 52(2), 159–167.

473 Okamura B, Feist SW (2011) Emerging diseases in freshwater systems. *Freshwater Biology*.  
 474 56(4), 627–637.

475 Parker JD, Torchin ME, Hufbauer RA, Lemoine NP, Alba C, Blumenthal DM, Bossdorf O,  
 476 Byers JE, Dunn AM, Heckman RW (2013) Do invasive species perform better in their new  
 477 ranges? *Ecology*. 94(5), 985–994.

478 Poulin R, Paterson RA, Townsend CR, Tompkins DM, Kelly DW (2011) Biological invasions  
 479 and the dynamics of endemic diseases in freshwater ecosystems. *Freshwater Biology*. 56(4),  
 480 676–688.

481 Prenter J, MacNeil C, Dick JTA, Riddell GE, Dunn AM (2004) Lethal and sublethal toxicity of  
 482 ammonia to native, invasive, and parasitised freshwater amphipods. *Water Research*. 38(12),  
 483 2847–2850.

484 Rodgers JC, Parker KC (2003) Distribution of alien plant species in relation to human  
485 disturbance on the Georgia Sea Islands. *Diversity and Distributions*. 9(5), 385–398.

486 Rosewarne et al., 2012

487 Roy HE, Lawson Handley LJ, Schönrogge K, Poland RL, Purse BV (2011) Can the enemy  
488 release hypothesis explain the success of invasive alien predators and parasitoids?  
489 *BioControl*. 56(4), 451–468.

490 Strauss A, White A, Boots M (2012) Invading with biological weapons: the importance of  
491 disease-mediated invasions S. Perkins, ed. *Functional Ecology*. 26(6), 1249–1261.

492 Strayer DL (2010) Alien species in fresh waters: ecological effects, interactions with other  
493 stressors, and prospects for the future. *Freshwater Biology*. 55, 152–174.

494 Sures B (2008) Interactions between parasites and pollutants in the aquatic environment.  
495 *Parasite*. 15(3), 434–438.

496 Tompkins DM, Dunn AM, Smith MJ, Telfer S (2011) Wildlife diseases: from individuals to  
497 ecosystems: Ecology of wildlife diseases. *Journal of Animal Ecology*. 80(1), 19–38.

498 Torchin ME, Lafferty KD, Dobson AP, McKenzie VJ, Kuris AM (2003) Introduced species and  
499 their missing parasites. *Nature*. 421(6923), 628–630.

500 Torchin ME, Lafferty KD, Kuris AM (2002) Parasites and marine invasions. *Parasitology*.  
501 124(07), 137–151.

502 Torchin ME, Lafferty KD, Kuris AM (2001) Release from parasites as natural enemies:  
503 increased performance of a globally introduced marine crab. *Biological Invasions*. 3(4), 333–  
504 345.

505 Torchin ME, Mitchell CE (2004) Parasites, pathogens, and invasions by plants and animals.  
506 *Frontiers in Ecology and the Environment*. 2(4), 183–190. 142

507 UK Technical Advisory Group on the Water Framework Directive (2014) Revised classification  
508 of aquatic alien species according to their level of impact. Available from:  
509 <http://www.wfduk.org/tagged/alien-species>

510 Van Kleunen M, Weber E, Fischer M (2010) A meta-analysis of trait differences between  
511 invasive and non-invasive plant species. *Ecology Letters*. 13(2), 235–245.

512 Vorburger C, Ribi G (1999) Aggression and competition for shelter between a native and an  
513 introduced crayfish in Europe. *Freshwater Biology*. 42(1), 111–119.

514 Yang YT, Lee DY, Wang Y, Hu JM, Li WH, Leu JH, Chang GD, Ke HM, Kang ST, Lin SS, Kou  
515 GH, Lo CF (2014) The genome and occlusion bodies of marine *Penaeus monodon* nudivirus



(PmNV, also known as MBV and PemoNPV) suggest that it should be assigned to a new nudivirus genus that is distinct from the terrestrial nudiviruses. BMC genomics, 15(1), 628.

## Tables and Figures:

Table 1: A breakdown of the crayfish datasets, including historical Cefas data (2007-2014) and primary data collected for the study in 2012.

Sampling sub-set		Number of individuals ( <i>A. pallipes</i> )	Number of individuals ( <i>P. leniusculus</i> )	Tests conducted
A	Seven-year Cefas dataset on crayfish disease	210 individuals from (25 populations)	818 individuals from (44 populations)	Full histological examination
B	Crayfish collected by Cefas during mortality events	123 individuals from (25 populations)	37 individuals from (5 populations)	<i>A. astaci</i> presence and prevalence using PCR
C	Field-based collections specifically to inform study	68 individuals from (3 populations) (Table 2)	128 individuals from (6 populations) (Table 2)	Full histological examination

Table 2: Location and composition of crayfish populations sampled as part of the 2012 survey. The results show the prevalence of the three parasites recorded during the study for *Austropotamobius pallipes* and *Pacifastacus leniusculus* populations.

Species	Site (coordinates)	n	<i>T. contejeani</i>	Bacilliform virus	<i>B. astaci</i>
<i>A. pallipes</i>	Wyke Beck, Leeds, West Yorkshire (53.8225, -1.4819)	24	0.37	0.58	0.25
<i>A. pallipes</i>	Clapham Beck, Clapham, North Yorkshire (54.118116, -2.391811)	33	0.09	0.80	0.13
<i>A. pallipes</i>	River Kent, Kendal, Cumbria (54.341219, -2.747489)	11	0.09	0.80	0
<i>P. leniusculus</i> (formerly mixed species)	Bookill Gill Beck, Long Preston, North Yorkshire	28	0	0	0

	(54.022255 - 2.242651)				
<i>P. leniusculus</i> (formerly mixed species)	Cawthorne Dike, Cawthorne, South Yorkshire (53.575938, - 1.555192)	33	0	0.6	0
<i>P. leniusculus</i> (formerly mixed species)	River Ure, West Tanfield, North Yorkshire (54.203132, - 1.589163)	9	0	0	0
<i>P. leniusculus</i>	River Clyde, Elvanfoot, Scotland (55.433032, - 3.649609)	22	0	0.2	0
<i>P. leniusculus</i>	Aske Estate, Richmond, North Yorkshire (54.424541, - 1.724253)	24	0	0	0
<i>P. leniusculus</i>	Loch Ken, Dumfries and Galloway, Scotland. (55.0116161, - 4.0593604)	32	0	0	0

527

528 Table 2: Results of mixed effects models with significant predictors of parasitism and  
529 prevalence of parasites in crayfish populations using the Cefas dataset. Predictors are the  
530 variables that remained in the minimum adequate model. Non-significant predictors are  
531 variables removed to reach the minimum adequate model.  $X^2$  reflects the result of log  
532 likelihood test comparing minimum adequate model to null model. The term “Ecostat” is the  
533 ecological status of the sub-catchment according to the water framework directive. The term  
534 “Chemstat” is the chemical status of the sub-catchment according to the water framework  
535 directive. The terms “Ba” = *Branchiobdella astaci*, “Ph” = *Psorospermium haeckeli*, “Tc” =  
536 *Thelohania contejeani*, “BV” = bacilliform virus (putative *Nudiviridae*).

537

Response variable	Significant predictor(s)	Non-significant predictors	Model $R^2$	$X^2$	P value
Model 1: Parasite diversity (Simpson's Index)	Crayfish species	Ecostat Chemstat	0.98	7.50	<0.05

Model 2: Prevalence of bacilliform virus (both species)	Crayfish species	Ecostat Chemstat Presence of Ph, Ba, Tc Prevalence of Ph, Ba, Tc	0.24	11.30	<0.001
Model 3: Response: Prevalence of bacilliform virus ( <i>A. pallipes</i> only).	Presence of <i>Tc</i>	Ecostat Chemstat Presence of Ph, Ba Prevalence of Ph, Ba, Tc	0.27	0.01	<0.001
Model 4: Presence of crayfish plague ( <i>Aphanomyces astaci</i> ) in <i>A. pallipes</i> .	Chemstat	Ecostat	0.21	4.20	<0.05




Figure 1: Histological micrographs identifying infectious agents of crayfish. A) Low magnification view of a *Pacifastacus leniusculus* hepatopancreatic epithelial cell infected with a bacilliform virus (solid pink viroplasm in centre surrounded by cell organelles). B) High magnification view of a hepatopancreatocyte with a virally infected nucleus (arrow). C) Melanised gill tissue of an *Austropotamobius pallipes* infected with *Branchiobdella astaci*. D) Cross-section through an individual *B. astaci* parasite infecting the gill of *A. pallipes*. E) Tail muscle tissue of *A. pallipes* heavily infected with *Thelohania contejeani*. The arrow marked 'h' shows healthy, striated muscle tissue. The arrow marked 'i' shows infected muscle tissue which has been replaced with spores. F) A longitudinal section through a *Psorospermium haeckeli* sporocyst in the connective tissue of an *A. pallipes* host.




Figure 2: Presence and mean prevalence of parasites across the three population compositions. Isolated *Austropotamobius pallipes* populations (n=3 populations), isolated *Pacifastacus leniusculus* populations (n=3 populations) and *P. leniusculus* populations with recent *A. pallipes* overlap (n=3 populations). *Thelohania contejeani* (Microsporidia), Bacilliform virus (putative *Nudiviridae*) and *Branchiobdella astaci* (Annelida).