



This is a repository copy of *The myriad of complex demographic responses of terrestrial mammals to climate change and gaps of knowledge : a global analysis*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/177547/>

Version: Accepted Version

---

**Article:**

Paniw, M., James, T.D. [orcid.org/0000-0003-1363-4742](https://orcid.org/0000-0003-1363-4742), Ruth Archer, C. et al. (15 more authors) (2021) The myriad of complex demographic responses of terrestrial mammals to climate change and gaps of knowledge : a global analysis. *Journal of Animal Ecology*, 90 (6). pp. 1398-1407. ISSN 0021-8790

<https://doi.org/10.1111/1365-2656.13467>

---

This is the peer reviewed version of the following article: Paniw, M, James, TD, Ruth Archer, C, et al. The myriad of complex demographic responses of terrestrial mammals to climate change and gaps of knowledge: A global analysis. *J Anim Ecol*. 2021; 90: 1398–1407., which has been published in final form at <https://doi.org/10.1111/1365-2656.13467>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

The myriad of complex demographic responses of terrestrial  
mammals to climate change and gaps of knowledge: A global  
analysis

Submission to *Journal of Animal Ecology* for consideration as *Review*

Maria Paniw<sup>\*1,2</sup>, Tamora D. James<sup>3</sup>, C. Ruth Archer<sup>4</sup>, Gesa Römer<sup>5,6</sup>, Sam Levin<sup>7,8</sup>, Aldo Compagnoni<sup>7,8</sup>, Judy Che-Castaldo<sup>9</sup>, Joanne M. Bennett<sup>7,8,10</sup>, Andrew Mooney<sup>11</sup>, Dylan Z. Childs<sup>3</sup>, Arpat Ozgul<sup>2</sup>, Owen R. Jones<sup>5,6</sup>, Jean H. Burns<sup>12</sup>, Andrew P. Beckerman<sup>3</sup>, Abir Patwary<sup>3,13</sup>, Nora Sanchez-Gassen<sup>14</sup>, Tiffany M. Knight<sup>§7,8,15</sup> & Roberto Salguero-Gómez<sup>§13</sup>

§Shared senior co-authorship

<sup>1</sup>Ecological and Forestry Applications Research Centre (CREAF), Campus de Bellaterra (UAB) Edifici C, ES-08193 Cerdanyola del Vallès, Spain

<sup>2</sup>Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

<sup>3</sup>Department of Animal and Plant Sciences, University of Sheffield. Western Bank, Sheffield S10 2TN, UK.

<sup>4</sup>Institute of Evolutionary Ecology and Conservation Genomics, University of Ulm, Albert-Einstein-Allee 11, 89081, Ulm, Germany.

<sup>5</sup>Interdisciplinary Centre on Population Dynamics (CPop), University of Southern Denmark. Odense, Denmark.

<sup>6</sup>Department of Biology, University of Southern Denmark. Campusvej 55, 5230 Odense M, Denmark.

<sup>7</sup>Institute of Biology, Martin Luther University Halle-Wittenberg, Am Kirchtor 1, 06108, Halle (Saale), Germany

<sup>8</sup>German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany

<sup>9</sup>Alexander Center for Applied Population Biology, Conservation & Science Department, Lincoln Park Zoo, Chicago, IL 60614-4712 USA

<sup>10</sup>Centre for Applied Water Science, Institute for Applied Ecology, Faculty of Science and Technology, University of Canberra, Australia

<sup>11</sup>School of Natural Sciences, Zoology, Trinity College Dublin, Ireland

<sup>12</sup>Department of Biology, Case Western Reserve University, Cleveland, Ohio, USA.

<sup>13</sup>Department of Zoology, University of Oxford. 11a Mansfield Road, Oxford, OX1 3SZ, United Kingdom.

<sup>14</sup>Nordregio, Holmamiralens Väg 10, 11149 Stockholm, Sweden

<sup>15</sup>Department of Community Ecology, Helmholtz Centre for Environmental Research – UFZ, Theodor-Lieser-Straße 4, 06120 Halle (Saale), Germany

\*Corresponding author: Maria Paniw, Estación Biológica de Doñana (EBD-CSIC), Seville, 41001 Spain

Email: [maria.paniw@ebd.csic.es](mailto:maria.paniw@ebd.csic.es)

ORCID:

Maria Paniw 0000-0002-1949-4448; Tamora D James 0000-0003-1363-4742; Gesa Römer 0000-0002-4859-5870; Sam Levin 0000-0002-3289-9925; Aldo Compagnoni 0000-0001-8302-7492; Judy Che-Castaldo 0000-0002-9118-9202; Joanne M Bennett 0000-0002-7883-3577, Dylan Z Childs 0000-0002-0675-4933; Arpat Ozgul 0000-0001-7477-2642; Owen R Jones 0000-0001-5720-4686; Jean H Burns 0000-0001-6140-9094; Andrew P Beckerman 0000-0002-4797-9143; Tiffany M Knight 0000-0003-0318-1567; Roberto Salguero-Gómez 0000-0002-6085-4433

## Keywords

Climate vulnerability; temperature extremes; comparative demography; demographic rates; population growth rate

## Abstract

1. Approximately 25% of mammals are currently threatened with extinction, a risk that is amplified under climate change. Species persistence under climate change is determined by the combined effects of climatic factors on multiple demographic rates (survival, development, reproduction), and hence, population dynamics. Thus, to quantify which species and regions on Earth are most vulnerable to climate-driven extinction, a global understanding of how different demographic rates respond to climate is urgently needed.

2. Here, we perform a systematic review of literature on demographic responses to climate, focusing on terrestrial mammals, for which extensive demographic data are available.
3. To assess the full spectrum of responses, we synthesize information from studies that quantitatively link climate to multiple demographic rates. We find only 106 such studies, corresponding to 87 mammal species. These 87 species constitute < 1% of all terrestrial mammals.
4. Our synthesis reveals a strong mismatch between the locations of demographic studies and the regions and taxa currently recognized as most vulnerable to climate change. Surprisingly, for most mammals and regions sensitive to climate change, holistic demographic responses to climate remain unknown. At the same time, we reveal that filling this knowledge gap is critical as the effects of climate change will operate via complex demographic mechanisms: a vast majority of mammal populations display projected increases in some demographic rates but declines in others, often depending on the specific environmental context, complicating simple projections of population fates.
5. Assessments of population viability under climate change are in critical need to gather data that account for multiple demographic responses, and coordinated actions to assess demography holistically should be prioritized for mammals and other taxa.

## **Introduction**

The ca. 6,400 extant mammal species have colonized virtually all terrestrial and most aquatic habitats (Jenkins *et al.* 2013). This evolutionary success has been facilitated by the wide range of mammalian life history strategies (Healy *et al.* 2019), which enable them to cope with vastly different climates (Promislow *et al.* 1990). These strategies include extreme examples like male semelparity in some Australian marsupials, characterized by short, explosive mating seasons (Fisher *et al.* 2013), or high behavioral and demographic plasticity in long-lived primates that buffers populations from the negative effects of environmental variation (Campos *et al.* 2017). This

tremendous variation in life history strategies can be captured by differences among organisms in their rates and timing of survival, development, and reproduction (Stearns 1992). It is these demographic rates that determine population growth and thus species persistence (Morris & Doak 2002). Therefore, understanding the effects of climate drivers on the viability of natural mammal populations requires a simultaneous consideration of multiple demographic rates (Paniw *et al.* 2019).

Important efforts have been made in the last decade to increase the amount of comparative data to understand the variation in demographic rates across mammals (Conde *et al.* 2019). These data have resulted in the broader availability of open-access demographic information on populations of mammals and other taxa (Jones *et al.* 2009; Salguero-Gómez *et al.* 2016) and have produced key life-history knowledge, for instance on lifespan and mortality schedules (Conde *et al.* 2019; Jones *et al.* 2014). However, we still lack a holistic understanding of how climate drivers simultaneously affect survival, development, and reproduction in mammals worldwide. This gap is at odds with an emerging consensus that interactions among demographic rates and biotic and abiotic drivers hinder simplistic projections of persistence under climate change (Benton *et al.* 2006; Urban *et al.* 2016). For instance, a negative effect of climate on a specific demographic rate does not necessarily cause a population to go extinct, when another demographic rate responds positively to climate, or when population dynamics are mediated by interactions among climate drivers and density-dependent feedbacks (Reed *et al.* 2013; Paniw *et al.* 2019). Consequently, it is important to assess whether the complexity of demographic responses to climate is widespread among mammals and whether research quantifying such complexity is available for regions most vulnerable to climate change or for the most vulnerable species.

Here, we synthesize our understanding regarding where, which, and how mammal populations respond to climate. We searched for literature linking multiple demographic rates to climatic drivers, thus capturing the complexity of demographic responses, for 5,728 terrestrial mammal species (Myhrvold *et al.* 2015). We then performed a rigorous review of studies that matched our search

criteria and linked data from the review to information on ecoregion and species' vulnerability to climate change (Beaumont *et al.* 2011; IUCN 2019). Our review allows us to explore (i) whether mammal demographic studies are conducted in ecoregions that are most vulnerable to projected increases in temperature and precipitation extremes (Q1: *Where?*); (ii) whether potential future demographic responses to climate change reflect species' extinction risk as determined by the IUCN Red List status of mammals (Q2: *Which species?*); and (iii) through which demographic processes projected changes in climate may show negative and/or positive effects on populations (Q3: *How?*).

## **Methods**

We obtained scientific names of 5,728 terrestrial mammal species from the Amniote database (Myhrvold *et al.* 2015). For each species, we searched SCOPUS for studies (published before January 2018) that quantified demographic-rate-climate relationships (for the full list of search terms, see Supporting information S1). We also confirmed, using a subset of species, that a search in Web of Science did not produce different results (Supporting information S1). From any study that matched our search terms, we extracted information on demographic-rate-climate relationships only if (i) the study linked at least two different demographic rates (*i.e.*, survival, development/growth, or reproduction) to a climatic driver (*i.e.*, any direct or indirect measure of temperature or precipitation); and (ii) the response of a demographic rate to a climatic driver was quantified using statistical methods (*i.e.*, excluding qualitative or descriptive studies). To standardize our search protocol and to assess potential global demographic consequences of climate change, we recorded local climatic factors such as winds or storms only when these factors were related to temperature and precipitation. In addition, we only considered studies on natural populations of terrestrial mammals, or partially terrestrial mammals (e.g., polar bears), because initial results showed that there were few climate-related population studies on fully aquatic mammals, which considered distinct climatic drivers that lack future projection information (e.g., ocean circulation indices).

From all studies meeting our criteria, we extracted the effects any climatic driver had on unstructured or age- or stage-specific demographic rates. We grouped these effects as positive (*i.e.*, increased rate), negative (*i.e.*, reduced rate), no effect, or context-dependent (e.g., increased rate at low population densities and no effect at high densities; see Supporting information S1 for details). We also recorded whether climate effects were measured at annual or seasonal scales and, when applicable, the factors that mediate these effects (e.g., density, biotic interactions). In studies that indicated future projections of drivers, we recorded whether drivers were projected to increase, decrease, or show context-dependent trends.

For the 68% of studies that did not report on climate projections (Figs. S1.4 & S1.5) but assessed demographic effects of direct temperature and precipitation measures, we approximated whether future averages and extremes in these drivers were likely to increase, decrease, or remain unchanged under global climate change (Supporting information S1). To do so, we obtained monthly average temperatures and precipitation data as well as maximum and minimum monthly temperatures from 1979-2013 for all relevant study locations using *climatologies at high resolution for the earth's land surface areas* (CHELSA; ca. 1 km<sup>2</sup> resolution) (Karger *et al.* 2017). The monthly scale corresponded to the finest scale used to aggregate climatic variables for demographic analyses in the reviewed studies. We also obtained projections of these variables for 2041-2060 assuming five diverging climate models (NorESM1-M; MPI-ESM-LR; GISS-E2-R; CMCC-CM; CESM1-BGC; Supporting information S1). We averaged historical and projected climate records for each month and each of the five climate models. We then calculated changes in each of the climate variables by subtracting the monthly average historical records from the projected values. We set the change to 0 if 95 % CI of the monthly averages across the five projection models crossed corresponding average historical values. Finally, we averaged these changes across all months to obtain a unique value of projected increase, decrease, or no change in total precipitation or average and extremes in temperature. We also calculated the standard deviation of the monthly averaged historical and projected climate records and assessed the changes in this deviation using the 95 %

CI as above, which gave us a unique value of increase, decrease, or no change in the variation of the above variables.

To quantify potential demographic effects of global changes in temperature and precipitation, we linked the changes in precipitation and temperature variables obtained from CHELSA to the reviewed studies in the following way: If a study assessed seasonal/annual averages or deviations in precipitation, mean temperature, or minimum/maximum temperature but did not project future changes in these variables, we assigned future changes using our CHELSA climate data – assuming that precipitation and temperature variables used in a given study would change in the same direction as the CHELSA climatic data. From this information, we then determined whether a demographic rate would potentially decrease (e.g., where a rate has a positive response to maximum temperature and maximum temperature projected to decrease) or increase (e.g., where a rate has a positive response to maximum temperature and maximum temperature projected to increase). Unless explicitly stated otherwise in a study, we assumed that demographic rates that were not affected by a climatic variable would not change in the future, and ones that showed context-dependent responses would also likely show context-dependent responses in the future.

A full list of extracted studies and a more detailed description of the extraction protocol and climate modelling can be found in Supporting information S1 and Table S1.1. We note that the multitude of methodological approaches used to study demographic responses (e.g. correlation analyses, structured demographic models, individual-based models) and the lack of a full statistical report in some studies (Gerstner *et al.* 2017) render a full meta-analytical approach of climate-demography relationships impractical.

We determined how well our studies provided knowledge in global ecoregions with high biodiversity and high vulnerability to climate change (Q1, *Where?*) using two approaches. First, we built on the results of Beaumont and colleagues (2011), which found that temperature, unlike precipitation, is projected to become more extreme in all of the 132 terrestrial ecoregions of exceptional biodiversity (i.e., in the “Global 200”, Pimm *et al.* 2014). Specifically, the authors classified temperature



vulnerability of the G200 ecoregions using, among other measures, the weighted average minimum monthly distance in temperatures (under the A2 climate model ensemble) from the mean of the 1961-1990 baseline. The higher the distance, the more vulnerable an ecoregion (Beaumont *et al.* 2011). We quantified the proportion of our studies that were conducted in these highly biodiverse ecoregions, and the proportion of these studies that assess the demographic effects of temperature increases. Further, we quantified and visualized whether demographic rates were reported to be decreasing (-), not changing (0), or increasing (+) under increasing temperatures across ecoregion temperature vulnerability scores.

Second, to assess how well our studies provided knowledge in areas projected to experience the highest changes in temperature and precipitation, regardless of whether they have high biodiversity, we obtained gridded global data on annual average surface air temperature and total precipitation anomalies, the latter driven by seasonal variation (Collins *et al.* 2013), for the period (2030-2049) compared to present day (1986-2005). Climatic anomalies were based on the Representative greenhouse-gas Concentration Pathway scenario 6.0 (NCAR 2012), which is similar to the previous SRES A2 climate model ensemble (Moss *et al.* 2008). We matched the geographic locations of the studied mammal populations to the gridded global anomalies.

To assess whether demographic responses to projected changes in climate agree with the International Union for Conservation of Nature and Natural Resources (IUCN) Red List status of mammals (Q2: *Which species?*), we obtained IUCN assessments (including threats) for all species identified in the literature review. We used the R package *rredlist* (Chamberlain 2020) to access the IUCN Red List database and extract available information on whether the species are listed in the database, and, if so, what status they are assigned to and whether climate change is listed as an existing or potential threat. Lastly, for each IUCN threat category, we quantified the diversity of projected positive (+), negative (-), and context-dependent (CD) climatic effects across different demographic rates and life-cycle stages modelled for the mammal populations (Q3: *How?*).

## Results

For most terrestrial mammals, we did not find studies on detailed demographic responses to climate. We extracted information on climate-demography relationships from 106 studies, for a total of 87 mammal species, that quantified simultaneous responses to climate in at least two different demographic rates. These studies span 12 biomes, with the exception of tropical and subtropical coniferous forests and mangroves (Fig. S1.1). Overall, more studies assess only the direct effects of precipitation ( $n = 42$ ) than the direct effects of temperature ( $n = 11$ ) (Fig. S1.2); and in eight of the 106 studies, only indirect effects are assessed via indices such as the North Atlantic Oscillation (NAO) or El Niño–Southern Oscillation (ENSO), while only seven studies assess the combined demographic effects of temperature, precipitation and climate indices (Fig. S1.2). A majority of studies (71%) only test demographic responses to seasonal ( $n = 40$ ) or annual ( $n = 36$ ) averages or sums in climate variables, while 16 % of studies assess responses to climate extremes (maxima, minima, or deviation from averages) as well as averages (Fig. S1.3). Few studies (10%) test how different climatic drivers interact with one another, approximately half (55%) test for the effects of density dependence on demographic rates, and an additional 27% test for effects of non-climatic drivers other than population density (e.g. predation, food availability). Studies that test for driver interactions typically find context-dependent climate effects in some demographic rates, with 15% of all reviewed studies reporting such effects.

In addressing “Q1: *Where?*”, our synthesis reveals that few demographic studies are conducted in ecoregions that are both biodiverse and highly vulnerable to climate change (Olson and Dinerstein 2002). We find that 42 out of the 106 demographic studies were conducted in one of the G200 ecoregions (Fig. 1). However, only 13 of these studies assess the demographic effects of temperature increases. In addition, no study has examined the responses of different demographic rates in ecoregions with the highest vulnerability scores (e.g., the Central Congo Basin; darkest red in Fig. 1). Only one study, which included three primate species, assessed temperature effects in

a G200 ecoregion with relatively high vulnerability to climate change; and found no effects (Fig. 1 insert) likely due to the primates' behavioural and physiological flexibility before climate variation (Campos *et al.* 2017). Similarly, our synthesis reveals that few demographic studies are conducted in ecoregions projected to have the greatest anomalies in annual temperature and precipitation (Fig. S1.6). Only one study (Hunter *et al.* 2010) assessed climate-demography relationships, of polar bears (*Ursus maritimus*), in an arctic region with highest projected increases in average temperature ( $> 2.5$  °C; Fig. S1.6a). Lastly, no study has examined demographic responses to precipitation in ecoregions with highest projected decreases in precipitation. Only two studies (Campos *et al.* 2017; Marshall *et al.* 2016), considering a banded mongoose and two primate populations, examined demographic responses to precipitation in regions in East Africa, where precipitation is projected to increase substantially ( $> 100$  mm; Fig. S1.6).

In addition to an ecoregion bias, in answering “Q2: *Which species?*”, we highlight that demographic analyses suffer from a taxonomic bias too. Studies linking multiple demographic rates to climatic drivers are primarily performed in regions with a relatively low mammal richness and on species that are not currently vulnerable to climate change (Fig. 2), based on IUCN classifications. Indeed, the IUCN has identified at least 17% of listed vertebrates to be sensitive to climate change, *i.e.*, decreasing in numbers or losing habitat under changes in temperature and precipitation regimes due to elevated atmospheric CO<sub>2</sub> levels (Marris 2008). Our synthesis reveals that only 4% of all mammals assessed as climate sensitive by the IUCN have detailed studies linking demography to climate (*i.e.*, 13% of studies we assessed). Interestingly, the proportion of demographic rates per study that may decline under global climate change ( $0.31, \pm 0.10$  S.E.) is highest for species that have been flagged by the IUCN as climate sensitive. However, this proportion is followed closely by species for which climate change is not considered a threat by the IUCN (Fig. 2 insert).

In addressing “Q3: *How?*”, our synthesis reveals that complex demographic responses to climate are prevalent. Only eight (7%) of the 106 studies report unidirectional (all positive) responses of

demographic rates to climatic drivers, while in 19 cases no effect of climate on any demographic rate is reported (Figs. S1.4 & S1.5). For the vast majority of species, effects of climate reported in the studies (79%) and projected under global change as presented in the studies or approximated in our analysis (75%) can be positive, negative or context-dependent depending on the population studied and the specific driver, demographic rate, and life-cycle stage considered (examples in Fig. 4). In addition, 13 studies assess the effects of climate on population growth rates in addition to underlying demographic rates (Fig. S1.4, Table S1.1). These examples show that population responses are not readily predictable from a single demographic rate when multiple climatic drivers and their interactions with biotic drivers affect demography. For instance, Lawler and co-authors (2009) show that survival, stage transitions, and reproduction all determine how the population growth rate of the long-lived Verreaux's sifakas (*Propithecus verreauxi verreauxi*) responds to annual rainfall.

## Discussion

Our synthesis shows that gaining a global understanding of the mammals that are most vulnerable to climate-driven extinction will require filling important knowledge gaps. Studies on climate effects across the whole life cycle of populations are needed in regions that are highly susceptible to climate change, many of which are biodiversity hotspots (Beaumont *et al.* 2011). Many of these ecoregions are not only threatened by increases in climate extremes (Beaumont *et al.* 2011; Stott 2016) but also face strong pressures on biodiversity from direct human activities (Venter *et al.* 2016). A majority of studies linking demographic rates to climate in vulnerable ecoregions and elsewhere nevertheless focus on demographic responses to changes in average climate and do not account for interactions with non-climate drivers. It is possible that publication bias plays a part in our observations of these omissions, as authors may, for instance, explore interactions but not mention this in their published study due to insignificant or inconclusive results (Jennions and Møller 2002). At the same time, population dynamics are typically driven by compound effects of interacting climatic and non-climatic drivers (Benton *et al.* 2006; Zarnetske *et al.* 2012), which are

projected to become more extreme under climate change (Zscheischler *et al.* 2018). A stronger focus on extremes and driver interactions is therefore required to improve our estimates of population viability.

Our analysis also reveals that studies assessing responses to climate across different demographic rates are required for most mammals considered climate-sensitive by the IUCN. These results support recent calls to gain a better understanding of the complex effects of climate on population viability (e.g., Doak and Morris 2010; Cahill *et al.* 2013; Román-Palacios and Wiens 2020), allowing this threat to be understood and potentially mitigated through conservation. On the other hand, given that a large number of mammals not considered to be climate sensitive may actually show negative demographic responses to climate change, our results support the need for current IUCN efforts to re-evaluate the importance of climate as an extinction threat to mammals (Foden and Young 2016). It is noteworthy that the majority of demographic studies on mammals do not project demographic effects of climate change. Our simplified and broad approximation of climate change for these studies allowed for global synthesis in this review, but we note that this approximation does not match the spatiotemporal resolution of the climatic drivers assessed in the relevant studies, averages over seasonal climatic patterns, and does not allow examination of interactions with other important factors, which can exacerbate or dampen large-scale climate trends (van de Pol *et al.* 2013). Thus, an important challenge for future studies is to project changes in local climatic variables that have been considered important drivers of demographic rates (Snover *et al.* 2013).

By focusing on studies that have assessed several demographic responses to climate, we necessarily limited the number of taxa in our review. In fact, we identified at least 111 more studies on 68 additional species that only assessed climatic effects on single demographic rates. We stress here that we do not question the validity of such studies when population dynamics can be accurately predicted from the changes in one demographic rate with a uniquely high contribution to the population growth rate. However, population responses to climate are typically determined

by the covariation among multiple demographic rates, which itself is often mediated by a myriad of interacting biotic and abiotic factors (e.g., Reed *et al.* 2013). For instance, impalas (*Aepyceros melampus*), which the IUCN characterizes as threatened by drought (Table S1.1), may show positive or negative responses in survival and reproductive success under rainfall scarcity depending on the seasonal patterning of rainfall and population density (Ogutu *et al.* 2012). Similarly, meerkats (*Suricata suricatta*), which currently face no threats according to the IUCN, show both positive and negative responses to precipitation across several critical demographic rates depending on social stage, group size, and temperature extremes (Ozgul *et al.* 2014; Paniw *et al.* 2019). Therefore, as a cooperative breeder, meerkats may be vulnerable to compound effects of increases in seasonal climatic extremes that decrease group densities (Paniw *et al.* 2019). Such complex demographic responses make it challenging to project species' fates under climate change because the future of populations cannot be accurately determined from single demographic rates (Urban *et al.* 2016).

As consistently negative demographic responses to climate change are only relatively common in species the IUCN flagged as climate sensitive, our results suggest that complexity of demographic responses may buffer populations from adverse climate effects (González-Suárez and Revilla 2013). One important aspect of such complexity is that different populations may show starkly different responses to climate, and less severe demographic effects in one population may compensate for climate-related losses of individuals in another population (Schindler *et al.* 2015; Villellas *et al.* 2015). In addition, dispersal can stabilize decreasing populations and allow mammals to track new suitable habitats, and may itself be strongly mediated by climate (Parmesan and Yohe 2003; Root 2003; Travis *et al.* 2013). However, these important spatial aspects of demographic complexity have thus far been rarely assessed, with only three of the 106 studies quantitatively linking dispersal (emigration or immigration), in addition to local survival or reproduction, to climate (Kreuzer and Huntly 2003; Ozgul *et al.* 2014; Wauters *et al.* 2004). Therefore, despite the challenges involved in collecting long-term demographic parameters, including dispersal, across

the entire life cycle (Foden and Young 2016), the mechanistic insights gained from such parameters will be invaluable to understand the drivers of biodiversity loss under climate change (Travis *et al.* 2013; Urban *et al.* 2016).

In conclusion, our current mechanistic knowledge on mammal responses to climate change would benefit from strategic studies that fill the knowledge gaps identified here. Our take-home message echoes recent calls for more integrative climate-demography studies in bird populations (Zurell 2017), along with numerous studies showing the benefits of such integrative research for evidence-based conservation (Møller *et al.* 2012). Other taxa, such as insects or amphibians, are far less well studied demographically than mammals or birds, but recent evidence suggests that demographic responses to climate change are complex and that understanding this complexity is key to conservation planning (Boggs 2016; Muths *et al.* 2017). Along with recent calls for a renewed global effort to collect natural-history information (Urban *et al.* 2016), we therefore advocate for a coordinated effort to collect and model multiple demographic responses to climate across various taxa.

## **Acknowledgments**

This work was supported by the working group proposal “sAPROPOS: Analysis of PROjections of POpulationS” funded to RS-G by sDiv, the Synthesis Centre of the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig (DFG FZT 118-202548816). MP was supported by an ERC Starting Grant (33785) and a Swiss National Science Foundation Grant (31003A\_182286) to AO; and by a Spanish Ministry of Economy and Competitiveness Juan de la Cierva-Formación grant FJCI-2017-32893. RS-G was also supported by a NERC grant (R/142195-11-1), and JC, ORJ, and RS-G by an NSF award (DBI-1661342). TDJ was supported by NERC (ACCE DTP; NE/L002450/1), with additional CASE funding from the World Parrot Trust. We thank the Alexander von Humboldt foundation (award to TMK) that supported a retreat to write a first version of this manuscript.

376

377 **Data availability and archiving**

378 The data that support the findings in this study are available in the Supporting information S1 and  
379 Table S1.1. The code that supports the findings in this study is available in the Supplementary  
380 Online Materials, `climate_change_analyses_mammal_review.R`. Upon acceptance, all  
381 data and code will be archived on *Dryad* and will be freely accessible.

382

383 **Author Contributions**

384 MP, TDJ, GR, and RS-G devised the overall manuscript. MP and TDJ designed the literature review  
385 protocol, which was then implemented by MP, TDJ, GR, CRA, SL, AM, JC, NSG, JMB, and AP.  
386 The climatic data were derived by AC. The first draft of the manuscript was written by MP and RS-  
387 G, and all co-authors contributed to the final manuscript. See Table S1.2 for further specifics  
388 regarding task contributions.

389 **References**

390 Beaumont LJ, Pitman A, Perkins S, *et al.* 2011. Impacts of climate change on the world's most  
391 exceptional ecoregions. *Proc Natl Acad Sci USA* **108**: 2306–11.

392 Benton TG, Plaistow SJ and Coulson TN. 2006. Complex population dynamics and complex  
393 causation: devils, details and demography. *Proc Biol Sci* **273**: 1173–81.

394 Boggs CL. 2016. The fingerprints of global climate change on insect populations. *Curr Opin*  
395 *Insect Sci* **17**: 69-73.

396 Cahill AE, Aiello-Lammens M, Fisher-Reid C, *et al.* 2013. How does climate change cause  
397 extinction? *Proc R Soc Biol Sci* **280**: 20121890.

398 Campos FA, Morris WF, Alberts SC, *et al.* 2017. Does climate variability influence the  
399 demography of wild primates? Evidence from long-term life-history data in seven species.  
400 *Glob Change Biol* **23**: 4907–21.



401 Chamberlain S. 2020. *rredlist: 'IUCN' Red List Client*. R package version 0.6.0. Available at:  
 402 <https://CRAN.R-project.org/package=rredlist>.

403 Collins MR, Knutti J, Arblaster JL, *et al.* 2013. *Long-term climate change: projections,*  
 404 *commitments and irreversibility*. In: Climate Change 2013: The Physical Science Basis.  
 405 Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental  
 406 Panel on Climate Change [Stocker TF, Qin D, Plattner G-K, *et al.* (eds.)]. Cambridge  
 407 University Press: Cambridge.

408 Conde DA, Staerk J, Colchero F, *et al.* 2019. Data gaps and opportunities for comparative and  
 409 conservation biology. *Proc Natl Acad Sci USA* **116**: 9658–64.

410 Doak DF and Morris WF. 2010. Demographic compensation and tipping points in climate-induced  
 411 range shifts. *Nature* **467**: 959–62.

412 Fisher DO, Dickman CR, Jones ME, *et al.* 2013. Sperm competition drives the evolution of  
 413 suicidal reproduction in mammals. *Proc Natl Acad Sci USA* **110**: 17910–14.

414 Foden WB and Young BE. 2016. *IUCN SSC Guidelines for Assessing Species' Vulnerability to*  
 415 *Climate Change*. UCN Cambridge, England and Gland, Switzerland.

416 Gerstner K, Moreno-Mateos D, Grevitch J, *et al.* 2017. Will your paper be used in a meta-  
 417 analysis? Make the reach of your research broader and longer lasting. *Methods Ecol Evol* **8**:  
 418 777–784.

419 González-Suárez M and Revilla E. 2013. Variability in life-history and ecological traits is a buffer  
 420 against extinction in mammals. *Ecol Lett* **16**: 242–51.

421 Healy K, Ezard THG, Jones OR, *et al.* 2019. Animal life history is shaped by the pace of life and  
 422 the distribution of age-specific mortality and reproduction. *Nat Ecol Evol* **3**: 1217–24.

423 Hunter CM, Caswell H, Runge MC, *et al.* 2010. Climate change threatens polar bear populations:

424 a stochastic demographic analysis. *Ecology* **91**: 2883-97.

425 IUCN (International Union for Conservation of Nature). 2019. The IUCN red list of threatened  
 426 species. <http://www.iucnredlist.org>. Viewed 25 Jan 2020.

427 Jenkins CN, Pimm SL and Joppa L N. 2013. Global patterns of terrestrial vertebrate diversity and  
 428 conservation. *Proc Natl Acad Sci USA* **110**: E2602–10.

429 Jennions M and Møller A. 2002. Publication bias in ecology and evolution: An empirical  
 430 assessment using the 'trim and fill' method. *Biol Rev*, **77**: 211-222.

431 Jones KE, Bielby J, Cardillo M, *et al.* 2009. PanTHERIA: a species-level database of life history,  
 432 ecology, and geography of extant and recently extinct mammals: Ecological Archives E090-  
 433 184. *Ecology* **90**: 2648–48.

434 Jones OR, Scheuerlein A, Salguero-Gómez R, *et al.* 2014. Diversity of ageing across the tree of  
 435 life. *Nature* **505**: 169–173.

436 Kreuzer MP and Huntly NJ. 2003. Habitat-specific demography: evidence for source-sink  
 437 population structure in a mammal, the pika. *Oecologia* **134**: 343-49.

438 Lawler RR, Caswell H, Richard AF, *et al.* 2009. Demography of Verreaux's sifaka in a stochastic  
 439 rainfall environment. *Oecologia* **161**: 491–504.

440 Marris E. 2008. Pre-emptive strike: outwitting extinction. *Nat Clim Change* **1**: 140–41.

441 Marshall HH, Sanderson JL, Mwanghuya F, *et al.* 2016. Variable ecological conditions promote  
 442 male helping by changing banded mongoose group composition. *Behav Ecol* **27**: 978–87.

443 Møller AP, Fiedler W, and Berthold P. 2012. *Effects of Climate Change on Birds*. Oxford  
 444 University Press: Oxford, England.

445 Morris WF and Doak DF. 2002. *Quantitative Conservation Biology: Theory and Practice of*  
 446 *Population Viability Analysis*. Oxford University Press: Oxford, England.

447 Moss R, Babiker M, Brinkman S, *et al.* 2008. *Towards New Scenarios for Analysis of Emissions,*  
448 *Climate Change, Impacts, and Response Strategies.* Geneva: Intergovernmental Panel on  
449 Climate Change.

450 Muths E, Chambert T, Schmidt BR *et al.* 2017. Heterogeneous responses of temperate-zone  
451 amphibian populations to climate change complicates conservation planning. *Sci Rep* **7**:  
452 17102.

453 Myhrvold NP, Baldrige E, Chan B, *et al.* 2015. An amniote life-history database to perform  
454 comparative analyses with birds, mammals, and reptiles: Ecological Archives E096-269.  
455 *Ecology* **96**: 3109.

456 NCAR GIS Program. 2012. Climate Change Scenarios, version 2.0. Community Climate System  
457 Model, June 2004 version 3.0. <http://www.cesm.ucar.edu/models/ccsm3.0/> was used to  
458 derive data products. NCAR/UCAR. URL: <http://www.gisclimatechange.org>. Accessed  
459 December 12, 2020.

460 Ogutu JO, Piepho HP and Kanga E. 2012. Dynamics of an insularized and compressed impala  
461 population: rainfall, temperature and density influences. *Open J Ecol* **5**: 1–17.

462 Olson DM and Dinerstein E. 2002. The Global 200: Priority ecoregions for global conservation.  
463 *Ann Mo Bot Gard* **89**: 199–224.

464 Ozgul A, Bateman AW, English S, *et al.* 2014. Linking body mass and group dynamics in an  
465 obligate cooperative breeder. *J Anim Ecol* **83**: 1357–66.

466 Paniw M, Maag N, Cozzi G, *et al.* 2019. Life history responses of meerkats to seasonal changes  
467 in extreme environments. *Science* **363**: 631–35.

468 Parmesan C and Yohe G. 2003. A globally coherent fingerprint of climate change impacts across  
469 natural systems. *Nature* **421**: 37-42.

470 Pimm SL, Jenkins CN, Abell R, *et al.* 2014. The biodiversity of species and their rates of  
471 extinction, distribution, and protection. *Science* **344**: 1246752.

472 Promislow DEL and Harvey PH. 1990. Living fast and dying young: A comparative analysis of life-  
473 history variation among mammals. *J Zool* **220**: 417–37.

474 Reed TE, Grøtan V, Jenouvrier S, *et al.* 2013. Population growth in a wild bird is buffered against  
475 phenological mismatch. *Science* **340**: 488–91.

476 Román-Palacios C and Wiens JJ. 2020. Recent responses to climate change reveal the drivers of  
477 species extinction and survival. *Proc Natl Acad Sci USA* **117**: 4211-17.

478 Root TL, Price JT, Hall KR, *et al.* 2003. Fingerprints of global warming on wild animals and plants.  
479 *Nature* **421**: 57-60.

480 Salguero-Gómez R, Jones OR, Archer CR, *et al.* 2016. COMADRE: a global data base of animal  
481 demography. *J Anim Ecol* **85**: 371–84.

482 Schindler DE, Armstrong JB, and Reed TE. 2015. The portfolio concept in ecology and evolution.  
483 *Frontiers Ecol Evol* **13**: 257-263.

484 Snover AM, Mantua NJ, Littell JS, *et al.* 2013. Choosing and using climate-change scenarios for  
485 ecological-impact assessments and conservation decisions. *Cons Biol* **27**: 1147-57.

486 Stearns SC. 1992. *The Evolution of Life Histories*. Oxford: Oxford University Press.

487 Stott P. 2016. How climate change affects extreme weather events. *Science* **352**: 1517-1518.

488 Travis JM, Delgado M, Bocedi G, *et al.* 2013. Dispersal and species' responses to climate  
489 change. *Oikos* **122**: 1532-40.

490 Urban MC, Bocedi G, Hendry AP, *et al.* 2016. Improving the forecast for biodiversity under  
491 climate change. *Science* **353**: aad8466.

- 492 Van de Pol M, Brouwer L, Brooker LC, et al. 2013. Problems with using large-scale oceanic  
493 climate indices to compare climatic sensitivities across populations and species. *Ecography*  
494 **36**: 249-55.
- 495 Venter O, Sanderson EW, Magrach A, et al. 2016. Sixteen years of change in the global  
496 terrestrial human footprint and implications for biodiversity conservation. *Nat Comm* **7**:  
497 12558.
- 498 Villellas J, Doak DF, García MB, and Morris WF. 2015. Demographic compensation among  
499 populations: what is it, how does it arise and what are its implications? *Ecol Lett* **11**: 1139-  
500 1152.
- 501 Wauters LA, Matthyssen E, Adriaensen F, and Tosi G. 2004. Within-sex density dependence and  
502 population dynamics of red squirrels *Sciurus vulgaris*. *J Anim Ecol* **73**: 11-25.
- 503 Zarnetske P L, Skelly DK and Urban MC. 2012. Biotic multipliers of climate change. *Science* **336**:  
504 1516–18.
- 505 Zscheischler J, Westra S, van den Hurk BJJM, et al. 2018. Future climate risk from compound  
506 events. *Nat Clim Change* **8**: 469–77.
- 507 Zurell D. 2017. Integrating demography, dispersal and interspecific interactions into bird  
508 distribution models. *J Avian Biol* **48**: 1505-16.

509 **Supplementary Information** is available in the online version of the paper.

510

## 511 **Figure legends**

512

513 **Figure 1.** Global distribution of 106 mammal studies (grey points) that have assessed multiple  
514 demographic responses to climate across the species' life cycles. Point size indicates number of  
515 relationships between climatic drivers and age/stage-specific demographic rates (survival,

development, and/or reproduction). Red-scale map background indicates projected climate-change vulnerability for the most biodiverse (G200) ecoregions, with redder colours indicating higher increase in extreme temperatures. Left insert shows number of demographic rates decreasing (-), not changing (0), or increasing (+) under increasing temperatures as function of ecoregion vulnerability. Shading on insert indicates total number of demographic rates linked to temperature per ecoregion vulnerability level.


**Figure 2.** Global distribution of mammals (points) with available information on climate-demography relationships. Point and bar colours indicate levels of threat assessment by the IUCN (No IUCN - species not assessed; No T - species assessed and currently faces no threats; No CT - climate change not considered a threat; CT - climate change considered a threat). Darker background on the map indicates higher mammal richness (number of species). Bottom-left insert displays the mean proportion of demographic rates per mammal population  $\pm$  S.E. (error bars) that may decrease under projected global climate change in different IUCN categories. Total number of populations with at least one decreasing rate per threat level are indicated above the bars. Species highlighted in Figure 4 are mapped here.

**Figure 3.** Summary potential demographic responses under projected global changes in temperature and precipitation across IUCN threat categories. Out of total number of populations reviewed per IUCN threat category (indicated next to bars), we show the proportion of populations where increasing or decreasing (+/-) responses to climate are projected for different ages/stages of the same demographic rate ("within"; e.g. pre-reproductive versus reproductive stage), or for different demographic rates ("among", e.g. survival versus reproduction). Categories include No IUCN - species not assessed; No T - species assessed and currently faces no threats; No CT - climate change is not considered a threat; CT - climate change is considered a threat.

**Figure 4.** Detailed potential demographic responses to climate change for 11 example species. Demographic rates include survival (S), probability of reproducing and reproductive output (R), and growth and development (G), which can show only positive (+), only negative (-), context-dependent (CD; both positive and negative, depending on interactions with other drivers), or no (0) responses in the future. The number of instances that responses are recorded include all stage-specific and climate driver-specific combinations. From top left to bottom right, the species include Soay sheep (*Ovis aries*), agile antechinus (*Antechinus agilis*), yellow-bellied marmot (*Marmota flaviventer*), meerkat (*Suricata suricatta*), pika (*Ochotona curzoniae*), long-tailed wattled bat (*Chalinolobus tuberculatus*), Milne-Edwards's sifaka (*Propithecus edwardsi*), northern muriqui (*Brachyteles hypoxanthus*), Saiga antelope (*Saiga tatarica*), impala (*Aepyceros melampus*), and black-flanked rock-wallaby (*Petrogale lateralis*).

560


561 **Figure 1**



562

563

564 **Figure 2**




565

566

567



569



570

