



UNIVERSITY OF LEEDS

This is a repository copy of *The role of low carbon and high carbon materials in carbon neutrality science and carbon economics*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/177572/>

Version: Accepted Version

---

**Article:**

Grainger, A [orcid.org/0000-0001-8803-6013](https://orcid.org/0000-0001-8803-6013) and Smith, G (2021) The role of low carbon and high carbon materials in carbon neutrality science and carbon economics. *Current Opinion in Environmental Sustainability*, 49. pp. 164-189. ISSN 1877-3435

<https://doi.org/10.1016/j.cosust.2021.06.006>

---

© 2021, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# **The Role of Low Carbon and High Carbon Materials in Carbon Neutrality Science and Carbon Economics**

Alan Grainger

School of Geography, University of Leeds, Leeds LS2 9JT, UK.

George Smith

Trinity College, Broad Street, Oxford OX1 3BH, UK.

## **Abstract**

Focusing on switching the energy inputs to economies from fossil fuels to renewable energy and neglecting material outputs gives an over-optimistic picture of achieving carbon neutrality. We propose a set of equations that integrate analysis of energy and materials, provide a framework for a new carbon neutrality science, and lead to three carbon neutrality conditions. The equations are applied to low carbon materials, such as metals, and high carbon materials, such as wood. Refining carbon is the key carbon dioxide (CO<sub>2</sub>) emissions source to minimize for steel and aluminium, but slow technological change could create a 'carbon neutrality gap' by 2050. This will increase in size unless forest expansion is accelerated to offset remaining CO<sub>2</sub> emissions. Principles of a new carbon economics are proposed and applied. Policy priorities include integrating energy and materials in carbon neutrality strategies, strengthening carbon reporting standards, establishing national wood products databases, increasing afforestation rates, and controlling deforestation.

Key words: carbon neutrality; net zero carbon emissions; Life Cycle Analysis; materials; bio-based economy; carbon economics

## Introduction

Carbon neutrality, in which emissions of carbon dioxide into the atmosphere are offset by its removal from the atmosphere, could become the leading global environmental goal of the 21st Century. Concern about changes in global climate caused by the rising atmospheric concentration of carbon dioxide (CO<sub>2</sub>) (and other greenhouse gases) (1) has led a growing number of countries to commit themselves to carbon neutrality (2). In 2019 the UK pledged to achieve this goal, which is also called net zero carbon emissions, by 2050 (3). It was followed in 2020 by Japan (4), and by China, whose target is 2060 (5). Later in 2020 the Secretary General of the United Nations called on all governments to “declare a Climate Emergency in their countries until carbon neutrality is reached” (6).

One constraint on realizing this goal is that national strategies and academic research have so far focused on switching the *energy inputs* to economies from fossil fuels to renewable sources, and neglected the *material outputs* of these economies. Whereas 35% of the 49 Gigatonnes of CO<sub>2</sub> equivalent (Gt CO<sub>2</sub> eq) of all annual greenhouse gas emissions are directly linked to energy production, industry accounts for 21% (7) and too little attention has been paid to reducing emissions from using fossil fuels in “energy-intensive industries” which produce iron and steel, aluminium, chemicals, cement and forest products (8-12). Society depends on these materials, e.g. to construct buildings and vehicles which account for another 6% and 14% of emissions, respectively (7). As these and other material uses influence *demand* for energy, concentrating on energy *supply* is rather one-sided, so integrating the analysis of materials and energy is crucial to gaining a better understanding of likely future trends in CO<sub>2</sub> emissions from using fossil fuels. Yet only 6% of the 2,265 journal papers that we analyse here focus on materials, and they have been published in multiple, weakly interacting, scientific literatures (13). To fill this gap, bridge these literatures and contribute to the carbon neutrality literature, in this paper we show how trends in producing and using materials could promote or delay the achievement of carbon neutrality.

Materials differ in how their manufacture depends on fossil fuels. *Low carbon materials*, such as metals, have a low carbon content but fossil fuels consumed in their production and use emit much CO<sub>2</sub>, so this dependence must be changed to reduce total CO<sub>2</sub> emissions. Wood products, on the other hand, are *high carbon materials* that are rich in carbon sequestered from the atmosphere by forests and need less energy for their production. Forests, and carbon capture and storage technologies that store carbon dioxide in underground reservoirs or as carbonated minerals (14), are central to achieving carbon neutrality by offsetting CO<sub>2</sub> emissions (15). If forests are to be sustainable carbon sinks they should be managed sustainably, their trees regularly felled and replanted, and the carbon in harvested trees stored for long periods in wood products. To transfer large amounts of CO<sub>2</sub> from the atmosphere to the Earth the world will actually depend *more* on carbon, not less, so it is misleading to talk about a “low carbon economy” (16) – or even a “post-carbon economy” (17). Here we compare the main sources of CO<sub>2</sub> emitted in producing and using low and high carbon materials, and show how to reduce them.

Scientists have struggled to keep pace in conceptualizing novel environmental concepts (such as sustainable development) that originate in the government arena or outside it in civil society (18), and the same applies to carbon neutrality. By using physical variables and equations, a new *carbon neutrality science* could integrate energy and materials analysis, provide a common language for all carbon literatures, test hypotheses about alternative *carbon transition paths* (19-20), and support the political process of carbon neutrality by measuring progress

towards achieving carbon neutrality (and later zero carbon emissions). This paper identifies ten existing equations in carbon literatures, and extends life cycle analysis (21) to propose eight more equations for the new science and three conditions for carbon neutrality. It also proposes seven principles of a new *carbon economics* which can explain how carbon transition paths arise and how to manage the emerging carbon economy.

This paper has five sections. The first reviews key literatures. The second proposes new carbon neutrality equations and conditions, which are used in the third and fourth sections to evaluate the roles of low carbon and high carbon materials, respectively, in achieving carbon neutrality. The last section proposes and applies some principles of carbon economics.

## Key literatures

Research into reducing anthropogenic CO<sub>2</sub> emissions has grown rapidly since the year 2000 and is reported in seven literatures: decarbonization, low carbon economy, carbon transition, low carbon development, carbon neutrality, zero net carbon emissions and carbon footprint (Fig. 1). This section identifies equations published in these literatures and the relative priorities which each literature gives to energy and materials. Generic insights are provided by literatures on the circular economy and life cycle analysis, so these are reviewed too.

### *Global carbon budget*

Empirical analysis in this paper is framed by two equations used by the Intergovernmental Panel on Climate Change to represent net CO<sub>2</sub> emissions into the atmosphere ( $C_n$ ):

$$1. \quad C_n = C_{\text{ffc}} - C_{\text{oc}} + (C_{\text{luc}} - C_{\text{terr}})$$

where  $C_{\text{ffc}}$  refers to emissions from fossil fuel use and cement production,  $C_{\text{oc}}$  ocean uptake,  $C_{\text{luc}}$  net land use change emissions and  $C_{\text{terr}}$  other terrestrial uptake (22); and

$$2. \quad C_n = (C_{\text{end}} + C_{\text{eno}} + C_{\text{ind}} + C_{\text{tran}} + C_{\text{bld}}) - C_{\text{oc}} + C_{\text{afolu}}$$

where  $C_{\text{end}}$  denotes direct emissions from energy production,  $C_{\text{eno}}$  other energy emissions, e.g. in blast furnaces,  $C_{\text{ind}}$  emissions from industry,  $C_{\text{tran}}$  emissions from transport,  $C_{\text{bld}}$  emissions from buildings, and  $C_{\text{afolu}}$  net emissions from agriculture, forestry and other land use (7).

### *Decarbonization*

*Decarbonization* is the process by which energy generation and national economies become less dependent on the fossil fuels whose use emits CO<sub>2</sub> (23). "Deep decarbonization" requires a sharp fall in emissions (24) or structural changes in some economic sectors (25).

A quarter of a sample of 2,265 papers published in international peer-reviewed journals between 1995 and 2020 that we found in a Google Scholar search have 'decarbonization' or related words in their titles. The shares of papers on energy (43%), transport (12%) and buildings (6%) are commensurate with their shares of all direct greenhouse gas emissions (7), but producing materials only accounts for 5% and other industrial processes 2% (Table 1).

Early research led to the "Kaya identity", which "decomposes" the underlying causes of trends in carbon dioxide emissions from energy use ( $C_{\text{en}}$ ) into several "drivers", namely population

(P), gross domestic product (GDP), *energy intensity of GDP* (energy (E) used per unit GDP), and *carbon intensity of energy* (carbon emitted per unit of energy) (26):

$$3. \quad C_{en} = P \times (GDP/P) \times (E/GDP) \times (C_{en}/E)$$

This identity is still widely used (27), but has been extended by including CO<sub>2</sub> emissions and sequestration linked to "non-energy factors" (28) to give net national terrestrial CO<sub>2</sub> emissions (C<sub>nt</sub>):

$$4. \quad C_{nt} = C_{en} + (T_{ag} + T_{res} + T_{seq}) + (P_{sh} + P_{av}) + (C_f + C_{gs}) + (N_{non} + N_{seq} + N_{cr})$$

These factors include "territorial emissions" within a country from agriculture (T<sub>ag</sub>) and cement and fertilizer production etc. (T<sub>res</sub>), and net sequestration from land use change (T<sub>seq</sub>); "production emissions" from shipping (P<sub>sh</sub>) and aviation (P<sub>av</sub>); "consumption emissions" in imports of food (C<sub>f</sub>) and other goods and services (C<sub>gs</sub>); other aviation albedo effects (N<sub>non</sub>), other forms of sequestration (N<sub>seq</sub>), and sequestration funded overseas (N<sub>cr</sub>).

Modelling research shows that, if decarbonization is rapid, the delay between the time when global CO<sub>2</sub> emissions peak (t<sub>1</sub>) and the time when atmospheric CO<sub>2</sub> concentration peaks (t<sub>2</sub>) is related to the ratio between the rates of change in CO<sub>2</sub> emissions before (m\*<sub>t</sub>) and after (m\*<sub>d</sub>) peak emissions (29):

$$5. \quad t_2 - t_1 = t_1 \left( \frac{-m^*_t}{m^*_d} \right)$$

### *Low carbon economy*

A *low carbon economy* is the outcome of a decarbonization process, and was first recognized officially by the UK government in 2003 (30). It initially described an ideal scenario for reducing national CO<sub>2</sub> emissions to a low level (16), and this definition is still used (31). Exactly how "low" emissions must become is rarely stated, but a more formal definition, an economy in which greenhouse gas emissions are "decoupled" from economic growth (32), implies that emissions eventually decline to zero.

A third of papers are economic studies and a quarter have a national focus (Table 1). Papers on energy (11%) again dominate those on materials (2%). The recent derivation in this literature of an equivalent to the Kaya identity (33) from the same original "IPAT" source model (34) illustrates the compartmentalization of carbon literatures.

### *Carbon transition*

A *carbon transition* is the development path along which a shift is made from high to low carbon emissions. It begins at the Carbon Peak and ends with zero carbon emissions (Fig. 2). While it resembles decarbonization it links the change in carbon dependence to societal development and has a more specific ending.

A generic national *carbon transition curve* was proposed in 1997 to describe how over time countries pass from rising CO<sub>2</sub> emissions, as they depend on first wood and then fossil fuels for energy, to falling emissions as they increasingly rely on renewable energy sources (19-20).

The shift from fossil fuel to renewable energy dependence is also called an “*energy transition*” (35).

The carbon transition concept was inspired by the generic U-shaped *forest transition curve*, in which national forest cover declines as a country develops, and agriculture expands, switching from net deforestation (and CO<sub>2</sub> emissions) to net reforestation (and carbon sequestration) at a point called the 'forest transition' (Fig. 3). First seen in the present industrialized countries (36), it is rarer elsewhere in the world (37), so tropical deforestation remains a major source of CO<sub>2</sub> emissions (38).

In the various literatures which study the role of forests in global climate change (39), the stock of carbon stored above ground in a forest (C) is calculated by multiplying its area (A) by the volume per unit area (V) and density (D) of wood in trees, a factor (b) to convert this mass into total biomass and a carbon fraction quantity (f) to convert biomass into carbon (40):

$$6. \quad C = A \times V \times D \times b \times f$$

The global net forest carbon flux (C<sub>nf</sub>) is typically calculated as the sum of net fluxes in boreal forest (C<sub>nb</sub>), temperate forest (C<sub>nte</sub>) and tropical forest (C<sub>nt</sub>). As tropical forest is changing rapidly, C<sub>nt</sub> typically combines fluxes for sequestration by intact forest (C<sub>tri</sub>), and the net flux for land use change (C<sub>trluc</sub>), which combines emissions from forest clearance and sequestration by forest subsequently regrowing (41):

$$7. \quad C_{nf} = C_{nb} + C_{nte} + (C_{trluc} - C_{tri})$$

The carbon transition and forest transition curves use time as the independent variable, but in the *Environmental Kuznets Curve* the concentrations of some air pollutants, such as sulphur dioxide, peak when countries reach a certain level of economic development represented by GDP per capita (42). Its application to CO<sub>2</sub> is controversial (43).

The largest share of papers (26%) focuses on energy, compared with just 5% on materials and 1% on forests (Table 1). Many papers use "transition" as a synonym for "achieving" a low carbon economy (44) or energy sector (45). Yet some take a "pathway" approach consistent with the original vision of a generic trend (19). A few of these papers equate "pathway" with a future strategy (46) or scenario (47). Most, however, use theoretical frameworks in which influences from regulations, markets and other sources link pathways in CO<sub>2</sub> emissions to underlying societal trends (48-49). Some decarbonization studies use a pathway approach too (50).

### *Low carbon development*

*Low carbon development* is an equitable “development pathway which can achieve economic and social development while tackling global climate change” (51). It is particularly relevant to low and medium income countries (52), which face challenges in cutting CO<sub>2</sub> emissions but could “leap-frog” large-scale use of fossil fuels when they become less dependent on fuelwood (53), which still accounts for half of world wood production (54).

This literature is also dominated by energy (29% of papers). Just 7% of papers focus on economic development and poverty reduction. None covers materials (Table 1).

### *Carbon neutrality*

*Carbon neutrality* is achieved when emissions of CO<sub>2</sub> into the atmosphere are offset by CO<sub>2</sub> removals from the atmosphere. It is a weaker goal than zero carbon emissions, but equally specific and easier to measure at global scale. As with the UN Sustainable Development Goal ("target") of land degradation neutrality (55), it combines control of environmental (in this case atmospheric) degradation and terrestrial restoration.

A country achieves carbon neutrality when its carbon transition curve is intersected by its *carbon sequestration curve*, so the *carbon neutrality gap* between the two curves becomes zero (Fig. 2). Even higher “negative emissions” (56) lead to *carbon negativity* (57), which can reduce atmospheric CO<sub>2</sub> concentration to a level where its warming effect is no longer a threat.

The concept of carbon neutrality originated when individuals purchased carbon offsets against their emissions (58). It was then adopted by communities and organizations (59), before spreading to national scale (60). Concise definitions are rather elusive.

Energy studies are again dominant, with 26% of papers, while materials only account for 7%. Consistent with how the concept originated, 24% have a local or urban focus (Table 1).

### *Zero net carbon emissions*

While *net zero carbon emissions* is now a common policy goal (3), the term *zero net carbon emissions* has until now proved more popular in scientific studies, so it is used here as the name for this literature, though both are regarded as synonyms for carbon neutrality (61-62).

The two terms actually have different meanings when applied to buildings. In *zero net carbon buildings* "the amount of carbon emissions associated with a building's... construction [ $C_e$  = embodied carbon, discussed below] and... operational energy [ $C_o$ ] on an annual basis is zero or negative", i.e.  $C_e + C_o \leq 0$  (63). In *net zero energy buildings*, on the other hand, the focus is on operational energy, and energy imported from the electricity grid ( $E_{imp}$ ) is less than or equal to energy supplied to the grid by renewable energy technologies attached to the building ( $E_{exp}$ ) (64), or  $E_{imp} - E_{exp} \leq 0$ .

This is the smallest of the carbon literatures, even when including papers with 'energy', rather than 'carbon', in their titles. Some 70% of papers focus on buildings and only 14% on energy (Table 1). None focuses on materials, but their use in constructing buildings is evaluated.

### *Circular economy*

The excessive transfer of carbon from the Earth to the atmosphere can be treated as a consequence of a “*linear economy*” (65), in which under the current market system human utility is equated with consumption. This makes it seem acceptable to deposit wastes like CO<sub>2</sub> into the atmosphere and the other *environmental sinks* of water and land, and to extract new natural resources without recycling old products to reduce depletion of remaining *resources stocks*.

A “*circular economy*” closes feedback loops left open in the linear economy; reduces material “throughput” to conserve natural resources and limits waste flows to those which the environment can assimilate; and gives the environment “three economic functions – as resource

supplier, as waste assimilator, and as a direct source of utility” (65). It recognizes the prices of all three functions, so that, for example, any disutility resulting from CO<sub>2</sub> accumulating in the atmosphere should be corrected.

The political process of the circular economy has greatly expanded since the year 2000, as has its associated academic literature (66-67). It identifies normative actions to modify *biophysical* flows of resources, energy and wastes to increase prosperity and conserve the environment. Societal equity is rather neglected (68). The circular economy is a “contested concept” (69), so its political process has generated many indicator systems (70) and 114 definitions, though most of the latter combine “reduce, reuse and recycle activities” (71).

The parallel scientific process launched by Pearce and Turner (65) is still embryonic (72). Yet expanding its theoretical base, e.g. by including the frequency (“*circularity*”) and length of time (“*longevity*”) of resource use (73), should be more productive than trying to extract meaning from multiple definitions and indicator systems.

### *Life cycle analysis*

Life cycle analysis evaluates the environmental impacts of products from their origin in natural resources stocks to their recycling or disposal as waste in environmental sinks (21). It can help to integrate materials and energy in carbon neutrality studies, having evolved from the “net energy analysis” method (74), and is consistent with flows of energy and materials in a circular economy (75). Studies with the most comprehensive “*system boundary*” include the energy “*embodied*” in a product, e.g. by mining, refining ores into metals, processing metals into products, and transport between these stages ( $E_e$ ); the “*operating energy*” consumed in using the product ( $E_o$ ); and the “*demolition energy*” ( $E_{de}$ ) required for its recycling or disposal. Ramesh et al. (76) calculate overall *life cycle energy* ( $E_l$ ) for constructing and using buildings by:

$$8. \quad E_l = (E_{ein} + E_{er}) + E_o + E_{de}$$

where  $E_{ein}$  is initial *embodied energy* and  $E_{er}$  recurring embodied energy expended in repairs.

The embodied energy per tonne of metal ( $E_e$ ) is expressed by Rankin (77) as:

$$9. \quad E_e = \frac{E_m(1+w)}{(gR_mR_bR_{p1}R_{p2})} + \frac{E_b}{(gR_bR_{p1}R_{p2})} + \frac{E_{p1}}{R_{p2}} + E_{p2}$$

where  $E_m$  is mining energy;  $w$  the extra mass of waste per tonne of mined ore;  $g$  the ore grade;  $R_m$  the mass of ore sent from the mine for pre-processing (benefication);  $E_b$  the beneficiation energy per tonne of ore;  $R_b$  the fraction of ore recovered;  $E_{pi}$  the energy used in two chemical processing stages ( $i = 1-2$ ) and  $R_{pi}$  the fraction of material recovered in each stage.

*Life cycle zero energy buildings* have an annualized life cycle energy ( $E_{la}$ ) of zero. As they export more energy to the grid than they consume, mean annual net energy use ( $E_{oa}$ ) offsets mean annual embodied energy ( $E_{ea}$ ) over a building's life (78), or  $E_{la} = E_{ea} + E_{oa} \leq 0$ .

To estimate “global warming potential”, “*embodied energy*” derived from fossil fuels can be converted into an equivalent amount of “*embodied carbon*”, as can overall *life cycle energy*, though any product has other environmental impacts too (79).



### *Carbon footprint*

The *carbon footprint* concept is framed by life cycle analysis and refers to the mass of CO<sub>2</sub> (or all greenhouse gases) emitted “directly or indirectly.. by an activity or.. accumulated over the life stages of a product” (80).

Carbon footprint (CF) is estimated differently from the *ecological footprint* index (80), and combines CO<sub>2</sub> emissions from all energy uses and other activities A<sub>i</sub> (i = 1.....p):

$$10. \quad CF = \sum_{i=1}^p A_i \times EF_i$$

where EF<sub>i</sub> is the *emission factor* for activity i (81).

This literature has grown rapidly since 2005 (Fig. 1) and accounts for half the papers in our sample. It is alone in having similar numbers of papers on energy (10%) and materials (9%) (Table 1).

### **Conditions for carbon neutrality**

A new carbon neutrality science could counter the neglect of materials in this field by using physical variables and equations to integrate the analysis of energy and materials - and renewable and non-renewable resources - and measure progress in achieving carbon neutrality. Here, using a circular economy framework (65), we build on life cycle analysis (21) to add more equations to those listed above and use them to propose three conditions for carbon neutrality (Table 2).

#### *A circular economy framework*

The use of natural resources to generate energy, manufacture and use materials and emit CO<sub>2</sub> as waste can be conceptualized by an ideal circular economy framework in which Human Capital (Labour, and more generally human skills and practices) and Human-Made Capital (productive Capital, such as machinery, and consumer goods) are linked to Natural Capital (comprising Renewable and Non-Renewable Resource Capital and Environmental Quality). A circular economy framework includes the full range of resource and waste flows, though contemporary "linear economies" deviate from this ideal because recycling within Human-Made Capital is limited, and so a lot of waste is deposited in environmental sinks in the Environmental Quality component (65). The flows in Figure 4 are aggregates of those for the life cycles of billions of products as they pass through the stages of resource extraction, refining, processing, operation, disposal and recycling etc. Recycling reduces the depletion of Non-Renewable and Renewable Resource Capital, but the latter is continually renewed through sequestration of CO<sub>2</sub> from the atmosphere and regular harvesting. Energy is expended in managing renewable resources sustainably so that harvests take full advantage of CO<sub>2</sub> inputs. The carbon flows on which this paper focuses are a subset of the larger number of flows of resources and wastes which determine whether an economy is circular or not.

#### *Minimize carbon dioxide emissions at all life cycle stages of materials and energy resources*

Our analysis of carbon neutrality begins with the *Energy Life Cycle Equation*. Expanding equation 8, the total amount of energy used per unit mass of a product containing material *i*

during a single life cycle ( $E_{li}$ ) - its *life cycle energy* - combines the: (a) *cultivation energy* expended in managing the growth of wood and other renewable resources ( $E_{ci}$ ); (b) *extraction energy* needed to remove the material from its natural state, e.g. by mining mineral ores or harvesting trees ( $E_{mi}$ ); (c) *refining energy* used to obtain the material in its pure form, e.g. by refining ores to produce metals or converting trees into sawnwood and other primary products ( $E_{ri}$ ); (d) *processing energy* which converts a material into a product ( $E_{pi}$ ); (e) *transport energy* expended in all stages from extracting the raw material to delivering products to end-users ( $E_{ti}$ ); (f) *operating energy* consumed when using the product ( $E_{oi}$ ); (g) *disposal energy* expended in disposing of a product ( $E_{di}$ ); (h) *recycling energy* needed to recycle the material so it can be converted into a new product ( $E_{rei}$ ); and (i) *heat energy* saved by using processing and other waste to substitute for other forms of energy ( $E_{hi}$ ):

$$11. \quad E_{li} = E_{cij} + E_{mij} + E_{rij} + E_{pij} + E_{tij} + E_{oij} + E_{dij} + E_{reij} - E_{hij}$$

The size of each category depends on the technology used ( $j = 1 \dots m$ ) (82).

If materials are produced using fossil fuels then each *energy category* has a corresponding *carbon category* in the *Carbon Life Cycle Equation* to represent its associated CO<sub>2</sub> emissions:

$$12. \quad C_{li} = C_{cij} + C_{mij} + C_{rij} + C_{pij} + C_{tij} + C_{oij} + C_{dij} + C_{reij} - C_{hij}$$

Here  $C_{li}$  is *life cycle carbon*. Refining carbon ( $C_{ri}$ ) can also include CO<sub>2</sub> emitted when refining raw materials, e.g. to convert limestone into cement. This equation integrates energy and materials using a common unit of carbon, rather than energy as in the *exergy* concept (83). It does not include CO<sub>2</sub> emissions associated with unsustainable management of forests and other renewable resources. These are included in  $C_{luc}$  in equation 1 and  $C_{afolu}$  in equation 2.

The ideal path to carbon neutrality, and later zero carbon emissions, minimizes *life cycle carbon* in a *Carbon Transition Path Equation* which applies to materials and energy resources and represents total CO<sub>2</sub> emitted annually in producing and using billions of products:

$$13. \quad \text{Minimize } \sum C_{li} = \sum (C_{cij} + C_{mij} + C_{rij} + C_{pij} + C_{tij} + C_{oij} + C_{dij} + C_{reij} - C_{hij})$$

So the *first condition* for achieving carbon neutrality is that minimizing the sum of emissions at all stages of the life cycles of materials and energy resources is necessary to accelerate peaking of CO<sub>2</sub> emissions and then passage through the carbon transition (Fig. 2, Table 2).

National equations can be expanded, as in equation 4, to include imports and exports. Actual minimization of  $C_{li}$  along a development path will depend on how the *carbon intensities* of the technologies ( $j$ ) used in each category decline (84) under “*lock-in*” constraints (85).

Materials differ in their natural and embodied carbon contents. "Low carbon materials", such as metals, are naturally low in carbon but high in embodied carbon. "High carbon materials", such as wood, are naturally high in carbon but low in embodied carbon. Cement is an "Extreme carbon material", as it is naturally high in carbon in its original form of limestone and high in embodied carbon too (so the 'natural' classification of cement refers to the natural raw material (limestone) before it assumes its commercial form) (Table 3). Grouping materials with similar properties in this way helps in estimating embodied carbon intensity in *carbon impact assessments* of new processes and products (86). While embodiment may seem like a virtual

association it is physically very real, since each product is associated with a mass of CO<sub>2</sub> in the atmosphere.

Referring to C<sub>li</sub> in equation 12, the life cycle carbon benefits B<sub>lc</sub> of substituting one material (2) for another (1) in a product are expressed by a *Displacement Equation*:

$$14. \quad B_{lc} = C_{11} - C_{12}$$

B<sub>lc</sub> is 'normalized' by dividing by the difference between the masses of material 2 in the original product (m<sub>o</sub>) and substitute product (m<sub>s</sub>) to give a "*Displacement Factor*" DF<sub>12</sub> (87-88):

$$15. \quad DF_{12} = \frac{(C_{11} - C_{12})}{(m_s - m_o)}$$

The energy saved by recycling material i, rather than refining new ore, is indicated by the *Recycling Ratio* R<sub>i</sub> between its recycling energy (E<sub>rei</sub>) and refining energy (E<sub>ri</sub>), i.e. R<sub>i</sub> = E<sub>rei</sub> / E<sub>ri</sub>.

#### *Maximize carbon dioxide removals*

Since the present net terrestrial carbon sink is relatively small (see below), the *second condition* for carbon neutrality complements the first by requiring the maximization of CO<sub>2</sub> removals (C<sub>s</sub>) e.g. through terrestrial sequestration (C<sub>sterr</sub>), carbon capture and storage (C<sub>sccs</sub>) etc. (Table 2):

$$16. \quad \text{Maximize } \sum C_s = \sum (C_{sterr} + C_{sccs} + \dots)$$

#### *Offset carbon dioxide emissions by carbon dioxide removals*

The *third condition* states that carbon neutrality is achieved by offsetting gross CO<sub>2</sub> emissions (C<sub>g</sub>) by CO<sub>2</sub> removals (C<sub>s</sub>) (Table 2). At global scale, for all countries k = 1...n:

$$17. \quad \sum C_{gk} = \sum C_{sk}$$

Proximity to carbon neutrality can be measured by an absolute *Carbon Neutrality Gap* CNG =  $\sum C_{gk} - \sum C_{sk}$ , which equals 0 at carbon neutrality; and by a relative *Carbon Neutrality Index* CNI =  $\sum C_{gk} / \sum C_{sk}$ , which equals 1.0 at carbon neutrality.

The relative speeds of national carbon transitions can be compared by using the *Decarbonization Ratio* (D<sub>i</sub>). This divides the number of years between the Carbon Peak year and the current year (Y<sub>post</sub>), by the number of years between the Carbon Peak year and the historical Chordal Equivalent year, which had the same emissions as the current year (Y<sub>pre</sub>) (Fig. 5):

$$18. \quad D_i = \frac{Y_{post}}{Y_{pre}}$$

D<sub>i</sub> is 1.0 for symmetrical carbon transition curves. It can complement cumulative CO<sub>2</sub> emissions (46) in comparing alternative future scenarios, and is related to the delay between the peaks in global CO<sub>2</sub> emissions and global CO<sub>2</sub> concentration (equation 5) (29).

Most industrialized countries have now passed their Carbon Peaks. A long-term international database (89) shows that a few have symmetrical carbon transition curves in which the rate at which CO<sub>2</sub> emissions rise before the Carbon Peak resembles the subsequent rate of decline, e.g. Romania (Carbon Peak 1989) has a  $D_i$  value of 1.0. Other countries have skewed curves (Fig. 6). Countries that reached their Carbon Peaks before 1990 generally have  $D_i$  values above 1.0, e.g. France's emissions peaked in 1979 and its  $D_i$  is 2.1, representing a right-skewed curve (Table 4). Many countries with later Carbon Peaks have left-skewed curves and  $D_i$  values below 1.0, e.g. Italy's emissions peaked in 2004 and its  $D_i$  is 0.3. This general decline in  $D_i$  with Carbon Peak year (Fig. 7) implies that transitions are more rapid in countries with recent Carbon Peaks. For example, France's emissions in 2014 were 57% of their peak value but this drop took 35 years, while Italy's emissions declined to 68% of their peak value in just 10 years. This confounds expectations that learning from experience (90) should accelerate emissions decline. Yet national CO<sub>2</sub> *half lives* are at least 20 years for most countries with Carbon Peak years before 1990, but below 20 years for countries whose emissions peaked more recently (Table 4). Most national carbon transitions, however, are far from complete.

Few evaluations have yet been made of the availability and quality of the statistical data needed to quantify the equations in this section reliably (91).

### **Low carbon materials**

Metals are naturally low in carbon, but "embody" much more of it owing to the fossil fuels used to produce them and convert them into finished products. This section identifies the sources of CO<sub>2</sub> emissions in manufacturing and using metals which are critical to minimizing emissions to meet the first carbon neutrality condition (Table 2 and equation 13). It shows that constraints on changing technologies for producing metals could, by sustaining demand for fossil fuels, limit the rate of CO<sub>2</sub> emissions decline, regardless of renewable energy supply.

Steel accounts for 94% of all metal production and 70% of all greenhouse gas emissions associated with this. Producing steel and aluminium accounts for over 13% of all CO<sub>2</sub> emissions except for those from land use change. Copper and zinc are next in order of annual production (Table 5) (77). Steel, aluminium and copper are also used to make wind turbines, solar photovoltaic cells and lithium ion batteries for electric vehicles (92), so demand for them should remain strong in switching to renewable energy sources (93). We focus here on high longevity products, such as those used in vehicles, not low longevity products, such as beverage cans, many of which are rapidly recycled with high circularity (94).

Data available for analysis are limited. Our sample of 2,265 journal papers in the seven carbon literatures includes 17 papers on steel, but only 4 on aluminium, 2 on copper and none on zinc (Supplementary Table S1). Another Google Scholar search found 27 life cycle analyses for steel, 11 for aluminium, 19 for copper and 8 for zinc (Table S2), but none has a "cradle-to-grave" *system boundary* extending from extraction to recycling or disposal. Only 28% cover primary metal production, and not all of these include every stage from "cradle-to-[refinery] gate" (95), e.g. some focus on refining carbon and others on pre-processing ores. Yet most evaluate environmental impacts other than CO<sub>2</sub> emissions, e.g. emitting acidic gases such as sulphur dioxide (96), which together affect the overall sustainability of development.

### *Embodied carbon*

An overview by the Commonwealth Scientific and Industrial Research Organization (CSIRO) in Australia provides a set of estimates of *embodied carbon* in leading metals which, in our view, is uniquely comprehensive and internally consistent. Its estimates are also conservative, in having a common *emissions factor* for energy use (equation 10) based on using electricity derived from black coal at 35% efficiency. At 1 kg CO<sub>2</sub> eq /kWh this emissions factor is 27% greater than for oil, 2.3 times that for natural gas, 15 times that for nuclear power, and 100 times that for hydropower. Steel (an alloy of iron and other metals and up to 2% carbon) has the lowest embodied carbon per tonne of any leading metal: 2.2 t CO<sub>2</sub>, or a tenth of aluminium's 21.8 t CO<sub>2</sub> (Table 6). Copper and zinc have intermediate values of 3.3 t CO<sub>2</sub> and 4.6 t CO<sub>2</sub>, respectively, when produced using their most common pyrometallurgical and electrolytic processes, respectively (77). With the exception of aluminium, all CSIRO estimates of embodied carbon are in the middle of the range of other available estimates (Fig. 8).

### *Extraction carbon*

Extraction carbon is only 0.2% of embodied carbon for aluminium and 2.5% for steel, but up to 59.4% for copper (Table 6), since much crushing and grinding ('benefication') is needed to concentrate copper ore before shipping it from the mine for smelting (101).

The range of estimates of extraction carbon helps to explain the spread of estimates of embodied carbon in copper in Figure 8, its main driver being the variation in ore grade. Copper from Australia's Mount Isa mine, whose current ore grade is 3.0% (96), has an embodied carbon of 3.0 t CO<sub>2</sub> t<sup>-1</sup>, which is near the CSIRO estimate for the same grade (Fig. 8). The higher mean of 6.0 t CO<sub>2</sub> t<sup>-1</sup> for all mines in Chile reflects the lower mean ore grade of 0.7% in that country (102). Another large Australian mine (Olympic Dam), which has a 2% grade, has an even higher estimate of 8.5 t CO<sub>2</sub> t<sup>-1</sup>, but also a larger emissions factor (0.89 kg CO<sub>2</sub> eq /kWh) than Mount Isa (0.36 kg CO<sub>2</sub> eq /kWh) (96).

### *Refining carbon*

Refining carbon accounts for over 99% of embodied carbon in aluminium, 98% in steel and 89% in zinc, but as little as 41% for copper owing to the dominance of extraction carbon (Table 6). The ranges of estimates for aluminium, steel and zinc in Figure 8 are mainly driven by the dependence of refining on fossil fuels.

CSIRO's estimate of 21.8 t CO<sub>2</sub> t<sup>-1</sup> is the highest for aluminium, as it relies only on coal energy (77). In contrast, a global mean of 16.5 t CO<sub>2</sub> t<sup>-1</sup> (103) is 56% dependent on fossil fuels; another global mean of 14.4 t CO<sub>2</sub> t<sup>-1</sup> (104) is 71% dependent; and a European mean of 8.2 t CO<sub>2</sub> t<sup>-1</sup> (105) is just 19% dependent.

The steel estimates in Figure 8 have a more compact distribution, near the centre of which is the CSIRO estimate of 2.14 t CO<sub>2</sub> t<sup>-1</sup> (77). The others are 2.46 t CO<sub>2</sub> t<sup>-1</sup> for Poland (106), 2.04 t CO<sub>2</sub> t<sup>-1</sup> for China (107), 1.69 t CO<sub>2</sub> t<sup>-1</sup> for Europe (108), and 1.60 t CO<sub>2</sub> t<sup>-1</sup> for Italy (109). Only the last estimate includes extraction carbon, so for comparability the other values of embodied carbon in Figure 8 have been adjusted using CSIRO's estimate of 0.05 t CO<sub>2</sub> t<sup>-1</sup>. All these estimates are for processes dependent on fossil fuels.

The zinc estimates are all for the leading hydrometallurgical process and influenced by fossil fuel dependency too. A Chinese estimate of 6.1 t CO<sub>2</sub> t<sup>-1</sup> (110) is based on 100% coal energy, as is the CSIRO estimate of 4.6 t CO<sub>2</sub> t<sup>-1</sup> (77), while a European estimate of 3.1 t CO<sub>2</sub> t<sup>-1</sup> (111) is just 46% fossil fuel dependent. A lower global mean of 2.7 t CO<sub>2</sub> t<sup>-1</sup> (112) covers a range of fossil fuel dependencies. This, as for other means in this section, made it difficult for the authors of these studies to reliably estimate emission factors.

### *Transport carbon*

All the CSIRO estimates in Table 6 assume integrated mining and processing operations and so exclude transport carbon. Of the four metals reviewed here, transport energy is least crucial for aluminium, as its refining energy is so high. Until production expanded in China and the Persian Gulf (see below), most refining took place close to cheap nuclear or hydropower sources, to minimize refining energy. The UK now has only one small smelter at Lochaber, which uses hydropower. A larger smelter in Anglesey closed in 2009 when the nearby Wylfa nuclear power plant was decommissioned (113).

### *Reducing embodied carbon*

The key to reducing embodied carbon in most metals is to cut their refining carbon by switching to improved technologies that use renewable electricity (shown in green in Figure 8).

Aluminium, copper and zinc are already produced by electrical smelting. The 'best available technology' estimate for aluminium smelters of 3.5 t CO<sub>2</sub> t<sup>-1</sup> (104) in Figure 8 is based on hydropower. Improving anodes and cathodes could lower this value (114), but the Hall-Héroult smelting technology needs updating too (94). Before the year 2000, most aluminium smelters used hydropower or nuclear power, but the share of world production in China and the Persian Gulf based on fossil fuels has subsequently risen from 10% to 66%, and difficulties in replacing this new capacity are a "lock-in" effect (85) that will delay cuts in CO<sub>2</sub> emissions.

Switching to solar power could halve the European mean of embodied carbon for zinc from 3.1 t CO<sub>2</sub> t<sup>-1</sup> to 1.5 t CO<sub>2</sub> t<sup>-1</sup> (111). For copper, harnessing solar power in Chile's extensive drylands could cut its mean refining carbon of 6.0 t CO<sub>2</sub> t<sup>-1</sup> more sharply to 2.2 t CO<sub>2</sub> t<sup>-1</sup> (102), but reducing energy use in ore beneficiation is also needed to cut extraction carbon (96).

Switching to electricity poses the greatest challenge for steel production, since the widely used blast furnace method relies on fossil fuels to supply heat *and* reduce iron ore. The global dominance of steel is another "lock-in" effect that will limit the decline in total CO<sub>2</sub> emissions from all metal production if technologies change slowly. Blast furnaces could be replaced by hydrogen reduction and electric arc furnaces, using hydrogen derived by 'decarbonizing' fossil fuels (115) or electrolysing water, but this technology may not be operational until 2040 (116). It will take longer to spread worldwide, and since hydrogen reduction is expensive a high carbon price is also needed to make it economic (117). The same applies to electrowinning, an electrolytic process still under development (118).

### *Recycling carbon*

Greater recycling of metals is therefore crucial to counter slow technological change in primary production, especially for aluminium (114), whose Recycling Ratio between recycling energy

and refining energy ( $R_i$ ) is only 0.05-0.08, depending on the scrap input (Table 6). Steel has a larger  $R_i$  of 0.43, though its recycling energy of 10 GJ/t resembles that of aluminium.

### *Reducing operating carbon*

Minimizing CO<sub>2</sub> emissions will also require trade-offs between changes in *operating carbon* and embodied carbon. Operating carbon for steel vehicles can be cut by substituting lighter metals, such as aluminium, magnesium and titanium (119). These have higher strength to weight ratios ( $S$ ) than steel - whose  $S$  value varies but is assumed here to be 80 kN-m/kg (Table 7) - but their refining carbon is higher too. Yet for a car made of 10% aluminium, for which  $S = 130$  kN-m/kg, the drop in lifetime operating carbon could be up to six times the refining carbon of aluminium (120). Higher refining costs make the lighter metals more expensive, e.g. a tonne of aluminium costs four times as much as steel, but the actual cost difference is less since the final product is lighter. Substitution will also be influenced by the carbon price and safety issues.

### *Legacy carbon emissions*

This analysis shows that future CO<sub>2</sub> emissions scenarios for achieving carbon neutrality that are based solely on trends in renewable energy supply are over-optimistic, since they neglect delays in reducing fossil fuel demand by the metals industry. The lock-in effects of current technologies (steel) and fossil fuel-dependent refining capacity (aluminium), combined with the fossil fuel needs of new technologies, could leave a sizeable amount of *legacy carbon emissions* by 2050, and contribute to a substantial *carbon neutrality gap* (Fig. 2). If the metals industries are to become carbon neutral by 2050, all factories might therefore need their own carbon capture and storage facilities.

### *Data issues*

Life cycle analysis has evolved considerably since Ayres (74) criticized the tendency for studies to use "non-comparable units of measurement": all of the studies reviewed here reported "global warming potential" in comparable units. Yet Ayres' other criticism, about the use of "data from unreliable sources that cannot be checked", is still relevant, since estimates for metals rely on large numbers of company and industry data sources to quantify multiple variables. Other constraints include basing emissions factors on regional electricity data (96), and (as noted above) the small number of life cycle analyses for manufacturing metals (95). International reporting standards for energy intensive industries would ensure greater transparency, but to be feasible this would require a general strengthening of the current weak international institutional framework of these industries (10). Intergovernmental Panel on Climate Change (IPCC) guidelines (121) are currently the leading international standard for reporting CO<sub>2</sub> emissions by industries, but so far they do not cover complete life cycles.

### **High carbon materials**

Wood is naturally high in carbon, comprising half of carbon by weight, but compared with metals it needs less fossil fuel carbon to convert it into wood products. This section examines the feasibility of expanding global forest area to maximize CO<sub>2</sub> removals and offset CO<sub>2</sub> emissions to comply with the second and third carbon neutrality conditions (Table 2), and of converting the resulting extra wood supply into products to serve as long-term carbon stores.

## *Expanding forest area and carbon sequestration*

Forests already remove a considerable amount of carbon from the atmosphere. Temperate forests are now net carbon *sinks* since most industrialized countries, such as the USA, passed through their forest transitions long ago, and are on the upward sloping arm of the forest transition curve (Fig. 3) (36). However, owing to continuing deforestation, most tropical countries, such as Brazil, are still net carbon *sources* on the downward sloping arm. Only a few, such as Vietnam, have undergone their transitions (37). It is currently estimated, based on equation 7, that forests are a small net carbon sink of 4.0-5.9 Gt CO<sub>2</sub> a<sup>-1</sup> (1.1-1.6 Gt C a<sup>-1</sup>) (38, 41). So forests and other terrestrial sinks only offset a fraction of the 32 Gt CO<sub>2</sub> eq a<sup>-1</sup> of greenhouse gases emitted outside the land use sector (7).

Expanding forests is therefore necessary to maximize CO<sub>2</sub> removals to comply with the second carbon neutrality condition, and with the third condition by bringing the *carbon sequestration curve* nearer to the *carbon transition curve* (Fig. 2), yet forests only account for 1% of all papers in the seven carbon literatures (Table 1). At an IPCC conference in 1990 it was estimated that about 500 million hectares (Mha) of new forest could offset the prevailing net annual increment in atmospheric CO<sub>2</sub> of 10.6 Gt CO<sub>2</sub> a<sup>-1</sup> (2.9 Gt C a<sup>-1</sup>) from all sources, and so achieve carbon neutrality - though this term was not used then. Some 620 Mha of degraded tropical land were considered suitable for this "*carbonforest*" (122). While this estimate neglected economic and social factors, e.g. whether landowners wish to convert land to forest, this gap was tackled by research in the 1990s (15). The estimate also used poor quality statistical data, but a recent survey with very high resolution satellite images found sufficient land worldwide (123) to meet a new IPCC target of establishing up to 950 Mha of new forest by 2050 to limit global warming to 1.5°C (62). This area is twice as high as in 1990 since the net annual rise in atmospheric CO<sub>2</sub> has doubled. The Bonn Challenge is equally ambitious in aiming to establish 350 Mha of new forest by 2030 for this and other environmental goals (124).

If 10 Mha of new carbonforest had been planted annually from 1990, and CO<sub>2</sub> emissions remained unchanged, the world could have been carbon neutral within the present 2050 deadline. Yet CO<sub>2</sub> emissions have continued rising, and the rate of establishment of new tropical forest plantations fell from 1.8 Mha a<sup>-1</sup> in the 1980s to 0.8 Mha a<sup>-1</sup> in the 1990s, and from 2010 to 2015 was still only 0.9 Mha a<sup>-1</sup> (125). Planting outside the tropics peaked at 4.7 Mha a<sup>-1</sup> between 2000 and 2005, and the global rate of 2.5 Mha a<sup>-1</sup> ha between 2010 and 2015 could not even offset the annual natural forest loss of 5.8 Mha a<sup>-1</sup> ha (mainly in the tropics), despite temperate natural forest expanding at 1.4 Mha a<sup>-1</sup> (Table 8). The 110 Mha of new forest plantations established since 1990 is a small proportion of the 500 Mha and 950 Mha targets, so a huge effort is required if forest expansion is to offset all remaining CO<sub>2</sub> emissions by 2050.

Most industrialized countries, even those with large forest areas, such as Canada, have large carbon neutrality gaps (the blue bars in Figure 9), because net terrestrial sequestration ( $C_{\text{luc}} - C_{\text{terr}}$  in equation 1) (green bars) cannot offset fossil fuel CO<sub>2</sub> emissions (grey bars). Many tropical countries, such as Indonesia, have large carbon neutrality gaps too, since CO<sub>2</sub> emissions from deforestation greatly outweigh forest sequestration. Only a few countries where deforestation has ended, such as Malaysia and the Philippines, have quite small carbon neutrality gaps (126).

Contrary to assumptions that allowing the carbonforest to grow indefinitely will maximize carbon uptake, research shows that more carbon is sequestered by adopting normal forestry



practices, e.g. harvesting on specific rotations and thinning between harvests (127). The optimum rotation varies with tree species and ecological zone, but could be up to 80 years in temperate areas and lower in the tropics. Another finding is that carbon sequestration should be planned from the start to maximize total carbon storage in forests *and* wood products (Fig. 10), and that expanding existing uses of these products and substituting for other materials are both important (128). The current net annual sink associated with these "*harvested wood products*" is estimated at only 0.3 Gt CO<sub>2</sub> a<sup>-1</sup> (0.09 Gt C a<sup>-1</sup> (129)), but this is based on limited data since only in the last ten years have countries been required to provide statistics on these products to the UN Framework Convention on Climate Change (129).

### *Embodied carbon in wood products*

Wood products compare favourably with metals in their values of the life cycle carbon categories in equation 12, leading to a mean Displacement Factor of 2.1 in equation 15 for the carbon benefits of substituting wood for other materials (87).

Few measurements of the *cultivation carbon* ( $C_{cij}$ ) of wood in forest plantations are reported, but a recent estimate is 0.01 t CO<sub>2</sub> t<sup>-1</sup> of Eucalyptus logs for both it and extraction carbon in China. The main source of emissions is using diesel oil in trucks and equipment (130).

The *extraction carbon* incurred in harvesting trees is itself only a small proportion of the total embodied carbon of wood products (131). A typical US mean of 0.04 t CO<sub>2</sub> t<sup>-1</sup> of dry wood (Table 9) (132) is similar to the extraction carbon for steel (Table 6).

Total *embodied carbon* varies with the type of wood product, but is much lower than for metals, e.g. the 0.20 t CO<sub>2</sub> t<sup>-1</sup> of sawn softwood (Table 9) (133) is a tenth of that for steel (2.1 t CO<sub>2</sub> t<sup>-1</sup>) (Table 6). 'Wood panels', such as plywood, are more carbon intensive (134), e.g. embodied carbon is typically 0.45 t CO<sub>2</sub> t<sup>-1</sup> for plywood, or twice that for sawn softwood (133). The estimates of embodied carbon in Table 9 all use an emissions factor of 0.06 t CO<sub>2</sub> /GJ, except that for particle board (0.05 t CO<sub>2</sub> /GJ), which is identified as 'uncertain' (133).

The embodied carbon of wood could be cut by using battery-operated equipment and vehicles in forest management, and electricity for processing. Forest industries have long generated heat from wood processing waste, and on average about half of the total embodied energy in wood products is *heat energy* ( $E_{hij}$  in equation 11) recovered in this way (Table 9) (133).

The number of life cycle analyses of wood products - like that for metals - is still relatively low (Table S1) (133). The first detailed evaluation for the USA, for example, was not published until 2005 (131). Estimates vary considerably with tree species and energy use characteristics (131) and also differ in their measurement units, which supports Ayres' (74) critique.

### *Using more wood in buildings*

The top priority in expanding existing uses of wood is to increase long-term carbon storage in solid wood products, such as furniture (with lifetimes of 10-35 years), and those used in buildings (with lifetimes of at least 30-50 years) (127). We do not discuss low longevity products, such as paper, which are rapidly recycled with high circularity (135), or burning wood as fuel, the carbon neutrality of which is highly controversial (136-138).

Wood is a crucial part of "zero net carbon houses" designed to minimize embodied carbon and operating carbon (63), but the relative shares of these two categories are uncertain. In one estimate, embodied carbon only accounts for 10-20% of total life cycle carbon in buildings worldwide (76). Wood is already plentiful in buildings, and timber frame houses are the norm in Canada, Japan, New Zealand etc. A typical 2,074 m<sup>2</sup> timber frame house in Canada, for example, has twice as much wood as a brick house of similar size (139). Further reducing the non-wood content of buildings, e.g. by using wooden instead of concrete floors (140), can cut CO<sub>2</sub> emissions even more.

Substituting for cement and concrete is vital, as their manufacture accounts for a tenth of all CO<sub>2</sub> emissions except for those from land use change (Table 5). Cement is an 'extreme carbon material' (Table 3) since: (a) much CO<sub>2</sub> is emitted when limestone (calcium carbonate) is mixed with clay and heated to a high temperature to form 'clinker' – the intermediate material in manufacturing cement, which is then used to make concrete (141); and (b) more CO<sub>2</sub> is emitted when burning fossil fuel to provide the heat. Refining carbon (C<sub>rij</sub>) in equation 12 includes both types of emissions.

No feasible complete substitute for clinker has been found, in attempts to cut CO<sub>2</sub> emissions described in 39 papers in the seven carbon literatures - ten more than for all metals combined (Table S1). Fossil fuels could be 'decarbonized' into hydrogen, to generate energy (115), and carbon, to substitute for concrete (142). Yet incremental reductions are more likely, e.g. fly ash from coal power stations and steel blast furnace slag can already account for up to 30% of cement, and as much as 15% of fine limestone can be added without heating (143). Slow technological change in the cement industry could leave a large amount of *legacy carbon emissions* to add to that of metals, widening the 2050 carbon neutrality gap even further.

### *Moving to a new wood economy*

Owing to their lower embodied carbon, substituting wood products for metals, e.g. for steel in telegraph poles (144) and railway sleepers (145), can cut total CO<sub>2</sub> emissions. However, as global forest area expands and more trees are harvested, wood can also substitute for materials currently derived from fossil fuels, replacing the fossil fuel economy by a new *wood economy*. Glesinger published a blueprint for this, called "The Coming Age of Wood" (146), as early as 1949. Wood comprises cellulose, hemicellulose and lignin, three polymers that are separated when making paper. Goldstein showed again in 1975 how cellulose could be converted directly into polymers, such as rayon and cellophane, and broken down by hydrolysis and fermentation into ethanol, which can then be converted into ethylene and butadiene, the monomers used to form the polymers of polyethylene and polybutadiene (147).

These early visions have been reimagined for the 21st Century as a "*bio-based economy*", which is "the sustainable, eco-efficient, transformation of renewable biological resources into food, energy and other industrial products" (148). Just as oil refineries separate petroleum into fractions, "*biorefineries*" could convert wood and other forms of biomass into many products (149), e.g. degradable bio-plastics (150), and carbohydrates to serve as 'carriers' for generating hydrogen fuel (151). New plants are needed to convert cellulose and hemicellulose from birch wood into ethanol (152), but existing plants for manufacturing ethylene, propylene and butylene from fossil fuels can be "retrofitted" to take poplar wood instead (153). A transition metal catalyst method has been devised to depolymerize intractable lignin (154), as part of a new "green chemistry" initiative to use renewable feedstocks in chemical industries

(155). Wood might also be converted into other high carbon materials (156), such as graphene (157), carbon nano-tubes (158) and carbon fibre.

Wider political and commercial support is essential to realize the "bio-based economy" vision, since at the moment the European Union (EU) is virtually alone in having a strategy for this (159-160). In every country, new forests will compete with other land uses for each hectare of land (161). Yet support for bio-based economies should increase as the carbon price rises, and awareness grows of the competitiveness of new wood industries (153) and the potential to integrate these with climate change mitigation (162).

### **The emerging carbon economy**

To achieve carbon neutrality it must be economically feasible to introduce new technologies to minimize the carbon embodied in low carbon materials and, by increasing the use of high carbon materials and other means, maximize the transfer of CO<sub>2</sub> from the atmosphere to the Earth. This will require the market economy to undergo a structural change to a new *carbon economy* that sets a sufficiently high carbon price. (For reasons stated above, the term "carbon economy" is preferred to "low carbon economy".) This section identifies three features of the emerging carbon economy and proposes seven principles of a new carbon economics to explain them.

#### *Features of the emerging carbon economy*

Three distinctive features of the carbon economy are that:

1. *It is evolving*, and so is not pre-planned.
2. *Its evolution is being shaped by a diversity of actors using various approaches.* Initially, *governments* tried to cut CO<sub>2</sub> emissions by introducing new *national* regulations, and 'market-based mechanisms', e.g. pollution taxes and marketable pollution permits (65). The 1997 Kyoto Protocol of the UN Framework Convention on Climate Change allowed industrialized country (or Annex 1) signatories to use both approaches to achieve binding targets for emissions reductions. Carbon taxes have been introduced in many countries but vary in their effectiveness, longevity and acceptability (163). Carbon emissions trading schemes began in the EU in 2005 and then spread to China and other countries (164).

While environmental economists regard market-based mechanisms as more efficient than policies in changing polluting behaviour (65), critics claim that carbon neutrality can only be achieved if these mechanisms are supported by new policies (165), e.g. to replace intransigent technologies, such as coal power stations (166), and improve government administration of emissions trading schemes to generate the higher carbon prices needed to cut CO<sub>2</sub> emissions sharply (164).

However, many "*sub-national actors*" are also changing their energy use and everyday practices, such as cycling instead of using cars (167), and this can be just as effective as regulations or taxes in reducing *demand* for fossil fuels (168). Carbon offset schemes began in the 1980s to allow individuals to reduce the impacts of their travel (58). They were later incorporated in the Clean Development Mechanism of the Kyoto Protocol, so that industrialized countries listed in Annex 1 of the Protocol could offset their CO<sub>2</sub> emissions by funding carbon sequestration projects in other countries (169).

3. *It is uneven in space and time.* A heterogeneous global mosaic of national policies and market-based mechanisms has formed as governments have failed to agree on a universal binding successor to the Kyoto Protocol. So carbon can “leak” as industries migrate from 'strong' carbon economies to 'weaker' ones, e.g. aluminium smelting has shifted to fossil fuel-dependent economies in China and the Persian Gulf (104). Policies can also be reversed, e.g. in 2012 the Spanish government ended renewable energy subsidies (170).

### *Seven principles of carbon economics*

While 8% of papers in the seven carbon literatures are economic studies (Table 1) there is currently no specific literature on ‘carbon economics’. To fill this gap, seven principles of a new *carbon economics* are proposed here to explain the evolution and operation of the carbon economy.

1. *Carbon economics studies the allocation of overabundant carbon in the atmosphere while other factors of production remain scarce.* This adapts a longstanding definition of economics as "the study of the allocation of scarce resources" - capital, labour and land - "which have alternative uses" (171). Carbon is allocated by sequestering atmospheric carbon and avoiding volatilization of terrestrial carbon.

2. *The ideal human economy is a circular economy.* In an ideal human (as opposed to ecological) economy, materials extracted from Resource Capital would circulate within Human-Made Capital through recycling, instead of being deposited as waste in environmental sinks (Fig. 4) (65).

3. *The human economy is an integral part of the biosphere.* This follows the *global ecosystem framework* of ecological economics (172), instead of an environmental economics framework which separates the human economy from the biosphere (65). Figure 11 therefore combines the global ecosystem and circular economy frameworks.

4. *A sustainable human economy should stay within the ultimate carrying capacity of the biosphere.* This is another principle of the global ecosystem framework, and once the critical upper limit is exceeded, “remaining Natural Capital [replaces Hu]man-Made Capital as the limiting factor” (173). This remaining Natural Capital is the "Critical Natural Capital", which is essential for human life support, e.g. global environmental cycles, like the carbon cycle, and ecosystem services provided by biodiversity (174).

The Earth's carbon carrying capacity threshold, measured by the level of CO<sub>2</sub> in the atmosphere, may have been passed in the 1970s (175), when the current steep rise in global temperature also began (176). In another estimate, which divides global carrying capacity into multiple “planetary boundaries”, the biodiversity and nitrogen cycle thresholds have been passed too (177).

5. *New corrective economic institutions are needed when the scale of the human economy ‘overshoots’ ultimate carrying capacity.* Daly proposed in 1974 that to correct for overshoot the current institutions governing the market economy, e.g. those for contracts and property rights, should be supplemented by new institutions to stabilize human population, stop resource depletion, and limit social inequality (172). *Institutions* are not organizations but repeated practices, or "enduring regularities of human action in situations structured by rules, norms and

shared strategies, as well as by the physical world" (178). Daly implied that governments should impose the new institutions – as was common at the time. Since then, however, politics has changed dramatically, and studying institutional change has become a major research priority in various disciplines, including ecological economics (179).

Daly's proposal can be extended by using an approach, developed independently by Williamson and Ostrom, in which any society comprises multiple levels of institutions. Williamson proposed that institutions framing the market economy are structured hierarchically, with the institutions of resource allocation through *markets* being embedded in government institutions for *contracts*, which are nested in general "rules of the polity" that frame *property rights*, and are themselves embedded in *societal norms* (180). For a seamless shift to a carbon economy the arrangement of its institutions should be consistent with that of institutions governing the market economy. Ostrom's framework allows for this since it is more generic than Williamson's. "Operational institutions" - the everyday practices of *individuals* that are varied easily - are nested in the "collective choice institutions" of *groups* that change more slowly, and are framed by "constitutional choice institutions", complying with national and international *laws*, that vary even more slowly and are nested in "metaconstitutional institutions", such as *societal norms*, that rarely change (181).

6. *New institutions to govern the carbon economy can be created at all spatial scales.* According to Rhodes, following a shift in metaconstitutional institutions, in the 1990s many industrialized countries began switching from the conventional "government style" of governing, in which government steered society, to a new "*governance style*", in which society is "self-steering". So governing now involves decentralized, multiscale and networked interactions between all groups in society; civil society has greater autonomy; and individual citizens and non-governmental organizations can create institutions with national and even global impacts (182).

Institutions governing the carbon economy are represented in Figure 11 by an *institutional matrix*, in which rows correspond to actors at global to local scales, and columns to different economic or policy sectors. Institutions are established when norms, rules and practices are initiated and reproduced at various spatial scales and spread to other scales. Multiple *levels* are not identical to multiple *scales*, so actors at any scale may be associated with multiple levels of institutions.

This explains the *second* (diversity) feature of the carbon economy, described above. New *operational institutions* are created and reproduced by the repeated practices of many individuals, e.g. when purchasing carbon offsets for their travel. *Collective choice* institutions are created when corporations redesign their operations to be carbon neutral. *Constitutional choice* institutions are created by governments which introduce market-based mechanisms, and by UN organizations which adopt new international agreements to tackle global climate change. These are all influenced by, and consolidate, the new global *societal norm* of carbon neutrality. A multi-level institutional approach is already used in carbon transition studies (49, 183)

Spatio-temporal unevenness, the *third* feature of the carbon economy, results from a diversity of actors creating different institutions at different scales, times and places. Some scales have a full complement of institutions while others have none. Each cell in the institutional matrix varies in its *density* of institutions (Fig. 11).

Unevenness can be reduced by filling gaps in the institutional matrix, e.g. by revising the Paris Agreement, or the World Trade Organization's international trade rules (184); and constructing partial sets of institutions, e.g. for energy intensive industries (10), or to prevent *carbon bias* in the carbon economy, so that conservation does not favour high carbon density forests over high biodiversity forests (185).

7. *Institutions within Human Capital co-evolve with changes in Natural Capital.* “Co-evolution” of the human economy and biosphere is fundamental to ecological economics (186), and explains the *first* feature of the carbon economy - its evolution. “Polycentricity” in institutional change, identified by Ostrom (181), is matched by polycentric change in Natural Capital. So incremental changes in the life cycle stages of materials to minimize CO<sub>2</sub> emissions into the atmosphere, and maximize CO<sub>2</sub> removals from the atmosphere, can occur in parallel with changes in carbon economy institutions, and carbon price changes resulting from these. Other adaptive changes should eventually occur in response to breaches of the other “planetary boundaries” (177) noted above as more rules of the ecological economy come into play.

Co-evolution is apparent in a *carbon economy box* in which products and processes become economic at carbon prices inversely proportional to their CO<sub>2</sub> emissions (Fig. 12). As in the *McKelvey box* of resource economics (187), low carbon intensity technologies should become more economic as the carbon demand curve shifts and the carbon price rises, e.g. increasing the present carbon price of \$28 tC<sup>-1</sup> to \$67 tC<sup>-1</sup> should make it economic to manufacture steel by hydrogen reduction (117).

#### *The need for better statistics*

Better statistical data are needed for carbon economics research, and for the transparent national reporting needed for fair international trade in carbon (188). This will be promoted if new international trade institutions set common standards to upgrade carbon accounting methods (189), and if all products carry *carbon barcodes* to record the categories of embodied carbon (equation 12) added at each life cycle stage.

## **Conclusions**

Compared with other global goals, such as sustainable development (190), and other carbon strategies, such as decarbonization (23) and a low carbon economy (16), carbon neutrality is unambiguous and easier to measure at global scale. Yet achieving it is currently constrained since strategies and research have focused on energy (191) and neglected materials.

To fill this gap, we have extended existing equations in carbon literatures (76) by deriving more equations to integrate the analysis of the roles of energy and materials in achieving carbon neutrality. These equations can be used to establish a framework for a new carbon neutrality science; provide a common language for all carbon literatures; and monitor progress in realizing carbon neutrality. They also lead to three conditions for carbon neutrality: minimize CO<sub>2</sub> emissions at all stages of the life cycles of materials and energy resources; maximize CO<sub>2</sub> removals; and offset CO<sub>2</sub> emissions by CO<sub>2</sub> removals.

Applying these equations to low carbon materials, such as metals, shows that for aluminium, steel and zinc, refining carbon is the key source of CO<sub>2</sub> emissions to cut to minimize emissions to comply with the first condition. Extraction carbon is also important for copper. Trading off

reductions in refining carbon and operating carbon will be another challenge. Yet owing to delays in introducing the best available technologies worldwide by 2050, a substantial amount of *legacy carbon emissions* could remain, and contribute to a sizeable *carbon neutrality gap*.

Wood is naturally high in carbon but low in embodied carbon, and so can reduce CO<sub>2</sub> emissions if substituted for metals. If global forest area is expanded, to comply with the second condition for carbon neutrality by maximizing CO<sub>2</sub> removals, more wood will become available to substitute for metals and other materials and provide new terrestrial carbon stores. So burning wood as fuel is indeed wasteful (138). Yet slow forest expansion since an Intergovernmental Panel on Climate Change conference in 1990 called for forest-based climate change mitigation (192) raises doubts about whether sufficient forest expansion can occur before 2050 to offset remaining CO<sub>2</sub> emissions and satisfy the third condition for carbon neutrality.

Insufficient minimization of CO<sub>2</sub> emissions and maximization of CO<sub>2</sub> removals could therefore leave a substantial carbon neutrality gap by 2050. However, carbon neutrality is sufficiently flexible to allow sub-optimal forest expansion to be compensated by: (a) faster than expected cuts in CO<sub>2</sub> emissions, and (b) greater use of carbon capture and storage, as the evolving carbon economy sets a higher carbon price to make carbon neutrality more economically feasible.

To fill another gap, this paper has proposed seven principles of carbon economics that build on circular economy, ecological economics and new institutionalism frameworks. These principles explain three features of the carbon economy: its gradual evolution, the various approaches adopted by a diversity of actors, and its spatio-temporal unevenness, which has led to 'carbon leakage'. The seventh principle - co-evolution of the human economy and the biosphere - suggests that the carbon price could eventually rise sufficiently to correct the imbalance between CO<sub>2</sub> emissions and removals, and counter critics of emissions trading (165).

Far more research is needed into the role of materials in carbon neutrality. Future research in carbon neutrality science could study actual transition paths to carbon neutrality and ultimately zero carbon emissions, and the role of harvested wood products as global carbon stores. More carbon neutrality equations could be devised to cover, for example, the carbon benefits of constructing vehicles using materials that are lighter and stronger than existing ones; the balance between longevity and circularity in different materials (73); and the impact of recycling alloys on the quality of materials (193). Research in carbon economics could expand its theoretical base, and study the spatial unevenness of the carbon economy in detail.

Our research leads to five policy recommendations for governments coming under pressure to publish detailed carbon neutrality strategies (194-196). First, these strategies should integrate the energy inputs to economies and materials outputs. Second, they should encourage stronger international carbon reporting standards for energy-intensive industries, to improve transparency in national reports of progress towards carbon neutrality. Third, they should establish national databases of carbon stored in harvested wood products, to improve estimates of terrestrial carbon stocks other than in ecosystems and agroecosystems. Fourth, they should promote faster forest expansion. Fifth, they should reinvigorate the Reducing Emissions from Deforestation and Degradation (REDD+) mechanism of the UN Framework Convention on Climate Change, so that deforestation no longer cancels out afforestation gains.

A new carbon neutrality science will have much to study in the coming decades as humanity strives to ensure the sustainability of Planet Earth. It also has much to contribute to planning effective strategies. We have set out some simple equations which can be used to assess the

merits (or otherwise) of a range of proposed strategies for approaching carbon neutrality. If such basic metrics had been employed in the past, serious mistakes might have been avoided, for example in the case of the aluminium industry, and in the deployment of early-generation biofuels. We can only hope for better, more analytical, approaches in the future.

### **Acknowledgements**

The authors wish to thank the Editor-in-Chief and two anonymous reviewers for recognizing the potential in our original paper and for encouraging us to develop it further.



## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

\* of special interest

\*\* of outstanding interest

1. Scripps Institution of Oceanography, 2020. The Keeling Curve. University of California, San Diego. <https://scripps.ucsd.edu/programs/keelingcurve/>.
2. Flagg J.A., 2015. Aiming for zero: what makes nations adopt carbon neutral pledges? *Environmental Sociology* **1**: 202-212.  
\* A good overview of early government commitments to carbon neutrality.
3. BEIS, 2019. UK becomes first major economy to pass net zero emissions law. Press release, 27 June 2019, Department for Business, Energy & Industrial Strategy, London.
4. Harding R., 2020. Japan to be carbon neutral by 2050, insists prime minister. *Financial Times*, 26 October 2020.
5. Shepherd C., 2020. China's carbon pledge revives hopes of a climate game change. *Financial Times*, 23 September 2020.
6. Guterres A., 2020. Net-zero emissions must be met by 2050 or COVID-19 impact on global economies will pale beside climate crisis, Secretary-General tells Finance Summit. Press release, 12 November 2020, United Nations, New York.
7. IPCC, 2014. *Mitigation of Climate Change*. Working Group III Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge.
8. Ahman M., Nilsson L.J. and Johansson B., 2017. Global climate policy and deep decarbonization of energy-intensive industries. *Climate Policy* **17**: 634-649.
9. Gerres T., Chaves-Ávila J.P., Linares P. and San Román T.G., 2019. A review of cross-sector decarbonisation potentials in the European energy intensive industry. *Journal of Cleaner Production* **210**: 371-381.
10. Oberthür S., Khandekar G. and Wyns T., 2020. Global governance for the decarbonization of energy-intensive industries: Great potential underexploited. *Earth System Governance*. doi.org/10.1016/j.esg.2020.100072.  
\*A stimulating account of governing options in the energy intensive industries.
11. Rissman J., Bataille C., Masanet E., Adene N. et al., 2020. Technologies and policies to decarbonize global industry: review and assessment of mitigation drivers through 2070. *Applied Energy* **266**: 114848.  
\*A good overview of future prospects for the energy intensive industries.

12. Wesseling J.H., Lechtenböhmer S., Åhman M., Nilsson L.J., Worrell E. and Coenen L., 2017 The transition of energy intensive processing industries towards deep decarbonization: characteristics and implications for future research. *Renewable and Sustainable Energy Reviews* **79**: 1303-1313.
13. Wimbadi R. and Djalante R., 2020. From decarbonization to low carbon development and transition: A systematic literature review of the conceptualization of moving toward net-zero carbon dioxide emission (1995–2019). *Journal of Cleaner Production* **256**: 120307.  
\* A good evaluation of various carbon literatures.
14. Sanna A., Dri M., Hall M.R. and Maroto-Valer M., 2012. Waste materials for carbon capture and storage by mineralisation (CCSM)–A UK perspective. *Applied Energy* **99**: 545-554.  
\* A good overview of carbon capture and storage technology.
15. Grainger A., Iverson L.R., Marland G.H. and Prasad A., 2019. Comment on “The global tree restoration potential”. *Science* 10.1126/science.aay8334 (2019).
16. Richels R. and Edmonds J., 1995. The economics of stabilizing atmospheric CO<sub>2</sub> concentrations. *Energy Policy* **23**: 373-378.
17. Wiseman J., T Edwards T. and K Luckins K., 2013. Post carbon pathways: A meta-analysis of 18 large-scale post carbon economy transition strategies. *Environmental Innovation and Societal Transitions* **8**: 76-93.
18. Grainger A., 2010. Reducing uncertainty about hybrid lay-scientific concepts. *Current Opinion in Environmental Sustainability* **2**: 444-451.
19. Grainger A., 1997. Compensating for opportunity costs in forest-based strategies to mitigate global climate change. *Critical Reviews in Environmental Science and Technology* **27**: S163-176.
20. Van Kooten C.G., Grainger A., Ley E., Marland G. and Solberg B., 1997. Conceptual issues related to carbon sequestration: uncertainty and time. *Critical Reviews in Environmental Science and Technology* **27**: S65-S82.
21. Lee J.J., O'Callaghan P. and Allen D., 1995. Critical review of life cycle analysis and assessment techniques and their application to commercial activities. *Resources, Conservation and Recycling* **13**: 37-56.  
\* An early overview of life cycle analysis.
22. Watson R.T. et al., 2000. Land Use, Land-Use Change and Forestry. IPCC/Cambridge University Press, Cambridge.
23. Nakicenovic N., 1996. Decarbonization: Doing more with less. *Technological Forecasting and Social Change* **51**:1-17.  
\* A seminal paper on decarbonization.
24. Blackburn C., Harding A. and Moreno-Cruz J., 2017. Toward deep-decarbonization: an energy-service system framework. *Current Sustainable/Renewable Energy Reports* **4**: 181–190

25. Geels F.W., Sovacool B.K., Schwanen T. and Sorrell S., 2017. Sociotechnical transitions for deep decarbonisation. *Science* **357**: 1242-1244.
26. Yamaji K., Matsushashi R., Nagata Y. and Kaya Y., 1991. An integrated system for CO<sub>2</sub>/Energy/GNP analysis: case studies on economic measures for CO<sub>2</sub> reduction in Japan. *Workshop on CO<sub>2</sub> Reduction and Removal: Measures for the Next Century*, 19 March 1991. International Institute for Applied Systems Analysis, Laxenburg, Austria.
27. Muradov N., 2013. Decarbonization at crossroads: the cessation of the positive historical trend or a temporary detour? *Energy and Environmental Science* **6**: 1060 -1073.  
\* A thought-provoking paper on decarbonization.
28. Harper P., 2016. Decomposition for decarbonisation: evaluation of decarbonisation programmes. *Science Progress* **99**: 235–261.
29. Seshadri A.K., 2016. Decarbonization rate and the timing and magnitude of the CO<sub>2</sub> concentration peak. *Global and Planetary Change* **146**: 22-29.
30. DTI, 2003. Our energy future - creating a low carbon economy. Department of Trade and Industry, London.
31. Ekins P., Anandarajah G. and Strachan N., 2011 Towards a low-carbon economy: scenarios and policies for the UK. *Climate Policy* **11**: 865-882.
32. Zhuang G., 2008. How will China move towards becoming a low carbon economy? *China and World Economy* **16**: 93-105.
33. Fu B., Wu M., Che Y., Wang M., Huang Y and Bai Y., 2015 The strategy of a low-carbon economy based on the STIRPAT and SD models. *Acta Ecologica Sinica* **35**: 76-82.
34. Ehrlich P.R. and Holdren J.P., 1971. Impact of population growth. *Science* **171**: 1212-1217.
35. Cherp A., Vinichenko V., Jewell J., Brutschin E. and Sovacool B., 2018. Integrating techno-economic, social-technical and political perspectives on national energy transitions: a meta theoretical framework. *Energy Research and Social Science* **37**:175-190.  
\*\*An interdisciplinary study which places energy transitions within the context of a diversity of social science theories.
36. Mather A.S., 1992. The forest transition. *Area* **30**: 117-124.
37. Barbier E.B., Burgess J.C. and Grainger A., 2010. The forest transition: towards a more comprehensive theoretical framework. *Land Use Policy* **27**: 98–107
38. Le Queré C., Andres R.J., Boden T., Conway T., Houghton R.A. et al., 2013. The global carbon budget 1959–2011. *Earth System Science Data* **5**: 165–185.
39. Grainger A., 2010. Uncertainty in constructing global knowledge about tropical forests. *Progress in Physical Geography* **34**: 811-844.

40. Kauppi, P.E., Ausubel, J.H., Fang, J., Mather A.S., Sedjo, R.A. and Waggoner, P.E. 2006 *Proceedings of the National Academy of Sciences USA* 103, 17574-17579.
41. Pan Y.D., Birdsey R.A., Fang J., Houghton R., Kauppi P.E. et al., 2011. A large and persistent carbon sink in the world's forests. *Science* **333**: 988–993.
42. Selden T.M. and Song D., 1994. Environmental quality and development: is there a Kuznets Curve for air pollution emissions? *Journal of Environmental Economics and Management* **27**:147-162.
43. Itkonen J.V.A., 2012. Problems estimating the carbon Kuznets curve. *Energy* **39**: 274-280.
44. Jagger N., Foxon T. and A Gouldson A., 2013. Skills constraints and the low carbon transition. *Climate Policy* **13**: 43-57.
45. Newbery D.M., 2016; Towards a green energy economy? The EU Energy Union's transition to a low-carbon zero subsidy electricity system—Lessons from the UK's Electricity Market Reform. *Applied Energy* **179**: 1321-1330.
46. Wang T. and Watson J., 2010 Scenario analysis of China's emissions pathways in the 21st century for low carbon transition. *Energy Policy* **38**: 3537-3546.
47. Cranston G.R. and Hammond G.P., 2010. North and south: regional footprints on the transition pathway towards a low carbon, global economy. *Applied Energy* **87**: 2945-2951.
48. Foxon T.J., Pearson P.J.G. and Arapostathis S., 2013. Branching points for transition pathways: assessing responses of actors to challenges on pathways to a low carbon future. *Energy Policy* **52**: 146-158.  
\*An interesting paper on carbon transitions.
49. Geels F.W., Berkhout F. and Van Vuuren D.P., 2016. Bridging analytical approaches for low-carbon transitions. *Nature Climate Change* **6**: 576–583.  
\*Another interesting paper on carbon transitions.
50. Mathy S., Criqui P., Knoop K., Fishedick M. and Samadi S., 2016. Uncertainty management and the dynamic adjustment of deep decarbonization pathways. *Climate Policy* **16**: S47-S62.
51. Urban, F., 2010. The pro-poor low carbon development and the role of growth. *International Journal of Green Economics* **4**: 82–93.
52. Mulugetta Y. and Urban F., 2010. Deliberating on low carbon development. *Energy Policy* **38**: 7546-7549.  
\*A brief introduction to low carbon development.
53. Van der Zwaan B., Kober T. Dalla Longa F., Van der Laane A. and Kramer G.J., 2018. An integrated assessment of pathways for low-carbon development in Africa. *Energy Policy* **117**: 387-395.

54. Bailis R., Drigo R., Ghilardi A. and Masera O., 2015. The carbon footprint of traditional woodfuels. *Nature Climate Change* **5**: 266–272.
55. Grainger A., 2015. Is land degradation neutrality feasible in dry areas? *Journal of Arid Environments* **112**:14-24.
56. Daggash H.A., Heuberger C.F. and MacDowell N., 2019. The role and value of negative emissions technologies in decarbonising the UK energy system. *International Journal of Greenhouse Gas Control* **81**:181-198.
57. Vanholme B., Desmet T., Ronsse F., Rabaey K, et al., 2013. Towards a carbon-negative sustainable bio-based economy. *Frontiers in Plant Science* **4**: 174.  
\*An interesting discussion of carbon negativity in relation to a bio-based economy.
58. Trexler M.C. and Kosloff L.H., 2006. Selling carbon neutrality. *Environmental Forum*, March/April: 34-39.
59. Oxtoby D.W., 2008. Biologists and carbon neutrality. *Bioscience* **58**: 382-383.
60. Pilpola S., Arabzadeh V., Mikkola J. and Lund P.D., 2012. Analyzing national and local pathways to carbon-neutrality from technology, emissions, and resilience perspectives - case of Finland. *Energies* **12**: 949.  
\*A recent study of energy transitions to carbon neutrality at national and city scales which reviews earlier national studies.
61. Kilbert C.J. and Fard M.M., 2012. Differentiating among low-energy, low-carbon and net-zero-energy building strategies for policy formulation. *Building Research and Information* **40**: 625-637.
62. Allen M., Babiker M., Chen Y., De Coninck H., Connors S. et al., 2018. *An IPCC Special Report on the Impacts of Global Warming of 1.5°C above Pre-Industrial Levels*. Intergovernmental Panel on Climate Change, Geneva.
63. UK Green Building Council, 2019. *Net Zero Carbon Buildings: A Framework Definition*. UK Green Building Council, London.
64. Cellura M., Guarino F., Longo S. and Mistretta M., 2014. Energy life-cycle approach in net zero energy buildings balance: operation and embodied energy of an Italian case study. *Energy and Buildings* **72**: 371-381.
65. Pearce D. and Turner R. K., 1990. *Economics of Natural Resources and Environment*. Harvester, London.
66. Stahel W.R., 2016; The circular economy. *Nature* **531**: 435-438.
67. Luis E.C. and Celma D., 2020. Circular economy. A review and bibliometric analysis. *Sustainability* **12**(16): 6381.

68. Geissdoerfer M., Savaget P., Bocken N.M.P. and Hultink E.J., 2017. The circular economy – a new sustainability paradigm? *Journal of Cleaner Production* **143**: 757-768.
69. Korhonen J., Nuur C., Feldmann A. and Birkie S.E., 2018. Circular economy as an essentially contested concept. *Journal of Cleaner Production* **175**: 544-552.
70. Saidani M., Yannou B., Leroy Y. and Cluzel F., 2019. A taxonomy of circular economy indicators. *Journal of Cleaner Production Volume* **207**: 542-559.
71. Kirchherr J., Reike D. and Hekkert M., 2017. Conceptualizing the circular economy: an analysis of 114 definitions. *Resources, Conservation and Recycling* **127**: 221-232
72. Korhonen J., Honkasalo A. and Seppälä J., 2018. Circular economy: the concept and its limitations. *Ecological Economics* **143**: 37-46.
73. Figge F., Thorpe A.S., Givry P., Canning L. and Franklin-Johnson E., 2018. Longevity and circularity as indicators of eco-efficient resource use in the circular economy. *Ecological Economics* **150**: 297-306.
74. Ayres R.U., 1995. Life cycle analysis: A critique. *Resources, Conservation and Recycling* **14**: 199-223.
75. Niero M. and Kalbar P.P., 2018. Coupling material circularity indicators and life cycle based indicators: A proposal to advance the assessment of circular economy strategies at the product level. *Resources, Conservation and Recycling* **140**: 305-312.
76. Ramesh T. et al., 2010. Life cycle analysis of buildings: an overview. *Energy and Buildings* **42**:1592-1600.  
\*An introduction to life cycle analysis from a building perspective.
77. Rankin W.J., 2011. *Minerals, Metals and Sustainability: Meeting Future Material Needs*. CRC Press, London.
78. Hernandez P. and Kenny P., 2010. From net energy to zero energy buildings: defining life cycle zero energy buildings (LC-ZEB). *Energy and Buildings* **42**: 815-821.
79. Shirazi A. and Ashuri B., 2020. Embodied Life Cycle Assessment (LCA) comparison of residential building retrofit measures in Atlanta. *Building and Environment* **171**: 106644
80. Galli A., Wiedmann T., Ercin E., Knoblauch D. and Ewing B., 2012. Integrating ecological, carbon and water footprint into a “footprint family” of indicators: definition and role in tracking human pressure on the planet. *Ecological Indicators* **16**: 100-112
81. Dormer A., Finn D.P., Ward P. and Cullen J., 2013. Carbon footprint analysis in plastics manufacturing. *Journal of Cleaner Production* **51**: 133-141.
82. Morini A.A., Ribeiro M.J. and Hotza D., 2019. Early-stage materials selection based on embodied energy and carbon footprint. *Materials and Design* **178**: 107861.

83. Valero A., Valero A., Calvo G., Ortego A and Ascaso S., 2018. Global material requirements for the energy transition. An exergy flow analysis of decarbonisation pathways. *Energy* **159**: 1175-1184.
84. Zhang M., Mu H., Ning Y. and Song Y., 2009. Decomposition of energy-related CO<sub>2</sub> emission over 1991–2006 in China. *Ecological Economics* **68**: 2122-2128.
85. Unruh G.C., 2002. Escaping carbon lock-in. *Energy Policy* **30**: 317-325.  
\*A seminal paper on lock-in.
86. Rydh C.J. and Sun M., 2005. Life cycle inventory data for materials grouped according to environmental and material properties. *Journal of Cleaner Production* **13**: 1258-1268.
87. Sathre R. and O'Connor J., 2010. Meta-analysis of greenhouse gas displacement factors of wood product substitution. *Environmental Science and Policy* **13**:104–114.  
\*Compiles Displacement Factors for substituting wood products for non-wood products.
88. Bergman R., Puettmann M., Taylor A. and Skog K.E., 2014. The carbon impacts of wood products. *Forest Products Journal* **64**: 220-231.
89. World Bank, 2020. CO2 emissions (kt), from the Carbon Dioxide Information Analysis Centre. World Bank, Washington DC. Downloaded from; <https://data.worldbank.org/indicator/EN.ATM.CO2E.KT>
90. Crossan M.M., Lane H.W. and White R.E., 1999. An organizational learning framework: from intuition to institution. *Academy of Management Review* **24**: 522-536.
91. Sabolic D., 2018. Can decarbonization policy results be detected by simplistic analysis of macro-level statistical data? *Technology in Society* **53**:103-109.  
\*An important paper on the statistical basis of carbon transition studies.
92. Giurco D., Dominish E., Florin N., Watari T. and McLellan B., 2019. Requirements for minerals and metals for 100% renewable scenarios. In: Teske S. (eds) *Achieving the Paris Climate Agreement Goals*: 437-457. Springer, Cham.
93. De Koning A., Kleijn R., Huppes G., Sprecher B., Engelen G. and Tukker A., 2018. Metal supply constraints for a low-carbon economy? *Resources, Conservation and Recycling* **129**: 202-208.
94. Liu G., Bangs C.E. and Müller D.B., 2011. Unearthing potentials for decarbonizing the US aluminum cycle *Environmental Science and Technology* **45**:9515-9522.
95. Segura-Salazar J., Lima F.M. and Tavares L.M., 2019. Life Cycle Assessment in the minerals industry: Current practice, harmonization efforts, and potential improvement through the integration with process simulation. *Journal of Cleaner Production* **232**: 174-192.
96. Memary R., Giurco D., Mudd G. and Mason L., 2012. Life cycle assessment: a time-series analysis of copper. *Journal of Cleaner Production* **33**: 97-108.  
\*An interesting life cycle analysis of copper in Australia.

97. USGS, 2016. *2015 Minerals Yearbook: Aluminium*. US Geological Survey, Washington DC.
98. USGS, 2017. *2015 Minerals Yearbook: Copper*. US Geological Survey, Washington DC.
99. USGS, 2018. *2015 Minerals Yearbook: Iron and Steel*. US Geological Survey, Washington DC.
100. USGS, 2017. *2015 Minerals Yearbook: Zinc*. US Geological Survey, Washington DC.
101. Farjana S.H., Huda N. and Mahmud M.A.P., 2019. Life cycle analysis of copper-gold-lead-silver-zinc beneficiation process. *Science of the Total Environment* **659**: 41-52.
102. Moreno-Leiva S., Díaz-Ferrán G. and Haas G., 2017 Towards solar power supply for copper production in Chile: Assessment of global warming potential using a life-cycle approach. *Journal of Cleaner Production* **164**: 242-249.
103. Nunez P. and Jones S., 2016. Cradle to gate: life cycle impact of primary aluminium production. *International Journal of Life Cycle Assessment* **21**: 1594–1604.
104. Saevarsdottir G., Kvannd H. and Welch B.J., 2020. Aluminum production in the times of climate change: The global challenge to reduce the carbon footprint and prevent carbon leakage. *Journal of the Minerals, Metals & Materials Society* **72**: 296–308.
105. Achillas C., Vlachokostas C. and Koroneos C., 2020. Life cycle thinking: towards the sustainable management of resources in aluminium production. *Euro-Mediterranean Journal for Environmental Integration* **5**:16.
106. Burchart-Korol D., 2013. Life cycle assessment of steel production in Poland: a case study. *Journal of Cleaner Production* **54**: 235-243.
107. Li F., Chu M., Tang J., Liu Z., Wang J. and Li S., 2021. Life-cycle assessment of the coal gasification-shaft furnace-electric furnace steel production process. *Journal of Cleaner Production* **287**: 125075.
108. Bhaskar A., Assadi M., Somehsaraei H.N.N., 2020. Decarbonization of the iron and steel industry with direct reduction of iron ore with green hydrogen. *Energies* **13**: 758.  
\*An interesting study of new on iron and steel technology.
109. Renzulli P.A., Notarnicola B., Tassielli G. and Arcese G., 2016 Life cycle assessment of steel produced in an Italian integrated steel mill. *Sustainability* **8**: 719.
110. Qi C., Ye L., Ma X., Yang D., and Hong J., 2017. Life cycle assessment of the hydrometallurgical zinc production chain in China. *Journal of Cleaner Production* **156**: 451-458.
111. Werder M. and Steinfeld A., 2000. Life cycle assessment of the conventional and solar thermal production of zinc and synthesis gas. *Energy* **25**: 395-409.



112. Van Genderen E., Wildnauer M., Santero N. and Sidi N., 2016. A global life cycle assessment for primary zinc production. *International Journal of Life Cycle Assessment* **21**: 1580–1593.
113. Gribben R., 2011. Chancellor too late to save Britain's aluminium smelters. *Daily Telegraph*, 11 December.
114. Das S., 2012. Achieving carbon neutrality in the global aluminium industry. *Journal of Metals* **64**:285-290.  
\*\*A pioneering analysis of carbon neutrality in the metals trade.
115. Steinberg M., 1999 Fossil fuel decarbonization technology for mitigating global warming. *International Journal of Hydrogen Energy* **24**: 771-777.  
\*An early study of decarbonization technology.
116. Toktarova A., Karlsson I., Rootzén. J., Göransson L., Odenberger M. and Johnsson F., 2020. Pathways for low-carbon transition of the steel industry—a Swedish case study. *Energies* **13**: 3840.
117. Gielen D., Saygin D., Taibi E. and Birat J.P., 2020. Renewables-based decarbonization and relocation of iron and steel making: A case study. *Journal of Industrial Ecology* **24**: 1113-1125.
118. Bataille C., Åhman M., Neuhoff K., Nilsson L.J. and Fishedick M., 2018. A review of technology and policy deep decarbonization pathway options for making energy-intensive industry production consistent with the Paris Agreement. *Journal of Cleaner Production* **187**: 960-973.
119. ARPA, 2013. *METALS Program Overview*. Advanced Research Projects Agency, Department of Energy, Washington DC.
120. Aluminium Federation, 2014. Aluminium and sustainability. Fact Sheet No. 3, Aluminium Federation, Halesowen.
121. IPCC, 2006. IPCC Guidelines for National Greenhouse Gas Inventories. Intergovernmental Panel on Climate Change, National Greenhouse Gas Inventories Programme, Technical Support Unit; Institute for Global Environmental Strategies: Hayama, Japan.
122. Grainger A., 1990. Modelling the impact of alternative afforestation strategies to reduce carbon emissions. In *Proceedings of the IPCC Conference on Tropical Forestry Response Options to Global Climate Change*, São Paulo, Brazil, January 1990: 120-133. Report No. 20P-2003, Office of Policy Analysis, US Environmental Protection Agency, Washington DC: 93-104.
123. Bastin F., Finegold Y., Garcia C., Mollicone D. et al., 2019. The global tree restoration potential. *Science* **365**: 76–79.  
\*A recent assessment, based on satellite data, of the potential for sequestering carbon by expanding forest area.

124. International Union for the Conservation of Nature. Bonn Challenge. <https://www.bonnchallenge.org/>.
125. Keenan R.J., Reams G.A., Achard F., De Freitas J., Grainger A. and Lindquist E., 2015. Dynamics of global forest area, results from FAO Global Forest Resources Assessment 2015. *Forest Ecology and Management* **352**: 9–20.
126. UNFCCC, 2020. Greenhouse Gas Inventory Data - Detailed Data by Party. UN Framework Convention on Climate Change, Bonn. [https://di.unfccc.int/detailed\\_data\\_by\\_party](https://di.unfccc.int/detailed_data_by_party).
127. Lizarralde I., Rodriguez F. and Bravo F., 2016. Innovation in the value chain of wood products: data, equations and life cycle analysis. In Bravo F. et al (eds.). *Managing Forest Ecosystems: The Challenge of Climate Change*: 1-18. Kluwer, Dordrecht.
128. Perez-Garcia J., Lippke B., Cornick J. and Manriquez C., 2005. An assessment of carbon pools, storage and wood products markets substitution using life cycle analysis results. *Wood and Fiber Science* **37**: 140-148.
129. Johnston C.M.T. and Radeloff V.C., 2019. Global mitigation potential of carbon stored in harvested wood products. *Proceedings of the National Academy of Sciences USA* **116**: 14526–14531.
- \*A recent evaluation of the role of harvested wood products in climate change mitigation.
130. Zhao Q., Ding S., Wen Z. and Toppinen A., 2019. Energy flows and carbon footprint in the forestry-pulp and paper industry. *Forests* **10**: 725. <https://doi.org/10.3390/f10090725>
131. Puettmann M.E. and Wilson J.B., 2005. Life-cycle analysis of wood products: cradle-to-gate LCI of residential wood building materials. *Wood and Fiber Science* **37**: 18-29.
132. Abbas D. and Handler R.M., 2018. Life-cycle assessment of forest harvesting and transportation operations in Tennessee. *Journal of Cleaner Production* **176**: 512-520.
133. Hammond G. and Jones C, 2011. *Embodied Carbon: The Inventory of Carbon and Energy (ICE)*. University of Bath/Building Services Research and Information Organization, Bath.
134. Kazulis V., Muizniece I., Zihare L. and Blumberga D., 2017. Carbon storage in wood products. *Energy Procedia* **128**: 558-563.
135. Dias A.C. and Arroja L., 2012. Comparison of methodologies for estimating the carbon footprint—case study of office paper. *Journal of Cleaner Production* **24**: 30-35.
136. Sedjo R.A. and Tian X., 2012. An investigation of the carbon neutrality of wood bioenergy. *Journal of Environmental Protection* **3**: 989-1000.
137. Nian V., 2016. The carbon neutrality of electricity generation from woody biomass and coal, a critical comparative evaluation. *Applied Energy* **179**: 1069-1080.

138. Repo A., Tuovinen J.P. and Liski J., 2015. Can we produce carbon and climate neutral forest bioenergy? *GCB Bioenergy* **7**: 253-262
139. Salazar J. and Meil J., 2009. Prospects for carbon-neutral housing: The influence of greater wood use on the carbon footprint of a single-family residence. *Journal of Cleaner Production* **17**:1563-1571.  
\*A valuable comparison of traditional and wood-intensive houses.
- 140 Mithraratne N. and B Vale B., 2004. Life cycle analysis model for New Zealand houses. *Building and Environment* **39**: 483-492.
141. Andrew R.M., 2018. Global CO<sub>2</sub> emissions from cement production, 1928–2017. *Earth System Science Data* **10**: 2213–2239.
142. Halloran J.W., 2007. Carbon-neutral economy with fossil fuel-based hydrogen energy and carbon materials. *Energy Policy* **35**: 4839-4846.
143. Habert G., Miller S.A., John V.M., Provis J.L. et al., 2020. Environmental impacts and decarbonization strategies in the cement and concrete industries. *Nature Reviews Earth & Environment* **1**: 559–573.
144. Bolin C.T. and Smith S.T., 2011. Life cycle assessment of pentachlorophenol-treated wooden utility poles with comparisons to steel and concrete utility poles. *Renewable and Sustainable Energy Reviews* **15**: 2475-2486.
145. Rempelos G., Preston J. and Blainey S., 2020 A carbon footprint analysis of railway sleepers in the United Kingdom. *Transportation Research D* **81**: 102285.
146. Glesinger E., 1949. *The Coming Age of Wood*. Simon and Schuster, New York.  
\*The original vision of a wood economy.
147. Goldstein I.S., 1975. Potential for converting wood into plastics. *Science* **189**: 847-852.  
\*\*A classic study of how to make the transition from a fossil fuel economy to a wood economy.
148. EC, 2006. Framework 7 Theme 2: Food, Agriculture, Fisheries and Biotechnology, 2007 Work Programme. European Commission, Brussels.
149. Garlapati V.K., Chandel A.K., Kumar S.P.J., Sharma S., Sevda S. et al., 2020. Circular economy aspects of lignin: towards a lignocellulose biorefinery. *Renewable and Sustainable Energy Reviews* **130**:109977.
150. Zheng J and Suh S., 2019. Strategies to reduce the global carbon footprint of plastics. *Nature Climate Change* **9**: 374–378.
151. Zhang Y.H.P. and Mielenz J.R., 2011. Renewable hydrogen carrier—carbohydrate: constructing the carbon-neutral carbohydrate economy. *Energies* **4**: 254-275.

152. Liao Y., Koelewijn S.F., Van den Bossche G. Van Aelst J., Van den Bosch S. et al., 2020. A sustainable wood biorefinery for low-carbon footprint chemicals production. *Science* **367**: 1385-1390.

\*A new study of bio-refineries.

153. Sharifzadeh M., Wang L. and Shah N., 2015. Integrated biorefineries: CO<sub>2</sub> utilization for maximum biomass conversion. *Renewable and Sustainable Energy Reviews* **47**: 151-161.

154. Kärkäs M.D., Matsuura B.S., Monos T.M., Magallanes G. and Stephenson C.R.J., 2016. Transition-metal catalyzed valorization of lignin: the key to a sustainable carbon-neutral future. *Organic and Biomolecular Chemistry* **14**: 1853-1914.

155. Anastas P. and Eghbali N., 2010. Green chemistry: principles and practice. *Chemical Society Reviews* **39**: 301-312.

156. Hata T, Vystavel T., Bronsveld P., DeHosson J. et al., 2004. Catalytic carbonization of wood charcoal. *Carbon* **42**: 961-964.

157. Li D. and Kaner R.B., 2008. Graphene-based materials. *Science* **320**: 1170-1171

158. Baughman R.H., Zakhidov A.A. and De Heer W.A., 2002. Carbon nanotubes-the route toward applications. *Science* **297**:787-792.

159. EC, 2012. Innovating for Sustainable Growth: A Bioeconomy for Europe. European Commission, Brussels.

160 Staffas L., Gustavsson M. and McCormick K., 2013. Strategies and policies for the bioeconomy and bio-based economy: an analysis of official national approaches. *Sustainability* **5**:2751-2769.

\*A valuable comparison of the emergence and growing use of the terms "bioeconomy" and "bio-based economy" in national and international policies.

161. Iverson L.R., Brown S., Grainger A., Prasad A. and Liu D., 1993. Carbon sequestration in South/Southeast Asia: an assessment of technically suitable forest lands using geographic information systems analysis. *Climate Research* **3**: 23-38.

162. Bugge M.M., Hansen T. and Klitkou A., 2016. What is the bioeconomy? A review of the literature. *Sustainability* **8**: 691.

\*A critical comparison of different visions of the "bioeconomy".

163. Roy J., Ghosh D., Ghosh A. and Dasgupta S., 2013. Fiscal instruments: crucial role in financing low carbon transition in energy systems. *Current Opinion in Environmental Sustainability* **5**: 261-269.

164. Ervine K., 2018. How low can it go? Analysing the political economy of carbon market design and low carbon prices. *New Political Economy* **23**: 690-710.

165. Meckling J., Sterner T. and Wagner G., 2017. Policy sequencing toward decarbonization. *Nature Energy* **2**: 918-922.

166. Rosenbloom D. and Rinscheid A., 2020. Deliberate decline: An emerging frontier for the study and practice of decarbonization. *Wiley Interdisciplinary Reviews* **11**: e669.
167. Hsu A., Höhne N., Kuramochi T., Vilariño V. and Sovacool B.K., 2020. Beyond states: harnessing sub-national actors for the deep decarbonisation of cities, regions, and businesses. *Energy Research & Social Science* **70**: 101738
168. Bernstein S. and Hoffmann M., 2018. The politics of decarbonization and the catalytic impact of subnational climate experiments. *Policy Sciences* **51**: 189–211.
169. Huang Y. and Barker T., 2012. The Clean Development Mechanism and low carbon development: A panel data analysis. *Energy Economics* **34**:1033-1040.
170. Alonso P.M., Hewitt R., Pacheco J.D., Bermejo L.R., Jimenez V.H. et al., 2016. Losing the roadmap: Renewable energy paralysis in Spain and its implications for the EU low carbon economy. *Renewable Energy* **89**: 680-694.
171. Robbins, L., 1935. *An Essay on the Nature and Significance of Economic Science*. 2nd edition, Macmillan, London.
172. Daly H.E., 1974. The economics of the steady state. *American Economic Review* **64**: 15-21.
173. Daly H.E., 1994. Operationalizing sustainable development by investing in natural capital. In Jansson A, Hammer M, Folke C. and Costanza R (eds.). *Investing in Natural Capital*: 22- 37. Island Press, Washington DC.
174. Pearce D., 1991. Introduction. In Pearce D. (ed.). *Blueprint 2*: 1-10. Earthscan Publications, London.
175. Grainger A., 2017. The prospect of global environmental relativities after an Anthropocene tipping point. *Forest Policy and Economics* **79**: 36-49.
176. Stocker T.F., Qin D., Plattner G.-K., Tignor M., Allen S.K. et al. (eds.), 2013. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge.
177. Rockstrom J., Steffen W., Noone K., Persson A., Chapin F.S. et al., 2009. A safe operating space for humanity. *Nature*: 472-475.
178. Crawford S. E. S. and Ostrom E., 1995. A grammar of institutions. *American Political Science Review* **89**: 582-600.
179. Spash C.L., 2020. A tale of three paradigms: realising the revolutionary potential of ecological economics. *Ecological Economics* **169**: 106518.
- \*A good evaluation of ecological economics at the crossroads.

180. Williamson O.E., 2000. The new institutional economics. *Journal of Economic Literature* **38**: 595-613.
181. Ostrom E., 1990. *Governing the Commons: The Evolution of Institutions for Collective Action*. Cambridge University Press, Cambridge.
182. Rhodes R.A.W., 1996. The new governance: governing without government. *Political Studies* **44**: 652-667.
183. Andrews-Speed P., 2016 Applying institutional theory to the low-carbon energy transition. *Energy Research & Social Science* **13**: 216-225
184. Dobson N.L., 2018. The EU's conditioning of the 'extraterritorial' carbon footprint: a call for an integrated approach in trade law discourse. *Review of European, Comparative & International Environmental Law* **27**: 75-89.
185. Miles L. and Kapos V., 2008. Reducing greenhouse gas emissions from deforestation and forest degradation: global land-use implications. *Science* **320**: 1454-1455.
186. Ozkaynak B., Adaman F. and Devine P., 2012 The identity of ecological economics: retrospects and prospects. *Cambridge Journal of Economics* **36**: 1123-1142.
187. Falkie T.V. and McKelvey J.E., 1975. Principles of the mineral resource classification system of the US. *USGS Bulletin* 1450A. Bureau of Mines and US Geological Survey. Washington DC.
188. Barido D.P. De L., Avila N. and Kammen D.M., 2020. Exploring the enabling environments, inherent characteristics and intrinsic motivations fostering global electricity decarbonization. *Energy Research & Social Science* **61**: 101343.
189. Kanemoto K. and Moran D., 2019. Carbon-footprint accounting for the next phase of globalization: status and opportunities. *One Earth* **1**: 35-38.
190. WCED, 1987. *Our Common Future*. Report of the World Commission on Environment and Development. United Nations, New York.
191. Shinnar R. and Citro F., 2008. Decarbonization: Achieving near-total energy independence and near-total elimination of greenhouse emissions with available technologies. *Technology in Society* **30**: 1-16.
192. IPCC, 1990. *The IPCC Response Strategies*. First Assessment Report of the Intergovernmental Panel on Climate Change. World Meteorological Organisation and UN Environment Programme, Geneva.
193. Steinmann Z.J.N, Huijbregts M.A.J. and Reijnders L., 2018. How to define the quality of materials in a circular economy? *Resources, Conservation and Recycling* **14**: 362-363.
194. Hodgson C., 2021. MPs allege lack of joined-up plan on emissions targets. *Financial Times*, 5 March 2021.

195. Deutch J., 2020. Is Net Zero Carbon 2050 possible? *Joule* **4**: 1-4.

196. Rogelj J., Geden O., Cowie A. and Reisinger A., 2021. Three ways to improve net-zero emissions targets. *Nature* **591**: 365-368.

\*A recent evaluation of policy options for carbon neutrality.

Table 1. Numbers of papers in international journals in seven carbon literatures by principal categories 1995-2020.

	Decarbonization	%	Low Carbon Economy	%	Carbon Transition	%	Low Carbon Development	%	Carbon Neutrality	%	Zero Net Carbon Emissions	%	Carbon Footprint	%	Total	%
Energy	228	43	17	11	33	26	35	29	36	26	11	14	106	10	466	21
Transport	65	12	8	5	8	6	1	1	13	9	1	1	79	7	176	8
Buildings	34	6	0	0	6	5	6	5	11	8	55	70	59	5	171	8
Materials	26	5	3	2	7	5	0	0	10	7	0	0	98	9	144	6
Land Use	0	0	2	1	0	0	0	0	4	3	0	0	345	31	351	15
Forests	5	1	1	1	1	1	0	0	6	4	0	0	3	0	16	1
Economy	47	9	49	32	24	19	8	7	0	0	0	0	46	4	174	8
Development	7	1	0	0	3	2	8	7	0	0	0	0	3	0	21	1
Tourism	5	1	6	4	0	0	0	0	0	0	0	0	35	3	46	2
Decarbonization technology	50	9	0	0	0	0	0	0	0	0	0	0	0	0	50	2
Industry	12	2	15	10	3	2	7	6	0	0	0	0	90	8	127	6
Organizations	1	0	0	0	2	2	0	0	6	4	0	0	37	3	46	2
Urban	19	4	5	3	6	5	30	25	18	13	1	1	42	4	121	5
Local	2	0	4	3	3	2	4	3	15	11	6	8	4	0	38	2
National	9	2	37	24	22	17	12	10	11	8	2	3	32	3	125	6
Others	24	4	5	3	11	9	10	8	8	6	3	1	133	12	194	9
Total	534	24	152	7	129	6	121	5	138	6	79	3	1,116	49	2,265	

NB. All categories include synonyms, and zero net carbon emissions publications also include those on zero net energy.



Table 2. Three conditions for achieving carbon neutrality.

1. Minimize carbon dioxide emissions when producing and using materials and energy sources, to accelerate peaking of emissions and passage through the carbon transition.
2. Maximize carbon dioxide removals.
3. Offset carbon dioxide emissions by carbon dioxide removals.

Table 3. Classifying materials by their natural and embodied carbon contents.

Natural carbon content	Embodied carbon content	
	Low	High
Low	VERY LOW CARBON Non-carboniferous stone	LOW CARBON Metals
High	HIGH CARBON Wood products	EXTREME CARBON Cement

Table 4. National carbon transition curve characteristics of 17 countries.

	Carbon Peak Year	Chordal Equivalent of 2014	Decarbonization Ratio	Percent peak emissions in 2014	CO <sub>2</sub> half life (years)
<i>Right skewed</i>					
United Kingdom	1971	<1960	na	64	59
Sweden	1970	<1960	na	47	37
France	1979	1962	2.1	57	41
Belgium	1979	1961	1.9	67	25
Hungary	1984	<1960	na	46	28
Poland	1987	1969	1.5	61	35
<i>Quasi-symmetrical</i>					
Bulgaria	1987	1964	1.2	46	25
Romania	1989	1963	1.0	33	19
<i>Left skewed</i>					
Denmark	1996	1961	0.5	46	17
Portugal	2002	1992	1.2	67	18
Finland	2003	1972	0.4	68	17
Italy	2004	1971	0.3	68	15
Austria	2005	1976	0.3	79	22
Greece	2005	1988	0.5	68	14
Ireland	2007	1996	0.6	76	15
Spain	2007	1996	0.6	65	10
Croatia	2007	1993	0.5	70	12

*Curves without peaks:* Luxembourg, Netherlands, Norway, Slovenia, Switzerland, USA.

NB. Values of the Decarbonization Ratio are not available (na) for the United Kingdom, Sweden and Hungary, because the Chordal Equivalent Year precedes the beginning of the database that we used. Data were obtained from the World Bank (89).

Table 5. Carbon dioxide emitted in the production of five key materials in 2005<sup>1</sup> and 2015<sup>2</sup>.

Material	Global Production (Mt/a <sup>1</sup> )		Global CO <sub>2</sub> emissions <sup>3</sup> (Mt.a <sup>1</sup> )		Percent all CO <sub>2</sub> emissions <sup>4</sup>	
	2005	2015	2005	2015	2005	2015
Zinc	10.5	13.7	47.0	61.3	0.16	0.17
Copper	15.6	23.0	60.0	88.5	0.20	0.25
Aluminium	38.0	57.5	830.0	1,255.9	2.81	3.49
Steel	924.0	1,620.0	2,000.0	3,506.5	6.78	9.75
<i>Total</i>	<i>988.1</i>	<i>1,714.2</i>	<i>2,937.0</i>	<i>4,912.2</i>	<i>9.95</i>	<i>13.66</i>
Cement	2,600.0	4,100.0	2,300.0	3,626.9	7.79	10.08

NB. <sup>1</sup>The 2005 estimates are from Rankin (77). <sup>2</sup>The 2015 production estimates are from the US Geological Survey (97-100), and emissions are calculated using the same conversion factors as in Rankin (77). <sup>3</sup>Conversion to emissions assumes that all electricity is generated by black coal. <sup>4</sup>CO<sub>2</sub> emissions apart from those for land use change are 29,508 Mt in 2005 and 35,977 Mt in 2015 (89).

Table 6. Key energy and carbon categories of four leading metals (77).

Metal	Extraction		Production Process	Percent all production	Refining		Embodied		Percent Refining Energy	Input	Recycling	
	Energy (GJ/t metal)	Carbon (tCO <sub>2</sub> /t metal)			Energy (GJ/t)	Carbon (tCO <sub>2</sub> /t metal)	Energy (GJ/t)	Carbon (tCO <sub>2</sub> /t metal)			Energy (GJ/t)	Ratio
Copper	19.60	1.93	pyro	80	13.42	1.32	33.02	3.25	40.6	Scrap No. 1	4.4	0.33
	18.17	1.74	hydro	20	46.29	4.42	64.46	6.16	71.8	Scrap No. 2	20.1	1.50
										Low grade	49.3	3.67
Zinc	5.46	0.52	electrolytic	90	42.98	4.09	48.44	4.61	88.7	New scrap	3.8	0.12
	4.96	0.46	ISP	10	30.89	2.88	35.85	3.34	86.2	Slab	22.0	0.71
Aluminium	0.36	0.04	electrolytic	100	211.15	21.77	211.51	21.81	99.8	Alloy	17.5	0.08
Steel	0.56	0.05	BF/BOF	70	22.14	2.14	22.70	2.19	97.5	Billets	9.7	0.44

NB. ISP = imperial smelting process; BF = blast furnace; BOF = basic oxygen furnace. The Recycling ratio (R<sub>i</sub>) is calculated by dividing Recycling Energy by Refining Energy, and for copper the lower of the two Refining Energy values is used.

Table 7. The physical properties of three light metals and steel (119).

	Aluminium	Magnesium	Titanium	Steel
Density (Kg/m <sup>3</sup> )	2,710	1,740	4,510	7,860
Strength to weight ratio (kN-m/kg)	130	158	120	80

Table 8. Trends in changes in the areas of natural forest and forest plantations 1990-2015 (125).

	1990s	2000-05 million hectares a <sup>-1</sup>	2005-10	2010-15	Total area change million hectares
Forest plantations					
Non-tropical	3.36	4.65	3.81	1.60	83.87
Tropical	0.84	1.26	1.49	0.86	26.52
World	4.20	5.91	5.30	2.46	110.38
Natural forest					
Non-tropical	-1.08	-1.36	-0.61	0.61	-17.61
Tropical	-10.39	-9.13	-8.10	-6.38	-221.91
World	-11.47	-10.48	-8.72	-5.77	-239.52

Table 9. Typical values of extraction energy and carbon for forestry in the USA (132), and of the embodied energy and carbon of selected wood products in the UK showing total energy expended using fossil fuels and timber waste (133).

Product	Extraction energy	Extraction carbon	Embodied energy		Embodied carbon
	(GJ/dry t)	(t CO <sub>2</sub> /dry t)	Total (GJ/t)	Fossil fuels (GJ/t)	(t CO <sub>2</sub> /t)
Sawn softwood	0.59	0.04	7.4	3.2	0.20
Sawn hardwood	0.59	0.04	10.4	4.1	0.24
Particle board	0.59	0.04	14.5	11.1	0.54
Plywood	0.59	0.04	15.0	7.9	0.45



Figure 1. Trends in journal publications on carbon neutrality, carbon transition, zero net carbon emissions, low carbon development, low carbon economy, decarbonization and carbon footprint in five year periods from 1995 to 2019, by initial year.




Figure 2. Carbon neutrality achieved at the intersection of the carbon transition curve and the carbon sequestration curve. The carbon transition begins at the Carbon Peak and ends with zero carbon emissions (based on (19)).




Figure 3. Forest transition curve (36), elaborated to show associated carbon fluxes, and examples of countries which are still to pass through their transitions, or close to doing so, and those that have passed through their transitions.




Figure 4. A circular economy framework (based on 65), which shows flows of resources between Renewable and Non-Renewable Resource Capital and (as energy resources and materials) the Productive Capital (P) and Consumer Goods Capital (C) in the human economy, together with CO<sub>2</sub> inputs from the atmosphere to Renewable Resource Capital and outputs of waste from using Productive Capital and Consumer Goods Capital. Flows are labelled using the names of the principal life cycle stages.




Figure 5. Evaluating the relative speeds of carbon transitions using the number of years before the Carbon Peak ( $Y_{pre}$ ) and the number of years after the Carbon Peak ( $Y_{post}$ ).




Figure 6. The carbon transition curves for four industrialized countries, showing trends in CO<sub>2</sub> emissions (Mt CO<sub>2</sub> a<sup>-1</sup>) over time (89), to illustrate different degrees of symmetry: symmetrical (Romania), right-skewed (France), left-skewed (Italy) and minimal change (Switzerland).




Figure 7. The relationship between the Decarbonization Ratio and the Carbon Peak Year for 17 industrialized countries.




Figure 8. Estimates of the potential (green) and current (black) embodied carbon of aluminium (103-105), copper (96, 102), steel (106-109) and zinc (110-112), with CSIRO estimates (77) in red.






Figure 9. The carbon neutrality gap (Mt CO<sub>2</sub> eq a<sup>-1</sup>) between greenhouse gas emissions outside the land use sector and net terrestrial sequestration in industrialized countries (in 2018) and in other countries (various dates) (126).




Figure 10. Carbon storage in forest and wood products pools with substitution for other products (here the use of concrete in house construction), showing two rotations of 80 years with two thinnings in each rotation of forests in the Pacific Northwest region of the USA (based on 128).




Figure 11. A carbon economy framework, combining the ecosystem and circular economy frameworks and a new institutionalism framework, and showing the transfer of materials between the biosphere and the Productive Capital (P) and Consumer Goods Capital (C) in the human economy as it expands towards ultimate carrying capacity. The institutional matrix (inset) shows variation in the density of institutions at different scales, according to the scales at which institutions originate in different economic or policy sectors. Critical Natural Capital (174) is that remaining above ultimate carrying capacity.





Figure 12. A carbon economy box showing the rise in the carbon price (117) needed to shift the carbon feasibility line downward from current (red and black) values of the embodied carbon of steel, to make potential steel technologies (green) economic. Locations of estimates: (1) Poland, (2) Australia, (3) China, (4) Europe, and (5) Italy (77, 106-109).



Supplementary Table S1. The subjects of 144 papers on “Materials” and 16 papers on “Forests” in the 2,265 papers in seven carbon literatures in Table 1.

<b>Materials</b>	
Aluminium	4
Copper	2
Iron and steel	17
Zinc	0
Other metals	6
Cement	39
Plastics	9
Chemicals	7
Energy intensive industries	10
Other general studies	24
Forest products	26
<i>Sawnwood/pallets</i>	7
<i>Wood panels</i>	2
<i>Paper</i>	5
<i>Cork</i>	3
<i>Synthetic chemicals</i>	9
<b>Forests</b>	
Carbon sequestration	10
Fuelwood	6

Supplementary Table S2. The scope of 65 individual life cycle analyses for four leading metals.

	Aluminium	Copper	Steel	Zinc	Percentage
Primary production	4	4	4	6	28
Metal products	1	4	1	2	12
Comparisons	1	2	11	0	22
Recycling	4	7	7	0	28
Other	1	2	4	0	11
Total	11	19	27	8	100