



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/177573/>

Version: Accepted Version

---

**Article:**

Dabrowski, KK, Dross, F, Jeong, J et al. (2022) Tree Pivot-Minors and Linear Rank-Width. SIAM Journal on Discrete Mathematics, 35 (4). pp. 2922-2945. ISSN: 0895-4801

<https://doi.org/10.1137/21M1402339>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Tree Pivot-Minors and Linear Rank-Width\*

Konrad K. Dabrowski<sup>1</sup>, François Dross<sup>2</sup>, Jisu Jeong<sup>3</sup>, Mamadou Moustapha Kanté<sup>4</sup>,  
O-joung Kwon<sup>5,6</sup>, Sang-il Oum<sup>6,7</sup>, and Daniël Paulusma<sup>1</sup>

<sup>1</sup>School of Computing, University of Leeds, UK

<sup>2</sup>Institute of Informatics, University of Warsaw, Poland

<sup>3</sup>Clova AI Research, NAVER Corp, Seongnam, Korea

<sup>4</sup>Université Clermont Auvergne, LIMOS, CNRS, Aubière, France

<sup>5</sup>Department of Mathematics, Incheon National University, Incheon, Korea

<sup>6</sup>Discrete Mathematics Group, Institute for Basic Science (IBS), Daejeon, Korea

<sup>7</sup>Department of Mathematical Sciences, KAIST, Daejeon, Korea

August 4, 2021

## Abstract

Tree-width and its linear variant path-width play a central role for the graph minor relation. In particular, Robertson and Seymour (1983) proved that for every tree  $T$ , the class of graphs that do not contain  $T$  as a minor has bounded path-width. For the pivot-minor relation, rank-width and linear rank-width take over the role of tree-width and path-width. As such, it is natural to examine if, for every tree  $T$ , the class of graphs that do not contain  $T$  as a pivot-minor has bounded linear rank-width. We first prove that this statement is false whenever  $T$  is a tree that is not a caterpillar. We conjecture that the statement is true if  $T$  is a caterpillar. We are also able to give partial confirmation of this conjecture by proving:

- for every tree  $T$ , the class of  $T$ -pivot-minor-free distance-hereditary graphs has bounded linear rank-width if and only if  $T$  is a caterpillar;
- for every caterpillar  $T$  on at most four vertices, the class of  $T$ -pivot-minor-free graphs has bounded linear rank-width.

To prove our second result, we only need to consider  $T = P_4$  and  $T = K_{1,3}$ , but we follow a general strategy: first we show that the class of  $T$ -pivot-minor-free graphs is contained in some class of  $(H_1, H_2)$ -free graphs, which we then show to have bounded linear rank-width. In particular, we prove that the class of  $(K_3, S_{1,2,2})$ -free graphs has bounded linear rank-width, which strengthens a known result that this graph class has bounded rank-width.

---

\*An extended abstract of this paper appeared in the proceedings of EuroComb 2019 [21]. Dabrowski and Paulusma were supported by the Leverhulme Trust (RPG-2016-258). This work was mainly done when Jeong was in KAIST. Kanté was supported by the French Agency for Research under the projects DEMOGRAPH (ANR-16-CE40-0028) and ASSK (ANR-18-CE40-0025). Kwon was supported by the National Research Foundation of Korea funded by the Ministry of Education (No. NRF-2018R1D1A1B07050294). Kwon and Oum were supported by the Institute for Basic Science (IBS-R029-C1). Dross was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme Grant Agreement 714704.

<sup>1</sup>E-mail addresses: [k.k.dabrowski@leeds.ac.uk](mailto:k.k.dabrowski@leeds.ac.uk) (Dabrowski), [francois.dross@gmail.com](mailto:francois.dross@gmail.com) (Dross), [jisujeong89@gmail.com](mailto:jisujeong89@gmail.com) (Jeong), [mamadou.kante@uca.fr](mailto:mamadou.kante@uca.fr) (Kanté), [ojoungkwon@gmail.com](mailto:ojoungkwon@gmail.com) (Kwon), [sangil@ibs.re.kr](mailto:sangil@ibs.re.kr) (Oum), [daniel.paulusma@durham.ac.uk](mailto:daniel.paulusma@durham.ac.uk) (Paulusma).

# 1 Introduction

In order to increase our understanding of graph classes, it is natural to consider some notion of “width” and to research what properties graph classes of bounded width may have. We say that a graph class has *bounded* width (for some specific width parameter) if there exists a constant  $c$  such that the width of every graph in the class is at most  $c$ . In particular, this type of structural research has been done in the context of graph containment problems, where the aim is to determine whether a graph  $H$  appears as a “pattern” inside some other graph  $G$ . Here, a pattern is defined by specifying a set of graph operations that may be used to obtain  $H$  from  $G$ . For instance, a graph  $G$  contains a graph  $H$  as a *minor* if  $H$  can be obtained from  $G$  via a sequence of vertex deletions, edge deletions and edge contractions.

Tree-width and its *linear* variant, path-width, are the best-known graph width parameters due to their relevance for graph minor theory [47]. Rank-width is another well-known parameter, introduced by Oum and Seymour [43]. The rank-width of a graph  $G$  expresses the minimum width  $k$  of a tree-like structure obtained by recursively splitting the vertex set of  $G$  in such a way that each cut induces a matrix of rank at most  $k$  (see Section 2 for a formal definition). Rank-width is more general than tree-width in the sense that every graph class of bounded tree-width has bounded rank-width, but there are classes for which the reverse does not hold, for example, the class of all complete graphs [17].

The notion of rank-width has important algorithmic implications, as many NP-complete decision problems are known to be polynomial-time solvable not only for graph classes of bounded tree-width, but also for graph classes of bounded rank-width; see [16, 25, 28, 32, 44] for a number of meta-theorems capturing such decision problems. Rank-width is equivalent to clique-width [43], another important and well-studied width parameter. Linear rank-width is a linearized variant of rank-width, known to be equivalent to linear clique-width (see, for example, [42]) and to be closely related to the trellis-width of linear codes [31]. We formally define the notions of rank-width and linear rank-width in Section 2.

The problem of determining whether a given graph has linear rank-width at most  $k$  for some given integer  $k$  is NP-complete (this follows from a result of Kashyap [31]). On the positive side, Jeong, Kim, and Oum [29] gave an FPT algorithm for deciding whether a graph has linear rank-width at most  $k$ . Ganian [26] and Adler, Farley, and Proskurowski [1] characterized the graphs of linear rank-width at most 1. Recently, Nešetřil et al. [39] showed that every class of bounded linear rank-width is linearly  $\chi$ -bounded. However, our knowledge on linear rank-width, the topic of this paper, is still limited.

## Motivation

To increase our understanding of rank-width and linear rank-width, we may want to verify if classical results for tree-width and path-width stay valid when we replace tree-width with rank-width and path-width with linear rank-width. The following two structural results, related to path-width and tree-width, form the core of the Graph Minor Structure Theorem. Here, a graph  $G$  is *H-minor-free* for some graph  $H$  if  $G$  does not contain  $H$  as a minor.

**Theorem 1.1** (Robertson and Seymour [45]). *For every tree  $T$ , the class of  $T$ -minor-free graphs has bounded path-width.*

**Theorem 1.2** (Robertson and Seymour [46]). *For every planar graph  $H$ , the class of  $H$ -minor-free graphs has bounded tree-width.*




Figure 1: A graph before and after pivoting an edge (the example is taken from [20]).

It is known that edge deletions and contractions may increase the rank-width and linear rank-width [15]. Hence, working with minors is not a suitable approach for understanding rank-width and linear rank-width. Therefore, Oum [40] proposed the notions of vertex-minors and pivot-minors, two closely related notions, which were called  $\ell$ -reductions and  $p$ -reductions, respectively, in [11]. Taking vertex-minors or pivot-minors does *not* increase the rank-width or linear rank-width of a graph [40].

To define the notions of a vertex-minor and a pivot-minor, we need some terminology. The *local complementation* at a vertex  $u$  in a graph  $G$  replaces every edge of the subgraph induced by the neighbours of  $u$  with a non-edge, and vice versa. The resulting graph is denoted by  $G * u$ . An *edge pivot* is the operation that takes an edge  $uv$ , first applies a local complementation at  $u$ , then at  $v$ , and then at  $u$  again. We denote the resulting graph  $G \wedge uv = G * u * v * u$ . It is known that  $G * u * v * u = G * v * u * v$  [40], and thus  $G \wedge uv = G \wedge vu$ . An alternative definition of the edge pivot operation is as follows. Let  $S_u$  be the set of neighbours of  $u$  that are non-adjacent to  $v$  and let  $S_v$  be the set of neighbours of  $v$  that are non-adjacent to  $u$ , whereas we denote the set of common neighbours of  $u$  and  $v$  by  $S_{uv}$ . We replace every edge between any two vertices in distinct sets from  $\{S_u \setminus \{v\}, S_v \setminus \{u\}, S_{uv}\}$  by a non-edge and vice versa. Afterwards, we delete every edge between  $u$  and  $S_u$  and add every edge between  $u$  and  $S_v$ . We also delete every edge between  $v$  and  $S_v$  and add every edge between  $v$  and  $S_u$ . We refer to Figure 1 for an example.

A graph  $H$  is a *vertex-minor* of a graph  $G$  if  $H$  can be obtained from  $G$  by a sequence of local complementations and vertex deletions. A graph  $H$  is a *pivot-minor* of a graph  $G$  if  $H$  can be obtained from  $G$  by a sequence of edge pivots and vertex deletions. Hence  $H$  is a vertex-minor of  $G$  if  $H$  is a pivot-minor of  $G$ , but the reverse is not necessarily true. A graph is  *$H$ -vertex-minor-free* if it contains no vertex-minor isomorphic to  $H$ , and similarly, a graph is  *$H$ -pivot-minor-free* if it contains no pivot-minor isomorphic to  $H$ .

It is natural to ask whether parallel statements of Theorems 1.1 and 1.2 exist for rank-width or linear rank-width in terms of vertex-minors or pivot-minors. Below we discuss the state-of-the-art for this research direction.

## Related Work

A *circle graph* is the intersection graph of chords on a circle, and it is known that the class of circle graphs is closed under taking vertex-minors. Bouchet [11] characterized circle graphs in terms of three forbidden vertex-minors. Oum [40] showed that the class of circle graphs has unbounded rank-width, and asked, as an analogue to Theorem 1.2 for the vertex-minor relation, whether for every circle graph  $H$ , the class of  $H$ -vertex-minor-free graphs has bounded

rank-width. Recently, Geelen et al. [27] gave an affirmative answer to this question.

**Theorem 1.3** (Geelen, Kwon, McCarty, and Wollan [27]). *For every circle graph  $H$ , the class of  $H$ -vertex-minor-free graphs has bounded rank-width.*

Every pivot-minor of a graph is also a vertex-minor. Hence, for every graph  $H$ , the class of  $H$ -vertex-minor-free graphs is contained in the class of  $H$ -pivot-minor-free graphs. This leads to the question whether we can strengthen Theorem 1.3 by replacing the vertex-minor relation with the pivot-minor relation. However, this is not the case. In order to see this, we first observe that bipartite graphs are closed under taking pivot-minors [40]. Hence, no bipartite graph contains a non-bipartite circle graph as a pivot-minor. Now consider the class  $\mathcal{G}$  of  $n \times n$  grids, which has unbounded rank-width. As  $\mathcal{G}$  is a subclass of bipartite graphs,  $\mathcal{G}$  is  $H$ -pivot-minor-free for every non-bipartite circle graph  $H$  (such as, for example,  $H = K_3$ ). Hence, for every non-bipartite graph  $H$ , the class of  $H$ -pivot-minor-free graphs has unbounded rank-width. This means we can only hope to strengthen Theorem 1.3 by considering bipartite circle graphs  $H$ , and Oum [41] conjectured the following analogue to Theorem 1.2 for the pivot-minor relation:

**Conjecture 1** (Oum [41]). *For every bipartite circle graph  $H$ , the class of  $H$ -pivot-minor-free graphs has bounded rank-width.*

So far, Conjecture 1 has been verified for bipartite graphs [40], circle graphs [41], and line graphs [41]. If Conjecture 1 holds for all graphs, then this would imply both Theorems 1.2 and 1.3 [40].

We now turn to linear rank-width, for which Kanté and Kwon [30] conjectured the following analogue to Theorem 1.1 for the vertex-minor relation:

**Conjecture 2** (Kanté and Kwon [30]). *For every tree  $T$ , the class of  $T$ -vertex-minor-free graphs has bounded linear rank-width.*

So far, Conjecture 2 has been verified on every class of graphs whose prime graphs, with respect to split decompositions, have bounded linear rank-width [30]. For example, prime distance-hereditary graphs have at most three vertices, and therefore, for every tree  $T$ , the class of  $T$ -vertex-minor-free distance-hereditary graphs has bounded linear rank-width. Moreover, Conjecture 2 holds for every path  $T$  [33].

## Our Focus and Results

We focus on the remaining analogue, namely the analogue to Theorem 1.1 for the pivot-minor relation. We first prove that we cannot hope for a result that holds for every tree  $T$ . A *caterpillar* is a tree that contains a path  $P$ , such that every vertex not on  $P$  has a neighbour in  $P$ .

**Theorem 1.4.** *If  $T$  is a tree that is not a caterpillar, then the class of  $T$ -pivot-minor-free distance-hereditary graphs has unbounded linear rank-width.*

Due to Theorem 1.4, we conjecture the following:

**Conjecture 3.** *For every caterpillar  $T$ , the class of  $T$ -pivot-minor-free graphs has bounded linear rank-width.*

In contrast, the aforementioned result of Kwon et al. [33] confirming Conjecture 2 if  $T$  is a path implies that Conjecture 2 *has* been confirmed if  $T$  is a caterpillar: every caterpillar  $T$  is a pivot-minor of some path  $P$  [34, Theorem 4.6] and consequently, if  $T$  is a caterpillar, then the class of  $T$ -vertex-minor-free graphs is contained in the class of  $P$ -vertex-minor-free graphs.

By the fact that every caterpillar  $T$  is a pivot-minor of some path  $P$  and the fact that every path  $P$  is a caterpillar by definition, we can also formulate Conjecture 3 as follows:

**Conjecture 3 (alternative formulation).** *For every path  $P$ , the class of  $P$ -pivot-minor-free graphs has bounded linear rank-width.*

We make two contributions to Conjecture 3. We first show, in Section 4, that Conjecture 3 holds for distance-hereditary graphs.

**Theorem 1.5.** *Let  $n \geq 3$  be an integer. Every  $P_n$ -pivot-minor-free distance-hereditary graph has linear rank-width at most  $2n - 5$ .*

Theorems 1.4 and 1.5, together with the fact that every caterpillar is a pivot-minor of some path, yields the following dichotomy.

**Corollary 1.6.** *For every tree  $T$ , the class of  $T$ -pivot-minor-free distance-hereditary graphs has bounded linear rank-width if and only if  $T$  is a caterpillar.*

If a graph  $G$  is  $P_4$ -pivot-minor-free, then  $G$  has no induced subgraph isomorphic to  $P_4$ . This implies that  $G$  is distance-hereditary. Hence, Theorem 1.5 has the following consequence:

**Corollary 1.7.** *Every  $P_4$ -pivot-minor-free graph has linear rank-width at most 3.*

Below we give a short alternative proof of Corollary 1.7 (without an explicit bound) after introducing a more general strategy. A graph  $G$  is  $H$ -free if  $G$  does not contain the graph  $H$  as an induced subgraph, and  $G$  is  $(H_1, \dots, H_p)$ -free for some set of graphs  $\{H_1, \dots, H_p\}$  if  $G$  is  $H_i$ -free for every  $i \in \{1, \dots, p\}$ . We can now try to obtain for a caterpillar  $T$ , a constant bound on the linear rank-width of a  $T$ -pivot-minor-free graph by adapting the following *general* strategy:

**Step 1.** Show that the class of  $T$ -pivot-minor-free graphs is a subclass of a class of  $(H_1, H_2)$ -free graphs for some graphs  $H_1$  and  $H_2$ .

**Step 2.** Show that this class of  $(H_1, H_2)$ -free graphs has bounded linear rank-width.

An advantage of this strategy is that it will lead to a stronger result that forms the start of a *systematic* study into boundedness of linear rank-width of  $(H_1, H_2)$ -free graphs. This would address Open Problem 7.5 in [22], which asks for such a result. We refer to Section 6 for a further discussion on this.

To illustrate our general strategy for the case where  $H = P_4$ , we can do as follows. In Step 1, we observe that every  $P_4$ -pivot-minor graph is  $(P_4, \text{dart})$ -free (see Figure 2 for an illustration of the dart). In Step 2, we use a result of Brignall, Korpelainen, and Vatter [14], who showed that a class of  $(P_4, H)$ -free graphs has bounded linear rank-width if and only if  $H$  is a threshold graph. Hence, as the dart is a threshold graph, the class of  $(P_4, \text{dart})$ -free graphs, and thus the class of  $P_4$ -pivot-minor-free graphs, has bounded linear rank-width.

Whether  $P_n$ -pivot-minor-free graphs have bounded linear rank-width for  $n \geq 5$  remains a challenging open question. In the remainder, we focus on the other tree  $T$  on four vertices besides the  $P_4$ , which is the *claw*  $K_{1,3}$  (the 4-vertex star). We will prove the following result.




Figure 2: The graphs claw,  $P_5$ , bull,  $W_4$ , dart, and  $\overline{BW_3}$ .

**Theorem 1.8.** *Every claw-pivot-minor-free graph has linear rank-width at most 59.*

As a consequence, we have verified (the original formulation of) Conjecture 3 for every caterpillar  $T$  on at most four vertices. Since every tree on at most four vertices is a caterpillar we have in fact shown that if  $T$  is a tree on at most four vertices, then the class of  $T$ -pivot-minor-free graphs has bounded linear rank-width.

We first explain how we perform Step 1. In our previous paper [20], we proved that a graph is claw-pivot-minor-free if and only if it is (bull, claw,  $P_5$ ,  $W_4$ ,  $\overline{BW_3}$ )-free; see Figure 2 for pictures of these forbidden induced graphs. It is readily seen that to prove boundedness of linear rank-width of some graph class  $\mathcal{G}$  one may restrict to connected graphs in  $\mathcal{G}$ . In [20] we showed that a graph  $G$  is (bull, claw,  $P_5$ )-free if and only if every component of  $G$  is  $3P_1$ -free (the graph  $3P_1$  consists of three isolated vertices). Hence, we derive the following result, in which we specify the graphs  $H_1$  and  $H_2$  of Step 1 as  $H_1 = 3P_1$  and  $H_2 = W_4$ .

**Theorem 1.9** (Dabrowski et al. [20]). *Let  $G$  be a connected graph. Then  $G$  is claw-pivot-minor-free if and only if  $G$  is  $(3P_1, W_4, \overline{BW_3})$ -free. In particular, the class of connected claw-pivot-minor-free graphs belongs to the class of  $(3P_1, W_4)$ -free graphs.*

As Step 2, we must prove that  $(3P_1, W_4)$ -free graphs have bounded linear rank-width. With an eye on a future classification of boundedness of linear rank-width for  $(H_1, H_2)$ -free graphs, we aim to prove boundedness for classes of  $(H_1, H_2)$ -free graphs that are as large as possible. For integers  $1 \leq i \leq j \leq k$ , let  $S_{i,j,k}$  denote the *subdivided claw*, which is the graph obtained from the claw by subdividing its three edges  $i - 1$  times,  $j - 1$  times and  $k - 1$  times, respectively; see also Figure 3 and note that  $S_{1,1,1} = K_{1,3}$ . The *complement* of a graph  $G$  is the graph  $\overline{G}$  with  $V(\overline{G}) = V(G)$  and  $E(\overline{G}) = \{uv \mid u, v \in V(G) \text{ with } u \neq v \text{ and } uv \notin E(G)\}$ . The graph  $K_3$  denotes the *triangle*. As  $3P_1 = \overline{K_3}$  and  $W_4 = \overline{P_1 + 2P_2}$  is an induced subgraph of the complement of  $\overline{S_{1,2,2}}$ , the class of  $(3P_1, W_4)$ -free graphs is contained in the class of  $(\overline{K_3}, \overline{S_{1,2,2}})$ -free graphs. We prove the following result in Section 5:

**Theorem 1.10.** *Every  $(K_3, S_{1,2,2})$ -free graph has linear rank-width at most 58.*

As observed in Lemma 2.2 in Section 2, complementing a graph may increase its linear rank-width by at most 1. Hence, Theorems 1.9 and 1.10 and this observation imply Theorem 1.8.




Figure 3: The graphs  $S_{1,2,2}$  and  $P_1 + 2P_2$ ; note that  $P_1 + 2P_2$  is an induced subgraph of  $S_{1,2,2}$ .

Dabrowski et al. [19] proved that the class of  $(K_3, S_{1,2,2})$ -free graphs has bounded rank-width. As every class of bounded linear rank-width has bounded rank-width, but the reverse is not necessarily true, Theorem 1.10 is a strengthening of their result. Moreover, Theorem 1.10 is tight in the sense that even the class of  $S_{1,2,3}$ -free bipartite graphs is known to have unbounded linear rank-width [4].

It remains to prove Theorems 1.4, 1.5 and 1.10, which we do in Sections 3, 4 and 5, respectively. We discuss future research in Section 6.

## 2 Preliminaries

In this paper, all graphs have no loops and no multiple edges. For a graph  $G$ , let  $V(G)$  and  $E(G)$  denote the vertex set and edge set of  $G$ , respectively. For  $S \subseteq V(G)$ , let  $G[S] = (S, \{uv \mid uv \in E, u, v \in S\})$  denote the subgraph of  $G$  induced by  $S$ . For convenience, we write  $G[v_1, v_2, \dots, v_m]$  for  $G[\{v_1, v_2, \dots, v_m\}]$ . A graph  $H$  is an *induced subgraph* of  $G$  if  $H = G[S]$  for some  $S \subseteq V(G)$ . For a vertex  $v \in V(G)$ , we let  $G - v$  be the graph obtained from  $G$  by removing  $v$ . For a set  $S \subseteq V(G)$ , we let  $G - S$  be the graph obtained from  $G$  by removing all vertices in  $S$ . For an edge  $e \in E(G)$ , we let  $G - e$  be the graph obtained from  $G$  by removing  $e$ . For a set  $F \subseteq E(G)$ , we let  $G - F$  be the graph obtained from  $G$  by removing all edges in  $F$ .

The set of neighbours of a vertex  $v$  in a graph  $G$  is denoted by  $N_G(v)$ . The size of  $N_G(v)$  is the *degree* of  $v$ . For a set  $A \subseteq V(G)$ , we let  $N_G(A)$  denote the set of all vertices in  $V(G) \setminus A$  that have a neighbour in  $A$ . Two vertices  $v$  and  $w$  in  $G$  are *twins* if  $N_G(v) \setminus \{w\} = N_G(w) \setminus \{v\}$ . We say that twins  $v$  and  $w$  are *false twins* if  $v$  is not adjacent to  $w$ . An edge  $e$  of a connected graph  $G$  is a *cut edge* if  $G - e$  is disconnected.

Let  $A$  and  $B$  be two disjoint vertex subsets of a graph  $G$ . We let  $G \times (A, B)$  be the graph obtained from  $G$  by taking a *bipartite complementation* between  $A$  and  $B$ , that is, by replacing each edge between a vertex of  $A$  and a vertex of  $B$  by a non-edge, and vice versa. We say that  $A$  is *complete* to  $B$  if  $a$  is adjacent to  $b$  for every  $a \in A$  and every  $b \in B$ , whereas  $A$  is *anti-complete* to  $B$  if  $a$  is not adjacent to  $b$  for every  $a \in A$  and every  $b \in B$ . If  $A$  is complete or anti-complete to  $B$ , then  $A$  is *trivial* to  $B$ . If  $A$  consists of one vertex  $v$ , then we may say that  $v$  is complete or anti-complete to  $B$ .

A set  $F$  of edges is a *matching* in a graph  $G$  if no two edges in  $F$  have a common end-vertex. A set  $S$  of vertices in a graph  $G$  is an *independent set* if no two vertices in  $S$  are adjacent, whereas  $S$  is a *clique* if every pair of vertices in  $S$  is adjacent. The complete graph  $K_n$  is the graph on  $n$  vertices that form a clique. The complete bipartite graph  $K_{n,m}$  is the bipartite graph with a bipartition  $(A, B)$  such that  $|A| = n$ ,  $|B| = m$ , and  $A$  is complete to  $B$ . The graph  $W_n$  is the graph on  $n + 1$  vertices that is obtained from a cycle on  $n$  vertices by adding one vertex that is made adjacent to all vertices in the cycle. The *length* of a path is the number of edges in the path.

For two graphs  $G$  and  $H$  on disjoint vertex sets, we let  $G + H$  be the disjoint union of  $G$  and  $H$ , which has  $V(G + H) = V(G) \cup V(H)$  and  $E(G + H) = E(G) \cup E(H)$ . We let  $pG$  denote the disjoint union of  $p$  copies of  $G$ . The *subdivision* of an edge  $uv$  in a graph removes the edge  $uv$  and introduces a new vertex  $w$  that is made adjacent (only) to  $u$  and  $v$ .

A graph  $H$  is the *1-subdivision* of a graph  $G$  if  $H$  is obtained from  $G$  by subdividing each edge of  $G$  exactly once. A graph  $G$  is *distance-hereditary* if for every connected induced subgraph  $H$  of  $G$  and every two vertices  $v, w$  in  $H$ , the distance between  $v$  and  $w$  in  $H$  is the same as the distance in  $G$ .

Let  $G$  be a graph with vertices  $x_1, \dots, x_n$ . Let  $A = A_G$  denote the *adjacency matrix* of  $G$ , that is, entry  $A_G(i, j) = 1$  if  $x_i$  is adjacent to  $x_j$  and  $A_G(i, j) = 0$  if  $x_i$  is not adjacent to  $x_j$ . For a subset  $X \subseteq V(G)$ , the matrix  $A[X, V(G) \setminus X]$  is the  $|X| \times |V(G) \setminus X|$  submatrix of  $A$  restricted to the rows of  $X$  and the columns of  $V(G) \setminus X$ . The *cut-rank function* of  $G$  is the function  $\text{cutrk}_G : 2^{V(G)} \rightarrow \mathbb{N}$  such that for each  $X \subseteq V(G)$ ,

$$\text{cutrk}_G(X) := \text{rank}(A_G[X, V(G) \setminus X]),$$

where we compute the rank *over the binary field*. A *linear ordering* of  $G$  is a permutation of the vertices of  $G$ . The *width* of a linear ordering  $(x_1, \dots, x_n)$  of  $G$  is defined as  $\max_{1 \leq i \leq n} \{\text{cutrk}_G(\{x_1, \dots, x_i\})\}$ . The *linear rank-width*  $\text{lrw}(G)$  of  $G$  is the minimum width over all linear orderings of  $G$ .

Let  $X$  and  $Y$  be two disjoint subsets of vertices of a graph. For an ordering  $(x_1, \dots, x_n)$  of the vertices of  $X$  and an ordering  $(y_1, \dots, y_m)$  of the vertices of  $Y$ , we define the ordering

$$(x_1, \dots, x_n) \oplus (y_1, \dots, y_m) := (x_1, \dots, x_n, y_1, \dots, y_m).$$

The cut-rank function is invariant under taking local complementation. This implies that the linear rank-width of a graph does not increase when taking its vertex-minor.

**Lemma 2.1** (Bouchet [10]; See Oum [40]). *If  $G$  is obtained from  $H$  by a sequence of local complementations, then  $\text{cutrk}_G(X) = \text{cutrk}_H(X)$  for all  $X \subseteq V(G)$ . So, if  $G$  is a vertex-minor of  $H$ , then  $\text{lrw}(G) \leq \text{lrw}(H)$ .*

We need three structural lemmas on linear rank-width. Recall that the complement of a graph  $G$  is the graph  $\overline{G}$  with  $V(\overline{G}) = V(G)$  and  $E(\overline{G}) = \{uv \mid u, v \in V(G) \text{ with } u \neq v \text{ and } uv \notin E(G)\}$ .

**Lemma 2.2.** *If  $G$  has linear rank-width  $k$ , then  $\overline{G}$  has linear rank-width at most  $k + 1$ .*

*Proof.* Let  $H$  be the graph obtained from  $G$  by adding a vertex  $a$  complete to  $V(G)$ . Observe that  $\overline{G} = (H * a)[V(G)]$ . Then,  $\text{lrw}(\overline{G}) \leq \text{lrw}(H * a) = \text{lrw}(H) \leq \text{lrw}(G) + 1$ . Here, the second step follows from Lemma 2.1, and the third step follows from the fact that adding one vertex to a graph may increase the linear rank-width by at most one.  $\square$

**Lemma 2.3.** *Let  $G$  be a graph and  $A$  and  $B$  be two disjoint vertex subsets of  $G$ . If  $G$  has linear rank-width  $k$ , then  $G \times (A, B)$  has linear rank-width at most  $k + 2$ .*

*Proof.* Let  $H$  be the graph obtained from  $G$  by adding two adjacent vertices  $a$  and  $b$  such that

- $N_H(a) \cap V(G) = A$  and  $N_H(b) \cap V(G) = B$ .

Observe that  $G \times (A, B) = (H \wedge ab)[V(G)]$ . Since adding two vertices may increase the linear rank-width by at most two, we have  $\text{lrw}(H \wedge ab) = \text{lrw}(H) \leq \text{lrw}(G) + 2$ . Therefore, we have  $\text{lrw}(G \times (A, B)) \leq \text{lrw}(G) + 2$ .  $\square$

Let  $I$  be a set of pairwise twins in a graph  $G$ . We define  $G//I$  as the graph obtained from  $G$  by removing all the vertices of  $I$  except one vertex.

**Lemma 2.4.** *Let  $G$  be a graph and  $I_1, I_2, \dots, I_m$  be pairwise disjoint subsets of  $V(G)$  such that each  $I_i$  is a set of pairwise twins in  $G$ . Then  $\text{lrw}(G) \leq \text{lrw}(G//I_1//I_2//\dots//I_m) + 1$ .*

*Proof.* Let  $H := G//I_1//I_2//\cdots//I_m$ . For each  $j \in \{1, \dots, m\}$ , let  $w_j$  be the vertex kept from  $I_j$  in  $H$ . Let  $I := \bigcup_{j \in \{1, \dots, m\}} I_j$ . Suppose that  $L_H$  is a linear ordering of  $H$  with the optimal width. We obtain a linear ordering  $L_G$  of  $G$  from  $L_H$  by replacing each  $w_j \in \{w_1, \dots, w_m\}$  with any linear ordering of  $I_j$ . Let  $L_G := (v_1, v_2, \dots, v_n)$ . We claim that  $L_G$  has width at most  $\text{lrw}(H) + 1$ . Let  $i \in \{1, \dots, n\}$ , and let  $L := \{v_j \mid 1 \leq j \leq i\}$  and  $R := V(G) \setminus L$ . It suffices to show that  $\text{rank}(A(G)[L, R]) \leq \text{lrw}(H) + 1$ . We will use the fact that

(\*) if two rows of a matrix  $M$  are the same, then the matrix obtained from  $M$  by removing one of these rows has the same rank as  $M$ , and the same argument holds for columns.

We observe that at most one set of  $I_1, I_2, \dots, I_m$  may have a vertex in both  $L$  and  $R$ . Let  $J_1 := \{i \in \{1, 2, \dots, m\} \mid I_i \cap L \neq \emptyset\}$  and  $J_2 := \{i \in \{1, 2, \dots, m\} \mid I_i \cap R \neq \emptyset\}$ , and let  $W_1 := \{w_i \mid i \in J_1\}$  and  $W_2 := \{w_i \mid i \in J_2\}$ . First assume that no set of  $I_1, I_2, \dots, I_m$  has a vertex in both  $L$  and  $R$ . By (\*), we have  $\text{rank}(A(G)[L, R]) = \text{rank}(A(G)[(L \setminus I) \cup W_1, (R \setminus I) \cup W_2])$ . As the partition  $((L \setminus I) \cup W_1, (R \setminus I) \cup W_2)$  is considered when computing the width of  $L_H$ , we have  $\text{rank}(A(G)[L, R]) \leq \text{lrw}(H)$ . Therefore, we may assume that there is a set  $I_p$  of  $I_1, \dots, I_m$  having a vertex in both  $L$  and  $R$ . Let  $x \in I_p \cap L$  and  $y \in I_p \cap R$ . By (\*), we have

$$\begin{aligned} & \text{rank}(A(G)[L, R]) \\ &= \text{rank}(A(G)[(L \setminus I) \cup (W_1 \setminus \{w_p\}) \cup \{x\}, (R \setminus I) \cup (W_2 \setminus \{w_p\}) \cup \{y\}]). \end{aligned}$$

Note that one of the partitions  $((L \setminus I) \cup W_1, (R \setminus I) \cup (W_2 \setminus \{w_p\}))$  and  $((L \setminus I) \cup (W_1 \setminus \{w_p\}), (R \setminus I) \cup W_2)$  is considered when computing the width of  $L_H$ . Since  $x, y, w_p \in I_p$ , the matrix

$$A(G)[(L \setminus I) \cup (W_1 \setminus \{w_p\}) \cup \{x\}, (R \setminus I) \cup (W_2 \setminus \{w_p\}) \cup \{y\}]$$

can be obtained from  $A(G)[(L \setminus I) \cup W_1, (R \setminus I) \cup (W_2 \setminus \{w_p\})]$  by adding one column corresponding to  $y$ , and it also can be obtained from  $A(G)[(L \setminus I) \cup (W_1 \setminus \{w_p\}), (R \setminus I) \cup W_2]$  by adding one row corresponding to  $x$ . This implies that  $\text{rank}(A(G)[L, R]) \leq \text{lrw}(H) + 1$ . We conclude that  $\text{lrw}(G) \leq \text{lrw}(H) + 1$ .  $\square$

A *path decomposition* of a graph  $G$  is an ordered family  $\mathcal{B} = \{B_1, \dots, B_r\}$  of subsets of  $V(G)$  satisfying the following:

1. For every  $v \in V(G)$  there exists a  $t \in \{1, \dots, r\}$  with  $v \in B_t$ .
2. For every  $uv \in E(G)$  there exists a  $t \in \{1, \dots, r\}$  with  $\{u, v\} \subseteq B_t$ .
3. For every  $v \in V(G)$ , the set  $\{t \in \{1, \dots, r\} \mid v \in B_t\}$  consists of consecutive integers.

The *width* of a path decomposition  $(P, \mathcal{B})$  is defined as  $\max\{|B_t| \mid t \in \{1, \dots, r\}\} - 1$ . The *path-width* of  $G$  is the minimum width among all path decompositions of  $G$ .

We finish this section by proving that every tree with linear rank-width 1 is a caterpillar (recall that a caterpillar is a tree that contains a path  $P$ , such that every vertex not on  $P$  has a neighbour in  $P$ ).

**Theorem 2.5** (Adler and Kanté [2]). *For every tree  $T$ , the linear rank-width of a tree  $T$  is equal to the path-width of  $T$ .*

**Theorem 2.6** (Takahashi, Ueno, and Kajitani [48]). *Let  $G$  be a tree and  $k$  be a positive integer. Then  $G$  has path-width at most  $k$  if and only if for every vertex  $v$ ,  $G - v$  has at most two connected components with path-width exactly  $k$  and all other connected components of  $G - v$  have path-width less than  $k$ .*




Figure 4: A tree  $F$ , the tree  $F'$  that is obtained from  $F$  by subdividing each edge of  $F$  once, and the graph  $H \in \mathcal{C}$  that is obtained from  $F'$  after applying a local complementation at every vertex of degree at least 3 in  $F'$ .

**Lemma 2.7.** *A tree has linear rank-width at most 1 if and only if it is a caterpillar.*

*Proof.* Let  $T$  be a tree. First suppose that  $T$  is not a caterpillar. Then  $T$  contains  $S_{2,2,2}$  as an induced subgraph. Thus,  $T$  contains a vertex  $v$  such that  $T - v$  contains at least three connected components each containing an edge. So,  $T - v$  contains three connected components having path-width at least 1. By Theorem 2.6,  $T$  has path-width at least 2, and by Theorem 2.5,  $T$  has linear rank-width at least 2.

Now suppose that  $T$  is a caterpillar. We prove by induction on  $|V(T)|$  that  $T$  has path-width at most 1. We may assume that  $T$  has at least two vertices. Note that for every vertex  $v$ ,  $T - v$  has at most two connected components having an edge, which are still caterpillars, and all the other connected components are isolated vertices. So,  $T - v$  has at most two connected components having path-width 1 by induction, and all the other connected components have path-width 0. Thus, by Theorem 2.6,  $T$  has path-width at most 1. This proves the claim. We now apply Theorem 2.5 to conclude that every caterpillar has linear rank-width at most 1.  $\square$

### 3 The Proof of Theorem 1.4

In this section, we prove Theorem 1.4, which states that for every tree  $T$  that is not a caterpillar, the class of  $T$ -pivot-minor-free graphs has unbounded linear rank-width.

Let  $\mathcal{C}$  be the class of graphs that can be obtained from the 1-subdivision of a tree by applying a local complementation at every vertex of degree at least 3. We give an example of a graph in  $\mathcal{C}$  in Figure 4. Our proof of Theorem 1.4 consists of the following parts:

1. we show that  $\mathcal{C}$  has unbounded linear rank-width;
2. we show that every graph in  $\mathcal{C}$  is a distance-hereditary graph with some additional properties needed to prove the third step; and
3. we show that every graph in  $\mathcal{C}$  is  $T$ -pivot-minor-free whenever  $T$  is a tree that is not a caterpillar.

We start with the following lemma that proves the first part.




Figure 5: For the graphs  $F'$  and  $H$  in Figure 4, the decompositions  $D_{F'}$  and  $D_H$  are the canonical split decompositions of  $F'$  and  $H$ , respectively. Dashed edges denote marked edges and each circle denotes a bag. Note that by the definition of a split, every bag contains at least three vertices.

**Lemma 3.1.** *The class  $\mathcal{C}$  has unbounded linear rank-width.*

*Proof.* Adler and Kanté [2] proved that trees have unbounded linear rank-width. Note that the 1-subdivision  $H$  of a graph  $G$  contains  $G$  as a vertex-minor: for every subdivided vertex in  $H$ , perform a local complementation and remove it; this yields  $G$ . So, by Lemma 2.1, the class of 1-subdivisions of trees also has unbounded linear rank-width. As local complementations do not change the linear rank-width of a graph by Lemma 2.1, this means that  $\mathcal{C}$  has unbounded linear rank-width.  $\square$

We will now prove that  $\mathcal{C}$  is a subclass of the class of distance-hereditary graphs with some additional useful properties. In order to do this, we need the notion of a canonical split decomposition of a graph [18], which we define below.

A vertex partition  $(X, Y)$  of a connected graph  $G$  is a *split* of  $G$  if  $|X| \geq 2$ ,  $|Y| \geq 2$ , and  $N_G(Y)$  is complete to  $N_G(X)$ . A connected graph  $G$  on at least five vertices is *prime* if it has no split. A connected graph  $D$  with a distinguished set of edges  $M(D)$  is a *marked graph* if  $M(D)$  is a matching and each edge in  $M(D)$  is a cut edge. An edge in  $M(D)$  is a *marked edge*, and every other edge of  $D$  is an *unmarked edge*. A vertex incident with a marked edge is a *marked vertex*, and every other vertex of  $D$  is an *unmarked vertex*. Each connected component of  $D - M(D)$  is a *bag* of  $D$ . If a marked edge  $e$  is incident with a vertex of a bag  $B$ , we say that  $B$  is *incident* with  $e$ . A bag  $B_1$  of  $D$  is a neighbour bag of a bag  $B_2$  of  $D$  if there is a marked edge incident with both  $B_1$  and  $B_2$ . The *decomposition tree* of  $D$  is the graph obtained from  $D$  by contracting each bag into a vertex.

If  $G$  has a split  $(X, Y)$ , we construct a marked graph  $D$  on the vertex set  $V(G) \cup \{x_1, y_1\}$  for some new vertices  $x_1$  and  $y_1$  such that

- for every two distinct vertices  $x, y$  with  $\{x, y\} \subseteq X$  or  $\{x, y\} \subseteq Y$ , the property  $xy \in E(G)$

holds if and only if  $xy \in E(D)$ ,

- $x_1y_1$  is a new marked edge,
- $X$  is anti-complete to  $Y$ ,
- $x_1$  is complete to  $N_G(Y)$  (with only unmarked edges) and has no neighbours in  $V(G) \setminus N_G(Y)$ ,
- $y_1$  is complete to  $N_G(X)$  (with only unmarked edges) and has no neighbours in  $V(G) \setminus N_G(X)$ .

The graph  $D$  is a *simple decomposition* of  $G$ . To obtain a split decomposition, we will recursively take a simple decomposition of a bag, and when we take a simple decomposition of a bag, all the marked vertices remain marked vertices. That is, a *split decomposition* of a connected graph  $G$  is a marked graph  $D$  defined inductively to be either  $G$  or a marked graph obtained from a split decomposition  $D'$  of  $G$  by replacing some bag  $B$  of  $D'$  by the bags of a simple decomposition  $B'$  of  $B$  and keeping all the marked edges between vertices of  $V(B)$  and vertices of  $V(D') \setminus V(B)$ . We give an example of a split decomposition in Figure 5.

For a marked edge  $xy$  of a marked graph  $D$ , the *recomposition of  $D$  along  $xy$*  is the marked graph  $(D \wedge xy) - \{x, y\}$ , where when we pivot  $xy$ , we add unmarked edges between  $N_D(x) \setminus \{y\}$  and  $N_D(y) \setminus \{x\}$ . This operation can be seen as merging two adjacent bags  $B_1$  and  $B_2$  into one bag  $B$  where the union of  $B_1$  and  $B_2$  was a simple decomposition of  $B$ . It is not hard to see that if  $D$  is a split decomposition of  $G$ , then  $G$  can be obtained from  $D$  by recomposing along all the marked edges.

A split decomposition  $D$  is *canonical* if each bag of  $D$  is either a prime graph, a star, or a complete graph, and for every marked edge  $xy$  in  $D$ , recomposing  $xy$  results in a split decomposition having a bag that is neither a prime graph, a star, nor a complete graph. We note that the split decompositions in Figure 5 are canonical. We say that a bag is a *star bag* if it is a star and a *complete bag* if it is a complete graph.

**Theorem 3.2** (Cunningham and Edmonds [18]). *Every connected graph has a unique canonical split decomposition, up to isomorphism.*

Bouchet [9] described how split decompositions change under applying local complementations. A vertex  $v$  in a split decomposition  $D$  *represents* an unmarked vertex  $x$  (or is a *representative* of  $x$ ) if either  $v = x$  or there is a path of even length from  $v$  to  $x$  in  $D$  starting with a marked edge such that marked edges and unmarked edges appear alternately in the path. Observe that for every vertex  $v$  in a split decomposition  $D$  of a graph  $G$ , there exists a vertex  $x \in V(G)$  such that  $v$  represents  $x$ . Two unmarked vertices  $x$  and  $y$  are *linked* in  $D$  if there is a path from  $x$  to  $y$  in  $D$  such that unmarked edges and marked edges appear alternately in the path.

A *local complementation* at an unmarked vertex  $x$  in a split decomposition  $D$ , denoted by  $D * x$ , is the operation that replaces each bag  $B$  containing a representative  $w$  of  $x$  with  $B * w$ .

**Lemma 3.3** (Bouchet [9]). *Let  $D$  be the canonical split decomposition of a connected graph  $G$ . If  $x$  is an unmarked vertex of  $D$ , then  $D * x$  is the canonical split decomposition of  $G * x$ .*

Let  $x$  and  $y$  be linked unmarked vertices in a split decomposition  $D$ , and let  $P$  be the path in  $D$  linking  $x$  and  $y$  where unmarked edges and marked edges appear alternately. Observe that

such a path is unique. The *pivoting on  $xy$  of  $D$* , denoted by  $D \wedge xy$ , is the split decomposition obtained as follows: for each bag  $B$  containing an unmarked edge  $vw$  of  $P$ , we replace  $B$  with  $B \wedge vw$ .

**Lemma 3.4** (Adler, Kanté, and Kwon [3]). *Let  $D$  be the canonical split decomposition of a connected graph  $G$ . If  $xy \in E(G)$ , then  $D \wedge xy$  is the canonical split decomposition of  $G \wedge xy$ .*

An *internal* edge of a tree is an edge that is not incident with a vertex of degree 1. We also need the following result due to Bouchet [9].

**Lemma 3.5** (Bouchet [9]). *A canonical split decomposition of a tree  $T$  can be constructed by replacing each internal edge of  $T$  by a path of length 3, the middle edge of the path being the marked edge.*

A bag of a split decomposition is a *branching bag* if it is incident with at least three marked edges. Let  $\mathcal{M}$  be the set of all distance-hereditary graphs in which each connected component admits a canonical split decomposition with the property that every branching bag is a complete bag. We will prove that  $\mathcal{C}$  is a subclass of  $\mathcal{M}$ , so every graph of  $\mathcal{C}$  is a distance-hereditary graph with the additional property that every branching bag in its canonical split decomposition is a complete bag. In order to do this we need one more lemma.

**Lemma 3.6** (Bouchet [9]). *A connected graph is distance-hereditary if and only if every bag of its canonical split decomposition is either a star or a complete graph.*

**Lemma 3.7.**  *$\mathcal{C}$  is a subclass of  $\mathcal{M}$ .*

*Proof.* Let  $G \in \mathcal{C}$ . By the definition of  $\mathcal{C}$ , we find that  $G$  is obtained from the 1-subdivision  $T$  of some tree by performing local complementations at vertices of degree at least 3. By Lemma 3.5, we can first construct the canonical split decomposition of  $T$ . By Lemma 3.3 we can then perform local complementations at corresponding unmarked vertices in the canonical split decomposition of  $T$  to obtain the canonical split decomposition of  $G$  (we refer again to Figure 5 for an example). By construction, every bag of the canonical split decomposition of  $G$  is a star or a complete bag. Hence, by Lemma 3.6, we find that  $G$  is a distance-hereditary graph. From our construction we also note that every branching bag is complete. We conclude that  $G \in \mathcal{M}$ .  $\square$

We will now prove that for every tree  $T$  that is not a caterpillar,  $\mathcal{M}$  is  $T$ -pivot-minor-free. In order to do this we first show that  $\mathcal{M}$  is closed under taking pivot-minors.

**Lemma 3.8.** *The class  $\mathcal{M}$  is closed under taking pivot-minors.*

*Proof.* Let  $G \in \mathcal{M}$ . It is sufficient to show the following:

- (1) for  $v \in V(G)$ ,  $G - v$  is in  $\mathcal{M}$ ,
- (2) for an edge  $wz \in E(G)$ ,  $G \wedge wz$  is in  $\mathcal{M}$ .

We may assume that  $G$  is connected. Let  $D$  be the canonical split decomposition of  $G$ .

We first show (2). Let  $wz$  be an edge of  $G$ . By Lemma 3.4,  $D \wedge wz$  is the canonical split decomposition of  $G \wedge wz$ . Let  $P$  be the path in  $D$  linking  $w$  and  $z$  where unmarked edges and marked edges appear alternately. By the definition of pivoting in a canonical split decomposition, we obtain  $D \wedge wz$  from  $D$  as follows: for each bag  $B$  containing an unmarked

edge  $xy$  of  $P$ , we replace  $B$  with  $B \wedge xy$ . It is easy to observe that if  $B$  is a complete bag, then  $B \wedge xy$  is again a complete bag, and if  $B$  is a star bag, then  $B \wedge xy$  is again a star bag. Therefore,  $G \wedge wz$  is again contained in  $\mathcal{M}$ .

It remains to prove (1). Suppose that  $v \in V(G)$  and let  $B$  be the bag containing  $v$ . We may assume that  $D$  has at least two bags, otherwise the statement follows immediately.

**Case 1.**  $B$  is a complete bag.

If  $|V(B)| \geq 4$ , then after removing  $v$  in  $G$ , we find that  $B$  is still a complete bag of size at least 3. So,  $D - v$  is a canonical split decomposition of  $G - v$ . If  $|V(B)| = 3$ , then  $B - v$  is merged with one of the neighbour bags of  $B$ . This process does not change the type of the neighbour bag. It is possible that the two neighbour bags  $B_1$  and  $B_2$  of  $B$  in  $D$  are star bags, and they can be merged after  $B - v$  is merged with a neighbour bag. In this case, each of  $B_1$  and  $B_2$  has at most two neighbour bags, and after merging  $B_1$  and  $B_2$ , it is again a star bag that has at most two neighbour bags. Thus,  $G - v$  is in  $\mathcal{M}$  again.

**Case 2.**  $B$  is a star bag and  $v$  is a leaf of  $B$ .

If  $|V(B)| \geq 4$ , then  $D - v$  is the canonical split decomposition of  $G - v$ . If  $|V(B)| = 3$ , then  $B - v$  is merged with one of the neighbour bags of  $B$ . This process does not change the type of the neighbour bag. There might be two cases where the two neighbour bags  $B_1$  and  $B_2$  of  $B$  in  $D$  are merged after  $B - v$  is merged with a neighbour bag. If  $B_1$  and  $B_2$  are complete bags, then the merged bag becomes a complete bag. If  $B_1$  and  $B_2$  are star bags, then each of them has at most two neighbour bags in  $D$ , and after merging, the new bag is again a star bag that has at most two neighbour bags. So,  $G - v$  is in  $\mathcal{M}$ .

**Case 3.**  $B$  is star bag and  $v$  is the center of  $B$ .

Then  $v$  is a cut vertex of  $G$ ; that is,  $G - v$  is disconnected. Furthermore, each component of  $G - v$  either consists of a single vertex, or it admits a split decomposition obtained from a connected component of  $D - V(B)$  by removing a leaf of a star bag or a vertex in a complete bag. Thus, each component of  $G - v$  is in  $\mathcal{M}$ , and thus,  $G - v$  is also in  $\mathcal{M}$ . This concludes the proof of the lemma.  $\square$

We also need a known characterization of graphs of linear rank-width at most 1 in terms of their canonical split decompositions.

**Lemma 3.9** (Kanté and Kwon [30]). *Let  $G$  be a connected graph with canonical split decomposition  $D$ . Then  $G$  has linear rank-width at most 1 if and only if  $G$  is distance-hereditary and the decomposition tree of  $D$  is a path.*

We are now ready to prove Theorem 1.4.

**Theorem 1.4 (restated).** *If  $T$  is a tree that is not a caterpillar, then the class of  $T$ -pivot-minor-free distance-hereditary graphs has unbounded linear rank-width.*

*Proof.* Let  $T$  be a tree that is not a caterpillar. As  $\mathcal{C} \subseteq \mathcal{M}$  by Lemma 3.7 and  $\mathcal{C}$  has unbounded linear rank-width by Lemma 3.1, it follows that  $\mathcal{M}$  has unbounded linear rank-width. Moreover,  $\mathcal{M}$  is a subclass of the class of distance-hereditary graphs. Hence, to prove the theorem it remains to show that every graph in  $\mathcal{M}$  is  $T$ -pivot-minor-free.

Suppose, for contradiction, that  $T$  is a pivot-minor of some graph  $H \in \mathcal{M}$ . As  $\mathcal{M}$  is closed under pivot-minors by Lemma 3.8, we find that  $T \in \mathcal{M}$ . Let  $L_T$  be a canonical split decomposition of  $T$ . Since  $T$  is a tree,  $L_T$  has no complete bags. So, by the definition of

$\mathcal{M}$ ,  $L_T$  has no branching bags, and thus, the decomposition tree of  $L_T$  is a path. Since  $T$  is distance-hereditary and the decomposition tree of  $L_T$  is a path,  $T$  has linear rank-width at most 1 by Lemma 3.9. Thus, by Lemma 2.7,  $T$  is a caterpillar, a contradiction.  $\square$

## 4 The Proof of Theorem 1.5

In this section, we prove Theorem 1.5, which states that  $P_n$ -pivot-minor-free distance-hereditary graphs have bounded linear rank-width.

To prove Theorem 1.5, we use the canonical split decomposition of a distance-hereditary graph, discussed in Section 3. A sequence  $B_1, B_2, \dots, B_n$  of distinct bags in a canonical split decomposition is a *path of bags* if for each  $i \in \{1, \dots, n-1\}$ ,  $B_{i+1}$  is a neighbour bag of  $B_i$ . As we will explain in the proof of Theorem 1.5, it follows from the definition of a canonical split decomposition, that at least half of the bags in a path of bags are star bags. By applying some pivot operations, we can extract a long path as a pivot-minor in this case. So, we may assume that the decomposition tree of the canonical split decomposition of a given graph has no long path. We use the following result which relates the path-width of a decomposition tree and the linear rank-width of the graph to conclude the theorem.

**Proposition 4.1** (Kanté and Kwon [30]). *Let  $D$  be the canonical split decomposition of a connected distance-hereditary graph  $G$ , and let  $T_D$  be the decomposition tree of  $D$ . Then  $\frac{1}{2} \text{pw}(T_D) \leq \text{lrw}(G) \leq \text{pw}(T_D) + 1$ .*

We also use a tight version of Theorem 1.1. We will use the fact that if a graph contains a minor isomorphic to  $P_n$ , then it also contains a subgraph isomorphic to  $P_n$ .

**Theorem 4.2** (Bienstock, Robertson, Seymour, and Thomas [6]). *For every tree  $T$  on  $n$  vertices, the class of  $T$ -minor-free graphs has path-width at most  $n - 2$ .*

We are now ready to prove Theorem 1.5.

**Theorem 1.5 (restated).** *Let  $n \geq 3$  be an integer. Every  $P_n$ -pivot-minor-free distance-hereditary graph has linear rank-width at most  $2n - 5$ .*

*Proof.* Let  $G$  be a distance-hereditary graph having no pivot-minor isomorphic to  $P_n$ . We will show that  $G$  has linear rank-width at most  $2n - 5$ . We may assume that  $G$  is connected. Let  $D$  be the canonical split decomposition of  $G$  and let  $T_D$  be its decomposition tree.

We claim that  $D$  has no path with  $2n - 4$  bags. Suppose that such a path of bags  $B_1, B_2, \dots, B_{2n-4}$  exists. As no two complete bags are neighbour bags in a canonical split decomposition, at most  $n - 2$  bags in the sequence are complete bags. Thus, there are at least  $n - 2$  star bags. Let  $B_{i_1}, B_{i_2}, \dots, B_{i_t}$  be the sequence of all star bags in  $B_1, B_2, \dots, B_{2n-4}$  where  $1 \leq i_1 < i_2 < \dots < i_t \leq 2n - 4$ . For convenience, we assign  $B_0 = B_{2n-3} = \emptyset$ .

We claim that for every  $k \in \{1, \dots, t\}$ , there is a graph  $G_k$  pivot-equivalent to  $G$  with a canonical split decomposition  $D_k$  such that

- (\*) for every  $k' \in \{1, \dots, k\}$ , the bag  $D_k[V(B_{i_{k'}})]$  is a star bag whose center has no neighbour in  $V(B_{i_{k'}-1}) \cup V(B_{i_{k'}+1})$ .

For  $k = 1$ , suppose that  $D$  does not satisfy the property (\*). Choose a vertex  $v'$  in  $B_{i_1}$  that has no neighbour in  $V(B_{i_1-1}) \cup V(B_{i_1+1})$ . Such a vertex exists, as each bag has at least three vertices. Let  $v$  be the vertex of  $G$  represented by  $v'$  (recall that for every marked vertex, there

is a vertex of  $G$  that it represents). We further choose a vertex  $w$  of  $G$  represented by the center  $w'$  of  $B_{i_1}$ . Since  $v'w'$  is an edge of  $B_{i_1}$ ,  $v$  is linked to  $w$  in  $D$ , and therefore,  $v$  is adjacent to  $w$  in  $G$ .

Note that for a star  $H$  with a leaf  $b$  and center  $a$ ,  $H \wedge ab$  is the star with vertex set  $V(H)$  whose center is  $b$ . Thus, in  $D \wedge vw$ ,  $v'$  becomes the center of  $B_{i_1} \wedge v'w'$ , which has no neighbour in  $V(B_{i_1-1}) \cup V(B_{i_1+1})$ . Thus,  $G_1 = G \wedge vw$  and  $D_1 = D \wedge vw$  satisfy  $(*)$ .

Now, assume that  $k > 1$  and the property  $(*)$  is satisfied for  $k - 1$ . If the center of  $D_{k-1}[V(B_{i_k})]$  has no neighbour in  $V(B_{i_k-1}) \cup V(B_{i_k+1})$ , then  $G_k = G_{k-1}$  and  $D_k = D_{k-1}$  satisfy  $(*)$ . So, we may assume that the center of  $D_{k-1}[V(B_{i_k})]$  has a neighbour in  $V(B_{i_k-1}) \cup V(B_{i_k+1})$ . We distinguish two cases.

**Case 1.** The center of  $D_{k-1}[V(B_{i_k})]$  has a neighbour in  $V(B_{i_k-1})$ .

Note that if  $D_{k-1}[V(B_{i_k-1})]$  is a star bag, then by the inductive hypothesis, its center has no neighbour in  $V(B_{i_k})$ . But this is not possible by the definition of a canonical split decomposition. Thus,  $D_{k-1}[V(B_{i_k-1})]$  is a complete bag. We choose a vertex  $v$  in  $G_{k-1}$  represented by a vertex in  $D_{k-1}[V(B_{i_k-1})]$  having no neighbour in  $V(B_{i_k-2}) \cup V(B_{i_k})$  and choose a vertex  $w$  in  $G_{k-1}$  represented by a vertex in  $D_{k-1}[V(B_{i_k})]$  having no neighbour in  $V(B_{i_k-1}) \cup V(B_{i_k+1})$ . Observe that  $v$  is adjacent to  $w$  in  $G_{k-1}$ , because the center of  $D_{k-1}[V(B_{i_k})]$  has a neighbour in  $V(B_{i_k-1})$ . Thus, in  $D_{k-1} \wedge vw$ , the bag induced by  $V(B_{i_k})$  is a star bag whose center has no neighbour in  $V(B_{i_k-1}) \cup V(B_{i_k+1})$ . As the bags on  $V(B_{i_1}), \dots, V(B_{i_{k-1}})$  are not changed by this pivot operation,  $G_k = G_{k-1} \wedge vw$  and  $D_k = D_{k-1} \wedge vw$  satisfy  $(*)$ .

**Case 2.** The center of  $D_{k-1}[V(B_{i_k})]$  has a neighbour in  $V(B_{i_k+1})$ .

We choose a vertex  $v$  in  $G_{k-1}$  represented by a vertex in  $D_{k-1}[V(B_{i_k+1})]$  having no neighbour in  $V(B_{i_k}) \cup V(B_{i_k+2})$  and choose a vertex  $w$  in  $G_{k-1}$  represented by a vertex in  $D_{k-1}[V(B_{i_k})]$  having no neighbour in  $V(B_{i_k-1}) \cup V(B_{i_k+1})$  and linked to  $v$  in  $D_{k-1}$ . Such a vertex  $w$  exists, because the center of  $D_{k-1}[V(B_{i_k})]$  has a neighbour in  $V(B_{i_k+1})$ , and thus, if  $D_{k-1}[V(B_{i_k+1})]$  is a star, then its center has a neighbour in  $V(B_{i_k})$ . This implies that  $v$  is adjacent to  $w$  in  $G_{k-1}$ . In  $D_{k-1} \wedge vw$ , the bag induced by  $V(B_{i_k})$  is a star bag whose center has no neighbour in  $V(B_{i_k-1}) \cup V(B_{i_k+1})$ . As the bags on  $V(B_{i_1}), \dots, V(B_{i_{k-1}})$  are not changed by this pivot operation,  $G_k = G_{k-1} \wedge vw$  and  $D_k = D_{k-1} \wedge vw$  satisfy  $(*)$ .

Hence, we have found that the claim holds.

Now, in  $D_t$ , let  $v_j$  be a vertex of  $G_t$  represented by the center of  $D_t[V(B_{i_j})]$  for each  $j \in \{1, \dots, t\}$ , and let  $v_0$  be a vertex of  $G_t$  represented by a leaf of  $D_t[V(B_{i_1})]$  which has no neighbour in  $V(B_{i_1+1})$ , and let  $v_{t+1}$  be a vertex of  $G_t$  represented by a leaf of  $D_t[V(B_{i_t})]$  which has no neighbour in  $V(B_{i_t-1})$ . It is not difficult to check that  $v_0v_1v_2 \cdots v_tv_{t+1}$  is an induced path of  $G_t$  on  $t + 2 \geq n$  vertices. This contradicts the assumption that  $G$  has no pivot-minor isomorphic to  $P_n$ . We conclude that  $D$  has no path with  $2n - 4$  bags  $B_1, B_2, \dots, B_{2n-4}$ .

The above means that the decomposition tree  $T_D$  has no path on  $2n - 4$  vertices. By Theorem 4.2,  $T_D$  has path-width at most  $2n - 6$ . By Proposition 4.1,  $G$  has linear rank-width at most  $2n - 5$ .  $\square$

## 5 The Proof of Theorem 1.10

In this section, we prove Theorem 1.10, which states that the class of  $(K_3, S_{1,2,2})$ -free graphs has linear rank-width at most 58. We prove the following statements in this order:

1. bipartite  $2P_2$ -free graphs, which form a subclass of bipartite  $(P_1 + 2P_2)$ -free graphs, have linear rank-width at most 1;
2. bipartite  $(P_1 + 2P_2)$ -free graphs, which form a subclass of bipartite  $S_{1,2,2}$ -free graphs, have linear rank-width at most 3;
3. bipartite  $S_{1,2,2}$ -free graphs have linear rank-width at most 3;
4. non-bipartite  $(K_3, C_5, S_{1,2,2})$ -free graphs have linear rank-width at most 3; and
5.  $(K_3, S_{1,2,2})$ -free graphs with an induced  $C_5$  have linear rank-width at most 58.

Note that Statements 3–5 cover all cases for proving Theorem 1.10. So, we first consider bipartite  $2P_2$ -free graphs.

**Lemma 5.1.** *Every bipartite  $2P_2$ -free graph has linear rank-width at most 1.*

*Proof.* Let  $G$  be a bipartite  $2P_2$ -free graph with bipartition  $(A, B)$ . It is well known [49] that a bipartite graph with bipartition  $(X_1, X_2)$  is  $2P_2$ -free if and only if it is a bipartite chain graph, that is, for each  $i \in \{1, 2\}$ , the neighbourhoods of the vertices in  $X_i$  can be ordered linearly with respect to the inclusion relation. We may assume that  $G$  is connected. Hence, as  $G$  is  $2P_2$ -free, we can define a sequence  $A_1, A_2, \dots, A_m$  of pairwise vertex-disjoint subsets of  $A$  such that

- $A_1 \cup A_2 \cup \dots \cup A_m = A$ ,
- each  $A_i$  is a maximal set of pairwise twins in  $G$ ,
- for integers  $i, j \in \{1, \dots, m\}$  with  $i < j$ ,  $N_G(A_i) \subsetneq N_G(A_j)$ .

If  $m = 1$ , then  $G$  is complete bipartite. In this case, we take a linear ordering  $L_1$  of  $A_1$  and a linear ordering  $L_2$  of  $V(G) \setminus A_1$  arbitrarily. It is not hard to see that  $L_1 \oplus L_2$  is a linear ordering of width at most 1. Hence  $G$  has linear rank-width at most 1.

Now suppose that  $m \geq 2$ . In this case,  $G$  is not complete bipartite. Notice that for each  $i \in \{2, \dots, m\}$ , there is a vertex  $v \in B$  that has a neighbour in  $A_i$  but does not have a neighbour in  $A_1 \cup \dots \cup A_{i-1}$ ; otherwise, vertices in  $A_{i-1} \cup A_i$  have the same neighbourhood in  $B$ , which contradicts the maximality of  $A_i$ . For each  $i \in \{2, \dots, m\}$ , let  $B_i := N_G(A_i) \setminus N_G(A_{i-1})$ , and let  $B_1 := N_G(A_1)$ . Since  $G$  is connected, we have  $B = B_1 \cup B_2 \cup \dots \cup B_m$ .

For each  $i \in \{1, \dots, m\}$ , let  $L_i^A$  be an ordering of  $A_i$  and  $L_i^B$  be an ordering of  $B_i$ . It is not difficult to check that the linear ordering  $L_1^B \oplus L_1^A \oplus L_2^B \oplus L_2^A \oplus \dots \oplus L_m^B \oplus L_m^A$  has width at most 1.  $\square$

We now consider bipartite  $(P_1 + 2P_2)$ -free graphs and show the following lemma.

**Lemma 5.2.** *Every bipartite  $(P_1 + 2P_2)$ -free graph has linear rank-width at most 3.*

*Proof.* Let  $G$  be a bipartite  $(P_1 + 2P_2)$ -free graph with bipartition  $(A, B)$ . Then  $G \times (A, B)$  is  $P_5$ -free. We may assume without loss of generality that  $G \times (A, B)$  is connected. Then, as  $G \times (A, B)$  is also bipartite,  $G \times (A, B)$  is readily seen to be  $2P_2$ -free. By Lemma 5.1,  $G \times (A, B)$  has linear rank-width at most 1. By Lemma 2.3,  $G = G \times (A, B) \times (A, B)$  has linear rank-width at most 3.  $\square$

We now consider  $S_{1,2,2}$ -free bipartite graphs and need two results by Lozin [35].

**Lemma 5.3** (Lozin [35]). *Every connected bipartite  $(S_{1,2,2}, P_7)$ -free graph is  $(P_1 + 2P_2)$ -free.*

**Lemma 5.4** (Lozin [35]). *Let  $G$  be a bipartite  $S_{1,2,2}$ -free graph with no twins. If  $G$  contains an induced  $P_7$ , then  $G$  is  $K_{1,3}$ -free (and thus  $G$  has maximum degree at most 2).*

**Proposition 5.5.** *Every bipartite  $S_{1,2,2}$ -free graph has linear rank-width at most 3.*

*Proof.* Let  $G$  be a bipartite  $S_{1,2,2}$ -free graph with bipartition  $(A, B)$ . We may assume that  $G$  is connected. If  $G$  is  $P_7$ -free, then by Lemma 5.3,  $G$  is  $(P_1 + 2P_2)$ -free, and by Lemma 5.2,  $G$  has linear rank-width at most 3. Thus, we may assume that  $G$  contains an induced subgraph isomorphic to  $P_7$ . Note that  $G$  is not a complete bipartite graph.

Let  $I_1, I_2, \dots, I_m$  be the vertex partition of  $G$  such that each  $I_i$  is a maximal set of pairwise twins in  $G$ . Since  $G$  is connected and bipartite, each  $I_i$  is contained in one of  $A$  or  $B$ . Let  $G_1 := G // I_1 // I_2 // \dots // I_m$ . Note that  $G_1$  is also connected. We claim that  $G_1$  has no twins. Note that  $G_1$  is not an edge, because  $G$  is not a complete bipartite graph. So,  $G_1$  has at least three vertices. Moreover,  $G_1$  still has an induced subgraph isomorphic to  $P_7$ , as  $P_7$  has no twins.

Suppose for contradiction that  $G_1$  has two twins  $v_1$  and  $v_2$ , and assume that  $v_1$  and  $v_2$  were identified from  $I_{i_1}$  and  $I_{i_2}$  for some  $i_1$  and  $i_2$ , respectively. Since each  $v_i$  has a neighbour and  $G_1$  has at least three vertices,  $v_1$  and  $v_2$  are in the same part of the bipartition, and thus  $I_{i_1}$  and  $I_{i_2}$  are contained in the same part of the bipartition of  $G$ . Thus  $I_{i_1}$  and  $I_{i_2}$  have the same neighbourhoods in  $G$ , contradicting the fact that they are maximal sets of pairwise twins in  $G$ . So,  $G_1$  has no twins. Then  $G_1$  has linear rank-width at most 2 because by Lemma 5.4 every vertex has degree at most 2. By Lemma 2.4, we conclude that  $G$  has linear rank-width at most 3.  $\square$

We now consider  $(K_3, C_5, S_{1,2,3})$ -free graphs and need the following result as a lemma.

**Lemma 5.6** (Dabrowski, Dross, and Paulusma [19]). *Let  $G$  be a connected  $(K_3, C_5, S_{1,2,3})$ -free graph that does not contain a pair of false twins. Then  $G$  is either bipartite or an induced cycle.*

**Proposition 5.7.** *Every non-bipartite  $(K_3, C_5, S_{1,2,2})$ -free graph has linear rank-width at most 3.*

*Proof.* Let  $G$  be a connected non-bipartite  $(K_3, C_5, S_{1,2,2})$ -free graph. By Lemma 5.6,  $G$  is a graph obtained from an induced cycle  $C = c_1 c_2 \dots c_k c_1$  by adding false twins. For each  $i \in \{1, \dots, k\}$ , let  $U_i$  be the maximal set of false twins containing  $c_i$  in  $G$ . As  $G // U_1 // U_2 // \dots // U_k$  is isomorphic to  $C$  and  $C$  has linear rank-width at most 2, by Lemma 2.4, we find that  $G$  has linear rank-width at most 3.  $\square$

We now consider  $(K_3, S_{1,2,2})$ -free graphs that contain an induced  $C_5$ . We first introduce some additional terminology and lemmas. A graph is *3-partite* if its vertex set can be partitioned into three independent sets. We need the following known result.

**Theorem 5.8** (Alecú et al. [5]). *Let  $G$  be a 3-partite graph on  $n$  vertices with vertex partition  $(V_1, V_2, V_3)$  such that*

- (a) *for every  $a \in V_1, b \in V_2, c \in V_3, G[a, b, c]$  is isomorphic to neither  $K_3$  nor  $3P_1$ ,*

(b)  $G[V_1 \cup V_2]$ ,  $G[V_1 \cup V_3]$ , and  $G[V_2 \cup V_3]$  are  $2P_2$ -free.

Then  $V(G)$  admits a linear ordering  $x_1, \dots, x_n$  and a labelling  $\ell : V(G) \rightarrow \{a, b, c\}$  such that  $x_i x_j \in E(G)$  for  $i < j$  if and only if  $(\ell(x_i), \ell(x_j)) \in \{(a, b), (b, c), (c, a)\}$ .

The following lemma follows from Theorem 5.8 after observing that each cut of the linear ordering  $x_1, \dots, x_n$  has cut-rank at most 3, because it has at most three different rows.

**Lemma 5.9.** *Let  $G$  be a 3-partite graph with vertex partition  $(V_1, V_2, V_3)$  such that*

- (a) *for every  $a \in V_1$ ,  $b \in V_2$ ,  $c \in V_3$ ,  $G[a, b, c]$  is isomorphic to neither  $K_3$  nor  $3P_1$ ,*
- (b)  *$G[V_1 \cup V_2]$ ,  $G[V_1 \cup V_3]$ , and  $G[V_2 \cup V_3]$  are  $2P_2$ -free.*

*Then  $G$  has linear rank-width at most 3.*

Let  $G$  be a graph and  $V_1, V_2, V_3$  be three pairwise disjoint independent sets of  $G$ . We denote the subgraph of  $G$  induced by  $V_1 \cup V_2 \cup V_3$  as  $G[V_1, V_2, V_3]$ . Moreover, if  $G[V_1, V_2, V_3]$  satisfies conditions (a) and (b) in Lemma 5.9, then we say that  $G[V_1, V_2, V_3]$  is *nice*.

We are now ready to prove the following result. We note that Brandstädt, Mahfud and Mosca [12] gave an alternative proof of the result from [19], which shows that  $(K_3, S_{1,2,2})$ -free graphs have bounded rank-width. Some parts of the proof of our result below are similar to parts of the proof of [12]. As we need to use slightly different arguments, we have chosen to keep our proof self-contained. However, we explicitly indicate whenever there is overlap between our arguments and the ones used in [12].

**Proposition 5.10.** *Every  $(K_3, S_{1,2,2})$ -free graph that contains an induced  $C_5$  has linear rank-width at most 58.*

*Proof.* Let  $G$  be a  $(K_3, S_{1,2,2})$ -free graph that contains an induced subgraph  $C$  isomorphic to  $C_5$ . We may assume without loss of generality that  $G$  is connected. We write  $C = c_1 c_2 c_3 c_4 c_5 c_1$  and interpret subscripts modulo 5. Let  $U := V(G) \setminus V(C)$ .

Since  $G$  is  $K_3$ -free, every vertex in  $U$  has either no neighbours in  $C$  or exactly one neighbour in  $C$  or exactly two neighbours, which are not consecutive in  $C$ .

We claim that every vertex of  $U$  has a neighbour in  $C$ . This can be seen as follows. Suppose, for contradiction, that  $U$  contains a vertex that has no neighbour in  $C$ . As  $G$  is connected, this means that  $U$  contains two vertices  $w_1$  and  $w_2$ , such that  $w_1$  has a neighbour in  $C$  and  $w_2$  is adjacent to  $w_1$  but  $w_2$  has no neighbour in  $C$ . If  $w_1$  has exactly one neighbour  $c_i$  in  $C$ , then  $G[c_i, c_{i+1}, c_{i-1}, c_{i-2}, w_1, w_2]$  is isomorphic to  $S_{1,2,2}$ . If  $w_1$  has two neighbours  $c_{i-1}, c_{i+1}$  in  $C$ , then  $G[w_1, w_2, c_{i-1}, c_i, c_{i-2}, c_{i-3}]$  is isomorphic to  $S_{1,2,2}$ . However, both cases are not possible, as  $G$  is  $S_{1,2,2}$ -free. Hence, we conclude that every vertex in  $U$  has a neighbour in  $C$ .

Consequently, we can partition  $U$  into ten parts  $\{V_1, \dots, V_5, W_1, \dots, W_5\}$  such that for each  $i \in \{1, \dots, 5\}$ ,

- $V_i$  is the set of vertices whose unique neighbour in  $C$  is  $c_i$ , and
- $W_i$  is the set of vertices that are adjacent to  $c_{i-1}$  and  $c_{i+1}$ .

Each set in  $\{V_1, \dots, V_5, W_1, \dots, W_5\}$  is an independent set as  $G$  is  $K_3$ -free. We verify basic relations between these parts. Let  $i \in \{1, \dots, 5\}$ .

- (1)  $V_i$  is complete to  $V_{i-1} \cup V_{i+1}$ , and anti-complete to  $V_{i-2} \cup V_{i+2}$ .

Suppose, for contradiction, that there are  $a \in V_i$  and  $b \in V_{i+1}$  that are not adjacent. Then  $G[c_i, a, c_{i+1}, b, c_{i-1}, c_{i-2}]$  is isomorphic to  $S_{1,2,2}$ , a contradiction. Thus, there are no such vertices. This implies that  $V_i$  is complete to  $V_{i+1}$ , and by symmetry also complete to  $V_{i-1}$ . Suppose that there are  $a \in V_i$  and  $b \in V_{i+2}$  that are adjacent. Then  $G[c_i, c_{i+1}, a, b, c_{i-1}, c_{i-2}]$  is isomorphic to  $S_{1,2,2}$ , a contradiction. Hence,  $V_i$  is anti-complete to  $V_{i+2}$  and by symmetry also anti-complete to  $V_{i-2}$ .  $\diamond$

- (2)  $W_i$  is anti-complete to  $W_{i-2} \cup W_{i+2}$ .

This is because  $G$  is  $K_3$ -free.  $\diamond$

- (3)  $W_i$  is complete to  $V_i$ , and anti-complete to  $V_{i-1} \cup V_{i+1}$ .

Suppose that there are vertices  $a \in W_i$  and  $b \in V_i$  that are not adjacent to each other. Then  $G[c_{i-1}, a, c_i, b, c_{i-2}, c_{i-3}]$  is isomorphic to  $S_{1,2,2}$ , a contradiction. This implies that  $W_i$  is complete to  $V_i$ . As  $G$  is  $K_3$ -free,  $W_i$  is anti-complete to  $V_{i-1} \cup V_{i+1}$ .  $\diamond$

- (4)  $G[V_i \cup W_{i+2}]$ ,  $G[V_i \cup W_{i-2}]$ ,  $G[W_i \cup W_{i+1}]$  are  $2P_2$ -free.

Suppose that there are  $a_1, a_2 \in V_i$  and  $b_1, b_2 \in W_{i+2}$  such that  $a_1b_1, a_2b_2 \in E(G)$  and  $a_1b_2, a_2b_1 \notin E(G)$ . Then  $G[c_i, c_{i-1}, a_1, b_1, a_2, b_2]$  is isomorphic to  $S_{1,2,2}$ , a contradiction. So,  $G[V_i \cup W_{i+2}]$  is  $2P_2$ -free, and by symmetry  $G[V_i \cup W_{i-2}]$  is  $2P_2$ -free. Suppose that there are  $a_1, a_2 \in W_i$  and  $b_1, b_2 \in W_{i+1}$  such that  $a_1b_1, a_2b_2 \in E(G)$  and  $a_1b_2, a_2b_1 \notin E(G)$ . Then  $G[c_{i-1}, c_{i-2}, a_1, b_1, a_2, b_2]$  is isomorphic to  $S_{1,2,2}$ , a contradiction.  $\diamond$

- (5) For  $v \in V_i, w \in W_{i+2}, z \in W_{i-2}$ ,  $\{v, w, z\}$  is not an independent set.

Suppose that such  $v, w, z$  forming an independent set exist. Then  $G[c_i, v, c_{i+1}, w, c_{i-1}, z]$  is isomorphic to  $S_{1,2,2}$ , a contradiction.  $\diamond$

By Claims (4) and (5) and the fact that  $G$  is  $K_3$ -free, we deduce that for each  $i \in \{1, \dots, 5\}$ ,  $G[V_i, W_{i-2}, W_{i+2}]$  is a nice 3-partite graph. Moreover,  $V_i$  is complete to  $V_{i-1}, V_{i+1}, W_i$  and anti-complete to  $V_{i+2}, V_{i+3}, W_{i-1}, W_{i+1}$ , whereas  $W_i$  is anti-complete to  $W_{i-2}, W_{i+2}$ . By doing bipartite complementations between  $V_i$  and  $V_{i+1} \cup W_i$  for each  $i \in \{1, \dots, 5\}$ , we may assume that each edge not incident to a vertex of  $C$  belongs to  $G[V_i, W_{i+2}, W_{i-2}]$  for some  $i \in \{1, \dots, 5\}$ .

Our goal is now to compute a refined set of pairwise non-intersecting 3-partite graphs by doing a small number of bipartite complementations such that each new 3-partite graph satisfies conditions (a) and (b) of Lemma 5.9.

We first observe that each  $G[V_i, W_{i-2}, W_{i+2}]$  intersects only  $G[V_{i+1}, W_{i-1}, W_{i-2}]$  and  $G[V_{i-1}, W_{i+2}, W_{i+1}]$ . We now aim, for each  $i \in \{1, \dots, 5\}$ , to split  $V_i, W_{i-2}$  and  $W_{i+2}$  in such a way that we can construct the desired non-intersecting 3-partite graphs after some bipartite complementations. We will use the same construction as in [12], but the way we use the different sets differs from [12].

For each  $i \in \{1, \dots, 5\}$ , let us define the following partition of  $W_i$ :

$$\begin{aligned} W_i^- &= \{x \in W_i \mid x \text{ has a non-neighbour in } W_{i+1}\}, \\ W_i^+ &= \{x \in W_i \setminus W_i^- \mid x \text{ has a non-neighbour in } W_{i-1}\}, \\ W_i^* &= W_i \setminus (W_i^- \cup W_i^+). \end{aligned}$$

By definition,  $W_i^+ \cup W_i^*$  is complete to  $W_{i+1}$  and  $W_i^-$  is complete to  $W_{i+1}^*$ . We claim that  $W_i^-$  is also complete to  $W_{i+1}^-$ . Suppose that a vertex  $x \in W_i^-$  has a non-neighbour  $y$  in  $W_{i+1}^-$ .

Then, by definition  $y$  has a non-neighbour  $z$  in  $W_{i+2}$ . Therefore,  $G[c_{i+1}, z, x, c_{i-1}, c_{i+2}, y]$  is isomorphic to  $S_{1,2,2}$  because  $x$  is not adjacent to  $z$  by (2), a contradiction. Thus,  $W_i^-$  is complete to  $W_{i+1}^-$ .

We now show some relationships between the  $V_i$ 's and  $W_i$ 's, which were also proven in [12], but we add the proofs for completeness. Let  $i \in \{1, \dots, 5\}$ .

(a)  $V_i$  is anti-complete to  $W_{i+2}^+$  and  $W_{i-2}^-$ .

Suppose, for contradiction, that a vertex  $x \in V_i$  has a neighbour  $z \in W_{i+2}^+ \cup W_{i-2}^-$ . First suppose that  $z \in W_{i+2}^+$ . Then, by definition,  $z$  has a non-neighbour  $y \in W_{i+1}$ . However, now  $G[c_i, c_{i-1}, x, z, y, c_{i+2}]$  is isomorphic to  $S_{1,2,2}$ , a contradiction. Now suppose that  $z \in W_{i-2}^-$ . Then, by definition,  $z$  has a non-neighbour  $y \in W_{i-1}$ . However, now  $G[c_i, c_{i+1}, y, c_{i-2}, x, z]$  is isomorphic to  $S_{1,2,2}$ , another contradiction.  $\diamond$

(b) If a vertex  $x \in V_i$  has a neighbour in  $W_{i+2}^*$  (resp.  $W_{i-2}^*$ ), then  $x$  is complete to  $W_{i+2}^-$  and anti-complete to  $W_{i-2}$  (resp. complete to  $W_{i-2}^+$  and anti-complete to  $W_{i+2}$ ).

Suppose  $x \in V_i$  has a neighbour  $y$  in  $W_{i+2}^*$ . Because  $W_{i+2}^*$  is complete to  $W_{i+3} = W_{i-2}$ , we find that  $x$  is anti-complete to  $W_{i-2}$ . Suppose that  $x$  has a non-neighbour  $z$  in  $W_{i+2}^-$ . By definition, there is a vertex  $w \in W_{i-2}$  not adjacent to  $z$ , but adjacent to  $y$ . Then,  $G[y, x, w, c_{i-1}, c_{i+1}, z]$  is isomorphic to  $S_{1,2,2}$ , a contradiction.

Now suppose that  $x \in V_i$  has a neighbour  $y$  in  $W_{i-2}^*$ . Because by definition  $W_{i-2}^*$  is complete to  $W_{i-3} = W_{i+2}$ , we find that  $x$  is anti-complete to  $W_{i+2}$ . Suppose that  $x$  has a non-neighbour  $z$  in  $W_{i-2}^+$ . By definition,  $z$  has a non-neighbour  $w$  in  $W_{i+2}$ , which is adjacent to  $y$ . Again,  $G[y, x, w, c_{i+1}, c_{i-1}, z]$  is isomorphic to  $S_{1,2,2}$ , a contradiction.  $\diamond$

For each  $i \in \{1, \dots, 5\}$ , let

$$\begin{aligned} V_i^1 &= \{x \in V_i \mid x \text{ has a neighbour in } W_{i+2}^*\}, \\ V_i^2 &= \{x \in V_i \mid x \text{ has a neighbour in } W_{i-2}^*\}, \\ V_i^3 &= \{x \in V_i \mid x \text{ has no neighbour in } W_{i+2}^* \cup W_{i-2}^*\}. \end{aligned}$$

From (a), we know that for each  $x \in V_i$ , in the graph  $G[V_i \cup W_{i-2} \cup W_{i+2}]$ ,  $x$  can have neighbours in only  $W_{i+2}^- \cup W_{i+2}^*$  and in  $W_{i-2}^+ \cup W_{i-2}^*$ . From (b), every vertex in  $V_i^1$  is complete to  $W_{i+2}^-$  and anti-complete to  $W_{i-2}$ , and every vertex in  $V_i^2$  is complete to  $W_{i-2}^+$  and anti-complete to  $W_{i+2}$ . Thus,  $V_i^1, V_i^2, V_i^3$  are disjoint sets.

The common neighbours between a vertex  $x \in V_i$  and  $y \in V_{i+1}$  are in  $W_{i-2}^*$  and the common neighbours between  $x \in V_i$  and  $y \in V_{i-1}$  are in  $W_{i+2}^*$ . For each  $i \in \{1, \dots, 5\}$ , we define the following 3-partite graphs:

$$\begin{aligned} G_i^1 &= G[V_i^1, V_{i-1}^2, W_{i+2}^*], \\ G_i^2 &= G[V_i^3, W_{i+2}^-, W_{i-2}^+]. \end{aligned}$$

Now, we do some bipartite complementations. First we do bipartite complementations between  $W_i^+ \cup W_i^*$  and  $W_{i+1}$ , and between  $W_i^-$  and  $W_{i+1}^- \cup W_{i+1}^*$  for each  $i$ . The resulting graph has the property that the edges between  $W_i$  and  $W_{i+1}$  are always between  $W_i^-$  and  $W_{i+1}^+$ . Secondly, we do bipartite complementations between  $V_i^1$  and  $W_{i+2}^-$ , and  $V_i^2$  and  $W_{i-2}^+$  for each  $i$ . Lastly, we remove  $C$ . Observe that the edges of the remaining graph are all contained in one  $G_i^j$  for some  $i \in \{1, \dots, 5\}$  and  $j \in \{1, 2\}$ , and by definition the  $G_i^j$ 's are pairwise disjoint.

Because all of the graphs  $G_i^j$  are disjoint, the linear rank-width of the resulting graph is the maximum linear rank-width of its connected components. Since each connected component is a nice 3-partite graph, by Lemma 5.9, we conclude that the linear rank-width of the resulting graph is at most 3. We will now count the number of times we applied bipartite complementations. For each  $i \in \{1, \dots, 5\}$ , we did one bipartite complementation to keep only the edges between  $V_i$  and  $W_{i-2} \cup W_{i+2}$ , and then two bipartite complementations to remove the edges between  $W_i$  and  $W_{i+1}$  apart from those between  $W_i^-$  and  $W_{i+1}^+$ , and finally two bipartite complementations to remove the edges between  $V_i^1$  and  $W_{i+2}^-$ , and  $V_i^2$  and  $W_{i-2}^+$ , that is, in total five bipartite complementations, resulting in a total of 25 bipartite complementations. So, by Lemma 2.3, the graph  $G - V(C)$  has linear rank-width at most  $3 + 2 * 25 = 53$ . Because  $|V(C)| = 5$ , we conclude that the linear rank-width of  $G$  is at most 58.  $\square$

We are now ready to prove Theorem 1.10.

**Theorem 1.10 (restated).** *Every  $(K_3, S_{1,2,2})$ -free graph has linear rank-width at most 58.*

*Proof.* Let  $G$  be a  $(K_3, S_{1,2,2})$ -free graph. We may assume that  $G$  is connected. If  $G$  is bipartite, we use Proposition 5.5. If  $G$  is non-bipartite but  $C_5$ -free, we use Proposition 5.7. In the remaining case, we use Proposition 5.10.  $\square$

## 6 Concluding Remarks

In this paper we researched the relationship between pivot-minors and boundedness of linear rank-width. We first proved that for every tree  $T$  that is not a caterpillar, the class of  $T$ -pivot-minor-free graphs has unbounded linear rank-width. We then posed Conjecture 3, which states that an affirmative answer can be found whenever  $T$  is a caterpillar. We were only able to give an affirmative answer to this conjecture that holds for every caterpillar  $T$ , if the class of  $T$ -pivot-minor-free graphs is, in addition, also distance-hereditary. We also proved that the class of  $K_{1,3}$ -pivot-minor-free graphs has bounded linear rank-width. As a next step for proving Conjecture 3, it seems natural to consider the case where  $T = K_{1,r}$  for  $r \geq 4$ . We also proved Conjecture 3 for  $P = P_4$ . Since Conjecture 3 is equivalent to the alternative conjecture that for every path  $P$ , the class of  $P$ -pivot-minor-free graphs has bounded linear rank-width, the case where  $P = P_5$  is another interesting open case.

For obtaining our results (in particular, the case where  $T = K_{1,3}$ ) we followed a general strategy consisting of two steps. We believe this strategy is also useful for making further progress towards Conjecture 3. However, Step 1 of the strategy requires us to find a hereditary graph class (class of graphs that can be characterized by a set of forbidden induced subgraphs) that contains the class of  $T$ -pivot-minor-free graphs under consideration. In general, finding an appropriate hereditary graph class is a challenging task.

The fact that  $K_{1,3}$ -pivot-minor-free graphs have bounded linear rank-width follows from a stronger result that we showed, namely that  $(K_3, S_{1,2,2})$ -free graphs have bounded linear rank-width. Showing this stronger result will be useful for a systematic study on the boundedness of linear rank-width of  $(H_1, H_2)$ -free graphs. Such a classification already exists for  $H$ -free graphs, as observed in [22]: for a graph  $H$ , the class of  $H$ -free graphs has bounded linear rank-width if and only if  $H$  is a subgraph of  $P_3$  not isomorphic to  $3P_1$ . We note that similar classifications also exist for other width parameters: for the tree-width of  $(H_1, H_2)$ -free graphs [7], which was later generalized to a classification for tree-width of  $\mathcal{H}$ -free graphs, where  $\mathcal{H}$  is a finite

set of graphs [37], rank-width of  $H$ -free graphs (see [24]), rank-width of  $H$ -free bipartite graphs [23, 36, 38], and up to five non-equivalent open cases, rank-width of  $(H_1, H_2)$ -free graphs (see [8] or [22]), and for the mim-width of  $H$ -free graphs [13], whereas there is still an infinite number of open cases left for the mim-width of  $(H_1, H_2)$ -free graphs [13].

We leave a systematic study into boundedness of linear rank-width of  $(H_1, H_2)$ -free graphs for future research. Here, we only collect known results. The class of  $(H_1, H_2)$ -free graphs has bounded linear rank-width if

- one of  $H_1$  and  $H_2$  is a subgraph of  $P_3$  that is not isomorphic to  $3P_1$  [22],
- $(H_1, H_2) = (K_3, S_{1,2,2})$  and  $(3P_1, \overline{S_{1,2,2}})$  (Theorem 1.10)
- $(H_1, H_2) = (P_4, F)$  where  $F$  is a threshold graph [14].

The class of  $(H_1, H_2)$ -free graphs has unbounded linear rank-width if

- $(H_1, H_2) = (K_3, S_{1,2,3})$  or  $(3P_1, \overline{S_{1,2,3}})$  [4],
- $(H_1, H_2) = (P_4, F)$  where  $F$  is not a threshold graph [14],
- all known cases where  $(H_1, H_2)$ -free graphs have unbounded rank-width.

## References

- [1] Isolde Adler, Arthur M. Farley, and Andrzej Proskurowski. Obstructions for linear rank-width at most 1. *Discrete Appl. Math.*, 168:3–13, 2014.
- [2] Isolde Adler and Mamadou Moustapha Kanté. Linear rank-width and linear clique-width of trees. *Theoret. Comput. Sci.*, 589:87–98, 2015.
- [3] Isolde Adler, Mamadou Moustapha Kanté, and O-joung Kwon. Linear rank-width of distance-hereditary graphs I. A polynomial-time algorithm. *Algorithmica*, 78(1):342–377, 2017.
- [4] Bogdan Alecu, Mamadou Moustapha Kanté, Vadim V. Lozin, and Viktor Zamaraev. Between clique-width and linear clique-width of bipartite graphs. *Discrete Mathematics*, 343(8):111926, 2020.
- [5] Bogdan Alecu, Vadim V. Lozin, Dominique de Werra, and Viktor Zamaraev. Letter graphs and geometric grid classes of permutations: characterization and recognition. *Discrete Appl. Math.*, 283:482–494, 2020.
- [6] Dan Bienstock, Neil Robertson, Paul Seymour, and Robin Thomas. Quickly excluding a forest. *J. Combin. Theory Ser. B*, 52(2):274–283, 1991.
- [7] Hans L. Bodlaender, Nick Brettell, Matthew Johnson, Giacomo Paesani, Daniël Paulusma, and Erik Jan van Leeuwen. Steiner trees for hereditary graph classes: A treewidth perspective. *Theoret. Comput. Sci.*, 867:30–39, 2021.
- [8] Marthe Bonamy, Nicolas Bousquet, Konrad K. Dabrowski, Matthew Johnson, Daniël Paulusma, and Théo Pierron. Graph isomorphism for  $(H_1, H_2)$ -free graphs: an almost complete dichotomy. *Algorithmica*, 83(3):822–852, 2021.

- [9] André Bouchet. Transforming trees by successive local complementations. *J. Graph Theory*, 12(2):195–207, 1988.
- [10] André Bouchet. Connectivity of isotropic systems. In *Combinatorial Mathematics: Proceedings of the Third International Conference (New York, 1985)*, volume 555 of *Ann. New York Acad. Sci.*, pages 81–93. New York Acad. Sci., New York, 1989.
- [11] André Bouchet. Circle graph obstructions. *J. Combin. Theory Ser. B*, 60(1):107–144, 1994.
- [12] Andreas Brandstädt, Suhail Mahfud, and Raffaele Mosca. Bounded clique-width of  $(S_{1,2,2}$ , triangle)-free graphs. arXiv:1608.01820, 2016.
- [13] Nick Brettell, Jake Horsfield, Andrea Munaro, Giacomo Paesani, and Daniël Paulusma. Bounding the mim-width of hereditary graph classes. *J. Graph Theory*, to appear.
- [14] Robert Brignall, Nicholas Korpelainen, and Vincent Vatter. Linear clique-width for hereditary classes of cographs. *Journal of Graph Theory*, 84(4):501–511, 2017.
- [15] Bruno Courcelle. Clique-width and edge contraction. *Inform. Process. Lett.*, 114(1-2):42–44, 2014.
- [16] Bruno Courcelle, Johann A. Makowsky, and Udi Rotics. Linear time solvable optimization problems on graphs of bounded clique-width. *Theory Comput. Syst.*, 33(2):125–150, 2000.
- [17] Bruno Courcelle and Stephan Olariu. Upper bounds to the clique width of graphs. *Discrete Appl. Math.*, 101(1-3):77–114, 2000.
- [18] William H. Cunningham and Jack Edmonds. A combinatorial decomposition theory. *Canad. J. Math.*, 32(3):734–765, 1980.
- [19] Konrad K. Dabrowski, François Dross, and Daniël Paulusma. Colouring diamond-free graphs. *J. Comput. System Sci.*, 89:410–431, 2017.
- [20] Konrad K. Dabrowski, François Dross, Jisu Jeong, Mamadou Moustapha Kanté, O-joung Kwon, Sang-il Oum, and Daniël Paulusma. Computing small pivot-minors. *Proc. WG 2018, LNCS*, 11159:125–138, 2018.
- [21] Konrad K. Dabrowski, François Dross, Jisu Jeong, Mamadou Moustapha Kanté, O-joung Kwon, Sang-il Oum, and Daniël Paulusma. Tree pivot-minors and linear rank-width. *Proc. EuroComb 2019, Acta Mathematica Universitatis Comenianae*, 88(3):577–583, 2019.
- [22] Konrad K. Dabrowski, Matthew Johnson, and Daniël Paulusma. Clique-width for hereditary graph classes. *London Mathematical Society Lecture Note Series*, 456:1–56, 2019.
- [23] Konrad K. Dabrowski and Daniël Paulusma. Classifying the clique-width of  $H$ -free bipartite graphs. *Discrete Appl. Math.*, 200:43–51, 2016.
- [24] Konrad K. Dabrowski and Daniël Paulusma. Clique-width of graph classes defined by two forbidden induced subgraphs. *The Computer Journal*, 59(5):650–666, 2016.

- [25] Wolfgang Espelage, Frank Gurski, and Egon Wanke. How to solve NP-hard graph problems on clique-width bounded graphs in polynomial time. *Proc. WG 2001, LNCS*, 2204:117–128, 2001.
- [26] Robert Ganian. Thread graphs, linear rank-width and their algorithmic applications. *Proc. IWOCA 2010, LNCS*, 6460:38–42, 2011.
- [27] Jim Geelen, O-joung Kwon, Rose McCarty, and Paul Wollan. The grid theorem for vertex-minors. *Journal of Combinatorial Theory, Series B*, in press.
- [28] Michael U. Gerber and Daniel Kobler. Algorithms for vertex-partitioning problems on graphs with fixed clique-width. *Theoretical Computer Science*, 299(1):719–734, 2003.
- [29] Jisu Jeong, Eun Jung Kim, and Sang-il Oum. The “art of trellis decoding” is fixed-parameter tractable. *IEEE Trans. Inform. Theory*, 63(11):7178–7205, 2017.
- [30] Mamadou Moustapha Kanté and O-joung Kwon. Linear rank-width of distance-hereditary graphs II. Vertex-minor obstructions. *European J. Combin.*, 74:110–139, 2018.
- [31] Navin Kashyap. Matroid pathwidth and code trellis complexity. *SIAM J. Discrete Math.*, 22(1):256–272, 2008.
- [32] Daniel Kobler and Udi Rotics. Edge dominating set and colorings on graphs with fixed clique-width. *Discrete Applied Mathematics*, 126(2–3):197–221, 2003.
- [33] O-joung Kwon, Rose McCarty, Sang-il Oum, and Paul Wollan. Obstructions for bounded shrub-depth and rank-depth. *J. Combin. Theory Ser. B*, 149:76–91, 2021.
- [34] O-joung Kwon and Sang-il Oum. Graphs of small rank-width are pivot-minors of graphs of small tree-width. *Discrete Appl. Math.*, 168:108–118, 2014.
- [35] Vadim V. Lozin.  $E$ -free bipartite graphs. *Diskretn. Anal. Issled. Oper. Ser. 1*, 7(1):49–66, 103, 2000.
- [36] Vadim V. Lozin. Bipartite graphs without a skew star. *Discrete Mathematics*, 257(1):83–100, 2002.
- [37] Vadim V. Lozin and Igor Razgon. Tree-width dichotomy. arXiv:2012.01115, 2020.
- [38] Vadim V. Lozin and Jordan Volz. The clique-width of bipartite graphs in monogenic classes. *International Journal of Foundations of Computer Science*, 19(2):477–494, 2008.
- [39] Jaroslav Nešetřil, Patrice Ossona de Mendez, Roman Rabinovich, and Sebastian Siebertz. Classes of graphs with low complexity: The case of classes with bounded linear rankwidth. *European J. Combin.*, 91:103223, 2021.
- [40] Sang-il Oum. Rank-width and vertex-minors. *J. Combin. Theory Ser. B*, 95(1):79–100, 2005.
- [41] Sang-il Oum. Excluding a bipartite circle graph from line graphs. *J. Graph Theory*, 60(3):183–203, 2009.

- [42] Sang-il Oum. Rank-width: algorithmic and structural results. *Discrete Appl. Math.*, 231:15–24, 2017.
- [43] Sang-il Oum and Paul Seymour. Approximating clique-width and branch-width. *J. Combin. Theory Ser. B*, 96(4):514–528, 2006.
- [44] Michaël Rao. MSOL partitioning problems on graphs of bounded treewidth and clique-width. *Theoretical Computer Science*, 377(1–3):260–267, 2007.
- [45] Neil Robertson and P. D. Seymour. Graph minors. I. Excluding a forest. *J. Combin. Theory Ser. B*, 35(1):39–61, 1983.
- [46] Neil Robertson and P. D. Seymour. Graph minors. V. Excluding a planar graph. *J. Combin. Theory Ser. B*, 41(1):92–114, 1986.
- [47] Neil Robertson and P. D. Seymour. Graph minors. XX. Wagner’s conjecture. *J. Combin. Theory Ser. B*, 92(2):325–357, 2004.
- [48] Atsushi Takahashi, Shuichi Ueno, and Yoji Kajitani. Minimal acyclic forbidden minors for the family of graphs with bounded path-width. *Discrete Math.*, 127(1-3):293–304, 1994.
- [49] Mihalis Yannakakis. The complexity of the partial order dimension problem. *SIAM J. Algebraic Discrete Methods*, 3(3):351–358, 1982.