UNIVERSITY OF LEEDS

This is a repository copy of *Nondestructive diagnostics of magnesium deficiency based on distribution features of chlorophyll concentrations map on cucumber leaf*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/179313/>

Version: Accepted Version

Article:

Shi, J, Li, W, Zhai, X et al. (4 more authors) (2019) Nondestructive diagnostics of magnesium deficiency based on distribution features of chlorophyll concentrations map on cucumber leaf. *Journal of Plant Nutrition*, 42 (20). pp. 2773-2783. ISSN 0190-4167

<https://doi.org/10.1080/01904167.2019.1659332>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

1 Nondestructive diagnostics of magnesium deficiency based on
2 distribution features of chlorophyll concentrations map on cucumber leaf

3 Jiyong Shi^{a, c}, Wenting Li^a, Xiaodong Zhai^a, Zhiming Guo^{a, c}, Mel Holmes^{b, c}, Haroon
4 Elrasheid Tahir^a, Xiaobo Zou^{a, c, *}

5 ^a School of Food and Biological Engineering, Jiangsu University, Zhenjiang 212013,
6 China

7 ^b School of Food Science and Nutrition, the University of Leeds, Leeds LS2 9JT, United
8 Kingdom

9 ^c Joint Laboratory of China-UK on food nondestructive sensing, Jiangsu University,
10 Zhenjiang 212013, China

11 *Corresponding author. Tel: +86 511 88780201; Fax: +86 511 88780201

12 Email address: zou_xiaobo@ujs.edu.cn (Xiaobo Zou)

Abstract

A new and nondestructive method for diagnosing magnesium (Mg) deficiency based on chlorophyll concentration distribution features of cucumber leaves was proposed. Mg deficient cucumber plants and Control plants were grown under non-soil conditions with special nutrient supply. Cucumber leaves were employed to collect hyperspectral images using a **visible and near infrared** (VIS/NIR) hyperspectral imaging system (400-900 nm) and determine reference chlorophyll concentrations using high performance liquid chromatography (HPLC). An optimal chlorophyll concentration calibration model ($R_p = 0.9087$) was constructed and used to detect chlorophyll distribution maps of Mg deficient leaves and Control leaves. Results shown that chlorophyll content distributed more unevenly on Mg deficient leaves than Control leaves. The Standard Deviation (SD) value of the chlorophyll content at all the pixels on a chlorophyll distribution map was calculated for Mg deficient diagnostics. An Mg deficiency diagnostics model with satisfied performance (diagnostic rate 93.33%) was obtained. The result indicated the SD value of chlorophyll concentrations on the whole cucumber leaf could be employed to diagnose Mg deficiency nondestructively.

Keywords: diagnostics; magnesium; deficiency; distribution; chlorophyll; cucumber

1. Introduction

Magnesium (Mg) is an essential macronutrient element in plant growth and productivity, the shortage of Mg element causes imbalances in plant growth and drastically most importantly affect the quality and yield of agricultural products (Khaitov, 2018; Kwano et al., 2017; Ortas, 2018). Low level of Mg supplement often result in low chlorophyll production, which drastically impact the plant growth, flowering and fruit bearing (Farhat et al., 2016; Farzadfar et al., 2017; Saghaiesh et al., 2019). Moreover, Mg shortage also affects the production of chemical components that are connected with the product qualities, like appearance, aroma, taste and nutrition (Canizella et al., 2015; Gomez-Perez et al., 2018; Nikolic et al., 2014; Schurt et al., 2014). Although Mg deficiency diagnostics methods based on the Mg element content analysis with chemical methods, like inductively coupled plasma mass spectrometry, get good performance for diagnosing Mg deficient plants, the chemical analysis process is time-consuming, laborious and high cost (Guo et al., 2016; Nartvaranant, 2018). Therefore, more attention should be paid to the diagnostics methods that assess Mg status more efficiently.

Rapid and nondestructive diagnostics methods based on leaf chlorophyll analysis have been proposed to diagnose nutrients deficiencies in plants. It is well known that the shortage of Mg element results in low chlorophyll in plant leaves and low chlorophyll density regions appear on Mg deficient plant leaves, so chlorophyll density can be used reasonably as an indicator for Mg deficient diagnostics (Samborska et al.,

2018; Tatagiba et al., 2016). Many research demonstrated that reflectance in Vis/NIR spectrum is sensitive to the chlorophyll concentrations of leaf tissues (Cayuela et al., 2014; Kurenda et al., 2014; Schouten et al., 2014; Zhou et al., 2017). Therefore rapid and nondestructive diagnostics methods based on soil and plant analyzer development (SPAD) analysis, visible (Vis) spectral technology or near infrared (NIR) technology have been proposed to diagnose Mg deficiency (Marian et al., 2009; Shaahan et al., 1999). In these methods, one value was used to represent the chlorophyll level of the whole leaf sample. However, Mg deficiency causes a low chlorophyll density in small regions of a cucumber leaf at the early stage. The measurement of chlorophyll content on the whole cucumber leaf can improve the performance for diagnosis of Mg deficiency.

Hyperspectral imaging technology has been used to detect the chlorophyll concentration distribution map on cucumber leaves. Hyperspectral imaging acquire both spectral and spatial information from an object and produces a three dimensional data cube that includes the spectral data of each pixel. As the spectral data of pixels is sensitive to the quality compounds in biological products, hyperspectral imaging technology has been employed to determine the distribution of various quality indexes in food and agricultural products, like triterpene acids in loquat leaf (Shi et al., 2018), compounds (protein, carbohydrate, sialic acid) in edible bird's nest (Shi et al., 2017), chlorophyll in pepper leaves (Yu et al., 2016), tenderness in salmon (He et al., 2014), etc. These previous studies indicate that hyperspectral imaging technology could

determine the chlorophyll concentration distribution map on a cucumber leaf, and the chlorophyll distribution map can be used for nutrient diagnostics.

In this paper, the distribution features of chlorophyll content were employed to establish new diagnostics method for diagnosing Mg deficiency. Mg deficient leaf samples were cultured under non-soil conditions and Mg status of the leaf samples were confirmed by Mg element analysis. Cucumber leaves were employed to collect hyperspectral images using a VIS/NIR hyperspectral imaging system (400-900 nm) and determine reference chlorophyll concentrations using high performance liquid chromatography (HPLC). The chlorophyll distribution maps of Mg deficient leaves and Control leaves were detected nondestructively. The main objectives of this paper are to (1) find differences in chlorophyll content distribution maps of Mg deficient and Control leaf samples; (2) extract suitable chlorophyll distribution features for diagnosing Mg deficiency based on chlorophyll content distribution maps.

2. Materials and methods

2.1 leaf samples

Fresh leaf of Mg deficient cucumber plants and Control cucumber plants (*Cucumis sativus*, Biyu-3, FuNong seeds Co. Ltd., Shanghai, China) were used as analytical samples. Cucumber seeds were germinated in plug plant seed tray with perlite rock and distilled water. Seedlings were thinned to one plant/pot with a planting density of 20 cm × 20 cm. Complete nutrient solution with all necessary elements was applied to the

cucumber plants before the three-leaf stage, then all the cucumber plants were divided into two groups, Control group and Mg deficiency group. Control fertilization with all macronutrients was performed to plants in Control group, and no Mg fertilization (complete nutrient solution with Mg eliminated) was performed to plants in Mg deficiency group. The components of complete nutrient solution were NO_3 11.75 mmol L^{-1} , NH_4 1.0 mmol L^{-1} , H_2PO_4 1.25 mmol L^{-1} , K 6.5 mmol L^{-1} , Ca 2.75 mmol L^{-1} , Mg 1.0 mmol L^{-1} , SO_4 1.0 mmol L^{-1} , Fe 15 $\mu\text{mol L}^{-1}$, Mn 10 $\mu\text{mol L}^{-1}$, Zn 5 $\mu\text{mol L}^{-1}$, B 25 $\mu\text{mol L}^{-1}$, Cu 0.75 $\mu\text{mol L}^{-1}$, Mo 0.5 $\mu\text{mol L}^{-1}$. All the plants were grown under non-soil conditions (perlite rock) in green houses at Jiangsu University in China (32.11N, 119.27E). The amount of nutrient solution for each plant per day was 500ml for the first two weeks after transplanting and 800-1000ml until harvesting. The experiment was performed in three replications, and 60 Mg deficient plants and 60 Control plants were obtained.

2.2 Mg element determination

To detect the nutrient status of plants in Mg deficient group and Control group, Mg element concentrations in nutrient deficient groups and Control group were determined. 45 leaves at the first three nodes, the middle nodes, and the youngest nodes in Mg deficient and Control plants were collected for Mg elements determination. For each Mg element determination, 0.4g fresh leaf tissues without leaf veins was dried at 80°C to a constant weight. The dry matter was dissolved into 25 mL nitric and perchloric acid mixture, placed on a electric hot plate and heated at 200 °C for 60 mins. Then the digest

was analyzed using atomic absorption spectrophotometer (AA300, PerkinElmer Inc, USA) (Sofi et al., 2017).

2.3 Hyperspectral image collection

A line-scan reflectance hyperspectral imaging system with a spectral resolution of 2.8 nm was employed to collect raw hyperspectral image of cucumber leaf samples. The imaging system includes a spectrograph (ImSpector, V10E, Spectra Imaging Ltd., Finland), a high-performance camera (Bci4-1300, C-Cam Technologies, Belgium), an illumination unit consisting two fibre-optic light-guiding branches connected to a DC illuminator (DC-950A, Fiber Lite Illuminator, USA), a transitional stage operated by a stepper motor (Zolix SC300-1A, Zolix. Corp., China), an enclosure (450×600×750 mm) and a computer. Cucumber leaf sample was placed on the conveyor belt to be scanned line by line using the hyperspectral imaging system. In order to improve the signal-to-noise ratio, the speed of sample movement, exposure time and the distance of a sample from the camera were optimized and set to 10 mm/scan, 45 ms and 18 cm. White reference image (a white ceramic tile considering to be ~99% reflective) and dark reference image (an image taken when the light is off and the lens is covered) were acquired to correct the raw images. The three dimensional data cube of a cucumber leaf hyperspectral image was shown in Fig. 1.

Insert Fig. 1 here.

137

138 2.4 Chlorophyll concentration distribution map determination

139 As described in 2.3, the hyperspectral imaging data cube contains a spectrum with
140 a specific wavelength range for each pixel in a 2-dimensional image of the sample. At
141 the same time, researchers demonstrated there are good relationship between leaf
142 spectra and chlorophyll concentrations (Cayuela et al., 2014; Kurenda et al., 2014).
143 Therefore, it possible to estimate the distribution of chlorophyll concentrations on the
144 whole cucumber leaf based on the spectra data of each pixel and the chlorophyll
145 concentration calibration model. Basically, chlorophyll concentration distribution map
146 determination contains three steps, (1) building chlorophyll calibration models, (2)
147 testing the calibration models, and (3) determining chlorophyll distribution map.

148

149 2.4.1 Building calibration models

150 A calibration set and a prediction set were composed of 120 fresh cucumber
151 leaves. After hyperspectral image acquisition, spectral data of cucumber leaves in the
152 calibration set and prediction set was extracted by defining an **region of interest** (ROI)
153 (50×50pixels) in the center region of hyperspectral images. **Partial least squares**
154 (PLS), **interval partial least squares** (iPLS), **simulated annealing - Interval partial least**
155 **squares** (SA-iPLS) were employed to build chlorophyll calibration models based on
156 the extracted spectral data and reference chlorophyll concentrations determined using

high performance liquid chromatography (HPLC) (Shi et al., 2016; Shi et al., 2012).

The root mean square error of cross-validation (RMSECV), the correlation coefficient in the calibration set (R_c), the root mean square error of prediction (RMSEP) and the correlation coefficient in the prediction set (R_p) (Shi et al., 2012) were calculated to evaluate the established calibration models.

2.4.2 Testing the calibration models

An independent testing set was composed of 40 fresh cucumber leaves. After hyperspectral image data acquisition, the spectral data (430-960nm) were extracted and substituted in the established PLS, iPLS, SA-iPLS chlorophyll concentration calibration models to predict the chlorophyll concentrations of the samples in the testing set. Then the reference chlorophyll concentrations of the testing samples were detected using HPLC. Finally, the root mean square error (RMSET) and the correlation coefficient of the testing set (R_t) were calculated to test the established calibration models.

2.4.3 Estimating chlorophyll distribution map

The flowchart of estimating a chlorophyll distribution map was shown in Fig. 2. After hyperspectral image acquisition, data cube of all pixels were extracted and

converted to spectra curves. The spectra data of every pixel was substituted in the established chlorophyll concentration calibration model to predict chlorophyll concentrations for each pixel. Then chlorophyll concentrations for each pixel were ranked into a 2-D matrix according to the coordinates of the pixels. Finally, chlorophyll distribution map can be obtained by converting the 2-D matrix to a 2-D figure, as shown in Fig. 2.

Insert Fig. 2 here.

2.5 Diagnostics of Mg deficiency based on chlorophyll distribution map

Leaf samples in Mg deficiency group and Control group were used to detect chlorophyll concentration distribution maps using the methods described in above sections. The chlorophyll distribution maps of Mg deficient leaves were compared with those of Control leaves. Based on the results of comparison, chlorophyll distribution features were extracted and used to develop new Mg deficiency diagnostic methods.

3. Results and discussion

3.1 Mg nutrient status of cucumber plants

Mg content in Control plants growing with complete nutrient solution and Mg deficiency group growing with no Mg nutrient solution were analyzed using the method

described in section 2.1. 15 cucumber leaves at the first three nodes, 15 cucumber leaves at the middle nodes and 15 cucumber leaves at the highest nodes in plants of Mg deficiency group and Control group were collected for Mg analysis. The results of Mg element analysis for Mg deficiency group and Control group were shown in Fig. 3. In Fig. 3, it shows that the averages of Mg element at the first three nodes, middle nodes and the highest nodes in plants of Mg deficiency group were 0.81 mg/g, 1.15 mg/g and 1.39 mg/g, respectively. While in the plants of Control group, the averages of Mg element at the first three nodes, middle nodes and the highest nodes were 1.04 mg/g, 1.26 mg/g and 1.43 mg/g, respectively. It could be found that the averages of Mg elements increase from the lowest nodes to the highest nodes. It could also be found that the average of Mg element at the first three nodes in Mg deficiency group is obviously lower than the averages of the other nodes. This results show that leaves at the first three nodes in Mg deficiency group have entered into Mg deficient status.

Insert Fig. 3 here.

3.2 Chlorophyll concentration distribution maps of Mg deficient leaves

3.2.1 Building calibration models

As described in section 2.4.1, a calibration set and a prediction set contained 120 fresh cucumber leaves were used to build chlorophyll content calibration models.

After hyperspectral image collection and HPLC analysis, the hyperspectral cucumber

images and reference chlorophyll content were obtained. Chemometrics methods PLS, iPLS, SA-iPLS were employed to build chlorophyll content calibration models using the hyperspectral image data and reference chlorophyll content. The performance of the PLS, iPLS, SA-iPLS models was shown in table 1. The R_c for PLS, iPLS and SA-iPLS calibration models were 0.8149, 0.8472 and 0.9165, respectively. The R_p for PLS, iPLS and SA-iPLS calibration models were 0.7928, 0.8294 and 0.9087, respectively. The results shown that the SA-iPLS calibration model got better performance than the rest two calibration models.

Insert Table 1 here.

3.2.2 Testing the calibration models

As described in section 2.4.2, an independent testing set contained 40 fresh cucumber leaves was used to test the established chlorophyll content calibration models. After hyperspectral image data acquisition, spectra data of the samples in testing set was extracted and substituted in the established PLS, iPLS, SA-iPLS models, so the predicted chlorophyll content of the samples in testing set was calculated. Then the samples in the testing set were used to determine reference chlorophyll content. Finally, the root mean square error of testing (RMSET) and the

correlation coefficient in the testing set (R_t) were calculated to test the established calibration models, as shown in Table 1. It could be found that the RMSET and R_t based on SA-iPLS were was 2.12 mg/g and 0.8938, respectively. The results mean that chlorophyll calibration model based on SA-iPLS gets better performance for predicting chlorophyll content of an unknown leaf sample.

3.2.3 Estimating distribution map

Mg deficient cucumber leaves and Control group cucumber leaves were used to collect hyperspectral images, then data cube of all pixels were extracted to estimate chlorophyll distribution maps, as described in Section 2.4.3. The chlorophyll distribution maps of cucumber leaves in Control group and Mg deficiency group are shown in Fig.4. The pixels with high chlorophyll content appeared along the main leaf veins on the distribution maps of Control group and Mg deficiency group. However, the pixels between the main leaf veins got lower chlorophyll content on the chlorophyll distribution maps of Mg deficiency group than those of Control group. This result made that chlorophyll content distributed less evenly on Mg deficient leaves than that on Control leaves. Results indicated that the Standard Deviation (SD) value of the chlorophyll content at all the pixels on a chlorophyll distribution map could be used as indicator for diagnosing Mg deficiency.

Insert Fig. 4 here.

3.3 Diagnostics of Mg deficiency based on chlorophyll distribution map

A calibration set and a prediction set containing 120 fresh Mg deficient leaves and Control leaves picked from the first nodes of cucumber plants were used to develop an Mg deficiency diagnostic method. After hyperspectral image acquisition, chlorophyll distribution maps of the 120 cucumber leaves were detected using the procedures described in section 3.2. The SD of chlorophyll content of all pixels in a chlorophyll distribution map was calculated for diagnostics of Mg deficiency, as the results in section 3.2.3 indicated that the homogeneity of chlorophyll content at the pixels of Mg deficient leaves was worse than that of Control leaves. The SD values of the Mg deficient and Control leaves in the calibration set were shown in Fig. 5.

Fig. 5 shows the SD values of the chlorophyll content at all pixels of the Mg deficient cucumber leaves and Control cucumber leaves. The SD values of chlorophyll concentrations in Mg deficient leaves were higher than 2.25 mg/g, whereas most SD values of Control leaves were lower than 2.25 mg/g. Therefore, 2.25 mg/g was defined as a threshold for detecting Mg deficiency, as shown in Fig. 5. According to this threshold, only one sample from Control group was misclassified into Mg deficiency group, as shown in Fig. 5 (a). After applying this threshold to an independent set (30 Mg deficiency leaves and 30 Control leaves), a diagnostic rates of 93.33% was achieved, as shown in Fig. 5 (b). Control leaves at the first node enter into aging status, in which chlorophyll in some small regions decreases slowly, early than the leaves at higher nodes. Chlorophyll decreasing in these leaves can increase their SD values and result

in the misclassification. Mg deficient leaves at the third nodes enter into Mg deficiency, which decreases chlorophyll in some small regions, lately than the leaves at lower nodes. Chlorophyll decreasing in these leaves can decrease their SD values and result in the misclassification.

Insert Fig. 5 here.

4. Conclusion

An optimal chlorophyll content calibration model based on the hyperspectral images of cucumber leaves and the reference chlorophyll content of cucumber leaves was established. Chlorophyll distribution maps of Mg deficient leaves and Control leaves were obtained through calculating the chlorophyll content at each pixel by substituting its hyperspectral signal to the optimal chlorophyll content calibration model. Compared with the chlorophyll concentration distribution map, the main feature of Mg deficient chlorophyll distribution map is that chlorophyll content decreases in the regions between main leaf veins. The SD value of the chlorophyll content at all the pixels in Mg deficient and Control leaves were extracted and used as the indicator for diagnosing Mg deficiency. Result shown that a diagnostics model with good performance (diagnostic rate 93.33%) were established for measurement of Mg deficiency in cucumber plants. The result indicated that the extracted chlorophyll distribution feature could be employed to diagnose Mg deficiency.

299

300 Acknowledgment

301 The authors gratefully acknowledge the financial support provided by the National Key
302 Research and Development Program of China (2017YFC1600805), the National
303 Natural Science Foundation of China (31772073, 60901079), the Natural Science
304 Foundation of Jiangsu Province (BE2016306, BK20130505), China Postdoctoral
305 Science Foundation (2016M600379), Natural Science Foundation of the Jiangsu
306 Higher Education Institutions of China (16KJB550002), Jiangsu Planned Projects for
307 Postdoctoral Research Funds (1601080B), Six Talent Peaks Project in Jiangsu Province
308 (GDZB-016), and Priority Academic Program Development of Jiangsu Higher
309 Education Institutions (PAPD).

310

311 References

- 312 Canizella, B. T., Moreira, A., Moraes, L., Fageria, N. K. 2015. Efficiency of magnesium
313 use by common bean varieties regarding yield, physiological components, and
314 nutritional status of plants. *Communications in Soil Science and Plant Analysis*.
315 46:1376-1390.
- 316 Cayuela, J. A., Yousfi, K., Martinez, M. C., Garcia, J. M. 2014. Rapid determination of
317 olive oil chlorophylls and carotenoids by using visible spectroscopy. *Journal of the*
318 *American Oil Chemists Society*. 91:1677-1684.
- 319 Farhat, N., Elkhouni, A., Zorrig, W., Smaoui, A., Abdelly, C., Rabhi, M. 2016. Effects
320 of magnesium deficiency on photosynthesis and carbohydrate partitioning. *Acta*
321 *Physiologiae Plantarum*. 38:145.
- 322 Farzadfar, S., Zarinkamar, F., Hojati, M. 2017. Magnesium and manganese affect
323 photosynthesis, essential oil composition and phenolic compounds of *Tanacetum*
324 *parthenium*. *Plant Physiology and Biochemistry*. 112:207-217.
- 325 Gomez-Perez, L., Valdez-Aguilar, L. A., Cadena-Zapata, M., Cartmill, D. L., Cartmill,
326 A. D., Benavides-Mendoza, A. 2018. Biomass and accumulation of potassium, calcium,
327 and magnesium in gladiolus as affected by heat units and corm size. *Communications*
328 *in Soil Science and Plant Analysis*. 49:344-357.
- 329 Guo, W. L., Nazim, H., Liang, Z. S., Yang, D. F. 2016. Magnesium deficiency in plants:
330 An urgent problem. *Crop Journal*. 4:83-91.
- 331 He, H. J., Wu, D., Sun, D. W. 2014. Potential of hyperspectral imaging combined with

332 chemometric analysis for assessing and visualising tenderness distribution in raw
 333 farmed salmon fillets. *Journal of Food Engineering*. 126:156-164.

334 Khaitov, B. 2018. Effects of Rhizobium inoculation and magnesium application on
 335 growth and nodulation of soybean (*Glycine max* L.). *Journal of Plant Nutrition*.
 336 41:2057-2068.

337 Kurenda, A., Zdunek, A., Schluter, O., Herppich, W. B. 2014. VIS/NIR spectroscopy,
 338 chlorophyll fluorescence, biospeckle and backscattering to evaluate changes in apples
 339 subjected to hydrostatic pressures. *Postharvest Biology and Technology*. 96:88-98.

340 Kwano, B. H., Moreira, A., Moraes, L., Nogueira, M. A. 2017. Magnesium-manganese
 341 interaction in soybean cultivars with different nutritional requirements. *Journal of Plant*
 342 *Nutrition*. 40:372-381.

343 Marian, W., Gabriel, F., Krystyna, Z., Elzbieta, S. 2009. Early diagnostics of
 344 macronutrient deficiencies in three legume species by color image analysis. *Computers*
 345 *and Electronics in Agriculture*. 65:125-132.

346 Nartvaranant, P. 2018. Determination suitable sample method of rose apple (*Syzygium*
 347 *jambos* L.) for evaluation of plant nutrient status. *Journal of Plant Nutrition*. 41:2741-
 348 2748.

349 Nikolic, N., Borisev, M., Pajevic, S., Zupunski, M., Topic, M., Arsenov, D. 2014.
 350 Responses of wheat (*Triticum aestivum* L.) and maize (*Zea mays* L.) plants to cadmium
 351 toxicity in relation to magnesium nutrition. *Acta Botanica Croatica*. 73:359-373.

352 Ortas, I. 2018. Influence of potassium and magnesium fertilizer application on the yield

353 and nutrient accumulation of maize genotypes under field conditions. *Journal of Plant*
354 *Nutrition*. 41:330-339.

355 Saghaiesh, S. P., Souri, M. K., Moghaddam, M. 2019. Effects of different magnesium
356 levels on some morphophysiological characteristics and nutrient elements uptake in
357 Khatouni melons (cucumis melo var. Inodorus). *Journal of Plant Nutrition*. 42:27-39.

358 Samborska, I. A., Kalaji, H. M., Sieczko, L., Goltsev, V., Borucki, W., Jajoo, A. 2018.
359 Structural and functional disorder in the photosynthetic apparatus of radish plants under
360 magnesium deficiency. *Functional Plant Biology*. 45:668-679.

361 Schouten, R. E., Farneti, B., Tijssens, L., Alarcon, A. A., Woltering, E. J. 2014.
362 Quantifying lycopene synthesis and chlorophyll breakdown in tomato fruit using
363 remittance VIS spectroscopy. *Postharvest Biology and Technology*. 96:53-63.

364 Schurt, D. A., Lopes, U. P., Duarte, H., Rodrigues, F. A. 2014. Effect of Magnesium
365 on the Development of Sheath blight in Rice. *Journal of Phytopathology*. 162:617-620.

366 Shaahan, M. M., El-Sayed, A. A., El-Nour, E. A. A. A. 1999. Predicting nitrogen,
367 magnesium and iron nutritional status in some perennial crops using a portable
368 chlorophyll meter. *Scientia Horticulturae*. 82:339-348.

369 Shi, J. Y., Chen, W., Zou, X. B., Xu, Y. W., Huang, X. W., Zhu, Y. D., Shen, T. T.
370 2018. Detection of triterpene acids distribution in loquat (*Eriobotrya japonica*) leaf
371 using hyperspectral imaging. *Spectrochimica Acta Part A-Molecular and Biomolecular*
372 *Spectroscopy*. 188:436-442.

373 Shi, J. Y., Hu, X. T., Zou, X. B., Zhao, J. W., Zhang, W., Holmes, M., Huang, X. W.,

374 Zhu, Y. D., Li, Z. H., Shen, T. T., Zhang, X. L. 2017. A rapid and nondestructive
 375 method to determine the distribution map of protein, carbohydrate and sialic acid on
 376 Edible bird's nest by hyper-spectral imaging and chemometrics. *Food Chemistry*.
 377 229:235-241.

378 Shi, J. Y., Hu, X. T., Zou, X. B., Zhao, J. W., Zhang, W., Huang, X. W., Zhu, Y. D., Li,
 379 Z. H., Xu, Y. W. 2016. A heuristic and parallel simulated annealing algorithm for
 380 variable selection in near-infrared spectroscopy analysis. *Journal of Chemometrics*.
 381 30:442-450.

382 Shi, J. Y., Zou, X. B., Zhao, J. W., Holmes, M. 2012. Near infrared quantitative analysis
 383 of total flavonoid content in fresh Ginkgo biloba leaves based on different wavelength
 384 region selection methods and partial least squares regression. *Journal of Near Infrared
 385 Spectroscopy*. 20:295-305.

386 Shi, J. Y., Zou, X. B., Zhao, J. W., Wang, K. L., Chen, Z. W., Huang, X. W., Zhang, D.
 387 T., Holmes, M. 2012. Nondestructive diagnostics of nitrogen deficiency by cucumber
 388 leaf chlorophyll distribution map based on near infrared hyperspectral imaging. *Scientia
 389 Horticulturae*. 138:190-197.

390 Sofi, J. A., Rattan, R. K., Kirmani, N. A., Chesti, M. H., Bisati, I. 2017. Diagnosis and
 391 recommendation integrated system approach for major and micronutrient diagnostic
 392 norms for apple (*Malus Domestica* Borkh) under varying ages and management
 393 practices of apple orchards. *Journal of Plant Nutrition*. 40:1784-1796.

394 Tatagiba, S. D., DaMatta, F. M., Rodrigues, F. A. 2016. Magnesium decreases leaf

395 scald symptoms on rice leaves and preserves their photosynthetic performance. *Plant*
396 *Physiology and Biochemistry*. 108:49-56.

397 Yu, K. Q., Zhao, Y. R., Zhu, F. L., Li, X. L., He, Y. 2016. Mapping of chlorophyll and
398 spad distribution in pepper leaves during leaf senescence using visible and near-infrared
399 hyperspectral imaging. *Transactions of the ASABE*. 59:13-24.

400 Zhou, H., Fang, H., Pan, J., Jiang, L. J., He, Y., Shag, Y. N. 2017. Chlorophyll content
401 research of haematococcus pluvialis based on immersed Visible/Near-Infrared
402 spectroscopy. *Spectroscopy and Spectral Analysis*. 37:3375-3378.