

This is a repository copy of *Cyanidiophyceae (Rhodophyta) tolerance to precious metals: metabolic response to Cl<sub>4</sub>K<sub>2</sub>Pd and AuCl<sub>4</sub>K*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/179816/>

Version: Accepted Version

---

**Article:**

Sirakov, Maria, Palmieri, Maria Palmieri, Iovinella, Manuela et al. (5 more authors)

(Accepted: 2021) *Cyanidiophyceae (Rhodophyta) tolerance to precious metals: metabolic response to Cl<sub>4</sub>K<sub>2</sub>Pd and AuCl<sub>4</sub>K*. *Plants*. ISSN 2223-7747 (In Press)

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

Communication

# Cyanidiophyceae (Rhodophyta) Tolerance to Precious Metals: Metabolic Response to $\text{Cl}_4\text{K}_2\text{Pd}$ and $\text{AuCl}_4\text{K}$

Maria Sirakov <sup>1</sup>, Maria Palmieri <sup>2</sup>, Manuela Iovinella <sup>3,\*</sup>, Seth J. Davis <sup>3</sup>, Milena Petriccione <sup>4</sup>, Maria Rosa Di Cicco <sup>2</sup>, Mario De Stefano <sup>2</sup> and Claudia Ciniglia <sup>2</sup>

<sup>1</sup> Department of Biology and Evolution of Marine Organisms, Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Naples, Italy; maria.sirakov@szn.it

<sup>2</sup> DISTABIF, Università degli Studi di Caserta “L. Vanvitelli”, Via Vivaldi, 43, 81100 Caserta, Italy; maria.palmieri@unicampania.it (M.P.); mariarosa.dicicco@unicampania.it (M.R.D.C.); mario.destefano@unicampania.it (M.D.S.); claudia.ciniglia@unicampania.it (C.C.)

<sup>3</sup> Department of Biology, University of York, York YO10 5DD, UK; mi676@york.ac.uk; seth.davis@york.ac.uk

<sup>4</sup> Council for Agricultural Research and Economics (CREA)—Research Centre for Olive, Citrus and Tree Fruit, Via Torrino 3, 81100 Caserta, Italy; milena.petriccione@crea.gov.it

\* Correspondence: mi676@york.ac.uk; Tel.: +39-0823274582

**Abstract:** Cyanidiophyceae are polyextremophilic red algae adapted to live in geothermal and volcanic sites with a high concentration of heavy and rare metals and can mobilize and adsorb metals selectively. In this work, we assessed the capacity of 3 strains of *Galdieria* (*G. maxima*, *G. sulphuraria*, *G. phlegrea*) and one strain of *Cyanidium caldarium* to tolerate different concentrations of rare metal as  $\text{Cl}_4\text{K}_2\text{Pd}$  and  $\text{AuCl}_4\text{K}$  by monitoring algal growth in cultures exposed to the metals and investigating algae potential oxidative stress induced by them.

**Keywords:** metal tolerance; *Galdieria sulphuraria*; precious metal;s metabolic response

**Citation:** Sirakov, M.; Palmieri, M.; Iovinella, M.; Davis, S.J.; Petriccione, M.; Di Cicco, M.R.; De Stefano, M.; Ciniglia, C. Cyanidiophyceae (Rhodophyta) Tolerance to Precious Metals: Metabolic Response to  $\text{Cl}_4\text{K}_2\text{Pd}$  and  $\text{AuCl}_4\text{K}$ . *Plants* **2021**, *10*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor: Firstname Last-name

Received: date

Accepted: date

Published: date

**Publisher’s Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

In the last decade, there was a remarkable and growing demand for recovering elements and energetic resources from waste streams [1]. In particular, high levels of concern were directed towards Rare Earth Elements (REEs) due to their extensive use in superconductors, catalysts, and the electronic industry. Conversely, the discharge in the environment and the suitability of recycling REE from e-waste are relevant topics because of their hazard for the environment and health, besides their economic value. These issues become evident to both governments and electronic industries, which are increasingly prone to develop new methods to remove REEs from the environment and possibly recycle them back into a “closed-loop economy” production cycle [2–6] while simultaneously achieving energy optimization goals [7,8]. Recently, biological methods were developed to ensure the recovery of small quantities of these metals from wastewater systems [5], using mainly bacteria [9–12] or plants known for their ability to immobilize heavy metals in the cell wall and compartmentalization in vacuoles [13]. Interestingly, polyextremophilic algae have intrinsic properties that make them capable of selective removal and concentration of metals, thanks to their adaptation to live in geothermal and volcanic sites [14–17]. Geothermal fluids leach out of the hot volcanic rocks and are enriched by enormous amounts of minerals and metals, including lithium, sulphur, boric acid, and precious metals such as gold, platinum, palladium, and silver [18].

Cyanidiophyceae, unicellular red algae, survive in extreme conditions, very low pH (0.0–3.0) and high temperatures (37–55 °C), and colonize acid and hydrothermal sites but also rocks and muddy soil around hot ponds [19]. They are divided into 3 genera: *Cyanidioschyzon*, *Cyanidium* and *Galdieria*, which differ in size, cellular shape, and growth conditions. *Cyanidioschyzon merolae*, the only species belonging to the *Cyanidioschyzon*


genus, differs from the other two taxa in the lack of a cell wall and division by binary fission [20]. *Cyanidioschyzon*, *Cyanidium* and *Galdieria* can grow both on ammonia and nitrate. *Cyanidioschyzon* and *Cyanidium* species are obligatory autotrophs, while *Galdieria* ones can grow auto-, mixo-, and heterotrophically and tolerate high concentrations of salts [21], thus making *Galdieria* more suitable for biotechnological applications [14]. The ability of *Galdieria sulphuraria* of recovering REEs was already assessed [5,22] and confirmed by an approved patent [23]. In this report, we focused in deep for the first time on the ability of different *Galdieria* species (*Galdieria maxima*, *Galdieria sulphuraria*, *Galdieria phlegrea*) and *Cyanidium caldarium* to tolerate different concentrations of REEs (such as palladium-Cl<sub>4</sub>K<sub>2</sub>Pd and gold-AuCl<sub>4</sub>K). We also investigated the metabolic response and possible oxidative stress induced by these metals by monitoring superoxide dismutase (SOD), catalase (CAT), ascorbate peroxidase (APX) activities.

## 2. Results


Polyextremophilic microalgae, such as Cyanidiophyceae, have a high intrinsic capacity to uptake metals, involving active and passive mechanisms [5]. Heavy, rare, or precious metals can influence algae physiology in various ways, likely inhibiting different physiological processes. To evaluate the suitability of Cyanidiophyceae other than *G. sulphuraria* for biotechnological application to recover REEs effectively, we tested the tolerance to Cl<sub>4</sub>K<sub>2</sub>Pd and AuCl<sub>4</sub>K by monitoring the growth and metabolic response of 4 different taxa, exposed to each of these metals at a concentration in the range 1–10 g/L. As more deeply discussed in Section 4, the growth was evaluated after 4 days (96 h) since the single metal exposure. The results were expressed in the form of Maximum Growth Rate (MGR).

As shown in Figure 1, the presence of AuCl<sub>4</sub>K significantly reduced cellular duplication in *G. maxima* at all the concentrations tested; Cl<sub>4</sub>K<sub>2</sub>Pd did not negatively affect cell growth, and no statistical difference was recorded between MGR in control and tests (Figure 1A). Regarding *G. phlegrea*, both metals induced a trend of reduction in growth rate at both concentrations (Figure 1B). Viceversa, in *G. sulphuraria*, AuCl<sub>4</sub>K reduced cell growth at the maximum concentration, while the MGR appeared not affected by Cl<sub>4</sub>K<sub>2</sub>Pd, as shown by the MGR values at 10 g/L comparable to control. A decrease in growth rate was recorded at lower concentration (1 g/L); presumably, the highest amount of palladium was beneficial for the growth of this strain, or even the improvement of the cell duplication should be interpreted as a defence of the algal strain. Finally, *C. caldarium* showed a high tolerance to Cl<sub>4</sub>K<sub>2</sub>Pd, whereas AuCl<sub>4</sub>K significantly inhibited cell duplication as metal concentration increased (Figure 1D). The highest concentration of palladium (10 g/L) improved the growth, and in *G. maxima* and *G. sulphuraria*, the MGR values outperformed the controls. Subsequently, it was decided to evaluate the ROS scavenging activities of SOD, CAT and APX in all algae tested in the presence of Cl<sub>4</sub>K<sub>2</sub>Pd and AuCl<sub>4</sub>K at a concentration of 1 g/L, after an incubation period of 24 h. The reason behind this choice was that the antioxidant activity could be considered a measure of the cell effectiveness in responding to the impact of metals, increasing their tolerance as a protective mechanism necessary to remove ROS before they can damage sensitive parts of the cellular machinery. In particular, the SOD, which catalyses the dismutation of O<sub>2</sub><sup>-</sup> (singlet oxygen) to O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub>, was defined as the first cellular defence against ROS production. Meanwhile, CAT catalyses the production of H<sub>2</sub>O from the degradation of H<sub>2</sub>O<sub>2</sub> and ROOH, respectively. Finally, APX reduces H<sub>2</sub>O<sub>2</sub> to H<sub>2</sub>O using the ascorbate as an electron donor. The strain/metal-specific metabolic responses were quite diverse, as shown in Figure 2. Indeed, APX activity significantly increased only in *G. maxima* in response to Cl<sub>4</sub>K<sub>2</sub>Pd, while in the presence of AuCl<sub>4</sub>K, all the enzymatic activities appeared reduced (Figure 2A). Concerning the other strains, in *G. phlegrea*, all the enzymes tested activity decreased in the presence of Cl<sub>4</sub>K<sub>2</sub>Pd and increased in the presence of AuCl<sub>4</sub>K (Figure 2B); in *G. sulphuraria*, both metals induced an

enzymatic activity decrease (Figure 2C). Finally, we observed a significant increase of all the enzymes in *C. caldarium* in the presence of  $\text{Cl}_4\text{K}_2\text{Pd}$ , but not in  $\text{AuCl}_4\text{K}$  (Figure 2D).



**Figure 1.** Evaluation of metal tolerance through MGR monitoring after 4 days (96 h). Maximum growth rate in presence of different concentrations of palladium ( $\text{Cl}_4\text{K}_2\text{Pd}$ -orange-left panel) and gold ( $\text{AuCl}_4\text{K}$ -blue-right panel), for the species *G. maxima* (A), *G. phlegrea* (B), *G. sulphuraria* (C) and *C. caldarium* (D). Error bars represent standard deviation of three replicates; (\*) =  $p$ -value  $\leq 0.000000001$  calculated by  $T$ -test.



**Figure 2.** Evaluation of enzymatic activities after metals exposure. Relative units represent: enzymatic activity as units/g of dry weight (SOD); enzymatic activity as nmol H<sub>2</sub>O<sub>2</sub>/g of fresh weight (CAT); enzymatic activity as μmol ascorbate/g of fresh weight (APX). Enzymatic activities were monitored in *G. maxima* (A), *G. phlegrea* (B), *G. sulphuraria* (C) and *C. caldarium* (D), treated with 1 g/L of palladium (Cl<sub>4</sub>K<sub>2</sub>Pd—orange bars) and gold (AuCl<sub>4</sub>K—blue bars) after 96 h. Mean (± SD) was calculated from three replicates. (\*) = *p*-value ≤ 0.05 calculated by *T*-test.

### 3. Discussion

A significant increase of the enzyme compared to the control suggests a high scavenging activity of the singlet oxygen in peroxide of hydrogen, which can be expressed as an evident tolerance of these algae to the metal under examination. An increase in the activity of both antioxidant enzymes is necessary to reduce the concentrations of both singlet oxygen and hydrogen peroxide, minimizing the risks. In general, the modulation of antioxidant enzymes is an essential adaptive response to counteract adverse conditions; in fact, maintaining a high antioxidant capacity in the cells can be correlated with increased tolerance against different types of environmental stress [24].

Our results indicate that rare and precious metals can be tolerated by all the strains tested, even if there is a clear higher tolerance to  $\text{Cl}_4\text{K}_2\text{Pd}$  vs.  $\text{AuCl}_4\text{K}$  considering growth rates. Comparing the growth rate in the presence of the different concentrations of the metals, it seems clear that the growth and the metabolism of *G. phlegrea* appear to be more affected by the presence of both metals, showing a decrease of both growth and metabolic responses. The contribution to the oxidative equilibrium of the examined extremophile microalgae and the induction of antioxidant enzymes could result from the adaptation of the cell to the development of intracellular ROS. However, there is no clear correlation between any enzymatic activity and the better performing growth of the other 3 strains tested.

Although metals generally induce inhibition in microalgal growth, several reports also suggest their positive roles. It is known that metals at small concentrations are useful for microalgal metabolism since they participate in the synthesis of proteins involved in photosynthesis, nitrogen assimilation, phosphorous acquisition,  $\text{CO}_2$  fixation and DNA transcription [25]. Algae can develop efficient defence mechanisms to counteract the toxicity and improve their survival, even at high metal concentrations [26]. One of the defence strategies is the accumulation of the metals, which consists of the metal adsorption on the cell surface (biosorption), followed by their entry into the cell protoplast (bioaccumulation). When metals are accumulated inside the cell, the algae activate molecular mechanisms as other defence strategies to reduce their toxicity [26]. *G. sulphuraria* can survive in harsh environments rich in heavy and rare metals by detoxifying and transforming them into less toxic derivatives [27]. The defence strategies developed by algae to prevent the toxic effect of some metals represent a good opportunity for biotechnological purposes. The study from Ju et al. (2016) showed the ability of *G. sulphuraria* to recover both  $\text{Cl}_4\text{K}_2\text{Pd}$  and  $\text{AuCl}_4\text{K}$  inefficiently [5]. However, the authors did not test this strain's tolerance to growth in the presence of these metals. In contrast, we consider that tolerance and growth capacity is an essential parameter to take in account for the biotechnological application, such as REEs recovering.

### 4. Materials and Methods

#### 4.1. Strains Cultivation

The algal strains used in this study belong to the algal collection of the University of Campania "L. Vanvitelli" derived from the University of Naples ([www.acuf.net](http://www.acuf.net)), namely ACUF 3.4.5 (*G. maxima*), ACUF 7.6.21 (*G. phlegrea*), ACUF 9.2.11 (*G. sulphuraria*) and ACUF 626 (*C. caldarium*). All the strains were inoculated in Allen medium containing  $(\text{NH}_4)_2\text{SO}_4$  as nitrogen source, at pH 1.5 by adding  $\text{H}_2\text{SO}_4$  [28] and cultivated at 37 °C, kept mixed on an orbital shaker under a photon irradiance of 150  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  with 16/8-light/dark cycle provided by cool-light fluorescent lamps (Philips TLD30w/55). Cell densities of the algal cultures were assessed, recording the Optical Density (OD) at 750 nm with a spectrophotometer (Bausch & Lomb Spectronic 20).

#### 4.2. Experimental Procedure

Microalgal cultures at exponential phase were inoculated into fresh Allen medium enriched with  $\text{Cl}_4\text{K}_2\text{Pd}$  and  $\text{AuCl}_4\text{K}$  at concentrations ranging from 1 to 10 g/L. Growth

rates were calculated within 96h using spectrophotometric measurements of the optical density (OD 550 nm, Bausch & Lomb Spectronic 20), which were then used in the following equation for the Maximum Growth Rate (MGR):

$$\text{MGR (1/d)} = (\text{Ln}(N_t) - \text{Ln}(N_0)) / ((t - t_0))$$

where:

$N_t$  is the optical density at the final time

$N_0$  is the optical density at the initial time

$T$  is the final time (days)

$T_0$  is the initial time (days)

All analyses were performed in triplicates.

#### 4.3. Enzyme Extraction and Assays

Algal cultures grown in the presence of the minimal dose of palladium and gold (1 g/L) were harvested by centrifugation at 14,000 rpm for 10 min after 96 h of exposure. The algal pellets were washed using  $\text{KH}_2\text{PO}_4$  (0.1 M pH7.8) followed by centrifugation at 12,000 rpm for 4 min at 4 °C, twice. Proteins were extracted, homogenizing the sample with liquid nitrogen using a mortar and a pestle. The obtained powder was resuspended in 3 mL of Lysis Buffer ( $\text{KH}_2\text{PO}_4$  0.5 M pH7.8, DTT 2 mM, EDTA 1 mM, PMSF 1 mM, PEG 1.25 mM) and centrifuged at 14,000rpm for 20 min at 4 °C. The supernatant was used for measurement after Bradford quantification.

SOD (EC 1.15.1.1) activity was assayed by the photochemical inhibition nitroblue tetrazolium (NBT) method [6]. The reaction mixture contained 50 mM sodium phosphate buffer (pH 7.8), 13 mM methionine, 75 mM NBT, 0.1 mM EDTA and 30  $\mu\text{L}$  of enzyme extract and 2 mM riboflavin. The reaction was started by switching on the light (two 15 W fluorescent lamps) for 15 min, and the absorbance was measured at 560 nm. Two samples without the enzymatic extract and illumination were used as controls. One SOD unit was defined as the amount of enzyme corresponding to 50% inhibition of the NBT reduction. The enzyme activity was expressed as units per 1 mg of protein ( $\text{U mg}^{-1}$  protein).

CAT (EC 1.11.1.6) activity was assayed according to Aebi (1984) [29], with minor modifications. The  $\text{H}_2\text{O}_2$  decrease was determined after the reaction of the extract in the presence of 50 mM potassium phosphate buffer (pH 7.0) containing 20 mM  $\text{H}_2\text{O}_2$ . The reaction was monitored, measuring the decrease in the absorbance at 240 nm for 100 s. The CAT activity was calculated according to the molar extinction coefficient of  $\text{H}_2\text{O}_2$  ( $39.4 \text{ mM}^{-1} \text{ cm}^{-1}$ ) and expressed as  $\text{nmol H}_2\text{O}_2 \text{ min}^{-1} \text{ mg}^{-1}$  protein.

APX (EC 1.11.1.1) activity was assayed according to Nakano and Asada (1981) [30]. The ascorbate oxidation was determined using the reaction mixture containing 50 mM potassium phosphate buffer (pH 7.0), 0.1 mM EDTA- $\text{Na}_2$ , 0.5 mM ascorbic acid and 100  $\mu\text{L}$  of crude enzyme extract. The reaction started by adding 0.1 mM  $\text{H}_2\text{O}_2$ , monitoring the decreasing absorbance at 290 nm for 100 s. The APX activity was calculated according to the molar extinction coefficient of ascorbate ( $2.8 \text{ mM}^{-1} \text{ cm}^{-1}$ ) and is expressed as  $\text{nmol di H}_2\text{O}_2 \text{ min}^{-1} \text{ mg}^{-1}$  protein.

Each condition for each experimental approach was tested 3 times independently.

## 5. Conclusions

Our observations strongly suggest that other strains than *G. sulphuraria* can be used to recover REEs due to their high tolerance to precious and heavy metals. Nevertheless, further studies will be necessary to clarify the biological mechanisms underlying the tolerance capacity of *Cyanidiophyceae* and their strategies to respond to metal toxicity for future biotechnological applications.

**Author Contributions:** Conceptualization, M.S. and M.I.; Formal analysis, M.P. (Maria Palmieri), M.P. (Milena Petriccione) and M.R.D.C.; Methodology, M.P. (Maria Palmieri), M.P. (Milena Petriccione) and M.R.D.C.; Supervision, C.C.; Visualization, S.J.D., M.D.S. and C.C.; Writing—original draft, M.S. and M.I.; Writing—review & editing, M.S. and M.I. All authors have read and agreed to the published version of the manuscript.

**Funding:**

**Institutional Review Board Statement:**

**Informed Consent Statement:**

**Data Availability Statement:**

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Aebi H. Catalase in vitro. *Method Enzym* **1984** *105*,121–126
2. Albertano P., Ciniglia C., Pinto G., Pollio A. The taxonomic position of *Cyanidium*, *Cyanidioschyzon* and *Galdieria*: an update. *Hydrobiologia* **2000**; *433*,137-143.
3. Allen M.M. & Stanier RY. Selective isolation of blue-green algae from water and soil *J Gen Microbiol* **1968**; *51*, 203-209
4. Arshadi M, Mousavi SM, Rasoulnia P. Enhancement of simultaneous gold and copper recovery from discarded mobile phone PCBs using *Bacillus megaterium*: RSM based optimization of effective factors and evaluation of their interactions. *Waste Management* **2016** *57*, 158-167
5. Awasthi AK & Li J. An overview of the potential of eco-friendly hybrid strategy for metal recycling from WEEE. *Resources, Conservation & Recycling* **2017**, *126*, 228-239
6. Beauchamp C, Fridovich I Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Anal Biochem Review* **1971**, *44*, 76–287
7. Bourcier WL, Lin M, Nix G. Recovery of Minerals and Metals from Geothermal Fluids. In: SME Annual Meeting Cincinnati, OH, United States February 24, 2003 UCRL-CONF-2005; 215135.
8. De Luca P, Taddei R & Varano L. *Cyanidioschyzon merolae*: a new alga of thermal acidic environments *Webbia* **1978**, *33*(1), 37-44
9. Doemel W, & Brock ML. The physiological ecology of *Cyanidium caldarium* *J Gen Microbiol* **1971**, *67*, 17-32
10. Güzel Y, Rainer M, Mirza MR, Messner CB, Bonn GK. Highly selective recovery of phosphopeptides using trypsin-assisted digestion of precipitated lanthanide-phosphoprotein complexes *Analyst* **2013**, *138*, 2897-2905
11. Homosomi Y, Baba Y, Kubota F, Kamiya N, Goto M. Biosorption of rare earth elements by *Escherichia coli* *J.Chem Eng Jpn* **2013**, *46*:450-454
12. Jiang M, Ohnuki T, Kozai NM, Tanaka K, Suzuki Y, Sakamoto F, Kamiishi E, Utusnomiya S. The Biological nano-mineralization of Ce phosphate by *Saccharomyces cerevisiae* *Chem Geol* **2010**, *277*, 61-9
13. Ju X, Igarashi K, Miyashita S, Mitsuhashi H, Inagaki K, Fujii S, Sawada H, Kuwabara T, Minoda A. [Effective and selective recovery of gold and palladium ions from metal wastewater using a sulfothermophilic red alga, \*Galdieria sulphuraria\*](#). *Bioresour Technol* **2016**, *211*, 759-64.
14. Kuroda K, Ueda M. Engineering of microorganisms towards recovery of rare metal ions *Appl Microbiol Biotechnol* **2010**, *87*, 53-60
15. Latorre M, Cortés MP, Travisany D, Di Genova A, Budinich M, Reyes-Jara A, Hödar C, González M, Parada P, Roberto A, Bobadilla-Fazzini, Combiazo V, Maass A. The bioleaching potential of a bacterial consortium. *Biores Technol* **2016**, *218*, 659-666
16. Matsunaga T, Takeyama H, Nakao T, Yamazawa A. Screening of marine microalgae for bioremediation of cadmium-polluted seawater *J Biotechnol* **1999** *70*, 33-38
17. Mehta SK, Gaur JP. Use of algae for removing heavy metal ions from wastewater: progress and prospects. *Crit Rev Biotechnol* **2005**; *25*, 113-152
18. Minoda A & Chuo-ku. Agent for selective metal recovery method, and metal elution method. Patent Application Publication, US 2019/0024209 A1, **2019**
19. Minoda A, Sawada H, Suzuki S, Miyashita S, Inagaki K, Yamamoto T, Tsuzuki M. . Recovery of rare earth elements from the sulfothermophilic red alga *Galdieria sulphuraria* using aqueous acid. *Appl Microbiol Biotechnol* **2015** *99*(3),1513-9
20. Nakano Y, Asada K Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiol* **1981** *22*, 867–880
21. Okamoto OK, Pinto E, L., Latorre R, Bechara EJH, Colepicolo P. Antioxidant modulation in response to metal-induced oxidative stress in algal chloroplasts. *Arch Environ Contam Toxicol* **2001** *40*, 18-24
22. Sharma SS, Dietz KJ, Mimura T. Vacuolar compartmentalization as indispensable component of heavy metal detoxification in plants. *Plant, cell & environment* **39** (5), 1112-1126, 2016

23. Sirakov M, Toscano E, Iovinella M, Davis S, Petriccione M, Ciniglia C. Tolerance and metabolic responses of Cyanidiphytina (Rhodophyta) towards exposition to  $\text{Cl}_4\text{K}_2\text{Pd}$  and  $\text{AuCl}_4\text{K}$ . *Phycol int* **2018** 1 (1)
24. Wang J, Chen C. Biosorbents for heavy metal removal and their future *Biotechnol Adv* **2009** 27, 195-226
25. Wang M, Tan Q, Chiang F, et al. Recovery of rare and precious metals from urban mines—A review[J]. *Front. Environ. Sci. Eng.*, **2017**, 11(5): 1.

Zhang L, Dong H, Liu Y, Bian L, Wang X, Zhou Z, Huang Y. Bioleaching of rare earth elements from bastnaesite-bearing rock by actinobacteria. *Chem Geol* **2018** 483 544-557

1. Di Cicco, M.R.; Spagnuolo, A.; Masiello, A.; Vetromile, C.; Nappa, M.; Lubritto, C. Energetic and environmental analysis of a wastewater treatment plant through static and dynamic monitoring activities. *Int. J. Environ. Sci. Technol.* **2020**, 17, 4299–4312.
2. Yüksel Güzel; Matthias Rainer; R. Mirza, M.; B. Messner, C.; K. Bonn, G. Highly selective recovery of phosphopeptides using trypsin-assisted digestion of precipitated lanthanide–phosphoprotein complexes. *Analyst* **2013**, 138, 2897–2905.
3. HOSOMOMI, Y.; BABA, Y.; KUBOTA, F.; KAMIYA, N.; GOTO, M. Biosorption of Rare Earth Elements by *E. coli*. *J. Chem. Eng. JAPAN* **2013**, 46, 13we031.
4. Jiang, M.; Ohnuki, T.; Kozai, N.; Tanaka, K.; Suzuki, Y.; Sakamoto, F.; Kamiishi, E.; Utsunomiya, S. Biological nano-mineralization of Ce phosphate by *Saccharomyces cerevisiae*. *Chem. Geol.* **2010**, 277, 61–69.
5. Ju, X.; Igarashi, K.; Miyashita, S. ichi; Mitsushashi, H.; Inagaki, K.; Fujii, S. ichiro; Sawada, H.; Kuwabara, T.; Minoda, A. Effective and selective recovery of gold and palladium ions from metal wastewater using a sulfophilic red alga, *Galdieria sulphuraria*. *Bioresour. Technol.* **2016**, 211, 759–764.
6. Kuroda, K.; Ueda, M. Engineering of microorganisms towards recovery of rare metal ions. *Appl. Microbiol. Biotechnol.* **2010**, 87, 53–60.
7. di Cicco, M.R.; Spagnuolo, A.; Masiello, A.; Vetromile, C.; Lubritto, C.; Nappa, M.; Corbo, G. Energy monitoring of a wastewater treatment plant in salerno, campania region (southern italy). In *Advances in Science, Technology and Innovation*; Springer, Cham, 2019; pp. 107–111.
8. Vetromile, C.; Spagnuolo, A.; Petraglia, A.; Masiello, A.; di Cicco, M.R.; Lubritto, C. Pre- and post-operam comparison of the energy consumption of a radio base station under energy efficiency actions. *Energy Build.* **2021**, 236, 110772.
9. Arshadi, M.; Mousavi, S.M.; Rasoulnia, P. Enhancement of simultaneous gold and copper recovery from discarded mobile phone PCBs using *Bacillus megaterium*: RSM based optimization of effective factors and evaluation of their interactions. *Waste Manag.* **2016**, 57, 158–167.
10. Latorre, M.; Cortés, M.P.; Travisany, D.; Di Genova, A.; Budinich, M.; Reyes-Jara, A.; Hödar, C.; González, M.; Parada, P.; Bobadilla-Fazzini, R.A.; et al. The bioleaching potential of a bacterial consortium. *Bioresour. Technol.* **2016**, 218, 659–666.
11. Wang, M.; Tan, Q.; Chiang, J.F.; Li, J. Recovery of rare and precious metals from urban mines—A review. *Front. Environ. Sci. Eng.* **2017**, 11, 1–17.

12. Zhang, L.; Dong, H.; Liu, Y.; Bian, L.; Wang, X.; Zhou, Z.; Huang, Y. Bioleaching of rare earth elements from bastnaesite-bearing rock by actinobacteria. *Chem. Geol.* **2018**, *483*, 544–557.
13. Sharma, S.S.; Dietz, K.J.; Mimura, T. Vacuolar compartmentalization as indispensable component of heavy metal detoxification in plants. *Plant Cell Environ.* **2016**, *39*, 1112–1126.
14. Cicco, M.R. di; Palmieri, M.; Altieri, S.; Ciniglia, C.; Lubritto, C. Cultivation of the Acidophilic Microalgae *Galdieria phlegrea* with Wastewater: Process Yields. *Int. J. Environ. Res. Public Heal.* **2021**, *Vol. 18*, Page 2291 **2021**, *18*, 2291.
15. Matsunaga, T.; Takeyama, H.; Nakao, T.; Yamazawa, A. Screening of marine microalgae for bioremediation of cadmium-polluted seawater. *J. Biotechnol.* **1999**, *70*, 33–38.
16. Mehta, S.K.; Tripathi, B.N.; Gaur, J.P. Enhanced sorption of Cu<sup>2+</sup> and Ni<sup>2+</sup> by acid-pretreated *Chlorella vulgaris* from single and binary metal solutions. *J. Appl. Phycol.* **2002**, *14*, 267–273.
17. Wang, J.; Chen, C. Biosorbents for heavy metals removal and their future. *Biotechnol. Adv.* **2009**, *27*, 195–226.
18. Bourcier, W.L.; Lin, M.; Nix, G. Recovery of Minerals and Metals from Geothermal Fluids. *2003 SME Annu. Meet.* **2005**, 19.
19. Doemel, W.N.; Brock, T.D. The Physiological ecology of *Cyanidium caldarium*. *J. Gen. Microbiol.* **1971**, *67*, 17–32.
20. De Luca, P.; Taddei, R.; Varano, L. “*Cyanidioschyzon merolae*”: a new alga of thermal acidic environments. *Webbia* **1978**, *33*, 37–44.
21. Albertano, P.; Ciniglia, C.; Pinto, G.; Pollio, A. The taxonomic position of *Cyanidium*, *Cyanidioschyzon* and *Galdieria*: An update. *Hydrobiologia* **2000**, *433*, 137–143.
22. Minoda, A.; Sawada, H.; Suzuki, S.; Miyashita, S. ichi; Inagaki, K.; Yamamoto, T.; Tsuzuki, M. Recovery of rare earth elements from the sulfothermophilic red alga *Galdieria sulphuraria* using aqueous acid. *Appl. Microbiol. Biotechnol.* **2015**, *99*, 1513–1519.
23. Minoda, A. Agent for selective metal recovery, metal recovery method, and metal elution method 2019, 1–24.
24. Okamoto, O.K.; Pinto, E.; Latorre, L.R.; Bechara, E.J.H.; Colepicolo, P. Antioxidant Modulation in Response to Metal-Induced Oxidative Stress in Algal Chloroplasts. *Arch. Environ. Contam. Toxicol.* **2001**, *40*, 18–24.
25. Miazek, K.; Iwanek, W.; Remacle, C.; Richel, A.; Goffin, D. Effect of metals, metalloids and metallic nanoparticles on microalgae growth and industrial product biosynthesis: A review. *Int. J. Mol. Sci.* **2015**, *16*, 23929–23969.
26. Priyadarshini, E.; Priyadarshini, S.S.; Pradhan, N. Heavy metal resistance in algae and its application for metal nanoparticle synthesis. *Appl. Microbiol. Biotechnol.* **2019**, *103*, 3297–3316.

- 
27. Schönknecht, G.; Chen, W.; Ternes, C.M.; Barbier, G.G.; Shrestha, R.P.; Stanke, M.; Bräutigam, A.; Baker, B.J.; Banfield, J.F.; Garavito, R.M.; et al. Gene Transfer from Bacteria and Archaea Facilitated Evolution of an Extremophilic Eukaryote. *Science* (80-. ). **2013**, *339*, 1207–1210.
  28. Allen, M.B. Studies with cyanidium caldarium, an anomalously pigmented chlorophyte. *Arch. Mikrobiol.* **1959**, *32*, 270–277.
  29. Aebi, H. Catalase in Vitro. *Methods Enzymol.* **1984**, *105*, 121–126.
  30. Nakano, Y.; Asada, K. Hydrogen Peroxide is Scavenged by Ascorbate-specific Peroxidase in Spinach Chloroplasts. *Plant Cell Physiol.* **1981**, *22*, 867–880.