

This is a repository copy of *Linear gyrokinetic stability of a high β non-inductive spherical tokamak*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/179896/>

Version: Accepted Version

Article:

Patel, Bhavin, Dickinson, David orcid.org/0000-0002-0868-211X, Roach, C. M. et al. (1 more author) (2021) Linear gyrokinetic stability of a high β non-inductive spherical tokamak. Nuclear Fusion. 016009. ISSN: 1741-4326

<https://doi.org/10.1088/1741-4326/ac359c>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Linear gyrokinetic stability of a high β non-inductive spherical tokamak

B.S. Patel^{1,2}, D. Dickinson¹, C.M. Roach² and H.R. Wilson¹

¹ University of York, Heslington, York, YO10 5DD, UK

² Culham Centre for Fusion Energy, Abingdon OX14 3DB, UK

E-mail: bhavin.s.patel@ukaea.uk

March 2021

Abstract. Spherical tokamaks (STs) have been shown to possess properties desirable for a fusion power plant such as achieving high plasma β and having increased vertical stability. To understand the confinement properties that might be expected in the conceptual design for a high β ST fusion reactor, a 1GW ST plasma equilibrium was analysed using local linear gyrokinetics to determine the type of micro-instabilities that arise. Kinetic ballooning modes (KBMs) and micro-tearing modes (MTMs) are found to be the dominant instabilities. The parametric dependence of these linear modes was determined and from the insights gained, the equilibrium was tuned to find a regime marginally stable to all micro-instabilities at $\theta_0 = 0.0$. This work identifies the most important micro-instabilities expected to generate turbulent transport in high β STs. The impact of such modes must be faithfully captured in first principles based reduced models of anomalous transport that are needed for predictive simulations.

1. Introduction

Understanding the confinement of tokamak plasmas in reactor relevant regimes is critical in the design of future fusion power plants. Scaling laws, such as the ITER98 IPB(y,2) [1] and the Petty08 [2] are often used as a metric for the quality of confinement. Both of these laws describe the confinement of existing conventional tokamaks equally well but extrapolate quite differently to reactor relevant regimes. For example, for the ITER baseline scenario, the 98 scaling predicts $\tau_{98} = 3.7\text{s}$, but Petty predicts an increased value of $\tau_{\text{Petty}} = 4.6\text{s}$. Global scalings are fitted to data from existing tokamaks not in the reactor regime, and there are inevitably unknown uncertainties in their extrapolation, especially if the turbulent regime is different. Global scalings can be used for ballpark estimates in confinement, but reliable predictions require high fidelity models for the neoclassical and turbulent transport and should be based on first principle models. Gyrokinetics has proven to be a useful tool in modelling turbulence and its associated transport and has been able to match experimental fluxes with reasonable success [3–5]. Applying linear gyrokinetics to a high β ST regime will provide an insight into the turbulent modes that may arise. This work aims to identify the linear micro-stability

properties of a conceptual equilibrium for a high β burning ST reactor and highlight the necessary physics that needs to be captured in a transport model. This will be critical for the development of STEP [6].

A potential non-inductive operating point is identified in Section 2, where a plasma equilibrium has been obtained using SCENE [7], a fixed boundary Grad-Shafranov solver. The reasoning behind this equilibrium design is briefly discussed but further details can be found in [8]. In Section 3, the dominant instabilities of the $\rho_\psi \equiv \psi/\psi_{\text{LCFS}} = 0.5$ surface was determined using linear GS2 [9]. It was found that KBMs and MTMs were the prevalent modes at the ion scale illustrating the electromagnetic nature of the turbulence that needs to be captured in a transport model. At deep electron scales the equilibrium was found to be stable. Section 4 explores the impact of different equilibrium parameters, such as the kinetic gradients and plasma β , on these modes. With the insights gained here, Section 5 perturbs the global plasma parameters and finds a more stable equilibrium. An equilibrium marginally stable to all the micro-instabilities is found, which suggests that operating at neoclassical levels of transport and thus high levels of confinement, may be possible. The feasibility of achieving such an equilibrium is briefly discussed. Appendix A compares the linear predictions made by CGYRO and GS2 for this equilibrium showing them to be in good agreement and Appendix B explores the cause of stability in the deep electron scale region.

2. 1GW ST operating point

This plasma equilibrium was chosen to produce a fusion power over 1GW as this was assumed to be sufficient to produce net electricity. In SCENE several global plasma parameters are specified such as the current going through the central column (which sets the vacuum magnetic field), the total toroidal plasma current and the plasma boundary. Furthermore, the temperature and density profiles are specified, though no transport models inform these profiles. The feasibility of the profiles could be tested by calculating the heat and particle sources required for a given transport model.

SCENE solves for the equilibrium whilst self-consistently calculating the neoclassical currents such as the bootstrap and diamagnetic current, returning the amount of auxiliary current needed to provide the total plasma current requested. The shape of the auxiliary current profile is prescribed and can be set to be consistent with the anticipated auxiliary current sources.

The major radius, R_{maj} , was set via a constraint on the power going through the separatrix P_{sep} . Assuming that 80% of the heating power is radiated away, similar to FNSF [10] and requiring that $P_{\text{sep}}/R_{\text{maj}} < 20\text{MW m}^{-1}$, similar to the ITER [11], $R_{\text{maj}} = 2.5\text{m}$ was set. The spherical tokamak power plant (STPP) is an existing design of a high β ST [12]. Using its centre column design [13] which had a rod current, $I_{\text{rod}} = 30\text{MA}$, generates 2.4T on axis for this design. To allow for sufficient space for this centre column the minor radius a was set to 1.5m, resulting in an aspect ratio of




Figure 1: Profiles for the baseline equilibrium showing the a) safety factor, b) electron temperature where $T_e = T_i$ and c) electron density. d) The flux surface contours for the baseline equilibrium.

$A = 1.67$.

With the plasma current set to $I_p = 21\text{MA}$, the prescribed pressure leads to $\beta_N = 5.5$, which satisfies MHD stability constraints with the q profile shown in Figure 1a. Note, that this was designed with $q_{\min} > 2$ to avoid the 2/1 NTM and sawteeth modes. Increasing I_p further would increase the requirements on the auxiliary current drive system. The elongation was set to $\kappa = 2.8$, inline with results from NSTX [10] and the triangularity of $\delta = 0.55$ to match STPP. Using the Europed model [14], a temperature pedestal height of 5.3keV at $\rho_\psi = 0.92$ was found for utilising the assumption that the width of the pedestal scales like $0.1(\beta_\theta^{\text{ped}})^{1/2}$, consistent with MAST data [15]. This work set the temperature pedestal height to 5keV located at $\rho_\psi = 0.9$, which happens to be similar to the ITER pedestal [16]. The core temperature was assumed to be 28keV with the $T_e = T_i$. A small amount of density peaking was specified such that $n_{e,\text{ped}}/n_{e0} = 0.9$, similar to ITER. The density was then scaled up such that $P_{\text{fus}} > 1\text{GW}$, leading to a Greenwald fraction of $f_{\text{GW}} = 0.52$. These profiles are shown in Figures 1b and 1c. The

Parameter	Value	Motivation
R_{maj} (m)	2.5	Power through the separatrix
a (m)	1.5	Allow space for centre column
R_0 (m)	3.15	Magnetic axis position - output from SCENE
I_{rod} (MA)	30.0	STPP centre column
I_p (MA)	21.0	Trade-off between confinement/stability and current drive
I_{aux} (MA)	8.2	Output from SCENE
P_{fus} (MW)	1100	Net electric device
P_{aux} (MW)	94	NUBEAM
κ	2.8	Limits based off of NSTX data
δ	0.55	Same as STPP
H_{98}, H_{Petty}	1.35, 0.94	Output given required P_{aux}
$T_{e0}, \langle T_e \rangle$ (keV)	28.0, 14.8	Assumption
$n_{e0}, \langle n_e \rangle (\times 10^{20} \text{m}^{-3})$	1.72, 1.54	Ensure $P_{\text{fus}} = 1.1\text{GW}$ given T_e assumption
f_{GW}	0.52	Ensure $P_{\text{fus}} = 1.1\text{GW}$ given T_e assumption
l_i	0.27	Maximise elongation
β_N	5.5	Output from SCENE - MHD stable
q_0	2.51	Avoid sawteeth/NTM/internal kink modes

Table 1: Basic global plasma parameters for this baseline operating point and the reasoning behind them.

resulting equilibrium is outlined in Table 1, with Figure 1d illustrating the flux surfaces.

The auxiliary current profile was set to be primarily off axis which reduced the internal inductance l_i to help with vertical stability, with a small amount necessary on axis to fill the hole in the bootstrap current. From Table 1, it can be seen that 8.2MA of auxiliary current was necessary for a non-inductive scenario. Using NUBEAM [17] an NBI configuration was identified capable of driving the required current. For the on axis current, an 8MW 1MeV beam was necessary and 86MW 500keV for the off axis current. This totals to an auxiliary power of $P_{\text{aux}} = 94\text{MW}$, which requires a $H_{98} = 1.35$ and $H_{\text{Petty}} = 0.94$. This H_{98} has been seen before in STs [18] and is very close to the Petty prediction suggesting that this is a reasonable starting point. This equilibrium is further described in [8].


2.1. Local equilibrium for micro-stability analysis

To begin with the dominant micro-instabilities will be examined using local linear gyrokinetics on a core flux surface where $\rho_\psi = 0.5$. Once the nature of the modes is identified, the parametric dependence of these modes will be determined which, in turn, will inform design choices that will help stabilise the equilibrium.


A Miller parameterisation [19] was used to model this equilibrium, with the parameters outlined in Table 2. The fit of the flux surface and poloidal field are shown

Parameter	$\rho_\psi = 0.5$
r/a	0.66
R_{maj}/a	1.79
$n_e (\times 10^{20} \text{m}^{-3})$	1.51
T_e (keV)	12.2
B_0 (T)	2.16
B_{unit} (T)	7.52
ρ_s (m)	0.0021
a/L_n	0.43
a/L_T	2.77
Δ	-0.57
q	4.30
\hat{s}	0.78
κ	3.03
s_κ	-0.14
δ	0.45
s_δ	0.19
β_e	0.15
$\beta_{e,\text{unit}}$	0.012
$\nu_{ee}(c_s/a)$	0.017
$\gamma_{\text{dia}}(c_s/a)$	0.08

Table 2: Local plasma and Miller parameters for the $\rho_\psi = 0.5$ surface of the equilibrium in Table 1.



(a)



(b)

Figure 2: Miller fit shown for $\rho_\psi = 0.5$ surface showing the a) flux surface contour and b) the poloidal field

in Figures 2a and 2b respectively, showing that Miller fit to the local equilibrium is reasonable. Several simulations were conducted using the full numerical equilibrium and it was found to agree with the Miller simulations.

2.1.1. Numerical set-up The gyrokinetics code GS2 was used to examine the micro-stability of this plasma [9]. Convergence tests in GS2 indicated that 128 poloidal (θ) grid

points, 8 energy grid points and 16 un-trapped grid points in pitch angle \dagger was sufficient to resolve the modes seen here. Two species were simulated, a thermal deuterium and electron species. It should be highlighted that no fast ions were included in this work, but must be examined in future work as they may have a significant impact on the micro-stability [20, 21]. Furthermore, the impact of including tritium and impurities is also left as future work.

Although there will be neutral beam injection, the driven rotation is expected to be low due to the high energy beams [8]. However, there will be a contribution to the $E \times B$ shearing rate from the pressure gradient. This diamagnetic flow shear, γ_{dia} , can be calculated from force balance. In this work Equation A10 from [22] is used to calculate γ_{dia} , shown in Equation 1.

$$\gamma_{\text{dia}} = \frac{1}{B} \left(\frac{\partial \psi}{\partial \rho} \right)^2 \left(\frac{1}{pn_i e (1 + \eta_i)} \left(\frac{\partial p}{\partial \psi} \right)^2 - \frac{1}{n_i e} \frac{\partial^2 p}{\partial \psi^2} \right) \quad (1)$$

The normalisations used here was that from CGYRO [23] where the bi-normal wavenumber $k_y = \frac{nq}{r}$, where n is the toroidal mode number and r is the minor radius of the flux surface. k_y is normalised to Larmor radius $\rho_s = \frac{c_s}{eB_{\text{unit}}/m_D c}$. Here $c_s = \sqrt{\frac{T_e}{m_D}}$ and $B_{\text{unit}} = \frac{q}{r} \frac{\partial \psi}{\partial r}$. This also leads to a different definition of normalised β where $\beta_{e,\text{unit}} = \frac{n_e T_e}{B_{\text{unit}}^2 / 2\mu_0}$. It should be noted that this isn't the normalisation used in GS2, but was chosen to allow for easier comparison to CGYRO in Appendix A. The traditional normalising field is the $B_0 = f/R_{\text{maj}}$ with $\beta_e = \frac{n_e T_e}{B_0^2 / 2\mu_0}$.

3. Identifying the important instabilities

3.1. Dominant instabilities at $\rho_\psi = 0.5$

This section will examine the unstable modes at $\theta_0 = 0.0$. Due to the up-down symmetry of the equilibrium, it is possible to force an odd or even parity ϕ eigenfunction in GS2. This allows for usage as a pseudo-eigensolver to determine the dominant odd and even eigenmode. Conventionally, even parity ϕ modes are called “twisting parity” and odd parity are called “tearing parity”. However, an odd parity ϕ eigenfunction doesn't guarantee that the mode will tear the field line, so in this work we will exclusively refer to the parity as even or odd. For all the figures in this work, a hollow data point will correspond to an even parity ϕ eigenfunctions and a filled data point to an odd parity mode. The figure markers for each type of mode will be kept consistent throughout this work, to easier allow for identification of the different modes.

The dominant odd and even parity instabilities of the $\rho_\psi = 0.5$ surface at an initial ballooning angle of $\theta_0 = 0$ are shown in Figure 3. Three different types of modes were found; in the very long wavelength region when $k_y \rho_s < 0.6$, micro-tearing modes (MTMs) were found which will be illustrated by filled orange triangles. At slightly shorter wavelengths, but still at the ion scale where $0.1 < k_y \rho_s < 2.0$, kinetic ballooning

\dagger In GS2 the number of trapped grid points in pitch angle is given by $n_\theta/2 + 1$.

modes (KBMs) were found, denoted by hollow blue circles. Finally as the electron scale is approached between $3.0 < k_y \rho_s < 6.0$, another MTM is found shown by filled green squares, though this will be shown to be different in nature to the low k_y MTMs. The eigenfunctions of these modes are shown in Figure 4.

From Figure 3, it can be seen that dominant mode transitions from the MTMs to the KBMs, with the MTMs being subdominant above $k_y \rho_s = 0.14$. Co-existing MTMs and KBMs have been seen before in simulations of JET, DIII-D, MAST and NSTX plasmas [21, 24–27]. Nonlinear simulations will be necessary to determine their relative contributions to the total turbulent transport.




Figure 3: Growth rate and frequencies of the dominant odd and even parity micro-instabilities as a function of $k_y \rho_s$ with $\theta_0 = 0$ for the $\rho_\psi = 0.5$ surface of the baseline equilibrium calculated by GS2. The hollow marker denotes an even parity eigenfunction and a filled marker an odd eigenfunction. The figure markers for each type of mode will be kept consistent throughout this work, to easier allow for identification of different modes. Note the log scale.

3.1.1. Low k_y MTM In this work an MTM will refer to any mode with a group velocity in the electron diamagnetic flow direction that has field line tearing. MTMs are generally identified with an even parity $A_{||}$ eigenfunction which is symmetric about $\theta = 0$ (when $\theta_0 = 0.0$) with ϕ being odd. However this does not guarantee field line tearing and a more precise definition can be used that quantifies this. A mode is tearing if the perturbation results in a field line that does not return to the equilibrium flux surface.

This can be characterised using the following equation [28–30]

$$C_{\text{tear}} = \frac{|\int A_{\parallel} dl|}{\int |A_{\parallel}| dl} \quad (2)$$

which $\int dl$ corresponds to an integral along the field line. $C_{\text{tear}} > 0$ corresponds to some tearing of the field line. This low k_y MTM has a $C_{\text{tear}} = 0.7$ indicating that this mode tears the equilibrium magnetic flux surfaces.

There are several mechanisms that can drive an MTM. The first is from a parallel thermal force arising from the different frictional forces experienced by electrons travelling in opposite directions along a temperature gradient. This generates a parallel current which in turn generates a perturbation in A_{\parallel} [31]. It was found that this mechanism could be examined in different limits and in the collisionless regime ($\nu_{ee} < \omega$) where this time dependent thermal force vanishes [32]. For this equilibrium, $\nu_{ee} = 0.017c_s/a$, which is well below the mode frequency where $\omega \sim 0.5c_s/a$, suggesting this mechanism should not be relevant.

Another mechanism proposed by Catto & Rosenbluth is where electrons close to the trapped-passing boundary can easily scatter across it, which increases the effective collisionality allowing for a destabilising current driving the tearing instability. This is valid when $\nu_{ee} < \epsilon\omega$ [33] where $\epsilon = a/R_{\text{maj}}$ is the inverse aspect ratio. This condition is satisfied in this collisionality regime.

Both of these mechanisms require a finite collisionality. However, local gyrokinetic simulations have found MTMs not described by these two mechanisms [34, 35].

In an MTM, the perturbed A_{\parallel} corresponds to a magnetic island forming at the rational surface. Particles stream freely along the perturbed field and undergo radial excursions about the rational surface. Radial transport is enhanced at large amplitudes when the islands on neighbouring rational surfaces overlap. The higher velocity of the electrons results in significant electron heat transport and is given as a potential reason for the electron heat transport often being found to dominate over ion heat transport in STs [36–38].

The eigenfunctions of the MTM at $k_y\rho_s = 0.1$ is shown in Figure 4a. The eigenfunctions have been normalised to the maximum value of ϕ , indicating that the A_{\parallel} fluctuation is significantly larger than the electrostatic fluctuations at $\theta = 0$. Furthermore, these MTMs were found unstable when B_{\parallel} fluctuations were excluded which is typical for these modes. The eigenfunction at $k_y\rho_s = 0.35$ is shown in Figure 4b and it is significantly less extended in the parallel direction compared to the $k_y\rho_s = 0.1$ mode shown in Figure 4a, as would be expected if $k_{\parallel} \propto k_y$.

The eigenfunctions exhibit two scales at work here, a broad oscillation in θ , and a much narrower oscillation in θ which corresponds to a single poloidal revolution due to the equilibrium variation. In the A_{\parallel} eigenfunction, there is a central peak at $\theta = 0.0$ and then the other peaks along the field line occur at the same $\theta \bmod 2\pi$, at the top and bottom of the flux surface. This can be seen clearer in Figure 5d. These extended modes are generated by the parallel electron dynamics as the ions would not be able




Figure 4: Eigenfunctions for the different modes seen across the k_y spectrum for the $\rho_\psi = 0.5$ surface. The amplitude of each mode have been normalised with respect to the maximum value of ϕ

to travel that far down the field line in the mode period due to their lower velocity. The extended nature of the mode in ballooning space required a parallel domain from $-71\pi \rightarrow 71\pi$, corresponding to $k_x = k_y \hat{s} \theta = 26.1$. Here $\hat{s} = \frac{r}{q} \frac{\partial q}{\partial r}$. Even linearly resolving such modes becomes computationally expensive. Similar extended MTM eigenfunctions have been seen in simulations of MAST and NSTX discharges [22, 39].

Interestingly, these low k_y MTMs were unstable with only A_{\parallel} (both ϕ and B_{\parallel} turned off), adiabatic ions, and without contributions from the trapped particles. This indicates that the important physics lies within the passing electrons. Moreover, these MTMs required collisions for instability. Furthermore, if the ∇B and curvature drifts were turned off the mode went stable even if the drive for these MTMs was increased, suggesting it wouldn't exist in a slab geometry. This differentiates it from the MTMs derived in Drake *et al* [32] which assumed no toroidal effects, though Rafiq *et al* [40] performed a similar derivation including these effects.

3.1.2. Low k_y KBM The KBM eigenfunction, shown in Figure 4c, is significantly less extended compared to the low k_y MTM, only extending a single poloidal revolution. The $B_{||}$ perturbation has a significant amplitude and without $B_{||}$, these KBMs are not driven unstable which has been seen before in high β NSTX-like simulations [41]. The KBMs do not cause field line tearing as they had $C_{\text{tear}} = 0.0$. KBMs drive similar levels of electron and ion heat transport as well as significant particle transport differentiating them from MTMs [42].

3.1.3. High k_y MTM From $3.0 < k_y \rho_s < 7.0$, a higher $k_y \rho_s$ MTMs is seen with the eigenfunction at $k_y \rho_s = 4.2$ shown in Figure 4d. This mode is also tearing as they had $C_{\text{tear}} = 0.5$. These are much less extended than the MTMs seen at lower k_y . Again, $B_{||}$ has little effect on these modes.

These MTMs are fundamentally different to the low k_y MTMs as they will be shown to be unstable in the collisionless regime, which have been found previously [35, 43, 44], though the driving mechanisms are not completely understood.

Above $k_y \rho_s = 7.0$ all the modes were found to be stable which suggests that ETG-like modes may not be significant for this equilibria. The cause of the stability is examined in more detail in Appendix B.

3.2. Impact of flow shear

3.2.1. Low k_y modes Flow shear has been shown to stabilise turbulence by shearing the modes. Linearly this corresponds to making the radial wavenumber, k_x , of the modes time dependent. The effectiveness of flow shear stabilisation can thus be determined by examining how the stability varies with k_x and in GS2 this is done with the parameter $\theta_0 \equiv \frac{k_x}{k_y \bar{s}}$ [45, 46]. If the mode is stable at different θ_0 , then when flow shear advects the mode it can be moved into a stabilising region, reducing its impact on the transport.

A scan was done from $\theta_0 = -\pi \rightarrow \pi$ at $k_y \rho_s = 0.35$ to determine the impact of flow shear on the low k_y modes. Even or odd solutions only exist when the system is up-down symmetric which is only true when $\theta_0 = 0$. Therefore, only the dominant mode can be examined with GS2 initial value simulations when $\theta_0 \neq 0$.

Figure 5a illustrates the sensitivity of the growth rates and mode frequency to θ_0 at $k_y \rho_s = 0.35$. It is clear that the KBM is highly ballooning and only dominates in an extremely narrow range of $\theta_0 \sim 0$. This suggests that KBMs are highly susceptible to flow shear stabilisation and that flow shear will act to increase their critical gradient.

However, Figure 5a shows the MTM growth rate is largely unaffected by variations in θ_0 suggesting that the MTMs will be more resilient against flow shear stabilisation. The eigenfunction for $\theta_0 = \pi$ is shown in Figure 5b; this can be compared with Figure 4b showing the MTM at the same $k_y \rho_s$ with $\theta_0 = 0.0$ where the similar structure is evident. This MTM is clearly insensitive to curvature as the mode is equally unstable at the inboard side compared to the outboard side. This indicates that the drifts reversal, which can occur at the outboard side, may not impact these modes. The central peak




Figure 5: a) Growth rate and frequency of dominant mode at $k_y \rho_s = 0.35$, over a scan in θ_0 . The KBM only dominates around $\theta_0 = 0$, while MTMs take over as the dominant mode for $|\theta_0| > 0.15$ with eigenvalue that is independent of θ_0 . b) The MTM eigenfunction when $\theta_0 = \pi$ and it can be seen that the peak in $A_{||}$ occurs at $\theta = \pi$. c) A close up of the eigenfunctions for the MTMs for a range of θ_0 . d) Full eigenfunction of the MTMs at various θ_0 on a vertical log plot.

in $A_{||}$ occurs at $\theta = \pi$ for $\theta_0 = \pi$ in Figure 5b. Figure 5c overlays the eigenfunctions at different θ_0 around $\theta = 0$ which shows that the central peak moves across θ as θ_0 is increased. Figure 5d shows the full eigenfunction, and the extended tail region all have a similar decaying envelope, indicating that the physics along these tails is not impacted by the central region, which is qualitatively similar to the extended electron tails seen recently in [47]. It is noted that compared to the $\theta_0 = 0.0$ case the relative amplitude of the tails is smaller than for the cases with finite θ_0 .

These results demonstrate that there is a competition between KBMs and MTMs in this region. A small amount of flow shear will reduce the impact the KBM has on the transport as the mode will spend very little time in the θ_0 region that is KBM unstable. However, the MTM which persists across θ_0 , will likely contribute significantly to the




Figure 6: Effective growth rate when including flow shear in the baseline equilibrium where the vertical black line represents the diamagnetic flow shear γ_{dia} .

fluxes regardless of flow shear. This is illustrated in Figure 6, where $\gamma_{E \times B}$ is included in a linear simulation by allowing θ_0 to vary in time. The vertical dashed black line shows diamagnetic flow shear level γ_{dia} [22]. With no flow shear the effective growth rate of a $\theta_0 = 0.0$ mode is that of the KBM with $\gamma = 0.093c_s/a$. This is stabilised to some extent by a small amount of flow shear, but the effective growth rate remains at $\gamma \approx 0.04c_s/a$, which is due to the MTM. On inclusion of flow shear, the eigenmode becomes that of the MTM, which dominates over the KBM for most of the θ_0 range apart from a narrow region around $\theta_0 = 0.0$.

3.2.2. High k_y modes This type of scan was then repeated for the high k_y MTM. Figure 7a illustrates this MTM has a narrow peak in γ around $\theta_0 = 0$, indicating that it is highly ballooning (like the KBM) and will also be stabilised by a small amount of flow shear.

However, a different tearing parity mode appears at $\theta_0 = \pm\pi$ that rotates in the ion diamagnetic direction but has an odd parity ϕ about $\theta_0 = \pi$. The eigenfunction is shown in Figure 7b and it has $C_{\text{tear}} = 0.9$, making it more tearing than the MTM situated around $\theta_0 = 0$. This is labelled as an iMTM (grey filled hexagons) and a tearing ion direction mode has not been reported in the literature before (to the best of the authors' knowledge), especially one that has a maximum growth rate on the inboard side. This highlights the exotic nature of this equilibrium.

Between $0.1\pi < |\theta_0| < 0.7\pi$, both modes are stable. Including flow shear will move the modes through the stable region, resulting in a lower overall growth rate. Figure 7c shows that the inclusion of flow shear reduces the effective growth, even for very low values of $\gamma_{E \times B}$, such that at diamagnetic levels of flow shear the mode is close to stable. A γ/k_{\perp}^2 argument would suggest that the impact of these MTMs on the total transport may not be significant.

Reduced levels of transport might be anticipated in a local equilibrium where the low k_y MTMs could be stabilised given that the KBMs and high k_y MTMs will likely




Figure 7: a) θ_0 scan of the dominant mode at $k_y \rho_s = 4.2$, b) eigenfunction for the inboard iMTM which should be compared with the eigenfunction of the outboard MTM at the same k_y shown in Figure 4d. c) effective growth rate with $E \times B$ shear. The vertical black line shows γ_{dia} .

be stabilised by flow shear.

4. Parametric dependence of micro-instabilities

The rest of this work aims to understand what drives these modes such that the equilibrium can be re-designed to help to stabilise them. The parametric dependence of the linear modes will be determined which will help to identify which plasma parameters should be the focus of optimisation. The primary focus here will be to find actuators to stabilise the low k_y MTMs which appear to be the most dangerous from the transport point of view. The KBMs and high k_y MTMs can be largely ignored when optimising the equilibrium, so long as they aren't driven significantly more unstable.

This section will examine the impact of certain local parameters which were found to be the most significant in the stability properties of this equilibrium. The parameters

examined were a/L_{Te} , a/L_{Ti} , a/L_n , ν_{ee} , $\beta_{e,\text{unit}}$, $\beta'_{e,\text{unit}}$, q and \hat{s} . This will then guide global changes that can be made to the equilibrium

4.1. Impact of kinetic profiles

In SCENE, the density and temperature profiles are prescribed, meaning the main transport assumption is quantified via the H_{98} and H_{Petty} scaling laws. This sub-section will investigate the impact of the kinetic profiles on the micro-stability. Changing the kinetic profiles will change the kinetic gradients a/L_n and a/L_T , so the impact of these needs to be quantified to identify desirable operating scenarios. In this section the electron and ion temperature gradient will be independently changed to explore their impact on the KBMs and MTMs. Next the density gradient of the electrons and ions will be scanned together such that quasi-neutrality is maintained.

In the gyrokinetic equation the pressure gradient influences the local equilibrium in two ways: firstly the impact on local magnetic equilibrium including magnetic drifts and local shear via the current profile; secondly its impacts on the profile gradients providing the linear drive terms. It is possible to define these terms independently allowing for the impact of those terms to be isolated. In GS2 the pressure gradient parameter, β' , affecting the magnetic geometry is prescribed independently of the kinetic profile gradients. A self-consistent pressure gradient would depend on the profile gradients, and satisfy: $\beta'_e \equiv -\beta_e a/L_p$ where $a/L_p = \sum_s \frac{n_s T_s}{n_{\text{ref}} T_{\text{ref}}} (a/L_{Ts} + a/L_{ns})^\dagger$. In the scans in this section we vary the kinetic profile gradients (inconsistently) at fixed magnetic geometry, to allow for the impact of the kinetic gradients to be isolated ‡ .

Changing the density and temperature profiles will also impact the collisionality as $\nu_* \propto n_s/T_s^2$, so a higher density, lower temperature scenario will have a higher collisionality. Furthermore, the impact of impurities and fast ions will be important. This work won't include an impurity species, but will briefly examine the impact Z_{eff} has on these modes. Examining the impact of fast ions such as fusion α 's is left as future work.

4.1.1. Electron Temperature gradient The impact of the electron temperature gradient was examined by scanning from $a/L_{Te} = 0 \rightarrow 7.0$ at $k_y \rho_s = 0.35$. Generally MTMs are driven unstable by the electron temperature gradient and KBMs are driven unstable by the total pressure gradient. The eigenvalues are shown in Figure 8a, with the reference equilibrium value shown with the vertical black dashed line.

There appears to be a critical gradient where the KBM is completely stable at $(a/L_{Te})_{\text{crit}}^{\text{KBM}} = 2.0$, which would correspond to $a/L_p = 5.63$. Its growth rate increases exponentially with a/L_{Te} which may lead to stiff transport and if γ becomes sufficiently

† This work is missing the contribution from the fast ions which may be significant at high β

‡ β' is related to the parameter α often used to define the pressure gradient with $\alpha = Rq^2\beta'$ in the infinite aspect ratio shifted circular geometry limit.




Figure 8: a) Examining the impact of plasma a/L_{Te} when $k_y \rho_s = 0.35$. The dominant even and odd instabilities are shown, with the KBM demonstrating much stiffer behaviour. Note the log scale in γ . The dashed orange line shows the analytic prediction ω_{MTM}^{CR} defined in the text. The vertical dashed black line shows the equilibrium value of a/L_{Te} . b) Eigenfunction of the ion direction odd parity mode (filled green stars) for $a/L_{Te} = 7.0$; despite having odd parity, the mode is not tearing as $C_{tear} = 0.0$.

large then the flow shear stabilisation may not be sufficient in suppressing the transport[†]. The mode frequency is dropping as the temperature gradient increases. A small drop in a/L_{Te} would allow for the MTM to become the dominant instability.

Looking at the dominant odd eigenmode, it can be seen that there's a critical gradient $(a/L_{Te})_{crit}^{MTM} = 1.0$. This critical gradient may be the limiting factor on the electron temperature profile as MTMs can drive significant electron heat flux. The mode frequency scales with the temperature gradient and this follows predictions made by Catto and Rosenbluth [33], that the mode frequency of an MTM is given by $\omega_{MTM}^{CR} = \omega_e^*[1 + \eta_e/2]$ where ω_e^* is the electron diamagnetic frequency defined as $\omega_e^* = k_y(a/L_{ne})$ and $\eta_e = (L_{ne}/L_{Te})$. The orange dashed line shows ω_{MTM}^{CR} , indicating that this scaling fits well.

The MTM growth rate displays a non-monotonic behaviour which has been seen before in MAST simulations [34] where the MTMs also had a mode frequency like ω_{MTM}^{CR} . This was thought to be related to a resonance with a drift frequency. If $|\omega|$ is increased sufficiently then this resonance is disturbed and the mode becomes damped. As observed previously for MTMs, instability only arises over a finite range in a/L_{Te} , which interestingly, is consistent with recent analytic calculations demonstrating that toroidal ETG modes can only be unstable over a finite range in $(\eta_e \omega_e^*)/\omega_{ke}$ [48]. Increasing a/L_{Te} reduces the growth rate at a given k_y , but pushes the MTM spectrum to longer wavelength as seen in Figure 9b.

[†] Of course as a/L_{Te} increases, so would $\beta'_{e,unit}$, which will be shown in Section 4.2.1 to be stabilising so the combined impact is not clear.

However, when $a/L_{Te} > 5.0$, an ion direction mode emerges that has an odd ϕ eigenfunction, shown in Figure 8b. Its frequency is tending towards the KBM frequency. This however, is not a tearing mode as it has $C_{\text{tear}} = 0.0$. Rather, this appears to be an odd parity KBM, which will be labelled as an oKBM and will be represented with the filled green stars as seen in Figure 8a. This oKBM is a higher order eigenstate of the KBM and has been seen before in steep-gradient simulations [49]. If this is an oKBM, it should be seen when scanning through a/L_{Ti} .

A scan was also done across $k_y \rho_s$ to see how the spectra changed with a/L_{Te} and the eigenvalues for the KBMs and MTMs are shown in Figures 9a and 9b respectively. It can be seen that as a/L_{Te} increases the KBM remains unstable at very low values of $k_y \rho_s$, suggesting a local model may fail at high gradients[†]. For the MTMs the non-monotonic behaviour persists across the k_y spectrum, with the $a/L_{Te} = 5.0$ having the lower growth rates than the $a/L_{Te} = 4.0$ scan.




Figure 9: A scan in k_y at several different values of a/L_{Te} showing the dominant a) even and b) odd mode. The colour denotes the mode type (KBM - blue, MTM - orange) and the line-style shows the electron temperature gradient. Note the log scale for the KBMs.

It is expected that the high k_y MTMs will be impacted by a/L_{Te} so a similar scan is shown in Figure 10 for $k_y \rho_s = 4.2$. Once again non-monotonic behaviour is seen with a critical electron temperature of $a/L_{Te} = 1.0$, similar to the lower $k_y \rho_s$ MTM seen earlier. Significant changes in a/L_{Te} would be required to stabilise this MTM, though given the sensitivity of the growth rate to θ_0 , the transport from these modes will be mitigated by flow shear.

4.1.2. Ion temperature gradient A similar scan was performed for the ion temperature gradient whilst keeping the other kinetic gradients fixed. For $k_y \rho_s = 0.35$, the KBM

[†] Once again, this would also increase β'_{eunit} which is stabilising so a consistent study should be required for each equilibrium.




Figure 10: Examining the impact of a/L_{Te} on the MTM seen at $k_y \rho_s = 4.2$. The vertical dashed black line shows the equilibrium value of a/L_{Te} .

and MTM eigenvalues are shown in Figure 11. The KBM has a similar critical gradient to the previous scan with $(a/L_{Ti})_{\text{crit}}^{\text{KBM}} = 2.0$, as expected if the relevant parameter is a/L_p . The KBM is strongly destabilised by a/L_{Ti} and the mode frequency in this case actually increases with a/L_{Ti} , which suggests that the KBM frequency scales like η_i/η_e .




Figure 11: Examining the impact of plasma a/L_{Ti} when $k_y \rho_s = 0.35$. The dominant even and odd instabilities are shown, with the KBM, once again, demonstrating much stiffer behaviour. The oKBM is again found at high a/L_{Ti} . The vertical dashed black line shows the equilibrium value of a/L_{Ti} .

Examining the low k_y MTM, its growth rate and mode frequency are largely unaffected by a/L_{Ti} as expected. Once again if the ion temperature gradient is pushed high enough then when $a/L_{Ti} > 5.5$ an oKBM appears at a similar threshold as the a/L_{Te} scan. The high k_y MTM is also unaffected by the ion temperature gradient.

It has been found in MTM driven transport, that 98% of the heat transport can occur in the electron channel [27]. This suggests that a/L_{Ti} will predominantly be determined by the balance between the neoclassical transport and the electron-ion exchange power, assuming the KBMs is suppressed by flow shear.

4.1.3. Density gradient A density gradient scan was performed from $a/L_n = -1 \rightarrow 1$, corresponding to $a/L_p = 3.54 \rightarrow 7.54$, with the negative density gradient allowing for a similar lower value of a/L_p as the temperature gradient scans. It can be seen from Figure 12a that when a/L_n is increased the KBM is destabilised, which further supports that this is a pressure gradient driven mode. The KBM is stable when $a/L_n < 0$ corresponding to $a/L_p = 5.54$ which is similar to the critical value in the temperature gradient scan. The mode frequency is unaffected which supports the idea that ω_{KBM} scales with η_i/η_e which remains unchanged in this scan. When the density gradient is negative the KBM is stabilised and an electrostatic passing electron mode (ES-PEM shown by hollow red diamonds) appears; this is a scenario that should be avoided for a reactor, though may occur locally and transiently if pellet fuelling occurs at the edge. This mode was found to only require passing electrons and depended on both the electron and ion temperature gradient.

The low k_y MTM seems to be stabilised by a large $|a/L_n|$, which has been seen before on NSTX [39] and on MAST [26]. This again was thought to be due to the mode frequency changing and disrupting a resonance, though the peak growth rate here occurs at $\omega = 0.4c_s/a$ and in the a/L_{Te} scan it occurred at $\omega = 0.6c_s/a$.




Figure 12: Examining the impact of a/L_n when a) $k_y \rho_s = 0.35$ and b) $k_y \rho_s = 4.2$. The dominant even and odd instabilities are shown. The vertical dashed black lines shows the equilibrium value of a/L_n .

For the high k_y MTMs, the equilibrium happens to lie at the peak of the growth rate spectrum so increasing the density gradient would help to stabilise the mode, but the effect is not as large compared to the impact on the low k_y MTMs. Given the similar

behaviour as the low k_y modes, similar measures can be taken to stabilise these modes, noting that their impact on transport will not be as significant given the flow shear stabilisation.

4.1.4. Scans in a/L_T and a/L_n at fixed a/L_p It has been shown that increasing any of the kinetic gradients drives the KBM unstable. Assuming the KBM is driven by the total pressure gradient, if the total pressure gradient is kept fixed a similar growth rate should be seen. However, if the temperature gradient is exchanged for density gradient then it should be expected that the MTM will be stabilised. This should be doubly beneficial for the stability as the drive from a/L_{Te} is reduced and the stabilisation from a/L_n is being increased. A scan was performed at fixed a/L_p where the density gradient was changed from $a/L_n = 0 \rightarrow 1.5$ and $a/L_{Te} = a/L_{Ti}$ was set. The eigenvalues are shown in Figure 13. Note the linear scale here.

It can be seen that the MTM is indeed stabilised, and when $a/L_n > 1.0$, the MTM is completely stable. The critical gradient occurs at the same value as the pure density gradient scan, indicating that reducing the electron temperature gradient is not having a significant impact on the MTM growth rate. This is explained by the “levelling” out of the MTM growth rate seen in Figure 8a during the a/L_{Te} scan, indicating that small changes in a/L_{Te} will not impact these MTMs significantly.

Overall, this implies that a peaked density profile could be beneficial in reducing MTM-based transport. The KBM is also stabilised slightly indicating the growth rate is slightly sensitive to L_n/L_T .




Figure 13: a) A scan was performed at fixed a/L_p where the contribution of a/L_n was varied for $k_y \rho_s = 0.35$. $a/L_{Te} = a/L_{Ti}$ was enforced. Note the linear scale. The vertical dashed black line shows the equilibrium value of a/L_n .

4.1.5. Collision Frequency The kinetic profiles were specified to generate $P_{\text{fus}} = 1.1\text{GW}$, but it would be possible to operate at a higher density and lower temperature at the same fusion power. This would directly impact the collisionality of the plasma given that $\nu_* \propto n_e/T_e^2$.

MTMs are sometimes reported to be highly sensitive to the electron collision frequency, so a scan was conducted from $\nu_{ee} = 0 \rightarrow 0.14c_s/a$ and ν_{ei} was consistently changed assuming $T_e = T_i$. An electron collision frequency of $0.14c_s/a$ (which corresponds to $\nu_* = 0.05$) is rather high for reactor relevant conditions and would be approximately at the Greenwald limit for this device. Given the high temperatures of a reactor, it will be difficult to operate at a significantly higher collision frequency.




Figure 14: a) Impact of collision frequency showing the dominant odd and even parity mode for different values of ν_{ee} when $k_y \rho_s = 0.35$. b) Eigenfunction of the low k_y MTM when $\nu_{ee} = 0.14c_s/a$ showing the reduced extent in ballooning space. c) Growth rate and frequency for the $k_y \rho_s = 4.2$ MTM as a function of ν_{ee} . The vertical dashed black line shows the equilibrium value of ν_{ee} .

When examining the low k_y MTM, as shown in Figure 14a, it appears that γ^{MTM} increases with ν_{ee} , and is only unstable with a finite collision frequency. Additionally,

the collisions reduce the extent of the mode in ballooning space as shown in Figure 14b where $\nu_{ee} = 0.14c_s/a$, compared to the nominal collisionality ($\nu_{ee} = 0.017c_s/a$) eigenfunction shown in Figure 4b. This is due to the passing electrons undergoing a collision before they can propagate further along the field line. The reduced extent of the mode would make nonlinear simulations easier to resolve.

As the collision frequency is dropped towards 0, the MTM growth rate tends to 0, which is also consistent with the collisionality scaling recovered from MTM simulations for NSTX by Guttenfelder *et al* [39]. This suggests that the confinement will scale favourably as collisionality is reduced, aligned with previous confinement scaling laws in STs where $B\tau_e \propto \nu_{*e}^{-0.82}$ [50].

However, Figure 14a also shows that as the collision frequency is increased the KBM is stabilised and a similar feature has been seen in a hybrid TEM/KBM mode in simulations of NSTX [27]. There is a critical collision frequency at which the dominant mode switches from a KBM to an MTM and the KBM becomes stable for $\nu_{ee} > 0.12c_s/a$. Furthermore, this is consistent with NSTX observations of increasing anomalous ion transport at low ν^* , attributed to KBM-TEM hybrid modes [51].

Examining the impact of collisions on the high k_y MTM, it can be seen from Figure 14c that these are collisionless MTMs, highlighting the completely different nature of these modes compared to the longer wavelength MTMs. Collisionless MTMs have been seen before [35, 43, 44, 52], but their mechanism is not fully understood. These are weakly stabilised by collisions over the range of the scan, but it appears that the impact is not significant. The impact of Z_{eff} was also examined and was found to have a similar impact to the ν_{ee} for all 3 modes being examined due to its impact on the collision frequency as $\nu_{ei} \rightarrow Z_{\text{eff}}\nu_{ei}$ in the Lorentz collision operator.

4.2. Magnetic equilibria

Both MTMs and KBMs are inherently electromagnetic modes, so the total magnetic field will have a significant impact on the modes via β_e . Furthermore, there is evidence that both of these modes are impacted by both q and \hat{s} , which are set by the toroidal field and plasma current profile. Both of these were also inputs to SCENE, so it is possible to vary the assumptions about the magnetic equilibrium to see how to further stabilise these modes. This section will perform local equilibrium scans to assess the impact of β_e , β'_e , q and \hat{s} , which might provide a guide to the impact of changing the toroidal field and plasma current profile in a global Grad-Shafranov equilibrium from SCENE.

4.2.1. Impact of $\beta_{e,\text{unit}}$ and $\beta'_{e,\text{unit}}$ A scan was conducted in $\beta_{e,\text{unit}} = 0.0 \rightarrow 0.024^\dagger$ at $k_y\rho_s = 0.35$, where $\beta_{e,\text{unit}} = 0.024$ at $\rho_\psi = 0.5$ approximately corresponds to a 40% drop in the field compared to the reference case in Table 2. In a self-consistent scan both $\beta_{e,\text{unit}}$ and $\beta'_{e,\text{unit}}$ would be scanned together, but in the first scan shown in Figure 15a

[†] The subscript unit corresponds to the normalising magnetic field being $B_{\text{unit}} = \frac{q}{r} \frac{\partial \psi}{\partial r}$.

$\beta_{e,\text{unit}}$ alone is scanned with $\beta'_{e,\text{unit}}$ fixed. This effectively scans in the strength of the magnetic perturbations.

Looking at the even modes, when $\beta_{e,\text{unit}} = 0$, a weakly unstable electrostatic passing electron mode (ES-PEM) appears, which was seen in the negative density gradient scan in Figure 12. Above a critical value of $\beta_{e,\text{unit}} = 0.01$, the KBM mode becomes unstable, before it begins to saturate around $\beta_{e,\text{unit}} = 0.02$. Looking at the MTMs, again there is a critical $\beta_{e,\text{unit}} = 0.006$, below which the MTM is stable. This is driven unstable by $\beta_{e,\text{unit}}$, but when $\beta_{e,\text{unit}}$ increases sufficiently then the oKBM is seen, which is very sensitive to $\beta_{e,\text{unit}}$ and quickly begins to approach the KBM growth rate.




Figure 15: Examining the impact of a) $\beta_{e,\text{unit}}$ at fixed $\beta'_{e,\text{unit}} = 0.078$ and b) $\beta'_{e,\text{unit}}$ at fixed $\beta_{e,\text{unit}} = 0.012$ on the dominant even and odd parity micro-instabilities at $k_y \rho_s = 0.35$. The vertical dashed black lines shows the equilibrium values.

$\beta'_{e,\text{unit}}$ is used to calculate the equilibrium and has large impacts on the magnetic drifts and local magnetic shear. A scan was performed in $\beta'_{e,\text{unit}}$ at the fixed equilibrium $\beta_{e,\text{unit}}$ to isolate its impact. This is shown in Figure 15b where it can be seen that at low $\beta'_{e,\text{unit}}$, the KBM/oKBM are the dominant instabilities. These are quickly stabilised and the dominant odd mode switches from an oKBM into the MTM and the dominant even mode switches from a KBM to a weakly unstable ES-PEM. The MTM is not significantly affected by $\beta'_{e,\text{unit}}$ and the reason can be determined by examining the drifts.

Figure 16 illustrates how $\omega_{\nabla B}$ and ω_{curv} are modified by changing $\beta'_{e,\text{unit}}$. The total magnetic drift is approximately the sum of $\omega_{\nabla B}$ and ω_{curv} . Here a negative drift frequency corresponds to “good curvature” and positive is “bad curvature”. Increasing $\beta'_{e,\text{unit}}$ makes $\omega_{\nabla B}$ negative such that the combination of the two drifts becomes stabilising on the outboard side, resulting in “good curvature”. This effect is highly stabilising for highly ballooning KBMs that are driven by bad curvature on the outboard side. The MTMs at low k_y are considerably more extended in the parallel direction, and much less sensitive to drift reversal at the outboard mid-plane, with growth rates that are relatively insensitive to $\beta'_{e,\text{unit}}$.




Figure 16: The impact $\beta'_{e,\text{unit}}$ has on $\omega_{\nabla B}$ and ω_{curv} . The ∇B drift is negative for the high β' cases when $|\theta| < \pi$, which is stabilising for ballooning modes.

For the high k_y MTM, Figure 17a shows a scan of changing $\beta_{e,\text{unit}}$ at fixed $\beta'_{e,\text{unit}}$. Exceeding a critical $\beta_{e,\text{unit}} = 0.003$ was found to destabilise this mode, corresponding to a 70% increase in the field. Figure 17b shows how changing $\beta'_{e,\text{unit}}$ impacts the high k_y MTM which is very different to the low k_y MTMs which were largely unaffected by $\beta'_{e,\text{unit}}$, which is related to the MTMs respective sensitivity to θ_0 . At lower $\beta'_{e,\text{unit}}$ this MTM is stabilised, but another MTM branch appears for $\beta'_{e,\text{unit}} < 0.04$, likely also driven unstable by $\beta_{e,\text{unit}}$. At $\beta'_{e,\text{unit}} = 0.0$, this MTM has $C_{\text{tear}} = 0.3$ which is lower than that for the equilibrium $\beta'_{e,\text{unit}}$ at $\theta_0 = 0$ which had $C_{\text{tear}} = 0.5$. However, a scenario with a very low $\beta'_{e,\text{unit}}$ and high $\beta_{e,\text{unit}}$ requires a very low pressure gradient which is not desirable for a reactor as this would result in a lower bootstrap fraction.

To examine the relevance of this low $\beta'_{e,\text{unit}}$ MTM, a scan was done where $\beta'_{e,\text{unit}}$ and $\beta_{e,\text{unit}}$ were changed together i.e. $\beta'_{e,\text{unit}} = \beta_{e,\text{unit}} a / L_p$ was maintained throughout this scan. Figure 17c shows that at sufficiently low $\beta_{e,\text{unit}}$, the original MTM is stabilised and the new MTM does not appear. However, an ETG mode (hollow grey upside-down triangle) appears when operating at lower $\beta_{e,\text{unit}}$ and $\beta'_{e,\text{unit}}$. This ETG is explored in more detail in Appendix B. But this suggests that this low $\beta'_{e,\text{unit}}$ MTM will not be relevant.

4.2.2. Safety factor profile The q profile has a large impact on both the MHD and the micro-stability of a reactor. Tailoring the q profile requires careful control of the auxiliary current profile so it may be difficult to optimise it for turbulence but this section highlights how it will impact these modes.




Figure 17: Examining the impact of changing a) $\beta_{e,unit}$ at fixed $\beta'_{e,unit} = 0.078$ and b) $\beta'_{e,unit}$ at fixed $\beta_{e,unit} = 0.012$. c) $\beta_{e,unit}$ and $\beta'_{e,unit}$ are changed together for the MTMs at $k_y \rho_s = 4.2$. Above $\beta_{e,unit} = 0.016$, the equilibrium was completely stable. The vertical dashed black lines shows the equilibrium values.

The $n = \infty$ ballooning stability boundary for ideal ballooning modes is often used as an initial indicator for the onset of KBMs [53] and in GS2 there is a module that can calculate this boundary [54]. Figure 18 shows how this ideal stability boundary changes with q , illustrating a somewhat complicated relation between the ideal ballooning mode and q . As q increases the stability boundary moves to high \hat{s} and lower $|\beta'|$. Therefore pushing to a higher q will make access to the second stability region easier. At sufficiently low q , the stability boundary gets pushed to higher $|\beta'|$ enabling the equilibrium to lie in the first stability region.

Evidently, the ideal ballooning mode is not sufficient in predicting the KBM threshold as it predicts this equilibrium to be stable, but it can give an idea of how the KBM will behave. Figure 19a shows a scan in q for the kinetic modes. For $q < 3.0$, when the equilibrium is in the first stability region, an ITG (hollow cyan pentagons) mode is dominant but as the equilibrium enter the second stability region, the KBM




Figure 18: $\hat{s} - |\beta'|$ diagram showing how the ideal ballooning stability boundary moves with q . The reference equilibrium value of \hat{s} and $|\beta'_{e,unit}|$ is shown by the green cross and has $q = 4.3$ (green curve).

becomes the dominant instability. As q is increased the KBM peaks and then becomes stabilised as the ideal boundary gets pushed further away from the equilibrium value. When the equilibrium is in the second stability region then increasing q stabilises the KBM and another ES-PEM appears, indicating that operating at high q may help to increase the KBM critical gradient. This will be useful if the MTMs can be suppressed.

A scan was also conducted with \hat{s} and Figure 19b shows how the KBM is destabilised by increasing \hat{s} , consistent with the ideal ballooning mode behaviour.




Figure 19: Dominant even parity mode eigenvalues when changing a) q and b) \hat{s} for $k_y \rho_s = 0.35$. The vertical dashed black lines shows the equilibrium values.

Next, the impact \hat{s} and q have on the low k_y MTMs is explored, and it can be expected that increasing q will be destabilising for this MTM given that $\nu_* \propto q$. To

isolate the impact on the MTM and avoid the oKBM being driven unstable, the q scan was run with $\nu_{ee} = 0.14c_s/a$. Figure 20a illustrates how there appears to be a linear relationship between q and the MTM growth rate, consistent with a ν_* scaling. The magnetic shear dependency displays non-monotonic behaviour as shown in Figure 20b.




Figure 20: Dominant odd parity mode eigenvalues when changing a) q (at $\nu_{ee} = 0.14c_s/a$) and b) \hat{s} for $k_y\rho_s = 0.35$. The vertical dashed black lines shows the equilibrium values.

The impact of \hat{s}/q is often examined as it is related to Landau damping and field line bending. For the MTM seen here it appears that \hat{s}/q has a non-monotonic behaviour, so the impact on the turbulent transport will be difficult to predict. This has been seen before on MAST [26]. However, there are examples where \hat{s}/q is stabilising for MTMs like in DIII-D [25] and counter-examples where \hat{s}/q tends to be destabilising, such as on NSTX [39] attributed to higher field line bending. This further highlights the complicated behaviour of MTMs so further work is needed to understand the dependence on the q -profile.

Figure 21a shows a scan in q for the high k_y MTMs, and it can be seen that the growth rate has a peak close to the local equilibrium value of q , so increasing or decreasing can reduce the growth rate. A higher B_ϕ device will likely have a higher q , so the combination of lower $\beta_{e,\text{unit}}$ and higher q indicates that this mode will be stabilised by a higher B_ϕ . At very high q an iMTM mode with $C_{\text{tear}} = 0.7$ was found, though it exists for a very narrow window. Looking at the behaviour with \hat{s} in Figure 21b, there is a non-monotonic dependence similar to the lower k_y MTM. However, it seems to be a weak dependence so it will likely not have a significant impact on this MTM.

4.3. Radial surfaces at $\rho = 0.3$ and $\rho = 0.85$

This analysis was repeated for two different flux surfaces at $\rho_\psi = 0.3$ and 0.85 and similar KBMs and MTMs were found, indicating that these modes and the trends reported here exist robustly across the whole equilibrium. These are discussed in more detail in [8].




Figure 21: Impact of a) q and b) \hat{s} on the MTMs at $k_y \rho_s = 4.2$. The vertical dashed black lines shows the equilibrium values.

5. Options for stabilising the equilibrium

Section 4 has indicated potential avenues for stabilising the dominant micro-instabilities based on local equilibrium scans. Here these insights are used to modify the global equilibrium using SCENE with two options being pursued. Firstly, the impact of increased toroidal field is examined. To maintain the same q , I_p would need to be increased. However, I_p was fixed to avoid the need for additional auxiliary current drive. Secondly, the impact of a more peaked density profile is considered.

5.1. Impact of higher toroidal field at constant plasma current

Increasing B_ϕ by going to higher I_{rod} at constant I_p , will reduce β_e and increase q . These changes have opposing impacts on the stability of the low k_y MTMs: reducing β_e is stabilising, but increasing q is destabilising.

To investigate the impact of higher field, SCENE was used to consistently generate equilibria with a higher toroidal field. The value of I_{rod} was increased from 30MA \rightarrow 50MA in 5MA increments, whilst the density and temperature profiles were kept fixed. The total plasma current was fixed as well as the shape of the auxiliary current profile. As I_{rod} increases, there was a small increase in the bootstrap current but this was offset by a small reduction in the diamagnetic current such that the total auxiliary current remained roughly constant. This results in a near linear increase in the safety factor profile with I_{rod} . Furthermore, most of the local equilibrium parameters stayed within 1% of the baseline value. The main parameters that did change for $\rho_\psi = 0.5$ surface were

- $\beta_{e,\text{unit}} : 0.012 \rightarrow 0.004$
- $\beta'_{e,\text{unit}} : -0.08 \rightarrow -0.03$
- $q : 4.3 \rightarrow 7.3$

- $\hat{s} : 0.78 \rightarrow 0.70$

Figure 22a illustrates how the growth rate and frequency of the $k_y \rho_s = 0.35$ mode varies over the scan in I_{rod}^\dagger , with $\beta_{e,\text{unit}}$ shown by the dashed red line. Firstly it is important to note that the KBM and two different MTMs were found in all of these equilibria, suggesting that these will exist over a range of different high β ST equilibria. The KBM is weakly stabilised by the higher rod current. It seems that the stabilisation from the lower $\beta_{e,\text{unit}}$ and higher q is counteracted by the lower $\beta'_{e,\text{unit}}$. For the MTM there is a very weak stabilisation as the increased q counteracts the lower $\beta_{e,\text{unit}}$. If I_{rod} is sufficiently increased, as $\beta_{e,\text{unit}} \rightarrow 0$, these modes should become stable, but $I_{\text{rod}} = 50\text{MA}$ is pushing the limits of engineering so going further beyond this is unlikely to be feasible.




Figure 22: Examining the impact of a higher field device by increasing I_{rod} for a) $k_y \rho_s = 0.35$ and b) $k_y \rho_s = 4.2$ at $\rho_\psi = 0.5$.


Examining the impact on the high k_y MTM, Figure 22b shows that operating at higher field stabilises the mode and when $I_{\text{rod}} > 40\text{MA}$, consistent with the $\beta_{e,\text{unit}}$, $\beta'_{e,\text{unit}}$ scan conducted in Figure 17c. However, the ETG seen there is not destabilised in this scan as discussed further in Appendix B.

5.2. Impact of density peaking


The results of the Section 3.2 suggest that the flow shear will wipe out the KBMs and the high k_y MTMs found so far. It was also demonstrated that increasing the density gradient was beneficial in stabilising the low k_y MTMs. If the a/L_T is reduced whilst increasing a/L_n without changing the fusion power, then it may be possible to design an equilibrium in which the majority of the linear instabilities are stable.

Using this information a new equilibrium was designed in SCENE with a more peaked density profile. An operating point was examined where $n_{e0}/\langle n_e \rangle$, which was


† Note this is not at fixed toroidal mode number n as ρ_s changes with B



(a)



(b)



(c)

Figure 23: Comparison of a) temperature, b) density and c) safety factor profile for the baseline case in Table 1 and peaked density case in Table 4.

equal to 1.12 in the baseline operating point, was increased to $n_{e0}/\langle n_e \rangle = 1.58$ at fixed $P_{\text{fus}} = 1.1 \text{ GW}$. The pedestal height and width were kept the same. To reduce the temperature gradient drive, the core temperature for both species was dropped from 28 keV to 20 keV . The auxiliary current profile was kept fixed. The resulting equilibrium parameters are outlined in Table 4. The temperature, density and safety factor profiles are illustrated in Figures 23a, 23b and 23c respectively.

The parameters are broadly similar to the baseline case, with the exception of the density and temperature values. The Miller and plasma parameters for the $\rho_\psi = 0.5$

Parameter	Baseline	Peaked n
R_{maj} (m)	2.5	2.5
a (m)	1.5	1.5
R_0 (m)	3.15	3.05
I_{rod} (MA)	30.0	30.0
I_p (MA)	21.0	21.0
I_{aux} (MA)	8.2	7.1
P_{fus} (MW)	1100	1100
P_{aux} (MW)	94	83
κ	2.8	2.8
δ	0.55	0.55
H_{98}	1.35	1.26
H_{Petty}	0.94	0.83
T_{e0} (keV)	28.0	22.0
$\langle T_e \rangle$ (keV)	14.8	14.1
n_{e0} (10^{20}m^{-3})	1.72	2.19
$\langle n_e \rangle$ (10^{20}m^{-3})	1.54	1.38
f_{GW}	0.52	0.52
l_i	0.27	0.38
β_N	5.5	5.13
q_0	2.51	2.07

Table 4: Comparing the plasma parameters for the baseline and the peaked density operating points.

Parameter	Baseline	Peaked n
r/a	0.66	0.67
R_{maj}/a	1.79	1.83
n_{e20} (m^{-3})	1.51	1.36
T_e (keV)	12.2	12.8
B_0 (T)	2.16	2.18
B_{unit} (T)	7.52	7.34
ρ_s (m)	0.0021	0.0022
a/L_n	0.43	1.40
a/L_T	2.77	2.34
Δ	-0.57	-0.48
q	4.30	3.14
\hat{s}	0.78	0.97
κ	3.03	2.87
s_κ	-0.14	-0.11
δ	0.45	0.34
s_δ	0.19	0.21
β_e	0.15	0.14
$\beta_{e,\text{unit}}$	0.012	0.013
$\nu_{ee}(c_s/a)$	0.017	0.013
$\gamma_{\text{dia}}(c_s/a)$	0.080	0.055

Table 5: Local plasma and Miller parameters for the $\rho_\psi = 0.5$ surface of the baseline and peaked density equilibrium.

surface are detailed in Table 5. It can be seen that many of the parameters are very similar to the baseline parameters in Table 2, with the notable exception of the higher a/L_n and lower a/L_T as expected. However, a/L_p has increased overall from 6.40 to 7.48, so it may be expected that the KBMs will be driven more unstable, though as seen in Section 4.2.1, this is counteracted by the increased $\beta'_{e,\text{unit}}$. \hat{s}/q has increased, due to the differences in the bootstrap and diamagnetic current profiles, with this design having a higher bootstrap fraction due to the peaked density profile. Following the trends seen in Figures 19a and 19b, this is expected to destabilise the KBMs but its impact on the MTMs is less clear.

The dominant instabilities are presented in Figure 24. Similar to the baseline operating point, in the low k_y region KBMs and extended MTMs are found. However, the peak MTM growth rate is $\gamma_{\text{MTM}} = 0.004c_s/a$, significantly lower than the baseline operating point where it was $\gamma_{\text{MTM}} = 0.04c_s/a$. The KBM peak growth rate has increased from $\gamma_{\text{KBM}} = 0.09c_s/a$ to $\gamma_{\text{KBM}} = 0.19c_s/a$, but these modes are still narrow in θ_0 so it will be shown that again these are wiped out by flow shear. The higher k_y

MTMs are also seen and have a lower growth rate reducing from $1.2c_s/a$ to $0.9c_s/a$, and it is expected that the flow shear will help to stabilise these modes as well. Finally above $k_y\rho_s = 6$, the equilibrium was found to be completely stable, similar to the previous case. It should be noted that these modes peak at a lower k_y in Figure 24 compared to the baseline case in Figure 3 and this is expected as ρ_s has not significantly changed but the value of q is lower compared to the baseline case in Table 2. Specifically, given that $k_y = \frac{nq}{r}$, to find a given toroidal mode number n at lower q , a lower k_y is needed. The low k_y MTM here was unstable down to $n = 3$.




Figure 24: Dominant odd and even mode for the $\rho_\psi = 0.5$ surface of the peaked density equilibrium without flow shear outlined in Table 5.

A simulation was run for this case with $E \times B$ flow shear at the diamagnetic level of $\gamma_{\text{dia}} = 0.055c_s/a$. The effective growth rate is shown in Figure 25 and it was found that above $k_y\rho_s = 0.2$ the equilibrium was completely stable. Below this value, the MTMs did cause a very slowly growing mode with a growth rate $\mathcal{O}(10^{-3})$. While nonlinear simulations are required to determine the level of driven transport, nevertheless this work hints that density peaking is favourable and the plasma may be able to operate close to neoclassical levels of transport. The impact of impurities and fast ions needs to be examined as they may drive the MTM more unstable, increasing the effective growth rate. In particular, density peaking can lead to impurity accumulation in the core of conventional tokamaks, which would be detrimental to the performance of a reactor. While quantifying the level of impurity accumulation and impact on radiative losses is important, this equilibrium suggests a possible pathway to a high performance high β ST. The open question is how to generate peaking in the equilibrium density profile.

The level of density peaking will be set by particle transport and fuelling. The




Figure 25: Effective growth rate for the $\rho_\psi = 0.5$ surface of the peaked density equilibrium with diamagnetic levels of flow shear.

transport for each species can be decomposed into a diffusive term D and a convective term V , shown in Equation 3.

$$\Gamma = D \frac{1}{n} \frac{\partial n}{\partial r} + V \quad (3)$$

The convective particle pinch term from the low k_y MTMs can be estimated using a quasi-linear argument to determine whether it leads to outward or inward particle transport. If the saturated turbulence has a quasi-linear character, then many attributes of the turbulent transport can be inferred from the properties of the linear instabilities. Assuming the quasi-linear turbulence is dominated by a particular linear mode, it is possible to determine the ratio of the convective pinch and diffusivity terms, V/D , shown in Equation 3. By introducing a trace Deuterium species and modifying the trace density gradient to find the point where $\Gamma \rightarrow 0$, the ratio of V/D can be determined. Examining $k_y \rho_s = 0.35$ when $n_{\text{trace},D}/n_e = 0.05$, Figure 26a illustrates that zero particle flux occurs at $a/L_n = -1.56$ meaning $V/D = -a/L_n/a = 1.56/1.5 = 1.04m^{-1}$. This means the particle convection is outwards making density peaking even more difficult. The particle flux is normalised to the linear electron heat flux and is $\mathcal{O}(10^{-5})$ so the effect may be small but will not help achieve a peaked density profile. A similar scan for the KBMs shows that they also have an outwards convection, shown in Figure 26b. The relative size of the particle flux is $\mathcal{O}(10^{-2})$, larger than the MTMs but still quite small overall. These quasi-linear ideas suggest that peaked density is unlikely to be produced through transport from these KBMs and MTMs. The generation of peaked density profiles will require either other mechanisms contributing to the particle transport, or a core localised particle source.




Figure 26: Linear particle flux of a trace deuterium when scanning through the trace density gradients for the a) MTM and b) KBM at $\rho_s = 0.35$. The particle flux has been normalised to the electron heat flux.

6. Conclusions

An equilibrium has been presented from a potential operating point of a high β spherical tokamak. Local micro-instability studies of the thermal plasma (neglecting fast ions) find that KBMs and MTMs are the dominant micro-instabilities. Furthermore, the equilibria examined here were found to be stable in the electron scale range where $k_y \rho_s > 6$.

Two different types of MTMs were found, a collisional MTM when $k_y \rho_s < 0.6$ and a collisionless MTM between $3.0 < k_y \rho_s < 6.0$. The KBMs and high k_y MTMs were highly ballooning, and unstable only in a very narrow range of $\theta_0 \sim 0.0$, such that diamagnetic levels of flow shear was sufficient in suppressing these modes. The low k_y MTM, however, was unaffected by flow shear and thus is expected to be the dominant source of transport for this equilibrium. These low k_y MTM were very extended along the field line in ballooning space and require resolving electron scales radially indicating a potential challenge for nonlinear simulations.

The parametric dependency of these modes on different equilibrium parameters was determined and it was found that the low k_y MTMs can be stabilised by the raising density gradient, suggesting that a peaked density profile would be beneficial to confinement. Moreover, the low k_y MTMs were destabilised by increasing ν_* , consistent with previous energy confinements scalings from ST experiments [38, 39, 50], indicating that a high temperature/low density device will benefit the confinement. Increasing the magnetic field was found to have little impact on the normalised growth rate of MTMs

at a fixed value of $k_y \rho_s$, owing to competing effects from reduced β and increasing ν_* . Nonlinear simulations with the full spectrum are of course required to rigorously estimate the transport, but if transport from a given mode followed the mixing length estimate, the effective thermal diffusivity from such a mode would still be proportional to $1/B^2$. The overall trends for growth rates with a/L_n , a/L_{Te} , β , q , ν_{ee} of the low k_y MTMs seen here are qualitatively similar to MTMs seen in the other work by [22, 35, 39, 55] on NSTX and MAST equilibria.

An optimised equilibrium was generated with a more peaked density profile and flatter temperature profile. With flow shear, the equilibrium was found to be marginally stable, indicating that operating close to neoclassical levels of transport may be feasible if such an equilibrium can be accessed. The convective pinch term was found to be outwards for both low k_y modes, suggesting that transport from MTMs and KBMs is unlikely to generate the required peaking without a central source of particle fuelling.

To quantify the level of transport driven from each of these different modes requires nonlinear simulations. These are required to test and develop reduced physics models that are capable of describing the transport generated by the various instabilities contributing to the nonlinear turbulence. Nonlinear calculations are also required to improve our understanding as to how experimental actuators will influence transport in high β STs. This will identify the modes that contribute most strongly to transport, and may provide insights in how this turbulence can be controlled.

7. Acknowledgements

The author would like to thank Francis Casson, Walter Guttenfelder, Emily Belli and Jeff Candy for useful discussions that assisted this work.

The views and opinions expressed herein do not necessarily reflect those of the European Commission. This work was supported by the Engineering and Physical Sciences Research Council [EP/L01663X/1, EP/R034737/1, EP/T012250/1]. This work used the ARCHER UK National Supercomputing Service (<http://www.archer.ac.uk>). We acknowledge the CINECA award under the ISCRA initiative, for the availability of high performance computing resources and support. This research used resources of the National Energy Research Scientific Computing Center (NERSC), a U.S. Department of Energy Office of Science User Facility located at Lawrence Berkeley National Laboratory, operated under Contract No. DE-AC02-05CH11231.

Appendix A. Cross code validation

Given the highly exotic nature of this equilibrium, the predictions made by GS2 were compared to CGYRO to ensure that these modes can be consistently found across codes. CGYRO is currently unable to force an odd or even eigenmode so only the dominant instability can be examined. The results for the $\rho_\psi = 0.5$ surface outlined in Table 2 are shown in Figure A1.




Figure A1: Comparison of the dominant eigenvalue predictions made by GS2 and CGYRO of the equilibrium outlined in Table 2.

It can be seen that both codes are in good agreement for this case, with CGYRO finding these MTMs with the extended mode structure, which indicates that the gyrokinetic equation is being correctly solved and that these modes are likely to appear in high β STs like STEP.

Appendix B. Stability of electron scale region

All the equilibria examined so far were found to be stable when $k_y \rho_e \sim \mathcal{O}(1)$. The STPP reactor design was also found to be stable to electron scale modes [12]. ETG-like instabilities are generally found at the electron scale in other tokamaks. To understand why they are stable here, different plasma parameters were scanned as done in the Section 3.2. This section examines the un-optimised case in Table 2 for $\rho_\psi = 0.5$.

Appendix B.1. Impact of kinetic profiles

Given that ETG is expected, the impact of a/L_{Te} was examined. Scans were for $a/L_{Te} = 5.0, 6.0 \& 7.0$ for $k_y \rho_s = 15 \rightarrow 70$. The equilibrium value of $a/L_{Te} = 2.77$ is well below this temperature gradient range, illustrating the large increase in drive necessary to find any ETG. Figure B1a shows how an unstable mode can be found if the drive is sufficiently high. It can be seen that increasing a/L_{Te} does drive the mode seen here unstable, consistent with ETG. This shows that a significant increase in temperature gradient would be necessary for this region of k_y -space to be unstable. The eigenfunction for $k_y \rho_s = 35$ at $a/L_{Te} = 7.0$ is shown in Figure B1b. The $A_{||}$ component is very small when normalised to ϕ suggesting it is not significant for these ETG modes.

To examine why the critical a/L_{Te} is high for this equilibrium other parameters




Figure B1: a) Growth rate spectrum for high values of k_y at different a/L_{Te} . The equilibrium value was $a/L_{Te} = 2.77$. b) Eigenfunctions for $k_y \rho_s = 35$ when $a/L_{Te} = 7.0$.

will be examined. The stability at $k_y \rho_s = 35$ will be investigated more carefully.

Appendix B.2. Impact of magnetic equilibrium

Appendix B.2.1. Impact of $\beta_{e,unit}$ and $\beta'_{e,unit}$ Earlier it was shown that an ETG appeared when reducing $\beta_{e,unit}$ and $\beta'_{e,unit}$. As they were both reduced, the high k_y MTM was stabilised and an ETG was destabilised, so it is expected that they will have an impact on the stability.

Figure B2a shows how the growth rate changes for the 3 different temperature gradients when changing $\beta_{e,unit}$. $\beta_{e,unit}$ is found to be destabilising to these high gradient ETG modes, up to a critical value $\beta_{e,unit} = 0.02$, when there is a rollover and at sufficiently high $\beta_{e,unit}$ these ETG are stabilised. However, at the equilibrium value of a/L_{Te} , the mode was completely stable regardless of the $\beta_{e,unit}$. A similar scan was conducted with $\beta'_{e,unit}$ and this appears to be a cause of the stability. Increasing $\beta'_{e,unit}$ strongly stabilises the ETG mode, such that when $\beta'_{e,unit} = 0$ the ETG mode is unstable even at the equilibrium a/L_{Te} . This has been seen before by Roach *et al* [56] and is attributed to high β' causing drift reversal at the outboard mid-plane. This all indicates a potential disadvantage with operating at higher field. The resulting lower $\beta'_{e,unit}$ could destabilise the ETG modes and have a detrimental effect on electron thermal transport.

Appendix B.2.2. Safety factor profile A critical gradient formula for ETG-like turbulence was found by Jenko *et al* [57] as follows

$$\left(\frac{a}{L_{Te}}\right)_{crit}^{ETG} \propto \left(1.3 + 1.9 \frac{\hat{s}}{q}\right) \quad (B.1)$$

This was generated from simulations of a low β , conventional aspect ratio tokamak. To examine its validity for this ST equilibrium, scans were performed in a/L_{Te} at




Figure B2: Impact of a) $\beta_{e,\text{unit}}$ and b) $\beta'_{e,\text{unit}}$ at different a/L_{Te} with $k_y\rho_s = 35$.

different values of q and \hat{s} to find the critical gradient. Equation B.1 suggests that the critical gradient will increase with \hat{s}/q but Figures B3a and B3b indicate this is not the case here. Performing a linear fit to these growth rates, an estimate of the critical gradient can be made and this is shown in Figure B4. It is clear that the scaling of Equation B.1 does not describe this regime. Rather, operating at low \hat{s}/q is actually beneficial to these modes.




Figure B3: Impact of a) q and b) \hat{s} with different a/L_{Te} at $k_y\rho_s = 35$.

This indicates that the scaling laws developed by Jenko are not applicable in this regime such that they cannot be utilised. Though this may not be of a major concern as the ETG was found to be stable for this equilibrium as well as the optimised equilibrium.

Appendix B.3. Higher field device

A scan was done looking at the impact of the higher I_{rod} devices on these ETG, as was conducted in Section 5.1. When examining the impact on the higher gradient cases at




Figure B4: Critical a/L_{Te} threshold for various a) q and b) \hat{s} at $k_y \rho_s = 35$.

$k_y \rho_s = 35$, Figure B5 shows that operating at higher field does destabilise these ETG modes, indicating the destabilisation from $\beta'_{e,unit}$ is overcoming the stabilisation from lower $\beta_{e,unit}$ and higher q . Nevertheless, for $a/L_{Te} = 2.77$, the modes are stable, so for this case it is not a concern. But it does highlight that operating at higher field may destabilise ETG modes.




Figure B5: Examining the impact of I_{rod} on the ETG modes for 3 higher temperature gradients at $k_y \rho_s = 35$. Note the reference equilibrium value is $a/L_{Te} = 2.77$.

References

- [1] EDA ITER et al. “Plasma confinement and transport”. In: *Nuclear Fusion* 39.12 ITER physics basis (1999), pp. 2175–2249.

- [2] CC Petty. “Sizing up plasmas using dimensionless parameters”. In: *Physics of Plasmas* 15.8 (2008), p. 781.
- [3] TL Rhodes et al. “L-mode validation studies of gyrokinetic turbulence simulations via multiscale and multifield turbulence measurements on the DIII-D tokamak”. In: *Nuclear Fusion* 51.6 (2011), p. 063022.
- [4] F Van Wyk. “Subcritical turbulence in the mega ampere spherical tokamak”. In: *arXiv:1703.03397* (2017).
- [5] NT Howard et al. “Quantitative comparison of experimental impurity transport with nonlinear gyrokinetic simulation in an Alcator C-Mod L-mode plasma”. In: *Nuclear Fusion* 52.6 (2012), p. 063002.
- [6] HR Wilson et al. “STEP - on the pathway to fusion commercialization”. In: *Commercialising Fusion Energy*. 2053-2563. IOP Publishing, 2020, 8-1 to 8–18.
- [7] HR Wilson. *SCENE-simulation of self-consistent equilibria with neoclassical effects*. Tech. rep. UKAEA Government Division, 1994.
- [8] Bhavin Patel. “Confinement physics for a steady state net electric burning spherical tokamak”. PhD thesis. University of York, 2021.
- [9] Michael Barnes et al. *GS2*. Feb. 2021. DOI: 10.5281/zenodo.4461680. URL: <https://bitbucket.org/gyrokinetics/gs2/>.
- [10] JE Menard et al. “Fusion nuclear science facilities and pilot plants based on the spherical tokamak”. In: *Nuclear Fusion* 56.10 (2016), p. 106023.
- [11] RJ Goldston, ML Reinke, and JA Schwartz. “A new scaling for divertor detachment”. In: *Plasma Physics and Controlled Fusion* 59.5 (2017), p. 055015.
- [12] HR Wilson et al. “Integrated plasma physics modelling for the Culham steady state spherical tokamak fusion power plant”. In: *Nuclear fusion* 44.8 (2004), p. 917.
- [13] GM Voss et al. “Toroidal field coil design for the spherical tokamak power plant”. In: *Fusion engineering and design* 48.3-4 (2000), pp. 407–418.
- [14] S Saarelma et al. “Integrated modelling of H-mode pedestal and confinement in JET-ILW”. In: *Plasma Physics and Controlled Fusion* 60.1 (2017), p. 014042.
- [15] A Diallo et al. “Dynamical evolution of pedestal parameters in ELMy H-mode in the National Spherical Torus Experiment”. In: *Nuclear Fusion* 51.10 (2011), p. 103031.
- [16] PB Snyder et al. “Pedestal stability comparison and ITER pedestal prediction”. In: *Nuclear Fusion* 49.8 (2009), p. 085035.
- [17] A Pankin et al. “The tokamak Monte Carlo fast ion module NUBEAM in the National Transport Code Collaboration library”. In: *Computer Physics Communications* 159.3 (2004), pp. 157–184.
- [18] JE Menard et al. “Physics design of the NSTX upgrade”. In: *37th European Physical Society Conference on Plasma Physics, Dublin, Ireland, online: <http://ocs.ciemat.es/EPS2010PAP/pdf>* P. Vol. 2. 2010.

- [19] RL Miller et al. “Noncircular, finite aspect ratio, local equilibrium model”. In: *Physics of Plasmas* 5.4 (1998), pp. 973–978.
- [20] J Chowdhury and BF McMillan. “Low n electromagnetic modes in spherical tokamaks”. In: *Plasma Physics and Controlled Fusion* 63.7 (2021), p. 075017.
- [21] Neeraj Kumar et al. “Turbulent transport driven by kinetic ballooning modes in the inner core of JET hybrid H-modes”. In: *Nuclear Fusion* 61.3 (2021), p. 036005.
- [22] DJ Applegate et al. “Microstability in a “MAST-like” high confinement mode spherical tokamak equilibrium”. In: *Physics of plasmas* 11.11 (2004), pp. 5085–5094.
- [23] J Candy, EA Belli, and RV Bravenec. “A high-accuracy Eulerian gyrokinetic solver for collisional plasmas”. In: *Journal of Computational Physics* 324 (2016), pp. 73–93.
- [24] DR Hatch et al. “Microtearing turbulence limiting the JET-ILW pedestal”. In: *Nuclear Fusion* 56.10 (2016), p. 104003.
- [25] X Jian et al. “Role of Microtearing Turbulence in DIII-D High Bootstrap Current Fraction Plasmas”. In: *Physical Review Letters* 123.22 (2019), p. 225002.
- [26] D Dickinson et al. “Kinetic instabilities that limit β in the edge of a tokamak plasma: a picture of an H-mode pedestal”. In: *Physical Review Letters* 108.13 (2012), p. 135002.
- [27] W Guttenfelder et al. “Progress in simulating turbulent electron thermal transport in NSTX”. In: *Nuclear Fusion* 53.9 (2013), p. 093022.
- [28] D Dickinson et al. “Towards the construction of a model to describe the inter-ELM evolution of the pedestal on MAST”. In: *Plasma physics and controlled fusion* 53.11 (2011), p. 115010.
- [29] DR Hatch. *Mode analyses of gyrokinetic simulations of plasma microturbulence*. 2010.
- [30] A Ishizawa et al. “Electromagnetic gyrokinetic simulation of turbulence in torus plasmas”. In: *Journal of Plasma Physics* 81.2 (2015).
- [31] AB Hassam. “Fluid theory of tearing instabilities”. In: *The Physics of Fluids* 23.12 (1980), pp. 2493–2497.
- [32] JF Drake and YC Lee. “Kinetic theory of tearing instabilities”. In: *The Physics of Fluids* 20.8 (1977), pp. 1341–1353.
- [33] PJ Catto and MN Rosenbluth. “Trapped electron modifications to tearing modes in the low collision frequency limit”. In: *The Physics of Fluids* 24.2 (1981), pp. 243–255.
- [34] DJ Applegate et al. “Micro-tearing modes in the mega ampere spherical tokamak”. In: *Plasma Physics and Controlled Fusion* 49.8 (2007), p. 1113.
- [35] D Dickinson et al. “Microtearing modes at the top of the pedestal”. In: *Plasma Physics and Controlled Fusion* 55.7 (2013), p. 074006.

- [36] CM Roach et al. “Confinement in START beam heated discharges”. In: *Nuclear fusion* 41.1 (2001), p. 11.
- [37] W Guttenfelder et al. “Electromagnetic transport from microtearing mode turbulence”. In: *Physical review letters* 106.15 (2011), p. 155004.
- [38] SM Kaye et al. “Confinement and local transport in the National Spherical Torus Experiment (NSTX)”. In: *Nuclear Fusion* 47.7 (2007), p. 499.
- [39] W Guttenfelder et al. “Scaling of linear microtearing stability for a high collisionality National Spherical Torus Experiment discharge”. In: *Physics of Plasmas* 19.2 (2012), p. 022506.
- [40] T Rafiq et al. “Microtearing modes in tokamak discharges”. In: *Physics of Plasmas* 23.6 (2016), p. 062507.
- [41] EA Belli and J Candy. “Fully electromagnetic gyrokinetic eigenmode analysis of high-beta shaped plasmas”. In: *Physics of Plasmas* 17.11 (2010), p. 112314.
- [42] DR Hatch et al. “A gyrokinetic perspective on the JET-ILW pedestal”. In: *Nuclear Fusion* 57.3 (2017), p. 036020.
- [43] AK Swamy et al. “Collisionless microtearing modes in hot tokamaks: Effect of trapped electrons”. In: *Physics of Plasmas* 22.7 (2015), p. 072512.
- [44] Chen Geng, David Dickinson, and Howard Wilson. “The physics of a small-scale tearing mode in collisionless slab plasmas”. In: *Plasma Physics and Controlled Fusion* 62.8 (2020), p. 085009.
- [45] EG Highcock. “The zero turbulence manifold in fusion plasmas”. In: *arXiv preprint arXiv:1207.4419* (2012).
- [46] CM Roach et al. “Gyrokinetic simulations of spherical tokamaks”. In: *Plasma Physics and Controlled Fusion* 51.12 (2009), p. 124020.
- [47] M. R. Hardman et al. *Extended electron tails in electrostatic microinstabilities and the nonadiabatic response of passing electrons*. 2021. arXiv: 2108.02822 [physics.plasm-ph].
- [48] Jason F Parisi et al. “Toroidal and slab ETG instability dominance in the linear spectrum of JET-ILW pedestals”. In: *Nuclear Fusion* 60.12 (2020), p. 126045.
- [49] HS Xie, ZX Lu, and B Li. “Kinetic ballooning mode under steep gradient: High order eigenstates and mode structure parity transition”. In: *Physics of Plasmas* 25.7 (2018), p. 072106.
- [50] M Valović et al. “Collisionality and safety factor scalings of H-mode energy transport in the MAST spherical tokamak”. In: *Nuclear Fusion* 51.7 (2011), p. 073045.
- [51] SM Kaye et al. “The dependence of H-mode energy confinement and transport on collisionality in NSTX”. In: *Nuclear Fusion* 53.6 (2013), p. 063005.
- [52] I Predebon and F Sattin. “On the linear stability of collisionless microtearing modes”. In: *Physics of Plasmas* 20.4 (2013), p. 040701.

- [53] RJ Groebner et al. “Limits to the H-mode pedestal pressure gradient in DIII-D”. In: *Nuclear fusion* 50.6 (2010), p. 064002.
- [54] JW Connor, RJ Hastie, and John Bryan Taylor. “High mode number stability of an axisymmetric toroidal plasma”. In: *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences* 365.1720 (1979), pp. 1–17.
- [55] T Rafiq et al. “Microtearing instabilities and electron thermal transport in low and high collisionality NSTX discharges”. In: *Physics of Plasmas* 28.2 (2021), p. 022504.
- [56] CM Roach, JW Connor, and S Janjua. “Trapped particle precession in advanced tokamaks”. In: *Plasma physics and controlled fusion* 37.6 (1995), p. 679.
- [57] F Jenko, W Dorland, and GW Hammett. “Critical gradient formula for toroidal electron temperature gradient modes”. In: *Physics of Plasmas* 8.9 (2001), pp. 4096–4104.