



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/179947/>

Version: Submitted Version

---

**Preprint:**

Chandra, V., Hwang, H.-C., Zakamska, N.L. et al. (Submitted: 2021) The SNIa runaway LP 398-9: detection of circumstellar material and surface rotation. [Preprint - arXiv] (Submitted)

<https://doi.org/10.48550/arXiv.2110.06935>

---

© 2021 The Author(s). For reuse permissions, please contact the Author(s).

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

## The SN Ia Runaway LP 398-9: Detection of Circumstellar Material and Surface Rotation

VEDANT CHANDRA,<sup>1,2</sup> HSIANG-CHIH HWANG,<sup>3,2</sup> NADIA L. ZAKAMSKA,<sup>2,3</sup> SIMON BLOUIN,<sup>4,5</sup> ANDREW SWAN,<sup>6</sup>  
THOMAS R. MARSH,<sup>7</sup> KEN J. SHEN,<sup>8</sup> BORIS T. GÄNSICKE,<sup>7</sup> J. J. HERMES,<sup>9</sup> ODELIA PUTTERMAN,<sup>9</sup> EVAN B. BAUER,<sup>1</sup>  
EVAN PETROSKY,<sup>2,10</sup> VIKRAM S. DHILLON,<sup>11,12</sup> STUART P. LITTLEFAIR,<sup>11</sup> AND RICHARD P. ASHLEY<sup>7,13</sup>

<sup>1</sup>Center for Astrophysics | Harvard & Smithsonian, 60 Garden St, Cambridge, MA 02138, USA

<sup>2</sup>Department of Physics & Astronomy, Johns Hopkins University, 3400 N Charles St, Baltimore, MD 21218, USA

<sup>3</sup>Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, USA

<sup>4</sup>Los Alamos National Laboratory, PO Box 1663, Los Alamos, NM 87545, USA

<sup>5</sup>Department of Physics and Astronomy, University of Victoria, Victoria, BC V8W 2Y2, Canada

<sup>6</sup>Department of Physics & Astronomy, University College London, Gower Street, London WC1E 6BT, UK

<sup>7</sup>Department of Physics, University of Warwick, Coventry, CV4 7AL, UK

<sup>8</sup>Department of Astronomy and Theoretical Astrophysics Center, University of California, Berkeley, CA 94720, USA

<sup>9</sup>Department of Astronomy & Institute for Astrophysical Research, Boston University, 725 Commonwealth Ave., Boston, MA 02215, USA

<sup>10</sup>Department of Physics, University of Michigan, 450 Church St, Ann Arbor, MI 48109, USA

<sup>11</sup>Department of Physics and Astronomy, University of Sheffield, Sheffield S3 7RH, UK

<sup>12</sup>Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain

<sup>13</sup>Isaac Newton Group of Telescopes, Apartado de Correos 321, Santa Cruz de La Palma, E-38700, Spain

### ABSTRACT

A promising progenitor scenario for Type Ia supernovae (SNeIa) is the thermonuclear detonation of a white dwarf in a close binary system with another white dwarf. After the primary star explodes, the surviving donor can be spontaneously released as a hypervelocity runaway. One such runaway donor candidate is LP 398-9, whose orbital trajectory traces back  $\approx 10^5$  years to a known supernova remnant. Here we report the discovery of carbon-rich circumstellar material around LP 398-9, revealed by a strong infrared excess and analyzed with follow-up spectroscopy. The circumstellar material is most plausibly composed of inflated layers from the star itself, mechanically and radioactively heated by the past companion's supernova. We also detect a 15.4 hr periodic signal in the UV and optical light curves of LP 398-9, which we interpret as surface rotation. The rotation rate is consistent with theoretical predictions from this supernova mechanism, and the brightness variations could originate from surface inhomogeneity deposited by the supernova itself. Our observations strengthen the case for this double-degenerate SNIa progenitor channel, and motivate the search for more runaway SNIa donors.

*Keywords:* White dwarf stars (1799), Type Ia supernovae (1728), Circumstellar dust (236), Stellar rotation (1629), Runaway stars (1417)

### 1. INTRODUCTION

Type Ia supernovae (SNeIa) are luminous transients that are valuable standard candles to measure cosmological parameters (e.g., Riess et al. 1998; Perlmutter et al. 1999; Riess et al. 2019), and play a crucial role in the chemical evolution of stellar populations (e.g., Tinsley 1979; Matteucci et al. 2009; Kirby et al. 2019). Despite the importance of SNeIa, their progenitor scenario is


uncertain, and is the subject of intense theoretical and observational efforts (e.g., Hillebrandt & Niemeyer 2000; Maoz et al. 2014). It is generally accepted that SNeIa originate from massive white dwarfs (WDs) that accrete matter from binary companions, but the nature and fate of the donor companion is unknown.

For several decades, the leading hypothesis was the ‘single degenerate’ scenario (e.g., Whelan & Iben 1973), in which a carbon–oxygen core (C/O) WD accretes matter from a nondegenerate star until it approaches the Chandrasekhar limit (Chandrasekhar 1931). However, it is challenging for the accreting WD to gain mass in the

first place (e.g., Nomoto 1982; Iben & Tutukov 1984), and there is a growing body of observational evidence that suggests most SNe Ia cannot originate from single-degenerate progenitors (e.g., Kerzendorf et al. 2009; Kasen 2010; Li et al. 2011; Bloom et al. 2012; Margutti et al. 2012; Woods et al. 2017). An alternative is the ‘double degenerate’ scenario in which two WDs could either merge and consequently explode (e.g., Iben & Tutukov 1984; Webbink 1984), or have one WD detonate after mass transfer (e.g., Bildsten et al. 2007). The explosion mechanism can be a double-detonation, during which a detonation in the outer helium shell induces a second detonation in the carbon core that is powerful enough to unbind the star (Taam 1980; Guillochon et al. 2010; Dan et al. 2011; Raskin et al. 2012; Pakmor et al. 2013; Shen & Bildsten 2014).

In the double-degenerate case, a double-detonation can occur during the mass transfer phase, well before the two WDs merge. In this ‘dynamically driven double-degenerate double-detonation’ ( $D^6$ ) scenario, the primary WD can explode well below the Chandrasekhar limit (Pakmor et al. 2013; Shen et al. 2018a,b; Tanikawa et al. 2018, 2019). Observational studies have independently found that a significant fraction of SNe Ia could originate from sub-Chandrasekhar mass WDs (Scalzo et al. 2014; Dhawan et al. 2017; Kirby et al. 2019; de los Reyes et al. 2020; Sanders et al. 2021). Recent theoretical work has shown that the  $D^6$  mechanism can reproduce the observational signatures of most SNIa (Shen et al. 2021a,b). In this scenario, the donor WD could survive the explosion of the primary if it occurs in the early stages of mass transfer (Shen & Schwab 2017). The binary orbit becomes spontaneously unbound after the supernova, ejecting the donor WD at the orbital velocity,  $\gtrsim 1000 \text{ km s}^{-1}$ .

Shen et al. (2018b) discovered three candidates that appear to be ‘runaway’ WD donors ejected from SNe Ia that exploded via the  $D^6$  mechanism. These stars are among the fastest unbound stars in the Galaxy, with estimated space velocities  $\gtrsim 1000 \text{ km s}^{-1}$ . Their radii are inflated by an order of magnitude compared to typical WDs, likely due to deposited energy from the SNIa ejecta. All three candidates in Shen et al. (2018b) have similar low-resolution optical spectra, with absorption signatures of carbon, oxygen, magnesium, and calcium. One of these candidates is LP 398-9, referred to as ‘D6-2’ in Shen et al. (2018b). Remarkably, LP 398-9’s inferred orbital trajectory extrapolated back  $9 \times 10^4$  years matches the on-sky position and distance of a known supernova remnant G70.0-21.5, strongly supporting its  $D^6$  origin (Fesen et al. 2015; Raymond et al. 2020).



**Figure 1.** Top: Location of LP 398-9 on the *Gaia* EDR3 color-magnitude diagram. The background sample consists of stars within 100pc of the Sun (Smart et al. 2021). Bottom: The *Gaia*-*WISE* color-color space, with the same background sample of nearby stars. LP 398-9 has a significant excess in the  $W1-W2$  color compared to other stars of a similar optical color. For comparison, we show the other two  $D^6$  candidates from Shen et al. (2018b). ‘D6-3’ does not have secure *WISE* photometry due to a crowded field, and is consequently absent from the bottom panel.

Here we present follow-up observations of LP 398-9 that reveal the presence of significant quantities of circumstellar material, as well as a 15.4 hr photometric period. We argue that both of these observables can be linked to the  $D^6$  origin of the system, strengthening the case for this SNIa progenitor channel. After summarizing our data and observations in §2, we present our analysis in §3. We present our results in §4, and discuss our findings in §5.

## 2. DATA


In this section we describe the archival data we collected for LP 398-9, as well as our own follow-up observations. In §2.1 we assemble the spectral energy distribution (SED) from archival data. In §2.2 we describe our light curves and follow-up photometry, and in §2.3 we describe our follow-up spectroscopy.

### 2.1. Spectral Energy Distribution

LP 398-9 has reliable archival photometry in the *GALEX* *NUV* (Martin et al. 2005; Million et al. 2016), Sloan *ugriz* (Fukugita et al. 1996; Gunn et al. 1998; Doi et al. 2010; Blanton et al. 2017; Ahumada et al. 2020), 2MASS *JH* (Skrutskie et al. 2006), and *WISE* *W1*, *W2*

(Wright et al. 2010; Mainzer et al. 2011; Mainzer et al. 2014) bands. The detections in *GALEX* *FUV*, 2MASS *Ks*, and *WISE* *W3*, *W4* are unreliable or absent. LP 398-9 also has secure astrometry from the *Gaia* space observatory Early Data Release 3 (EDR3; Prusti et al. 2016; Mignard et al. 2018; Brown et al. 2021; Lindegren et al. 2021), with  $\varpi/\sigma_\varpi \sim 18$ . We correct the observed photometry for interstellar extinction along the line of sight using 3D dust maps (Green 2018; Green et al. 2015, 2018a,b) queried at the *Gaia* inverse-parallax distance of 840 pc. We adopt the extinction law of Fitzpatrick & Massa (2007) with  $R_V = 3.1$ . The *Gaia* color-magnitude diagram and *Gaia*-*WISE* color-color diagram of LP 398-9 are shown in Figure 1. The lower panel shows LP 398-9’s infrared excess compared to other stars at a similar optical color. We confirmed that the other two D<sup>6</sup> candidates from Shen et al. (2018b) do not have a measurable IR excess: D6-1 has no *W1* – *W2* color excess, and D6-3 is in a crowded field and consequently does not have secure *WISE* photometry.

One potential contaminant of *WISE* imaging is source confusion due to the relatively coarse angular resolution of *WISE* (see e.g., Denny et al. 2020). In Figure 2 we compare the *J*-band image from 2MASS to the AllWISE image in channel *W1*. In the 2MASS image, there is no discernible background source within 6 arcseconds of LP 398-9. Furthermore, because the *WISE* data of LP 398-9 appears time-variable (§3.3), we exclude the possibility of a background blazar by finding no radio counterpart in the NRAO VLA Sky Survey (NVSS; Condon et al. 1998). Another method to verify the association between the *WISE* data and LP 398-9 is by comparing optical astrometry to the source position measured by *WISE*. In Figure 2 we display time-averaged measurements of the position of LP 398-9 on the sky as measured over 5 years of the NEOWISE mission (2014-2019), along with the reference position from AllWISE (2011). For additional details about the *WISE* astrometric solution, we defer to Cutri et al. (2012). We perform a trimmed least-squares linear regression (Rousseeuw & Van Driessen 2006; Cappellari et al. 2013) on the RA and Dec as a function of time to estimate the *WISE* proper motion. The parallax motion of  $\sim 1$  mas (known from *Gaia*) is negligible here. We obtain  $\Delta RA_{\text{WISE}} = 80 \pm 30$  mas/yr and  $\Delta Dec_{\text{WISE}} = 250 \pm 30$  mas/yr. This is consistent within  $1-\sigma$  with the *Gaia* EDR3 proper motion of LP 398-9 (Table 1). To ensure our method is robust, we compute the *WISE* proper motions of some nearby objects in the field and verify that they too match their *Gaia* data within uncertainties. This astrometric test confirms that the *WISE* photometric source is securely associated with LP 398-9.



**Figure 2.** Top left: 2MASS *J* band image from  $\approx 2001$ . Top right: AllWISE channel *W1* image ten years later, from  $\approx 2011$ . The centroid has clearly shifted by several arcseconds towards the NE direction. Bottom: Position of LP 398-9 on the sky during the course of 5 years of the NEOWISE mission. Each datapoint is the average of 20 individual measurements. The arrow indicates 1 year of the *Gaia* EDR3 proper motion, centred on the AllWISE coordinate. There is no bright contaminant within 6 arcseconds and the *WISE* proper motion is consistent with the optical proper motion from *Gaia*, supporting the fact that the IR excess and variability comes from LP 398-9 itself.

## 2.2. Light Curves

In addition to the time-averaged catalog measurements from AllWISE, we utilize individual exposures taken over the past nine years. We assemble the *WISE* light curve of LP 398-9 by querying the AllWISE (Wright et al. 2010) and NEOWISE (Mainzer et al. 2011; Mainzer et al. 2014) databases for single-exposure photometric measurements. There are 27 AllWISE datapoints from 2010 May 27 to 2010 November 10, and 178 NEOWISE datapoints from 2014 May 31 to 2019 November 10. We perform quality cuts to remove spurious detections with outlying RA/Dec offsets and undefined magnitude uncertainties. The resulting light curve has 168 datapoints over a 9 year baseline. The cadence of the observations is irregular and follows the *WISE* position-dependent scanning strategy, with a  $\approx 180$  day gap between successive observing runs. Each

run is  $\approx 1.5$  days long and consists of exposures taken  $\approx 1.5$  hours apart.

We queried archival photometry for LP 398-9 from the Zwicky Transient Facility (ZTF; Bellm et al. 2019; Masci et al. 2019). We assembled a light curve consisting of datapoints collected between 2020 January 13 and 2021 April 27 in the  $g$  and  $r$  photometric bands. We only used the most recent year of ZTF data to minimize the effect of long term brightness variations (discussed further in §3.3). To maximize our time coverage, we combined the  $g$  and  $r$  data after subtracting their respective median magnitudes (e.g., Burdge et al. 2020). We removed bad datapoints with ZTF quality flag `catflags`  $> 0$  and sharpness `sharp`  $> 0.5$  (e.g., Guidry et al. 2021). Our final ZTF  $gr$  lightcurve has 205 datapoints.

We observed LP 398-9 with HiPERCAM (Dhillon et al. 2016, 2018, 2021) on the 10.4-m Gran Telescopio Canarias (GTC) at the Observatorio del Roque de los Muchachos on the island of La Palma, Spain, on 2018 June 11 from 03:38 to 05:15 UTC. We utilized a customised high-throughput approximation of the Sloan  $ugriz$  filter set to obtain simultaneous 5-band photometry. A cadence of 1.222 sec was used for the  $g$ ,  $r$  and  $i$  arms, while every other readout was skipped on the  $u$  and  $z$  arms giving a 2.444 sec cadence. The weather was clear with seeing  $\sim 0.8''$ – $1.5''$ . The Moon was 80% illuminated. We debiased and flat-fielded the data using twilight sky flats. We performed comparative aperture photometry using apertures that tracked the target positions with radii set to 1.8 times the mean full width at half-maximum of the stellar profiles. We defer to Dhillon et al. (2021) for further details of the reduction procedures. We used *Gaia* EDR3 1798008691771518464 ( $G = 15.1$ ) as the main comparison star, and *Gaia* EDR3 1798020408442309248 ( $G = 16.5$ ) to confirm that the main comparison star was not itself variable. We also observed LP 398-9 with the Wide Field Camera on the 2.5-m Isaac Newton Telescope (INT), on 2018 August 2 from 00:17 to 04:44 UTC. We utilized a Sloan  $g$  filter with exposure times of 100 sec. We performed aperture photometry relative to a comparison star, and defer to Dhillon et al. (2021) for further details on the reduction procedures.

We observed LP 398-9 with the IO:O imager on the 2-m Liverpool Telescope (LT; Steele et al. 2004) at the Observatorio del Roque de los Muchachos, over the course of a month from 2021 July 8 to 2021 August 8. We obtained a total of 154 images in the Bessel  $B$  band (effective wavelength  $\simeq 4450 \text{ \AA}$ ) with an exposure time of 60 s, averaging approximately five images per night. We performed comparative aperture photometry against

the nearby star *Gaia* EDR3 1798008721834715648 ( $G = 13.8$ ).

We also analyzed observations of LP 398-9 obtained with the Space Telescope Imaging Spectrograph (STIS; Woodgate et al. 1998) of the *Hubble Space Telescope* (*HST*) over six consecutive orbits on 2020 October 3 (PI: Shen). The spectra are flux-calibrated and cover a wavelength range of 1600–3600  $\text{\AA}$  with the G230L grating. We use these spectra to derive an ultraviolet light curve that is analyzed in this paper. A full analysis of the STIS spectrum will be presented in a forthcoming publication. We took advantage of the photon-counting TIME-TAG mode to re-bin the spectra into 9-minute exposure times, and computed the average flux in the 1700–3100  $\text{\AA}$  range (e.g., Hermes et al. 2021). The resulting light curve has an effective central wavelength of 2520  $\text{\AA}$ , and contains 28 datapoints over  $\approx 8$  hours.

### 2.3. Spectroscopy

Shen et al. (2018b) obtained a low-resolution spectrum ( $R \approx 300$ ) of LP 398-9 from 3400 – 9000  $\text{\AA}$  with the Nordic Optical Telescope (NOT). They noted absorption lines indicating the presence of carbon, oxygen, magnesium, and calcium in the stellar photosphere, and measured a radial velocity  $20 \pm 60 \text{ km s}^{-1}$ .

We obtained a mid-resolution spectrum of LP 398-9 on 2020 November 14 using the Dual-Imaging Spectrograph (DIS) on the 3.5-m telescope at the Apache Point Observatory (PI: Chandra). We employed the B1200/R1200 gratings and a 1.5 arcsec slit width, providing a spectral resolution (Gaussian  $\sigma$ ) of 2  $\text{\AA}$  on the blue end and 0.5  $\text{\AA}$  on the red end, as estimated with sky emission lines. We obtained four 10-minute exposures bracketed by arc lamp exposures to ensure a reliable wavelength calibration. Our data covered a wavelength range of 4000–7000  $\text{\AA}$ , with a dichroic gap between 5200–6000  $\text{\AA}$ . We performed a standard data reduction—bias correction, flat-fielding, aperture extraction, and wavelength calibration—using the pipeline tools in IRAF (Tody 1986).

We compared the O I absorption line at  $\lambda 7002.23 \text{ \AA}$  with its theoretical rest wavelength to derive a radial velocity  $80 \pm 10 \text{ km s}^{-1}$ , which is consistent with the earlier measurement made by Shen et al. (2018b). We also used the  $\lambda 6300 \text{ \AA}$  O I sky emission line to verify that our absolute wavelength calibration is accurate to within  $5 \text{ km s}^{-1}$ . We therefore do not detect radial velocity variations that would indicate the presence of a close binary companion. As noted by Shen et al. (2018b), the low radial velocity is somewhat unexpected considering LP 398-9’s hypervelocity space motion, but could be a

statistical fluctuation due to small numbers and selection effects.

### 3. ANALYSIS

In this section we analyze our spectro-photometric observations of LP 398-9. We confirm the hypervelocity nature of LP 398-9 with the latest data from Gaia (§3.1). We model the spectral energy distribution of LP 398-9 and its excess infrared emission to derive the system’s parameters (§3.2). We present multi-band light curves that exhibit a 15.4 hr period, which we interpret as surface rotation (§3.3). Finally, we present follow-up spectroscopy that reveals the presence of circumstellar carbon (§3.4).

#### 3.1. Hypervelocity

We confirm the hypervelocity status of LP 398-9 using newly released astrometric data from Gaia EDR3 (Brown et al. 2021; Lindegren et al. 2021). Previously, Shen et al. (2018b) used data from Gaia DR2 to find a 99% credible interval of  $v_T = [700, 1500]$  km s<sup>-1</sup> for the tangential proper motion velocity of LP 398-9. We use the affine-invariant Markov Chain Monte Carlo (MCMC) sampler `emcee` (Foreman-Mackey et al. 2013; Foreman-Mackey et al. 2019) to sample the new astrometric data from Gaia EDR3 and consequently derive an updated posterior distribution for  $v_T$ . For the parallax, we employ the exponentially decreasing space density prior described in Bailer-Jones (2015), although its effect is likely minimal since the EDR3 parallax is quite precise ( $\varpi/\sigma_\varpi \sim 18$ ). We find a median tangential velocity  $v_T = 1030 \pm 60$  km s<sup>-1</sup>, with a 99% credible interval of  $v_T = [900, 1160]$  km s<sup>-1</sup>. This secures LP 398-9’s status as a hypervelocity star, despite its low apparent radial velocity.

#### 3.2. SED Model and IR Excess

To investigate the infrared excess of LP 398-9, we first fit a stellar model to the observed photometry at optical wavelengths, and then compare the predicted stellar flux in the infrared to the observed IR photometry. We fit the SED of LP 398-9 in the SDSS *ugriz* passbands using a bespoke grid of model spectra computed with the atmosphere code described in Blouin et al. (2018a,b). Since the non-detection of hydrogen lines in the spectrum of LP 398-9 rules out the presence of atmospheric hydrogen, we assume a helium-dominated atmosphere. The SED can also be significantly altered by the presence of metals due to line blanketing and changes in the atmospheric opacity. We therefore use the NOT spectrum of LP 398-9 observed by Shen et al. (2018b) to approximately constrain the metallicity during the fitting process. We adopt a surface gravity  $\log g = 5.5$

when computing the model spectra, and verify that neither the photometric nor spectroscopic fits are sensitive to this assumption.

We assume flat priors on the stellar parameters:  $2500 \leq T_{\text{eff}}/K \leq 10000$  and  $0.05 \leq R/R_\odot \leq 1$ . We adopt a Gaussian prior on the parallax  $\varpi/\text{mas}$  with a mean and standard deviation defined by the Gaia EDR3 astrometric measurement  $\pi = 1.19 \pm 0.06$  mas. To prevent any single band with underestimated uncertainties from dominating the fit, we implement a floor uncertainty of 0.03 magnitudes in all bands (e.g., Bergeron et al. 2019). We define a  $\chi^2$  log-likelihood (the  $\chi^2$  statistic multiplied by  $-0.5$ ) to compare the model fluxes to the observed fluxes and perform a preliminary fitting step to estimate the atmospheric metallicity of LP 398-9. First, we maximize the photometric  $\chi^2$  likelihood over the stellar parameters ( $T_{\text{eff}}$  and  $R$ ) assuming a negligible metallicity. Next, we vary the metallicity to match the Ca II H and K absorption lines on the archival NOT spectrum from Shen et al. (2018b), with other elements scaled accordingly to solar proportions. We repeat these steps until the stellar parameters and metallicity converge.

For the main fitting step, we fix the models to the best-fitting metallicity from the preliminary fitting step, and use the preliminary stellar parameters as initial values. We sample the posterior distributions of  $T_{\text{eff}}$ ,  $R$ , and  $\varpi$  using `emcee`. The stellar parameters are summarized in Table 1, along with uncertainties computed by taking the standard deviation of the respective MCMC samples. We adopt the posterior sample with the highest log-likelihood as our best-fitting stellar parameters for LP 398-9:  $T_{\text{eff}} = 7500 \pm 100$  K,  $R = 0.20 \pm 0.01 R_\odot$ .

We additionally repeated our entire analysis assuming oxygen-dominated and carbon-dominated atmospheres. The inferred  $T_{\text{eff}}$  and  $R$  are similar for the helium and oxygen atmospheres. A carbon-dominated atmosphere is ruled out, since it would require the presence of strong molecular C<sub>2</sub> absorption lines that are absent in our spectroscopic observations (discussed further in §3.4). The mass of LP 398-9 is quite uncertain, since the stellar structure is unknown and the star is in a temporary inflated state. However, given the D<sup>6</sup> origin of the system, the mass could plausibly lie in the 0.2–0.8  $M_\odot$  range (Shen 2015; Shen et al. 2018b). Assuming a uniform mass prior in this range, our fitted photometric radius implies  $\log g \approx 5.5 \pm 0.2$ . This is consistent with the Ca II H and K absorption lines on the archival NOT spectrum. We emphasize that the purpose of fitting the metallicity is to produce a self-consistent stellar model, not to precisely determine the metallicity of LP 398-9’s atmosphere. For example, the assumption of solar-scaled


**Table 1.** Parameters of LP 398-9

Parameter	Value
Gaia EDR3	
Source ID	1798008584396457088
RA (degrees)	324.61250
Dec. (degrees)	25.37374
G (mag)	16.97
BP–RP (mag)	0.41
$\varpi$ (mas)	$1.19 \pm 0.06$
Distance (pc)	$840 \pm 40$
$\mu_{\text{RA}}$ (mas/yr)	$98.28 \pm 0.07$
$\mu_{\text{DEC}}$ (mas/yr)	$240.18 \pm 0.06$
$v_T$ (km s $^{-1}$ )	$1030 \pm 60$
Stellar Model	
$T_{\text{eff}}$ (K)	$7500 \pm 100$
Radius ( $R_{\odot}$ )	$0.20 \pm 0.01$
Mass ( $M_{\odot}$ )	U [0.2, 0.8] (assumed)
$\log [g, \text{cm s}^{-2}]$	$5.5 \pm 0.2$
$v_R$ (km s $^{-1}$ )	$80 \pm 10$
Blackbody Infrared Model	
$T_{\text{bb}}$ (K)	$670 \pm 50$
$R_{\text{bb}}$ ( $R_{\odot}$ )	$5.4 \pm 0.9$
$M_{\text{dust}}$ ( $g$ )	$\sim 10^{20}$
Flat Disk Infrared Model ( $i = 0^{\circ}$ )	
$T_{\text{in}}$ (K)	$1300 \pm 200$
$T_{\text{out}}$ (K)	$\approx 300$

abundances is likely inappropriate in detail, but is sufficient for our purposes of solving the temperature and radius of LP 398-9.

To quantify and model the infrared excess around LP 398-9 we use the AllWISE catalog magnitudes in the *W1* and *W2* channels. We convert them to fluxes, and subtract the expected flux of LP 398-9 at these wavelengths using the stellar model described above. To this excess, we fit two models: a blackbody, and a flat disk model. The blackbody is parameterized by the temperature  $T_{\text{bb}}$  and a normalization factor that depends on the solid angle subtended on the sky. We use the *Gaia* EDR3 parallax of LP 398-9 to fix its distance, fitting for the remaining normalization parameter of  $R_{\text{bb}}$  in addition to  $T_{\text{bb}}$ .

For the disk model, we use the geometrically thin, optically thick debris disk model of Jura (2003). With a known distance, this model is further constrained by the



**Figure 3.** Spectral energy distribution of LP 398-9 with our best-fitting model for the WD and IR excess. The observed photometric bands are indicated on the top axis. The inset shows the 3890–4000 Å region of the archival NOT spectrum, with the best-fitting stellar model spectrum overlaid. Downward arrows indicate upper limits. Uncertainties are 1-sigma after applying our 0.03 mag floor, and in some cases are smaller than the marker size.

stellar temperature  $T_{\text{eff}}$  and radius  $R$ , with the free parameters being the two temperatures corresponding to the inner edge ( $T_{\text{in}}$ ) and outer edge ( $T_{\text{out}}$ ) of the disk. The inclination of the disk is difficult to constrain with the available observations, although it is probably low (more face-on) given the brightness of the IR excess compared to the star. Additionally, the temperature of the outer edge of the disk  $T_{\text{out}}$  is poorly constrained because observations at longer wavelengths are not available.

We fit the respective models using the nonlinear minimization algorithms in `scipy.optimize` (Virtanen et al. 2020). We repeat the fit on  $10^4$  Monte Carlo replicates of the data (adding the relevant Gaussian errors to the observed fluxes) to estimate the uncertainties on the parameters. The pure stellar, stellar + blackbody, and stellar + flat disk models are illustrated with the observed SED in Figure 3, and the best-fit IR model parameters are summarized in Table 1.

The blackbody temperature can be interpreted as the average temperature of an inflated, optically thin dust shell (Xu et al. 2018). Given the low temperature  $T_{\text{bb}} \approx 670$  K and large normalization radius  $R_{\text{bb}} \approx 5.4 R_{\odot}$ , we can rule out stellar or sub-stellar companions. A brown dwarf with  $T_{\text{eff}} \sim 700$  K would have a radius  $\sim 0.1 R_{\odot}$  (e.g., Sorahana et al. 2013), almost fifty times smaller


than the radius we derive from the normalization of the IR excess. Depending on the typical grain sizes,  $T_{\text{bb}} \approx 670$  K corresponds to dust located at orbital radii  $\approx 10$ – $30 R_{\odot}$ , or around 50–150 times the present radius of the star (Xu et al. 2018).

The flat disk model does not match the observed fluxes well, indicating that a geometrically thin, optically thick disk is insufficient to describe the material around LP 398-9. A low inclination ( $\approx 0^{\circ} - 30^{\circ}$ ) provides the best match to the *W1* and *W2* data. The inclination and inner disk temperature are degenerate – assumed inclinations of  $[0^{\circ}, 30^{\circ}, 60^{\circ}]$  result in best-fit  $T_{\text{in}} \approx [1300, 1600, 2000]$  K. Depending on the size and composition of the grains, circumstellar dust sublimates at temperatures between 1200–2000 K, and consequently any material inwards of the radius corresponding to  $T_{\text{in}}$  rapidly sublimates away, leaving a gap between the star and the debris disk (von Hippel et al. 2007; Rafikov & Garmilla 2012; Xu et al. 2018). Therefore, the inferred inner temperature  $T_{\text{in}} \approx 1300$  K is broadly consistent with dust sublimation temperatures, although the flat disk geometry seems unlikely. Our favored interpretation is a circumstellar dust shell model with a characteristic temperature of  $T_{\text{bb}} \approx 670$  K.

### 3.3. Periodic Photometric Variability

We searched the LT photometry of LP 398-9 for periodic variations using a Lomb-Scargle periodogram (Lomb 1976; Scargle 1982). We find a dominant peak at  $f = 1.55 \text{ day}^{-1}$ , corresponding to a period of  $\approx 15.4$  hours. We repeat our periodogram analysis on the ZTF photometry. Fitting the entire ZTF baseline from 2018 April 20 to 2021 April 27 yields a period 1% smaller than the LT period. However, this ZTF light curve has significant long-term brightness variations that contaminate the periodogram, requiring pre-whitening to remove trends. If we instead fit only the most recent ZTF campaign from 2020 January 6 to 2021 April 27 without pre-whitening, we derive a period within 0.03% of the LT period. We therefore use only the ZTF data from 2020 January 6 onwards in our subsequent analysis.


We adopt the peak of the LT periodogram as our assumed period. To estimate the uncertainty of this measurement, we computed 2500 Monte Carlo replicates of the LT light curve by adding Gaussian uncertainties, and computed half the difference between the 84th and 16th quantiles of the resulting period distribution. We derive a best-fit period  $P = 15.441 \pm 0.016$  hr. We phase-fold the STIS, LT, and ZTF light curves and display them in Figure 4. We fit a sinusoidal model to each phase-folded (un-binned) light curve to estimate the variability amplitudes. We find amplitudes of  $4.4 \pm 0.6\%$  for STIS,



**Figure 4.** Lomb-Scargle periodogram of the LT light curve of LP 398-9, along with the phase-folded HST-STIS, LT-*B*, and ZTF-*gr* light curves. The STIS datapoints are unbinned and consecutive from a single epoch, while the LT and ZTF datapoints are binned in phase. We overlay the sinusoidal models used to estimate the variability amplitude of each light curve.

$2.8 \pm 0.2\%$  for LT-*B*, and  $1.3 \pm 0.3\%$  for ZTF-*gr*. Figure 4 shows a slight color-dependent phase difference between the three datasets. However, simply changing the assumed period by as little as 0.3% (to  $P = 15.40317$ , for example) can erase this apparent phase difference. Therefore, the most plausible explanation is that this phase difference is non-physical, and is simply an artifact of systematic uncertainties in our photometry and period measurement.

Our 100-minute HiPERCAM *ugriz* observations are illustrated in Figure 5. The time baseline of this observation covers a fraction of our derived rotational period, and therefore represents a quasi-linear segment of the rotationally modulated flux variation. Both our HiPERCAM-*ugriz* and subsequent INT-*g* observations (not shown here) caught LP 398-9 in the dimming phase. We quantify the color dependence of the flux decline by fitting a simple linear regression to the light curves in each band using the trimmed least squares approach described in Cappellari et al. (2013). The rate of decline (slope) is strongly wavelength-dependent, with the




**Figure 5.** 100-minute HiPERCAM light curve of LP 398-9. The *ugriz* bands are arranged from top to bottom. We overlay a linear fit to each dataset with a black dashed line. The color-dependent trend is visually discernible, with the bluest *u* band declining twice as fast as the reddest *z* band. For each band we show the fitted slope of the flux decline in magnitudes per day.

bluest photometry declining twice as quickly as the reddest. The *g*-band slope is identical within uncertainties between the HiPERCAM and INT data. The trend of the HiPERCAM color-dependence (blue declining faster than red) is consistent with the trend in the variability amplitudes of the STIS, *LT-B*, and *ZTF-gr* light curves.

The color-dependence of the photometric variability could have several plausible explanations. A convective star spot with a temperature lower than the surrounding photosphere could produce such a signal, although it is challenging to constrain the temperature and size of the spot (e.g., Pelisoli et al. 2021). Alternatively, if a portion of the stellar surface is polluted with metals, there could be an opacity or line blanketing contrast as the star rotates. Additionally, light filtering through the circumstellar dust will be reddened by extinction.

As described in §2.1, we compiled a *WISE* light curve of LP 398-9 spanning 9 years, grouped into 14 approximately day-long observing runs spaced 180 days apart. Phase-folding the *WISE* light curve to the inferred stellar rotation period of 15.4 hr does not produce any coherent signal. This is expected, since the IR flux is dominated by emission from the surrounding circumstellar dust, which would greatly dilute any photometric variability from the star itself. However, the *WISE* data of LP 398-9 is overall more variable than the photometric uncertainties would indicate. We average the measurements in each observing run and illustrate the long-term *W1* and *W2* fluxes along with the *W1*–*W2* color in Figure 6. There is significant variation in both the *W1* and *W2* fluxes. The *W1*–*W2* color marginally




**Figure 6.** Long-term light curves from the duration of the *WISE* mission in the *W1* and *W2* bands, along with their difference. We average the individual measurements from each observing run to produce these datapoints. The insets show individual measurements from a single *WISE* observing run. The insets vertically span 1 magnitude, and horizontally span 2 days.

varies as well, potentially indicating a varying surface area or temperature of the circumstellar material. Recent evidence has shown that many WDs with debris disks are IR-variable (Swan et al. 2019, 2020). Those systems are thought to exhibit stochastic variation over long timescales due to collisional dust production and depletion (e.g., Kenyon & Bromley 2017a,b; Swan et al. 2021).

### 3.4. Spectroscopic Evidence of Circumstellar Carbon

In Figure 7 we illustrate the APO spectrum of LP 398-9, with several identified absorption lines of carbon, oxygen, magnesium, and calcium highlighted. We characterize the spectrum using theoretical models described in Blouin et al. (2018a,b). We fix the stellar parameters to the values computed from the SED fit in §3.2 and compute single-abundance spectra over a range of C and O to investigate the atmospheric abundances. We overlay in red a model spectrum with  $[C/He] = -5.0$ , and in blue a model with  $[O/He] = -1.0$ . The short-wavelength end of the spectrum demonstrates that this carbon abundance is already pushing the higher plausible limit, due to the development of strong  $C_2$  bands that are absent from our observed spectrum (Swan 1857; Johnson 1927). Yet, several narrow carbon lines remain unexplained by the stellar models (red arrows in Fig-



**Figure 7.** Mid-resolution spectrum of LP 398-9 from the 3.5-m telescope at the Apache Point Observatory, split into two wavelength regions. We indicate identified absorption lines of carbon, oxygen, magnesium, and calcium. We overlay in red a theoretical spectrum with the same stellar parameters as LP 398-9 and an atmospheric carbon abundance of  $[C/He] = -5.0$ , and independently in blue an oxygen abundance of  $[O/He] = -1.0$ . Red arrows indicate carbon absorption lines that cannot be explained by the stellar model, suggesting that they are formed in circumstellar material instead.

ure 7). The most plausible explanation for this non-photospheric absorption is the presence of circumstellar carbon around LP 398-9. Compared to the stellar photosphere, the circumstellar region has much lower pressures, and consequently contains less molecular  $C_2$ . This leaves more atomic C available to produce narrow absorption lines without creating large molecular bands on the spectrum.

## 4. RESULTS

### 4.1. Photometric Period

LP 398-9 exhibits sinusoidal photometric variation with a period  $P \approx 15.4$  hr. A stellar companion is quite unlikely, given the non-detection of radial velocity variations  $\gtrsim 10$  km s $^{-1}$ . The most natural explanation is that this photometric period corresponds to the rotational period of the star (e.g., Brinkworth et al. 2005; Hermes et al. 2021). There are several possible mech-

anisms that can produce rotationally modulated variability. WDs with exceptionally strong magnetic fields can have continuum opacity variations on their surfaces, producing photometric variations at the rotational period (e.g., Ferrario et al. 1997; Tremblay et al. 2015). However, the optical spectrum of LP 398-9 does not indicate a strong magnetic field, given the absence of visible Zeeman splitting. An alternative possibility is the formation of weakly magnetized starspots in the convective envelope of the star (e.g., Brinkworth et al. 2004; Gänsicke et al. 2020). These spots would be cooler than the surrounding regions, producing chromatic variations in the observed flux as they rotated in and out of view (e.g., Kilic et al. 2015; Reding et al. 2020).

Another interesting possibility is that the surface inhomogeneity of LP 398-9 is related to its origin as a D $^6$  SN Ia donor. When its companion exploded  $\sim 10^5$  years ago, LP 398-9 would have been blasted by a wave of ejecta. These radioactive ejecta would have been asymmetrically deposited on the surface of LP 398-9 (e.g., Bauer et al. 2019). In regular WDs, horizontal mixing can homogenize the surface distribution of pollutants on timescales of  $10^1 - 10^6$  years depending on the temperature and atmospheric composition (Cunningham et al. 2021). Since the stellar structure of LP 398-9 is quite uncertain, it is difficult to estimate the timescale with which diffusion is expected to homogenize the stellar surface. The vertical diffusion timescale is certainly long compared to regular WDs, since LP 398-9 is inflated to  $\approx 10$  times the typical WD radius. If the deposited surface asymmetry does survive to the present day, then it could produce variable line blanketing and continuum opacity, possibly producing the observed photometric signal (e.g., Brinkworth et al. 2004; Kilic et al. 2015; Maoz et al. 2015). If the surface abundances vary along the line of sight, this may be detectable with future time-resolved optical spectra.

Apart from the origin of the surface inhomogeneity, the rotational period of LP 398-9 can also be linked to the D $^6$  mechanism. Immediately prior to the SN explosion of the massive companion, the binary system would have been in an extremely compact, mass-transferring stage. The orbital velocity  $v_{\text{orb}}$  of the donor (mass  $M_2$ ) around the accreting primary (mass  $M_1$ ) is given by

$$v_{\text{orb}}^2 = \frac{GM_1}{a(1 + M_2/M_1)}, \quad (1)$$

where  $a$  is the orbital separation. LP 398-9 would have been ejected from the binary at  $v_{\text{orb}}$  when its companion exploded. The amount of low-velocity material left behind by the exploded companion is negligible, so LP 398-9 has probably not slowed down much since the supernova. This allows us to express the orbital period in

terms of the measured space velocity as

$$P_{\text{orb}} = \frac{2\pi a}{(1 + M_2/M_1)v_{\text{orb}}} = \frac{2\pi GM_1}{(1 + M_2/M_1)^2 v_{\text{orb}}^3}. \quad (2)$$

Assuming that the primary star that exploded as an SN is  $\approx 1.0 M_{\odot}$ , and that the donor mass is somewhere in the range  $M_2 = 0.2\text{--}0.8 M_{\odot}$ , this gives orbital periods at the time of SN detonation around  $P_{\text{orb}} = 3\text{--}7$  min, based on the  $\approx 1100 \text{ km s}^{-1}$  space velocity of LP 398-9.

Although the tidal quality factor of WDs is uncertain, it is reasonable to expect a degree of tidal synchronization between the spin and orbit of the WDs (e.g., Iben et al. 1998; Fuller & Lai 2011, 2012a,b, 2014; Yu et al. 2020, 2021). There are a handful of known tidally-distorted WDs in compact binaries, and the non-detection of significant tidal heating could imply that synchronization is near-perfect (Piro 2011; Benacquista 2011). Under the assumption that tidal locking keeps  $P_{\text{rot}} \approx P_{\text{orb}}$ , our reasoning implies a rotation period  $\approx 3\text{--}7$  minutes when the companion exploded. If LP 398-9 subsequently conserved its angular momentum, then the initial post-SN rotational period  $P_i$  and radius  $R_i$  and present-day  $P_f$ ,  $R_f$  can be related with

$$\frac{P_f}{P_i} = \left(\frac{R_f}{R_i}\right)^2 \quad (3)$$

We can set  $R_i \approx 0.01\text{--}0.02 R_{\odot}$  using the mass-radius relation for  $0.2\text{--}0.8 M_{\odot}$  WDs, and set  $R_f = 0.20 M_{\odot}$  from our SED fit in §3.2. Applying Equation 3, post-SN rotational periods  $P_i \approx 3\text{--}7$  minutes correspond to present-day rotational periods  $P_f \approx 5\text{--}50$  hours, which brackets our observed photometric period of 15.4 hr for LP 398-9 (§2.2). A larger initial donor radius (e.g., due to heating during mass transfer) corresponds to a shorter present-day rotational period. Conversely, angular momentum was probably not entirely conserved. Additional effects like magnetic braking and mass loss could have further slowed the rotational rate of the system over time. These rough estimates demonstrate that the observed rotational period can be plausibly linked to the D<sup>6</sup> scenario, assuming LP 398-9 mostly conserved its angular momentum after the SNIa explosion.

We emphasize here the differences and similarities between LP 398-9 and GD 492, another runaway star that was recently found to rotate with an 8.9 hr period (Hermes et al. 2021). GD 492 has a peculiar composition that suggests it is the partially burnt accretor left over from a Type Iax supernova (Raddi et al. 2018b,a, 2019). GD 492 also has a lower space velocity  $\simeq 850 \text{ km s}^{-1}$ , implying that its donor companion was an He-burning subwarf rather than another white dwarf (Bauer et al. 2019). In this scenario, GD 492 is rotating too slowly

to be the runaway subdwarf donor, and is most likely the burnt remnant of the accreting primary that underwent a failed supernova. Conversely, the higher space velocity of LP 398-9 — and its remarkable spectroscopic similarity to two other D<sup>6</sup> systems with space velocities  $\gtrsim 1500 \text{ km s}^{-1}$  — strongly supports a D<sup>6</sup> origin for this system. In this context, as detailed above, our detected rotation period points to LP 398-9 being a D<sup>6</sup> donor.

#### 4.2. Circumstellar Material

LP 398-9 has a strong IR excess that indicates the presence of significant quantities of circumstellar dust. Our optical spectrum shows narrow atomic carbon lines unexplained by the photospheric model, pointing to a circumstellar source of carbon. Our observations suggest a carbon-rich shell of circumstellar material inflated to more than an order of magnitude larger than present-day radius of the star.

The presence of carbon-rich circumstellar material can be plausibly explained by the D<sup>6</sup> origin of LP 398-9. The SNIa explosion of its stellar companion would have significantly polluted the atmosphere of LP 398-9, depositing thermal and radioactive energy and causing the atmosphere (dominated by carbon and oxygen) to puff up. The radius of LP 398-9 inferred from its luminosity and parallax is an order of magnitude larger than that of a typical WD, suggesting that it remains in a somewhat puffed phase like the other two runaways found in Shen et al. (2018b). Mechanical energy from the companion’s SNIa explosion could also have pushed the circumstellar material outwards to larger orbital radii. If some fraction of the inflated layers had detached from the star, it would appear today as an extended carbon-rich circumstellar shell. A low-velocity tail of the SNIa ejecta could also have been retained by LP 398-9 after the explosion.

Under the assumption of an optically thin dust shell surrounding LP 398-9, we can use our observations to approximate the total dust mass. We adopt the fitted black-body dust temperature and Gaia-inferred distance from Table 1. We assume a dust opacity  $\kappa(3.4 \mu\text{m}) = 500 \text{ cm}^2 \text{ g}^{-1}$  (Draine & Lee 1984; Draine 2003; Woitke et al. 2016), consistent with amorphous carbon grains. We estimate the dust mass using the relation from Hildebrand (1983):

$$M_{\text{dust}} = \frac{F(\lambda)D^2}{\kappa(\lambda)B(\lambda, T_d)} \quad (4)$$

Substituting the *WISE* *W1* flux as  $F(\lambda)$  provides an order-of-magnitude total dust mass  $M_{\text{dust}} \approx 10^{20} \text{ g} = 10^{-13} M_{\odot}$ . This dust mass is comparable to that of known dust disks around WDs (Jura 2003; Reach et al. 2005), although we stress that LP 398-9’s origin and

composition is quite different from those systems. For comparison, the rings of Saturn have a mass  $\approx 10^{22} g$  (Esposito 1993; Iess et al. 2019).

## 5. DISCUSSION

We have presented two lines of evidence tying LP 398-9 to the D<sup>6</sup> SNIa progenitor scenario: circumstellar material and surface rotation. We interpret the 15.4 hr photometric signal as a signature of rotationally modulated brightness variations, potentially stemming from surface inhomogeneities left over from the SN explosion  $\sim 10^5$  years ago. The rotational period itself can be explained by angular momentum conservation of LP 398-9 after its companion exploded, assuming the binary was tidally locked at the point of detonation. Several other effects could have also altered the rotational rate. Since tidal dissipation primarily occurs near the surface of the WD, the surface layers could preferentially synchronize without fully spinning up the core, invalidating our assumption of a rigid-body rotation (e.g., Goldreich & Nicholson 1989; Fuller & Lai 2012a,b, 2014). Additionally, LP 398-9 could have lost angular momentum in the time since the SN explosion. An obvious culprit could be the inflated stellar layers that today present as circumstellar dust tens of stellar radii away. The present-day total dust mass is a minuscule fraction of the stellar mass, making it an unlikely route for angular momentum to leave the system. However, there could have been episodes of further mass loss in the time since the supernova.

Although it is uncertain precisely what processes cause the D<sup>6</sup> donor WDs to inflate to their presently observed radii, this inflated phase is probably only temporary (Shen et al. 2018b; Bauer et al. 2019). Due to selection effects, we are most likely to find D<sup>6</sup> donors in their inflated state, since these are detectable out to a larger search volume. Our fitted stellar parameters imply a Kelvin-Helmholtz timescale  $\tau_{\text{KH}} \gtrsim 20$  Myr for LP 398-9, much longer than the flight time  $\approx 0.1$  Myr from the SN remnant G70.0-21.5. As LP 398-9 radiates away energy deposited by tidal heating and the SN itself, it will contract in size and gradually return to the WD cooling track. In the absence of any mechanism to remove angular momentum from the system, its rotation will speed up as it shrinks. Therefore, a plausible long-term outcome of the D<sup>6</sup> scenario is a class of rapidly rotating hypervelocity white dwarfs. These might have helium-dominated or carbon-dominated spectral types. However, such hypervelocity systems leave the Galaxy on short timescale  $\lesssim 10$  Myr, making it quite unlikely that we could detect them. Due to this selection effect,

we are overwhelmingly likely to find D<sup>6</sup> donors in their inflated state.

When an isolated WD has an observed IR excess, the usual explanation is circumstellar dust from a disrupted planetesimal (e.g., Debes & Sigurdsson 2002; Jura 2003; Farihi 2016; Veras 2021). While the existence of disrupted planetesimals around LP 398-9 is tantalizing, it is quite unlikely. The D<sup>6</sup> origin of LP 398-9 implies that it was until recently ( $\sim 10^5$  years ago) in a close binary system with another WD. During the final stages of binary evolution, the two WDs in the theorized D<sup>6</sup> scenario would be orbiting too close for a planet to maintain a stable orbit around LP 398-9. While it is plausible that the stars possessed a circumbinary planetary system, any surviving planetesimals must have occupied orbits wide enough to avoid engulfment during giant branch evolution, yet close enough to remain bound to LP 398-9 after the SN Ia of its companion. This scenario can be tested with future infrared spectroscopy — for example with the James Webb Space Telescope (Gardner et al. 2006) — that would reveal the temperature profile, composition, and geometry of the circumstellar material.

One question that remains is why LP 398-9 (‘D6-2’) is the only D<sup>6</sup> candidate with an infrared excess, out of the three candidates found by Shen et al. (2018b). ‘D6-1’ has secure *WISE* data with no detectable IR excess (Figure 1), while ‘D6-3’ does not have secure data in *WISE*. One possible explanation is that the other candidates are the products of older supernovae, and could have consequently lost their circumstellar shell over time. LP 398-9 is the only D<sup>6</sup> star with an associated SN remnant, suggesting that it was ejected recently enough that the remnant did not dissipate into the interstellar medium. Conversely, all three D<sup>6</sup> stars are clustered closely together on the color-magnitude diagram and have similar low-resolution spectra, perhaps indicating that they are in similar stages of their evolution. D6-1 and D6-3 should be monitored to search for rotationally modulated variations. If they are indeed the product of older supernovae that LP 398-9, then their surface composition may have homogenized enough to make their rotational signal undetectable.

Future observations of both LP 398-9 and the other D<sup>6</sup> candidates could aid in resolving these questions by placing firmer constraints on the time elapsed since the respective SNe Ia. Further spectroscopic observations of other D<sup>6</sup> candidates could search for carbon absorption indicative of circumstellar dust that is too cool to produce an IR excess in *WISE*. Additionally, more theoretical modelling could estimate the timescales over which ejected runaways cool and return to the WD cooling

track. Understanding these unique systems will shed light on the mechanism and after-effects of the double-degenerate channel for SNe Ia.

*Data Availability*—The corresponding author will gladly share any code and intermediate data products upon reasonable request.

## ACKNOWLEDGMENTS

VC acknowledges support from the James Mills Peirce fellowship at Harvard University, and the Institute for Data-Intensive Engineering and Science at Johns Hopkins University. VC and HCH were supported in part by Space@Hopkins. VC, HCH, and NLZ were supported in part by NASA-ADAP 80NSSC19K0581. NLZ acknowledges support from J. Robert Oppenheimer Visiting Professorship and the Bershadsky Fund at the Institute for Advanced Study. SB acknowledges support from the Laboratory Directed Research and Development program of Los Alamos National Laboratory (20190624PRD2) and from the Banting Postdoctoral Fellowships program, administered by the Government of Canada. AS acknowledges support from STFC grant ST/R000476/1. KJS is supported by NASA through the Astrophysics Theory Program (NNX17AG28G). The design and construction of HiPERCAM was funded by the European Research Council under the European Union’s Seventh Framework Programme (FP/2007-2013) under ERC-2013-ADG Grant Agreement no. 340040 (HiPERCAM). VSD and HiPERCAM operations are supported by STFC grant ST/V000853/1.

Based in part on observations obtained with the Apache Point Observatory 3.5-m telescope, which is owned and operated by the Astrophysical Research Consortium. Based in part on observations made with the Gran Telescopio Canarias (GTC), installed in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias, in the island of La Palma. Based in part on observations obtained with the Isaac Newton Telescope, operated on the island of La Palma by the Isaac Newton Group of Telescopes in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. The Liverpool Telescope is operated on the island of La Palma by Liverpool John Moores University in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias with financial support from the UK Science and Technology Facilities Council. This work has made use of observations made with the NASA/ESA *Hubble Space Telescope*, obtained from the Data Archive at the Space Telescope Science Insti-

tute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. These observations are associated with program #15871, and funding in-part provided by programs #15871 and #15918.

This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement. We gratefully acknowledge NASA’s support for construction, operation, and science analysis for the GALEX mission, developed in cooperation with the Centre National d’Etudes Spatiales of France and the Korean Ministry of Science and Technology. This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration. This publication also makes use of data products from NEOWISE, which is a project of the Jet Propulsion Laboratory/California Institute of Technology, funded by the Planetary Science Division of the National Aeronautics and Space Administration. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This research has made use of the NASA/IPAC Infrared Science Archive, which is funded by the National Aeronautics and Space Administration and operated by the California Institute of Technology. This research has made use of the VizieR catalogue access tool, CDS, Strasbourg, France. The original description of the VizieR service was published in 2000, A&AS 143, 23. This research has made extensive use of NASA’s Astrophysics Data System Bibliographic Services. Based on observations obtained with the Samuel Oschin 48-inch Telescope at the Palomar Observatory as part of the Zwicky Transient Facility project. ZTF is supported by the National Science Foundation under Grant No. AST-1440341 and a collaboration including Caltech, IPAC, the Weizmann Institute for Science, the Oskar Klein Center at Stockholm University, the University of Maryland, the University of Washington, Deutsches Elektronen-Synchrotron and Humboldt University, Los Alamos National Laboratories, the TANGO

Consortium of Taiwan, the University of Wisconsin at Milwaukee, and Lawrence Berkeley National Laboratories. Operations are conducted by COO, IPAC, and UW. Parts of the results in this work make use of the colormaps in the CMasher package.

*Software:* `numpy` (Harris et al. 2020), `scipy` (Virtanen et al. 2020), `emcee` (Foreman-Mackey et al.

2013; Foreman-Mackey et al. 2019), `matplotlib` (Hunter 2007), `lmfit` (Newville & Stensitzki 2018), `ltsfit` (Cappellari et al. 2013),

*Facilities:* Gaia, Sloan, WISE, ARC:3.5-m (DIS), GTC:10.4-m (HiPERCAM), Liverpool:2-m (IO:O), INT:2.5-m (WFC), FLWIO:2MASS, HST (STIS), GALEX, IRSA

## REFERENCES

- Ahumada, R., Prieto, C. A., Almeida, A., et al. 2020, The Astrophysical Journal Supplement Series, 249, 3, doi: [10.3847/1538-4365/ab929e](https://doi.org/10.3847/1538-4365/ab929e)
- Bailer-Jones, C. A. L. 2015, PASP, 127, 994, doi: [10.1086/683116](https://doi.org/10.1086/683116)
- Bauer, E. B., White, C. J., & Bildsten, L. 2019, The Astrophysical Journal, 887, 68, doi: [10.3847/1538-4357/ab4ea4](https://doi.org/10.3847/1538-4357/ab4ea4)
- Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019, Publications of the Astronomical Society of the Pacific, 131, 1, doi: [10.1088/1538-3873/aacbe](https://doi.org/10.1088/1538-3873/aacbe)
- Benacquista, M. J. 2011, Astrophysical Journal Letters, 740, 3, doi: [10.1088/2041-8205/740/2/L54](https://doi.org/10.1088/2041-8205/740/2/L54)
- Bergeron, P., Dufour, P., Fontaine, G., et al. 2019, The Astrophysical Journal, 876, 67, doi: [10.3847/1538-4357/ab153a](https://doi.org/10.3847/1538-4357/ab153a)
- Bildsten, L., Shen, K. J., Weinberg, N. N., & Nelemans, G. 2007, ApJL, 662, L95, doi: [10.1086/519489](https://doi.org/10.1086/519489)
- Blanton, M. R., Bershady, M. A., Abolfathi, B., et al. 2017, The Astronomical Journal, 154, 28, doi: [10.3847/1538-3881/aa7567](https://doi.org/10.3847/1538-3881/aa7567)
- Bloom, J. S., Kasen, D., Shen, K. J., et al. 2012, ApJL, 744, L17, doi: [10.1088/2041-8205/744/2/L17](https://doi.org/10.1088/2041-8205/744/2/L17)
- Blouin, S., Dufour, P., & Allard, N. F. 2018a, The Astrophysical Journal, 863, 184, doi: [10.3847/1538-4357/aad4a9](https://doi.org/10.3847/1538-4357/aad4a9)
- Blouin, S., Dufour, P., Allard, N. F., & Kilic, M. 2018b, The Astrophysical Journal, 867, 161, doi: [10.3847/1538-4357/aae53a](https://doi.org/10.3847/1538-4357/aae53a)
- Brinkworth, C. S., Burleigh, M. R., Wynn, G. A., & Marsh, T. R. 2004, Monthly Notices of the Royal Astronomical Society, 348, 33, doi: [10.1111/j.1365-2966.2004.07538.x](https://doi.org/10.1111/j.1365-2966.2004.07538.x)
- Brinkworth, C. S., Marsh, T. R., Morales-Rueda, L., et al. 2005, Monthly Notices of the Royal Astronomical Society, 357, 333, doi: [10.1111/j.1365-2966.2005.08649.x](https://doi.org/10.1111/j.1365-2966.2005.08649.x)
- Brown, A. G., Vallenari, A., Prusti, T., et al. 2021, Astronomy and Astrophysics, 649, 1, doi: [10.1051/0004-6361/202039657](https://doi.org/10.1051/0004-6361/202039657)
- Burdge, K. B., Prince, T. A., Fuller, J., et al. 2020, The Astrophysical Journal, 905, 32, doi: [10.3847/1538-4357/abc261](https://doi.org/10.3847/1538-4357/abc261)
- Cappellari, M., Scott, N., Alatalo, K., et al. 2013, MNRAS, 432, 1709, doi: [10.1093/mnras/stt562](https://doi.org/10.1093/mnras/stt562)
- Chandrasekhar, S. 1931, The Astrophysical Journal, 74, 81, doi: [10.1086/143324](https://doi.org/10.1086/143324)
- Condon, J. J., Cotton, W. D., Greisen, E. W., et al. 1998, AJ, 115, 1693, doi: [10.1086/300337](https://doi.org/10.1086/300337)
- Cunningham, T., Tremblay, P.-E., Bauer, E. B., et al. 2021, Monthly Notices of the Royal Astronomical Society, 503, 1646, doi: [10.1093/mnras/stab553](https://doi.org/10.1093/mnras/stab553)
- Cutri, R. M., Wright, E. L., Conrow, T., et al. 2012, Explanatory Supplement to the WISE All-Sky Data Release Products, Explanatory Supplement to the WISE All-Sky Data Release Products
- Dan, M., Rosswog, S., Guillochon, J., & Ramirez-Ruiz, E. 2011, ApJ, 737, 89, doi: [10.1088/0004-637X/737/2/89](https://doi.org/10.1088/0004-637X/737/2/89)
- de los Reyes, M. A. C., Kirby, E. N., Seitzzahl, I. R., & Shen, K. J. 2020, The Astrophysical Journal, 891, 85, doi: [10.3847/1538-4357/ab736f](https://doi.org/10.3847/1538-4357/ab736f)
- Debes, J. H., & Sigurdsson, S. 2002, The Astrophysical Journal, 572, 556, doi: [10.1086/340291](https://doi.org/10.1086/340291)
- Dennihi, E., Farihi, J., Fusillo, N. P. G., & Debes, J. H. 2020, The Astrophysical Journal, 891, 97, doi: [10.3847/1538-4357/ab7249](https://doi.org/10.3847/1538-4357/ab7249)
- Dhawan, S., Leibundgut, B., Spyromilio, J., & Blondin, S. 2017, A&A, 602, A118, doi: [10.1051/0004-6361/201629793](https://doi.org/10.1051/0004-6361/201629793)
- Dhillon, V., Dixon, S., Gamble, T., et al. 2018, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 10702, Ground-based and Airborne Instrumentation for Astronomy VII, ed. C. J. Evans, L. Simard, & H. Takami, 107020L, doi: [10.1117/12.2312041](https://doi.org/10.1117/12.2312041)

- Dhillon, V. S., Marsh, T. R., Bezawada, N., et al. 2016, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9908, Ground-based and Airborne Instrumentation for Astronomy VI, ed. C. J. Evans, L. Simard, & H. Takami, 99080Y, doi: [10.1117/12.2229055](https://doi.org/10.1117/12.2229055)
- Dhillon, V. S., Bezawada, N., Black, M., et al. 2021, MNRAS, 507, 350, doi: [10.1093/mnras/stab2130](https://doi.org/10.1093/mnras/stab2130)
- Doi, M., Tanaka, M., Fukugita, M., et al. 2010, Astronomical Journal, 139, 1628, doi: [10.1088/0004-6256/139/4/1628](https://doi.org/10.1088/0004-6256/139/4/1628)
- Draine, B. T. 2003, ARA&A, 41, 241, doi: [10.1146/annurev.astro.41.011802.094840](https://doi.org/10.1146/annurev.astro.41.011802.094840)
- Draine, B. T., & Lee, H. M. 1984, ApJ, 285, 89, doi: [10.1086/162480](https://doi.org/10.1086/162480)
- Esposito, L. W. 1993, AREPS, 21, 487, doi: [10.1146/annurev.astro.21.050193.002415](https://doi.org/10.1146/annurev.astro.21.050193.002415)
- Farihi, J. 2016, New Astronomy Reviews, 71, 9, doi: [10.1016/j.newar.2016.03.001](https://doi.org/10.1016/j.newar.2016.03.001)
- Ferrario, L., Vennes, S., Wickramasinghe, D. T., Bailey, J. A., & Christian, D. J. 1997, Monthly Notices of the Royal Astronomical Society, 292, 205, doi: [10.1093/mnras/292.2.205](https://doi.org/10.1093/mnras/292.2.205)
- Fesen, R. A., Neustadt, J. M., Black, C. S., & Koepfel, A. H. 2015, Astrophysical Journal, 812, 37, doi: [10.1088/0004-637X/812/1/37](https://doi.org/10.1088/0004-637X/812/1/37)
- Fitzpatrick, E. L., & Massa, D. 2007, ApJ, 663, 320, doi: [10.1086/518158](https://doi.org/10.1086/518158)
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, Publications of the Astronomical Society of the Pacific, 125, 306, doi: [10.1086/670067](https://doi.org/10.1086/670067)
- Foreman-Mackey, D., Farr, W., Sinha, M., et al. 2019, The Journal of Open Source Software, 4, 1864, doi: [10.21105/joss.01864](https://doi.org/10.21105/joss.01864)
- Fukugita, M., Ichikawa, T., Gunn, J. E., et al. 1996, The Astronomical Journal, 111, 1748, doi: [10.1086/117915](https://doi.org/10.1086/117915)
- Fuller, J., & Lai, D. 2011, Monthly Notices of the Royal Astronomical Society, 412, 1331, doi: [10.1111/j.1365-2966.2010.18017.x](https://doi.org/10.1111/j.1365-2966.2010.18017.x)
- . 2012a, Astrophysical Journal Letters, 756, 2, doi: [10.1088/2041-8205/756/1/L17](https://doi.org/10.1088/2041-8205/756/1/L17)
- . 2012b, Monthly Notices of the Royal Astronomical Society, 421, 426, doi: [10.1111/j.1365-2966.2011.20320.x](https://doi.org/10.1111/j.1365-2966.2011.20320.x)
- . 2014, Monthly Notices of the Royal Astronomical Society, 444, 3488, doi: [10.1093/mnras/stu1698](https://doi.org/10.1093/mnras/stu1698)
- Gänsicke, B. T., Rodríguez-Gil, P., Gentile Fusillo, N. P., et al. 2020, Monthly Notices of the Royal Astronomical Society, 499, 2564, doi: [10.1093/mnras/staa2969](https://doi.org/10.1093/mnras/staa2969)
- Gardner, J. P., Mather, J. C., Clampin, M., et al. 2006, SSRv, 123, 485, doi: [10.1007/s11214-006-8315-7](https://doi.org/10.1007/s11214-006-8315-7)
- Goldreich, P., & Nicholson, P. D. 1989, ApJ, 342, 1079, doi: [10.1086/167665](https://doi.org/10.1086/167665)
- Green, G. 2018, The Journal of Open Source Software, 3, 695, doi: [10.21105/joss.00695](https://doi.org/10.21105/joss.00695)
- Green, G. M., Schlafly, E. F., Finkbeiner, D. P., et al. 2015, Astrophysical Journal, 810, doi: [10.1088/0004-637X/810/1/25](https://doi.org/10.1088/0004-637X/810/1/25)
- Green, G. M., Schlafly, E. F., Finkbeiner, D., et al. 2018a, Monthly Notices of the Royal Astronomical Society, 478, 651, doi: [10.1093/mnras/sty1008](https://doi.org/10.1093/mnras/sty1008)
- Green, M. J., Hermes, J. J., Marsh, T. R., et al. 2018b, Monthly Notices of the Royal Astronomical Society, 477, 5646, doi: [10.1093/mnras/sty1032](https://doi.org/10.1093/mnras/sty1032)
- Guidry, J. A., Vanderbosch, Z. P., Hermes, J. J., et al. 2021, The Astrophysical Journal, 912, 125, doi: [10.3847/1538-4357/abee68](https://doi.org/10.3847/1538-4357/abee68)
- Guillochon, J., Dan, M., Ramirez-Ruiz, E., & Rosswog, S. 2010, ApJL, 709, L64, doi: [10.1088/2041-8205/709/1/L64](https://doi.org/10.1088/2041-8205/709/1/L64)
- Gunn, J. E., Carr, M., Rockosi, C., et al. 1998, The Astronomical Journal, 116, 3040, doi: [10.1086/300645](https://doi.org/10.1086/300645)
- Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, Nature, 585, 357, doi: [10.1038/s41586-020-2649-2](https://doi.org/10.1038/s41586-020-2649-2)
- Hermes, J. J., Putterman, O., Hollands, M. A., et al. 2021, The Astrophysical Journal Letters, 914, L3, doi: [10.3847/2041-8213/ac00a8](https://doi.org/10.3847/2041-8213/ac00a8)
- Hildebrand, R. H. 1983, QJRAS, 24, 267
- Hillebrandt, W., & Niemeyer, J. C. 2000, Annual Review of Astronomy and Astrophysics, 38, 191, doi: [10.1146/annurev.astro.38.1.191](https://doi.org/10.1146/annurev.astro.38.1.191)
- Hunter, J. D. 2007, Computing in Science and Engineering, 9, 90, doi: [10.1109/MCSE.2007.55](https://doi.org/10.1109/MCSE.2007.55)
- Iben, Jr., I., Tutukov, A. V., & Fedorova, A. V. 1998, The Astrophysical Journal, 503, 344, doi: [10.1086/305972](https://doi.org/10.1086/305972)
- Iben, I., J., & Tutukov, A. V. 1984, The Astrophysical Journal Supplement Series, 54, 335, doi: [10.1086/190932](https://doi.org/10.1086/190932)
- Iess, L., Militzer, B., Kaspi, Y., et al. 2019, Sci, 364, aat2965, doi: [10.1126/science.aat2965](https://doi.org/10.1126/science.aat2965)
- Johnson, R. C. 1927, RSPTA, 226, 157, doi: [10.1098/rsta.1927.0005](https://doi.org/10.1098/rsta.1927.0005)
- Jura, M. 2003, The Astrophysical Journal, 584, L91, doi: [10.1086/374036](https://doi.org/10.1086/374036)
- Kasen, D. 2010, ApJ, 708, 1025, doi: [10.1088/0004-637X/708/2/1025](https://doi.org/10.1088/0004-637X/708/2/1025)
- Kenyon, S. J., & Bromley, B. C. 2017a, The Astrophysical Journal, 850, 50, doi: [10.3847/1538-4357/aa9570](https://doi.org/10.3847/1538-4357/aa9570)
- . 2017b, The Astrophysical Journal, 844, 116, doi: [10.3847/1538-4357/aa7b85](https://doi.org/10.3847/1538-4357/aa7b85)
- Kerzendorf, W. E., Schmidt, B. P., Asplund, M., et al. 2009, ApJ, 701, 1665, doi: [10.1088/0004-637X/701/2/1665](https://doi.org/10.1088/0004-637X/701/2/1665)

- Kilic, M., Gianninas, A., Bell, K. J., et al. 2015, *Astrophysical Journal Letters*, 814, L31, doi: [10.1088/2041-8205/814/2/L31](https://doi.org/10.1088/2041-8205/814/2/L31)
- Kirby, E. N., Xie, J. L., Guo, R., et al. 2019, *The Astrophysical Journal*, 881, 45, doi: [10.3847/1538-4357/ab2c02](https://doi.org/10.3847/1538-4357/ab2c02)
- Li, W., Bloom, J. S., Podsiadlowski, P., et al. 2011, *Nature*, 480, 348, doi: [10.1038/nature10646](https://doi.org/10.1038/nature10646)
- Lindegren, L., Klioner, S. A., Hernández, J., et al. 2021, *Astronomy and Astrophysics*, 649, A2, doi: [10.1051/0004-6361/202039709](https://doi.org/10.1051/0004-6361/202039709)
- Lomb, N. R. 1976, *Ap&SS*, 39, 447, doi: [10.1007/BF00648343](https://doi.org/10.1007/BF00648343)
- Mainzer, A., Bauer, J., Grav, T., et al. 2011, *ApJ*, 731, 53, doi: [10.1088/0004-637X/731/1/53](https://doi.org/10.1088/0004-637X/731/1/53)
- Mainzer, A., Bauer, J., Cutri, R. M., et al. 2014, *Astrophysical Journal*, 792, 30, doi: [10.1088/0004-637X/792/1/30](https://doi.org/10.1088/0004-637X/792/1/30)
- Maoz, D., Mannucci, F., & Nelemans, G. 2014, *Annual Review of Astronomy and Astrophysics*, 52, 107, doi: [10.1146/annurev-astro-082812-141031](https://doi.org/10.1146/annurev-astro-082812-141031)
- Maoz, D., Mazeh, T., & McQuillan, A. 2015, *Monthly Notices of the Royal Astronomical Society*, 447, 1749, doi: [10.1093/mnras/stu2577](https://doi.org/10.1093/mnras/stu2577)
- Margutti, R., Soderberg, A. M., Chomiuk, L., et al. 2012, *ApJ*, 751, 134, doi: [10.1088/0004-637X/751/2/134](https://doi.org/10.1088/0004-637X/751/2/134)
- Martin, D. C., Fanson, J., Schiminovich, D., et al. 2005, *ApJL*, 619, L1, doi: [10.1086/426387](https://doi.org/10.1086/426387)
- Masci, F. J., Laher, R. R., Rusholme, B., et al. 2019, *Publications of the Astronomical Society of the Pacific*, 131, 1, doi: [10.1088/1538-3873/aae8ac](https://doi.org/10.1088/1538-3873/aae8ac)
- Matteucci, F., Spitoni, E., Recchi, S., & Valiante, R. 2009, *A&A*, 501, 531, doi: [10.1051/0004-6361/200911869](https://doi.org/10.1051/0004-6361/200911869)
- Mignard, F., Klioner, S. A., Lindegren, L., et al. 2018, *Astronomy and Astrophysics*, 616, A14, doi: [10.1051/0004-6361/201832916](https://doi.org/10.1051/0004-6361/201832916)
- Million, C., Fleming, S. W., Shiao, B., et al. 2016, *ApJ*, 833, 292, doi: [10.3847/1538-4357/833/2/292](https://doi.org/10.3847/1538-4357/833/2/292)
- Newville, M., & Stensitzki, T. 2018, *Non-Linear Least-Squares Minimization and Curve-Fitting for Python*, 65, doi: [10.5281/ZENODO.11813](https://doi.org/10.5281/ZENODO.11813)
- Nomoto, K. 1982, *The Astrophysical Journal*, 253, 798, doi: [10.1086/159682](https://doi.org/10.1086/159682)
- Pakmor, R., Kromer, M., Taubenberger, S., & Springel, V. 2013, *Astrophysical Journal Letters*, 770, doi: [10.1088/2041-8205/770/1/L8](https://doi.org/10.1088/2041-8205/770/1/L8)
- Pelisoli, I., Marsh, T. R., Dhillon, V. S., et al. 2021, 5, 1, <http://arxiv.org/abs/2108.11396>
- Perlmutter, S., Aldering, G., Goldhaber, G., et al. 1999, *ApJ*, 517, 565, doi: [10.1086/307221](https://doi.org/10.1086/307221)
- Piro, A. L. 2011, *Astrophysical Journal Letters*, 740, 2, doi: [10.1088/2041-8205/740/2/L53](https://doi.org/10.1088/2041-8205/740/2/L53)
- Prusti, T., De Bruijne, J. H., Brown, A. G., et al. 2016, *Astronomy and Astrophysics*, 595, doi: [10.1051/0004-6361/201629272](https://doi.org/10.1051/0004-6361/201629272)
- Raddi, R., Hollands, M. A., Gänsicke, B. T., et al. 2018a, *Monthly Notices of the Royal Astronomical Society: Letters*, 479, L96, doi: [10.1093/mnrasl/sly103](https://doi.org/10.1093/mnrasl/sly103)
- Raddi, R., Hollands, M. A., Koester, D., et al. 2018b, *The Astrophysical Journal*, 858, 3, doi: [10.3847/1538-4357/aab899](https://doi.org/10.3847/1538-4357/aab899)
- . 2019, *Monthly Notices of the Royal Astronomical Society*, 489, 1489, doi: [10.1093/mnras/stz1618](https://doi.org/10.1093/mnras/stz1618)
- Rafikov, R. R., & Garmilla, J. A. 2012, *ApJ*, 760, 123, doi: [10.1088/0004-637X/760/2/123](https://doi.org/10.1088/0004-637X/760/2/123)
- Raskin, C., Scannapieco, E., Fryer, C., Rockefeller, G., & Timmes, F. X. 2012, *ApJ*, 746, 62, doi: [10.1088/0004-637X/746/1/62](https://doi.org/10.1088/0004-637X/746/1/62)
- Raymond, J. C., Caldwell, N., Fesen, R. A., et al. 2020, *The Astrophysical Journal*, 888, 90, doi: [10.3847/1538-4357/ab5e84](https://doi.org/10.3847/1538-4357/ab5e84)
- Reach, W. T., Kuchner, M. J., von Hippel, T., et al. 2005, *The Astrophysical Journal*, 635, L161, doi: [10.1086/499561](https://doi.org/10.1086/499561)
- Reding, J. S., Hermes, J. J., Vanderbosch, Z., et al. 2020, *The Astrophysical Journal*, 894, 19, doi: [10.3847/1538-4357/ab8239](https://doi.org/10.3847/1538-4357/ab8239)
- Riess, A. G., Casertano, S., Yuan, W., Macri, L. M., & Scolnic, D. 2019, *The Astrophysical Journal*, 876, 85, doi: [10.3847/1538-4357/ab1422](https://doi.org/10.3847/1538-4357/ab1422)
- Riess, A. G., Filippenko, A. V., Challis, P., et al. 1998, *The Astronomical Journal*, 116, 1009, doi: [10.1086/300499](https://doi.org/10.1086/300499)
- Rousseeuw, P. J., & Van Driessen, K. 2006, *Data Mining and Knowledge Discovery*, 12, 29, doi: [10.1007/s10618-005-0024-4](https://doi.org/10.1007/s10618-005-0024-4)
- Sanders, J. L., Belokurov, V., & Man, K. T. F. 2021, *Monthly Notices of the Royal Astronomical Society*, 506, 4321, doi: [10.1093/mnras/stab1951](https://doi.org/10.1093/mnras/stab1951)
- Scalzo, R., Aldering, G., Antilogus, P., et al. 2014, *MNRAS*, 440, 1498, doi: [10.1093/mnras/stu350](https://doi.org/10.1093/mnras/stu350)
- Scargle, J. D. 1982, *ApJ*, 263, 835, doi: [10.1086/160554](https://doi.org/10.1086/160554)
- Shen, K. J. 2015, *Astrophysical Journal Letters*, 805, 1, doi: [10.1088/2041-8205/805/1/L6](https://doi.org/10.1088/2041-8205/805/1/L6)
- Shen, K. J., & Bildsten, L. 2014, *Astrophysical Journal*, 785, doi: [10.1088/0004-637X/785/1/61](https://doi.org/10.1088/0004-637X/785/1/61)
- Shen, K. J., Blondin, S., Kasen, D., et al. 2021a, *The Astrophysical Journal Letters*, 909, L18, doi: [10.3847/2041-8213/abe69b](https://doi.org/10.3847/2041-8213/abe69b)
- Shen, K. J., Boos, S. J., Townsley, D. M., & Kasen, D. 2021b, 1. <http://arxiv.org/abs/2108.12435>

- Shen, K. J., Kasen, D., Miles, B. J., & Townsley, D. M. 2018a, *The Astrophysical Journal*, 854, 52, doi: [10.3847/1538-4357/aaa8de](https://doi.org/10.3847/1538-4357/aaa8de)
- Shen, K. J., & Schwab, J. 2017, *The Astrophysical Journal*, 834, 180, doi: [10.3847/1538-4357/834/2/180](https://doi.org/10.3847/1538-4357/834/2/180)
- Shen, K. J., Boubert, D., Gänsicke, B. T., et al. 2018b, *The Astrophysical Journal*, 865, 15, doi: [10.3847/1538-4357/aad55b](https://doi.org/10.3847/1538-4357/aad55b)
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *The Astronomical Journal*, 131, 1163, doi: [10.1086/498708](https://doi.org/10.1086/498708)
- Smart, R. L., Sarro, L. M., Rybizki, J., et al. 2021, *Astronomy and Astrophysics*, 649, 1, doi: [10.1051/0004-6361/202039498](https://doi.org/10.1051/0004-6361/202039498)
- Sorahana, S., Yamamura, I., & Murakami, H. 2013, *ApJ*, 767, 77, doi: [10.1088/0004-637X/767/1/77](https://doi.org/10.1088/0004-637X/767/1/77)
- Steele, I. A., Smith, R. J., Rees, P. C., et al. 2004, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 5489, *Ground-based Telescopes*, ed. J. Oschmann, Jacobus M., 679–692, doi: [10.1117/12.551456](https://doi.org/10.1117/12.551456)
- Swan, A., Farihi, J., & Wilson, T. G. 2019, *Monthly Notices of the Royal Astronomical Society: Letters*, 484, L109, doi: [10.1093/mnrasl/slz014](https://doi.org/10.1093/mnrasl/slz014)
- Swan, A., Farihi, J., Wilson, T. G., & Parsons, S. G. 2020, *Monthly Notices of the Royal Astronomical Society*, 496, 5233, doi: [10.1093/mnras/staa1688](https://doi.org/10.1093/mnras/staa1688)
- Swan, A., Kenyon, S. J., Farihi, J., et al. 2021, *Monthly Notices of the Royal Astronomical Society*, doi: [10.1093/mnras/stab1738](https://doi.org/10.1093/mnras/stab1738)
- Swan, W. 1857, *Transactions of the Royal Society of Edinburgh*, 21, 411–429, doi: [10.1017/S0080456800032233](https://doi.org/10.1017/S0080456800032233)
- Taam, R. E. 1980, *ApJ*, 237, 142, doi: [10.1086/157852](https://doi.org/10.1086/157852)
- Tanikawa, A., Nomoto, K., & Nakasato, N. 2018, *The Astrophysical Journal*, 868, 90, doi: [10.3847/1538-4357/aae9ee](https://doi.org/10.3847/1538-4357/aae9ee)
- Tanikawa, A., Nomoto, K., Nakasato, N., & Maeda, K. 2019, *The Astrophysical Journal*, 885, 103, doi: [10.3847/1538-4357/ab46b6](https://doi.org/10.3847/1538-4357/ab46b6)
- Tinsley, B. M. 1979, *ApJ*, 229, 1046, doi: [10.1086/157039](https://doi.org/10.1086/157039)
- Tody, D. 1986, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 627, *Instrumentation in astronomy VI*, ed. D. L. Crawford, 733, doi: [10.1117/12.968154](https://doi.org/10.1117/12.968154)
- Tremblay, P. E., Fontaine, G., Freytag, B., et al. 2015, *Astrophysical Journal*, 812, 19, doi: [10.1088/0004-637X/812/1/19](https://doi.org/10.1088/0004-637X/812/1/19)
- Veras, D. 2021, arXiv, arXiv:2106.06550. <https://arxiv.org/abs/2106.06550>
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *Nature Methods*, 17, 261, doi: [10.1038/s41592-019-0686-2](https://doi.org/10.1038/s41592-019-0686-2)
- von Hippel, T., Kuchner, M. J., Kilic, M., Mullally, F., & Reach, W. T. 2007, *The Astrophysical Journal*, 662, 544, doi: [10.1086/518108](https://doi.org/10.1086/518108)
- Webbink, R. F. 1984, *The Astrophysical Journal*, 277, 355, doi: [10.1086/161701](https://doi.org/10.1086/161701)
- Whelan, J., & Iben, Icko, J. 1973, *The Astrophysical Journal*, 186, 1007, doi: [10.1086/152565](https://doi.org/10.1086/152565)
- Woitke, P., Min, M., Pinte, C., et al. 2016, *A&A*, 586, A103, doi: [10.1051/0004-6361/201526538](https://doi.org/10.1051/0004-6361/201526538)
- Woodgate, B. E., Kimble, R. A., Bowers, C. W., et al. 1998, *PASP*, 110, 1183, doi: [10.1086/316243](https://doi.org/10.1086/316243)
- Woods, T. E., Ghavamian, P., Badenes, C., & Gilfanov, M. 2017, *NatAs*, 1, 800, doi: [10.1038/s41550-017-0263-5](https://doi.org/10.1038/s41550-017-0263-5)
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, 140, 1868, doi: [10.1088/0004-6256/140/6/1868](https://doi.org/10.1088/0004-6256/140/6/1868)
- Xu, S., Rappaport, S., van Lieshout, R., et al. 2018, *Monthly Notices of the Royal Astronomical Society*, 474, 4795, doi: [10.1093/mnras/stx3023](https://doi.org/10.1093/mnras/stx3023)
- Yu, H., Fuller, J., & Burdge, K. B. 2021, *MNRAS*, 501, 1836, doi: [10.1093/mnras/staa3717](https://doi.org/10.1093/mnras/staa3717)
- Yu, H., Weinberg, N. N., & Fuller, J. 2020, *MNRAS*, 496, 5482, doi: [10.1093/mnras/staa1858](https://doi.org/10.1093/mnras/staa1858)