



This is a repository copy of *Sacral nerve stimulation versus the magnetic sphincter augmentation device for adult faecal incontinence: the SaFaRI RCT.*

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/181837/>

Version: Published Version

Article:

Jayne, D.G., Williams, A.E., Corrigan, N. et al. (16 more authors) (2021) Sacral nerve stimulation versus the magnetic sphincter augmentation device for adult faecal incontinence: the SaFaRI RCT. *Health Technology Assessment*, 25 (18). ISSN 1366-5278

<https://doi.org/10.3310/hta25180>

© Queen's Printer and Controller of HMSO 2021. This work was produced by Jayne et al. under the terms of a commissioning contract issued by the Secretary of State for Health and Social Care. This issue may be freely reproduced for the purposes of private research and study and extracts (or indeed, the full report) may be included in professional journals provided that suitable acknowledgement is made and the reproduction is not associated with any form of advertising. Applications for commercial reproduction should be addressed to: NIHR Journals Library, National Institute for Health Research, Evaluation, Trials and Studies Coordinating Centre, Alpha House, University of Southampton Science Park, Southampton SO16 7NS, UK.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>


Health Technology Assessment

Volume 25 • Issue 18 • March 2021

ISSN 1366-5278

Sacral nerve stimulation versus the magnetic sphincter augmentation device for adult faecal incontinence: the SaFaRI RCT

David G Jayne, Annabelle E Williams, Neil Corrigan, Julie Croft, Alison Pullan, Vicky Napp, Rachel Kelly, David Meads, Armando Vargas-Palacios, Adam Martin, Claire Hulme, Steven R Brown, Karen Nugent, Jen Lodge, David Protheroe, Sushil Maslekar, Andrew Clarke, Pasha Nisar and Julia M Brown



Sacral nerve stimulation versus the magnetic sphincter augmentation device for adult faecal incontinence: the SaFaRI RCT

David G Jayne^{1*}, Annabelle E Williams²,
Neil Corrigan³, Julie Croft³, Alison Pullan³,
Vicky Napp³, Rachel Kelly³, David Meads⁴,
Armando Vargas-Palacios⁴, Adam Martin⁴,
Claire Hulme⁵, Steven R Brown⁶, Karen Nugent⁷,
Jen Lodge⁸, David Protheroe⁹, Sushil Maslekar¹⁰,
Andrew Clarke¹¹, Pasha Nisar¹² and Julia M Brown³

¹Academic Surgery, Leeds Institute of Medical Research at St James's, University of Leeds, Leeds, UK

²Colorectal and General Surgery, Milton Keynes University Hospital, Milton Keynes, UK

³Clinical Trials Research Unit, Leeds Institute of Clinical Trials Research, University of Leeds, Leeds, UK

⁴Academic Unit of Health Economics, Leeds Institute of Health Sciences, University of Leeds, Leeds, UK

⁵Health Economics Group, Institute of Health Research, University of Exeter Medical School, Exeter, UK

⁶Department of Colorectal Surgery, Northern General Hospital, Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK

⁷University of Southampton, Southampton, UK

⁸Bowel Health and Pelvic Floor Dysfunction Community Urology and Colorectal Service (CUCS), Leeds Community Healthcare NHS Trust, Leeds, UK

⁹Department of Liaison Psychiatry, Leeds General Infirmary, Leeds and York Partnership NHS Foundation Trust, Leeds, UK

¹⁰St James's Hospital, Leeds Teaching Hospitals NHS Trust, Leeds, UK

¹¹Poole Hospital NHS Foundation Trust, Poole, UK

¹²St Peter's Hospital, Ashford and St Peter's Hospitals NHS Foundation Trust, Chertsey, UK

*Corresponding author

Declared competing interests of authors: David G Jayne reports grants from National Institute for Health Research (NIHR) Senior Investigator awards, and that he was a member of the NIHR Efficacy and Mechanism Evaluation (EME) Strategy Group and Prioritisation Group (2015–18) and NIHR Clinical Scientist Awards Panel (2015–2018). He currently sits on the NIHR Advanced Fellowship Panel (2018 to present) and is a member of the NIHR i4i Product Development Awards Committee (2019 to present). Julia M Brown reports grants from NIHR Senior Investigator awards and the NIHR Funding Committee during the conduct of the study, and declares membership of the Health Technology Assessment (HTA) Remit and Competitiveness Group (2016 to present), Clinical Trial Units funded by NIHR, the HTA Funding Committee Policy Group (2016 to present) and the HTA Clinical Evaluation and Trials Committee (2016 to present). Claire Hulme was a member of the NIHR HTA Commissioning Board (2013–17). Steven Brown was a member of the HTA Commissioning Board (2018–19). David Meads was a member of the NIHR HTA Elective and Emergency Specialist Care (EESC) Methods Group (2014–17) and NIHR HTA EESC Panel (2013–17). He is a member of a NIHR Programme Grants for Applied Research subpanel (2017 to present).

Published March 2021

DOI: 10.3310/hta25180

This report should be referenced as follows:

Jayne DG, Williams AE, Corrigan N, Croft J, Pullan A, Napp V, *et al.* Sacral nerve stimulation versus the magnetic sphincter augmentation device for adult faecal incontinence: the SaFaRI RCT. *Health Technol Assess* 2021;**25**(18).

Health Technology Assessment is indexed and abstracted in *Index Medicus/MEDLINE*, *Excerpta Medica/EMBASE*, *Science Citation Index Expanded (SciSearch®)* and *Current Contents®/Clinical Medicine*.

Health Technology Assessment

ISSN 1366-5278 (Print)

ISSN 2046-4924 (Online)

Impact factor: 3.370

Health Technology Assessment is indexed in MEDLINE, CINAHL, EMBASE, the Cochrane Library and Clarivate Analytics Science Citation Index.

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics (COPE) (www.publicationethics.org/).

Editorial contact: journals.library@nihr.ac.uk

The full HTA archive is freely available to view online at www.journalslibrary.nihr.ac.uk/hta. Print-on-demand copies can be purchased from the report pages of the NIHR Journals Library website: www.journalslibrary.nihr.ac.uk

Criteria for inclusion in the *Health Technology Assessment* journal

Reports are published in *Health Technology Assessment* (HTA) if (1) they have resulted from work for the HTA programme, and (2) they are of a sufficiently high scientific quality as assessed by the reviewers and editors.

Reviews in *Health Technology Assessment* are termed 'systematic' when the account of the search appraisal and synthesis methods (to minimise biases and random errors) would, in theory, permit the replication of the review by others.

HTA programme

Health Technology Assessment (HTA) research is undertaken where some evidence already exists to show that a technology can be effective and this needs to be compared to the current standard intervention to see which works best. Research can evaluate any intervention used in the treatment, prevention or diagnosis of disease, provided the study outcomes lead to findings that have the potential to be of direct benefit to NHS patients. Technologies in this context mean any method used to promote health; prevent and treat disease; and improve rehabilitation or long-term care. They are not confined to new drugs and include any intervention used in the treatment, prevention or diagnosis of disease.

The journal is indexed in NHS Evidence via its abstracts included in MEDLINE and its Technology Assessment Reports inform National Institute for Health and Care Excellence (NICE) guidance. HTA research is also an important source of evidence for National Screening Committee (NSC) policy decisions.

This report

The research reported in this issue of the journal was funded by the HTA programme as project number 12/35/07. The contractual start date was in January 2014. The draft report began editorial review in June 2019 and was accepted for publication in October 2020. The authors have been wholly responsible for all data collection, analysis and interpretation, and for writing up their work. The HTA editors and publisher have tried to ensure the accuracy of the authors' report and would like to thank the reviewers for their constructive comments on the draft document. However, they do not accept liability for damages or losses arising from material published in this report.

This report presents independent research funded by the National Institute for Health Research (NIHR). The views and opinions expressed by authors in this publication are those of the authors and do not necessarily reflect those of the NHS, the NIHR, NETSCC, the HTA programme or the Department of Health and Social Care. If there are verbatim quotations included in this publication the views and opinions expressed by the interviewees are those of the interviewees and do not necessarily reflect those of the authors, those of the NHS, the NIHR, NETSCC, the HTA programme or the Department of Health and Social Care.

© Queen's Printer and Controller of HMSO 2021. This work was produced by Jayne *et al.* under the terms of a commissioning contract issued by the Secretary of State for Health and Social Care. This issue may be freely reproduced for the purposes of private research and study and extracts (or indeed, the full report) may be included in professional journals provided that suitable acknowledgement is made and the reproduction is not associated with any form of advertising. Applications for commercial reproduction should be addressed to: NIHR Journals Library, National Institute for Health Research, Evaluation, Trials and Studies Coordinating Centre, Alpha House, University of Southampton Science Park, Southampton SO16 7NS, UK.

Published by the NIHR Journals Library (www.journalslibrary.nihr.ac.uk), produced by Prepress Projects Ltd, Perth, Scotland (www.prepress-projects.co.uk).

NIHR Journals Library Editor-in-Chief

Professor Ken Stein Professor of Public Health, University of Exeter Medical School, UK

NIHR Journals Library Editors

Professor John Powell Chair of HTA and EME Editorial Board and Editor-in-Chief of HTA and EME journals. Consultant Clinical Adviser, National Institute for Health and Care Excellence (NICE), UK, and Professor of Digital Health Care, Nuffield Department of Primary Care Health Sciences, University of Oxford, UK

Professor Andrée Le May Chair of NIHR Journals Library Editorial Group (HS&DR, PGfAR, PHR journals) and Editor-in-Chief of HS&DR, PGfAR, PHR journals

Professor Matthias Beck Professor of Management, Cork University Business School, Department of Management and Marketing, University College Cork, Ireland

Dr Tessa Crilly Director, Crystal Blue Consulting Ltd, UK

Dr Eugenia Cronin Senior Scientific Advisor, Wessex Institute, UK

Dr Peter Davidson Consultant Advisor, Wessex Institute, University of Southampton, UK

Ms Tara Lamont Senior Scientific Adviser (Evidence Use), Wessex Institute, University of Southampton, UK

Dr Catriona McDaid Senior Research Fellow, York Trials Unit, Department of Health Sciences, University of York, UK

Professor William McGuire Professor of Child Health, Hull York Medical School, University of York, UK

Professor Geoffrey Meads Emeritus Professor of Wellbeing Research, University of Winchester, UK

Professor James Raftery Professor of Health Technology Assessment, Wessex Institute, Faculty of Medicine, University of Southampton, UK

Dr Rob Riemsma Reviews Manager, Kleijnen Systematic Reviews Ltd, UK

Professor Helen Roberts Professor of Child Health Research, UCL Great Ormond Street Institute of Child Health, UK

Professor Jonathan Ross Professor of Sexual Health and HIV, University Hospital Birmingham, UK

Professor Helen Snooks Professor of Health Services Research, Institute of Life Science, College of Medicine, Swansea University, UK

Professor Ken Stein Professor of Public Health, University of Exeter Medical School, UK

Professor Jim Thornton Professor of Obstetrics and Gynaecology, Faculty of Medicine and Health Sciences, University of Nottingham, UK

Please visit the website for a list of editors: www.journalslibrary.nihr.ac.uk/about/editors

Editorial contact: journals.library@nihr.ac.uk

Abstract

Sacral nerve stimulation versus the magnetic sphincter augmentation device for adult faecal incontinence: the SaFaRI RCT

David G Jayne^{1*}, Annabelle E Williams², Neil Corrigan³, Julie Croft³, Alison Pullan³, Vicky Napp³, Rachel Kelly³, David Meads⁴, Armando Vargas-Palacios⁴, Adam Martin⁴, Claire Hulme⁵, Steven R Brown⁶, Karen Nugent⁷, Jen Lodge⁸, David Protheroe⁹, Sushil Maslekar¹⁰, Andrew Clarke¹¹, Pasha Nisar¹² and Julia M Brown³

¹Academic Surgery, Leeds Institute of Medical Research at St James's, University of Leeds, Leeds, UK

²Colorectal and General Surgery, Milton Keynes University Hospital, Milton Keynes, UK

³Clinical Trials Research Unit, Leeds Institute of Clinical Trials Research, University of Leeds, Leeds, UK

⁴Academic Unit of Health Economics, Leeds Institute of Health Sciences, University of Leeds, Leeds, UK

⁵Health Economics Group, Institute of Health Research, University of Exeter Medical School, Exeter, UK

⁶Department of Colorectal Surgery, Northern General Hospital, Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK

⁷University of Southampton, Southampton, UK

⁸Bowel Health and Pelvic Floor Dysfunction Community Urology and Colorectal Service (CUCS), Leeds Community Healthcare NHS Trust, Leeds, UK

⁹Department of Liaison Psychiatry, Leeds General Infirmary, Leeds and York Partnership NHS Foundation Trust, Leeds, UK

¹⁰St James's Hospital, Leeds Teaching Hospitals NHS Trust, Leeds, UK

¹¹Poole Hospital NHS Foundation Trust, Poole, UK

¹²St Peter's Hospital, Ashford and St Peter's Hospitals NHS Foundation Trust, Chertsey, UK

*Corresponding author d.g.jayne@leeds.ac.uk

Background: Preliminary studies using the FENIX™ (Torax Medical, Minneapolis, MN, USA) magnetic sphincter augmentation device suggest that it is safe to use for the treatment of adult faecal incontinence, but efficacy data are limited.

Objective: To compare FENIX with sacral nerve stimulation for the treatment of adult faecal incontinence in terms of safety, efficacy, quality of life and cost-effectiveness.

Design, setting and participants: Multicentre, parallel-group, unblinded, randomised trial comparing FENIX with sacral nerve stimulation in participants suffering moderate to severe faecal incontinence.

Interventions: Participants were randomised on an equal basis to either sacral nerve stimulation or FENIX. Follow-up occurred 2 weeks postoperatively and at 6, 12 and 18 months post randomisation.

Main outcome and measure: The primary outcome was success, defined as device in use and ≥ 50% improvement in Cleveland Clinic Incontinence Score at 18 months post randomisation. Secondary outcomes included complication rates, quality of life and cost-effectiveness. Between 30 October 2014 and 23 March 2017, 99 participants were randomised across 18 NHS sites

(50 participants to FENIX vs. 49 participants to sacral nerve stimulation). The median time from randomisation to FENIX implantation was 57.0 days (range 4.0–416.0 days), and the median time from randomisation to permanent sacral nerve stimulation was 371.0 days (range 86.0–918.0 days). A total of 45 out of 50 participants underwent FENIX implantation and 29 out of 49 participants continued to permanent sacral nerve stimulation. The following results are reported, excluding participants for whom the corresponding outcome was not evaluable. Overall, there was success for 10 out of 80 (12.5%) participants, with no statistically significant difference between the two groups [FENIX 6/41 (14.6%) participants vs. sacral nerve stimulation 4/39 (10.3%) participants]. At least one postoperative complication was experienced by 33 out of 45 (73.3%) participants in the FENIX group and 9 out of 40 (22.5%) participants in the sacral nerve stimulation group. A total of 15 out of 50 (30%) participants in the FENIX group ultimately had to have their device explanted. Slightly higher costs and quality-adjusted life-years (incremental = £305.50 and 0.005, respectively) were observed in the FENIX group than in the sacral nerve stimulation group. This was reversed over the lifetime horizon (incremental = –£1306 and –0.23 for costs and quality-adjusted life-years, respectively), when sacral nerve stimulation was the optimal option (net monetary benefit = –£3283), with only a 45% chance of FENIX being cost-effective.

Limitations: The SaFaRI study was terminated in 2017, having recruited 99 participants of the target sample size of 350 participants. The study is, therefore, substantially underpowered to detect differences between the treatment groups, with significant uncertainty in the cost-effectiveness analysis.

Conclusions: The SaFaRI study revealed inefficiencies in the treatment pathways for faecal incontinence, particularly for sacral nerve stimulation. The success of both FENIX and sacral nerve stimulation was much lower than previously reported, with high postoperative morbidity in the FENIX group.

Future work: Further research is needed to clarify the treatment pathways for sacral nerve stimulation and to determine its true clinical and cost-effectiveness.

Trial registration: Current Controlled Trials ISRCTN16077538.

Funding: This project was funded by the National Institute for Health Research (NIHR) Health Technology Assessment programme and will be published in full in *Health Technology Assessment*; Vol. 25, No. 18. See the NIHR Journals Library website for further project information.

Contents

List of tables	xi
List of figures	xiii
List of abbreviations	xv
Plain English summary	xvii
Scientific summary	xix
Chapter 1 Treatment options for faecal incontinence and rationale for SaFaRI clinical trial	1
Introduction	1
<i>Current treatment options</i>	1
<i>Rationale for the SaFaRI trial</i>	4
Methods	5
<i>Aim and objectives</i>	5
<i>Trial design</i>	6
<i>Participants</i>	6
<i>Interventions</i>	7
<i>Summary of protocol changes</i>	8
<i>Early trial closure</i>	8
<i>End points</i>	10
<i>Sample size</i>	10
<i>Randomisation</i>	10
<i>Blinding</i>	10
<i>Statistical methods</i>	11
Chapter 2 Results	13
Recruitment	13
Baseline data	13
Primary end point: device in use and $\geq 50\%$ improvement in Cleveland Clinic Incontinence Score at 18 months post randomisation	15
<i>Sensitivity analysis: additional covariates</i>	21
<i>Longitudinal analysis</i>	22
Secondary end point: device in use and $\geq 50\%$ improvement in Cleveland Clinic Incontinence Score at 12 months post randomisation	22
Secondary end point: device in use and $\geq 50\%$ improvement in Cleveland Clinic Incontinence Score at 6 months post randomisation	23
Secondary end point: intraoperative complications	24
Secondary end point: postoperative complications	24
Secondary end point: device explants	26
Secondary end point: Cleveland Clinic Incontinence Score	27
Secondary end point: EuroQol-5 Dimensions, five-level version	29
Secondary end point: faecal incontinence quality of life	30
Secondary end point: Obstructed Defecation Score	30
Secondary end point: Short Form questionnaire-12 items	30
Registered training cases	35

CONTENTS

Chapter 3 Health economics	37
Methods: trial	37
<i>Health utility and quality-adjusted life-year calculation</i>	37
<i>Resource use and costs</i>	37
<i>Analysis</i>	39
Methods: decision model	40
<i>Model type, cycle length, structure and health states</i>	40
<i>Parameter values: transition probabilities</i>	42
<i>Parameter values: health state utilities</i>	44
<i>Parameter values: costs</i>	44
<i>Validation</i>	45
<i>Analysis</i>	45
Results	46
<i>Trial analysis</i>	46
<i>Decision modelling</i>	48
Chapter 4 Discussion	53
Limitations	55
Future research	56
Chapter 5 Conclusions	57
Acknowledgements	59
References	61
Appendix 1	65

List of tables

TABLE 1 Summary of protocol changes	9
TABLE 2 Recruitment by site	13
TABLE 3 Minimisation factors	15
TABLE 4 Baseline characteristics	16
TABLE 5 Characteristics for participants included in primary analysis	20
TABLE 6 Number of successes at 18 months post randomisation	21
TABLE 7 Fixed effects of primary end point model	21
TABLE 8 Fixed effects of longitudinal model	22
TABLE 9 Number of successes at 12 months post randomisation	23
TABLE 10 Number of successes at 6 months post randomisation	23
TABLE 11 Number of patients experiencing at least one postoperative complication	24
TABLE 12 Time from operation to first complication	24
TABLE 13 Complications in FENIX arm	25
TABLE 14 Postoperative complications regression model	26
TABLE 15 Device status at 18 months post randomisation	26
TABLE 16 Reasons for explant	28
TABLE 17 Crude summaries of CCIS at baseline and at 6, 12 and 18 months post randomisation	28
TABLE 18 Crude summaries of EQ-5D-5L score and EQ-5D-5L VAS at baseline and at 6, 12 and 18 months post randomisation	29
TABLE 19 Crude summaries of FIQoL domains at baseline and at 6, 12 and 18 months post randomisation	31
TABLE 20 Crude summaries of ODS at baseline and at 6, 12 and 18 months post randomisation	33
TABLE 21 Crude summaries of SF-12 PCS and MCS components at baseline and at 6, 12 and 18 months post randomisation	34
TABLE 22 Device and procedure costs	38

LIST OF TABLES

TABLE 23	Parameter values: transition probabilities and event rates	42
TABLE 24	Parameter values: utilities and costs	44
TABLE 25	Cost breakdown	46
TABLE 26	Cost-effectiveness at 18 months	47
TABLE 27	Deterministic and probabilistic base-case cost-effectiveness results	49
TABLE 28	Deterministic one-way and scenario analyses	50
TABLE 29	Expected value of perfect information ($\lambda = \text{£}20,000$)	51
TABLE 30	Success defined by device in use and $\geq 50\%$ improvement in CCIS	65
TABLE 31	Success defined by $\geq 50\%$ improvement in CCIS	65
TABLE 32	Success defined by device in use	66
TABLE 33	Cross-tabulation of the components	66
TABLE 34	Fixed effects	67
TABLE 35	Fixed effects of EQ-5D-5L model	69
TABLE 36	Lifestyle domain model results: fixed effects	72
TABLE 37	Obstructed Defecation Score model results: fixed effects	77
TABLE 38	Fixed effect estimates of PCS model	79
TABLE 39	Complication costs (secondary care)	84
TABLE 40	Further treatment/surgery costs (secondary care)	86
TABLE 41	Medication costs	88
TABLE 42	Primary care costs	93
TABLE 43	Other service use	94
TABLE 44	Descriptive statistics	94

List of figures

FIGURE 1	The CONSORT flow diagram	14
FIGURE 2	Time from randomisation to surgery	18
FIGURE 3	Diagram of participants in primary analysis	19
FIGURE 4	Longitudinal model results	22
FIGURE 5	Time to explant: (a) FENIX MSA; and (b) SNS	27
FIGURE 6	Decision model structure	41
FIGURE 7	Cost-effectiveness acceptability curve	48
FIGURE 8	Markov model traces: (a) SNS; and (b) FENIX	49
FIGURE 9	Cost-effectiveness plane: SNS vs. FENIX	50
FIGURE 10	Cost-effectiveness acceptability curve	50
FIGURE 11	Cleveland Clinic Incontinence Score model results	68
FIGURE 12	EQ-5D score model results	69
FIGURE 13	EQ-5D VAS model results	70
FIGURE 14	Model residuals for EQ-5D-5L score	71
FIGURE 15	Model residuals for EQ-5D VAS	71
FIGURE 16	The FIQoL lifestyle model results	74
FIGURE 17	The FIQoL coping model results	74
FIGURE 18	The FIQoL depression model results	74
FIGURE 19	The FIQoL embarrassment model results	75
FIGURE 20	Model residuals for FIQoL: lifestyle	75
FIGURE 21	Model residuals for FIQoL: coping	76
FIGURE 22	Model residuals for FIQoL: depression	76
FIGURE 23	Model residuals for FIQoL: embarrassment	77
FIGURE 24	Obstructed Defecation Score model results	78
FIGURE 25	Model residuals for ODS	78

LIST OF FIGURES

FIGURE 26 Physical Component Summary model results	80
FIGURE 27 Mental Component Summary model results	80
FIGURE 28 Model residuals for SF-12: PCS	81
FIGURE 29 Model residuals for SF-12: MCS	81
FIGURE 30 Empirical probability plot: primary end point	82
FIGURE 31 Delta-betas: primary end point	82
FIGURE 32 Empirical probability plot: postoperative complications	83
FIGURE 33 Delta-betas: postoperative complications	83
FIGURE 34 Model residuals for CCIS end point	84

List of abbreviations

ABS	artificial bowel sphincter	MCS	Mental Component Summary
ASA	American Society of Anesthesiologists	MI	multiple imputation
BMI	body mass index	MSA	magnetic sphincter augmentation
BNF	<i>British National Formulary</i>	NICE	National Institute for Health and Care Excellence
CCIS	Cleveland Clinic Incontinence Score	NIHR HSC	National Institute for Health Research Horizon Scanning Centre
CEAC	cost-effectiveness acceptability curve	NMB	net monetary benefit
CI	confidence interval	ODS	Obstructed Defecation Score
CONSORT	Consolidated Standards of Reporting Trials	PCS	Physical Component Summary
CRF	case report form	PPI	patient and public involvement
CTRU	Clinical Trials Research Unit	PSA	probabilistic sensitivity analysis
DMEC	Data Monitoring and Ethics Committee	PSSRU	Personal Social Services Research Unit
EQ-5D	EuroQol-5 Dimensions	QALY	quality-adjusted life-year
EQ-5D-5L	EuroQol-5 Dimensions, five-level version	QoL	quality of life
EVPI	expected value of perfect information	RCT	randomised controlled trial
FI	faecal incontinence	SE	standard error
FIQoL	faecal incontinence quality of life	SF-12	Short Form questionnaire-12 items
GP	general practitioner	SF-6D	Short Form questionnaire-6 Dimensions
HRQoL	health-related quality of life	SNS	sacral nerve stimulation
ICER	incremental cost-effectiveness ratio	TSC	Trial Steering Committee
MAS	magnetic anal sphincter	USC	unexpected serious complication
		VAS	Visual Analogue Scale

Plain English summary

Faecal incontinence is a distressing condition for patients, and surgery is recommended if symptoms are having an effect on quality of life. One of the treatments recommended for faecal incontinence by the National Institute for Health and Care Excellence is sacral nerve stimulation, which aims to improve continence by stimulating the nerves to the back passage. A newer treatment involves surgery to implant a string of magnetic beads around the anal canal using the FENIX™ device (Torax Medical, Minneapolis, MN, USA). The aim of this study was to assess the benefits and risks of the FENIX device compared with sacral nerve stimulation.

The SaFaRI study aimed to recruit 350 participants with faecal incontinence, but was stopped early because of the manufacturer withdrawing the FENIX device for strategic reasons. In total, we recruited 99 participants. Fifty participants were allocated to receive the FENIX device and 49 participants were allocated to receive sacral nerve stimulation. The observed success rates with both devices were low: at 18 months following their entry into the study, 6 out of 41 (14.6%) participants in the FENIX group and 4 out of 39 (10.3%) participants in the sacral nerve stimulation group had the device both in use and producing a benefit. A total of 5 out of 50 (10.0%) participants allocated to receive the FENIX device did not have a device implanted, and 15 out of 45 (33.3%) participants who did have the FENIX device implanted needed to have it removed because of complications during the 18-month follow-up period. A total of 21 out of 49 (42.9%) participants allocated to receive sacral nerve stimulation did not have a permanent sacral nerve stimulation device implanted, and 0 of the 28 who did have a permanent sacral nerve stimulation device implanted needed to have it removed during the 18-month follow-up period. The costs associated with the FENIX device were higher because of a greater number of participants experiencing complications, meaning that the FENIX device is unlikely to be cost-effective in the treatment of faecal incontinence compared with sacral nerve stimulation.

Scientific summary

Background

Faecal incontinence is a distressing condition that affects between 5% and 10% of the adult population. Current treatment options include conservative measures with dietary modification and constipating agents, pelvic floor physiotherapy with or without biofeedback therapy, and surgical intervention for patients with moderate to severe symptoms. The most commonly used surgical intervention for patients with faecal incontinence resistant to medical treatment is sacral nerve stimulation. Sacral nerve stimulation involves a two-stage procedure, whereby a temporary stimulation phase is used to assess initial efficacy and if a $\geq 50\%$ reduction in weekly incontinence episodes or incontinence score is observed, then the patient may proceed to a permanent sacral nerve stimulation implant. Sacral nerve stimulation is recommended by the National Institute for Health and Care Excellence and, although short-term efficacy is reported to be good, the long-term efficacy as reported from a decision-to-treat perspective is only around 45–50%. Despite the high costs of the permanent implant, sacral nerve stimulation is believed to be cost-effective compared with the only other definitive treatment for faecal incontinence: a permanent colostomy.

More recently, a new device, FENIX™ (Torax Medical, Minneapolis, MN, USA), has been introduced to the market, consisting of a string of magnetic beads that is implanted around the anal canal to augment the anal sphincter. Initial results from small, single-centre studies have been promising, suggesting a $\geq 50\%$ improvement in incontinence in around 70% of participants, with a complication rate of 20% and a device explant rate of 10%. Only one previous randomised trial has been undertaken to evaluate the FENIX device, comparing it with the Acticon Neosphincter® (American Medical Systems, Minneapolis, MN, USA) and showing benefits in terms of shorter operating times and length of hospitalisation, and reduced costs.

Objectives

The objective was to undertake a randomised comparison of the safety, efficacy, and cost-effectiveness of the FENIX device with sacral nerve stimulation for the treatment of adults with moderate to severe faecal incontinence that is resistant to medical therapies. The primary outcome was success of the intervention (FENIX or sacral nerve stimulation) defined as the device in use and $\geq 50\%$ improvement in the participant-reported Cleveland Clinic Incontinence Score at 18 months post randomisation. The secondary outcomes included length of hospital stay, complications, reinterventions, constipation, quality of life and cost-effectiveness.

Methods

A multicentre randomised controlled trial was undertaken across 18 NHS hospital trusts involving colorectal surgeons who were members of the Association of Coloproctology of Great Britain and Ireland. It was estimated that 350 participants would be required to detect at least a 20% difference in the percentage of successes at 18 months post randomisation, where success was defined as the device in use and $\geq 50\%$ improvement in Cleveland Clinic Incontinence Score from baseline. Prior to randomising participants, all surgeons had to have performed a minimum of 10 permanent sacral nerve stimulation implants, observed a minimum of one FENIX procedure and performed two FENIX procedures under proctorship with data captured in the registration phase of the study. For inclusion in the study, participants had to be aged ≥ 18 years and to have been suffering from faecal incontinence

for more than 6 months with ≥ 2 incontinent episodes per week. The aetiology of incontinence was not specified and an anal sphincter defect of $\leq 180^\circ$, as assessed by endoanal ultrasound scan, was allowed. Participants were randomised equally to either FENIX or sacral nerve stimulation treatment, with minimisation factors including the treating surgeon, participant sex, severity of incontinence and degree of anal sphincter defect. The technique for FENIX implantation was standardised according to the manufacturer's guidance, whereas the technique of sacral nerve stimulation was left to the surgeon's normal practice. Data were captured at baseline, 2 weeks post intervention and at 6, 12 and 18 months post randomisation. The Cleveland Clinic Incontinence Score was used to assess incontinence and the Obstructed Defaecation Score was used to assess constipation. Quality of life was assessed by participant-reported questionnaires, using the faecal incontinence quality of life, EuroQol-5 Dimensions, five-level version and Visual Analogue Scale, and SF-12 questionnaires. In addition, participants completed the Health and Social Care Resource Use questionnaire.

All analyses were prespecified and conducted on an intention-to-treat basis. The results are reported with 95% confidence intervals and *p*-values for fixed effects. For all end points, missing outcome data were assumed to be missing at random, and the treatment effect was estimated via maximum likelihood estimation using all participants with non-missing outcome data for non-longitudinal end points. A sensitivity analysis of the primary end point was performed considering other covariates thought to be related to participant outcome. All modelling was performed using SAS® (SAS Institute Inc., Cary, NC, USA) version 9.4 glimmix procedure.

The economic evaluation was from the perspective of the social and health-care provider generating cost per quality-adjusted life-year over a lifetime horizon. We analysed trial cost and quality-adjusted life-year data and extrapolated these forward using a de novo decision-analytic model.

Results

The study was prematurely stopped in 2017 when the manufacturing company was bought by a multinational company and the FENIX device was withdrawn from the market. Between 30 October 2014 and 23 March 2017, 322 participants were assessed for eligibility; 23 participants were registered as training cases, and 99 participants were randomised into the study (50 FENIX and 49 sacral nerve stimulation). The baseline characteristics of the two groups were similar and in keeping with a population suffering from moderate to severe faecal incontinence. The median time from randomisation to FENIX implantation was 57.0 days (range 4.0–416.0 days). The median time from randomisation to temporary sacral nerve stimulation was 86.5 days (range 2.0–699.0 days) and the median time from randomisation to permanent sacral nerve stimulation was 371.0 days (range 86.0–918.0). Five out of 50 participants did not undergo FENIX implantation, and 5 out of 49 participants did not undergo temporary sacral nerve stimulation. A total of 32 participants continued to permanent sacral nerve stimulation, of whom three did not have a device implanted. For the primary end-point analysis, 19 participants had missing data, meaning that there was complete data available for analysis for 80 out of 99 (80.8%) participants.

Overall, there was success for 10 out of 80 (12.5%) participants, with no statistically significant difference between the two groups [FENIX 6/41 (14.6%) participants vs. sacral nerve stimulation 4/39 (10.3%) participants]. A longitudinal analysis using data obtained at 6, 12 and 18 months post randomisation did not show a statistically significant difference between the treatment groups, with no significant difference over time.

There were four intraoperative complications in four participants: three during FENIX implantation and one during implantation of a permanent sacral nerve stimulation. A total of 42 out of 85 participants experienced at least one postoperative complication: 33 out of 45 (73.3%) in the FENIX group and 9 out of 40 (22.5%) in the sacral nerve stimulation group. The adjusted odds ratio revealed a statistically

significant difference between the two treatments (11.21, 95% confidence interval 2.65–47.35; $p = 0.004$). A total of 15 out of 50 (30%) of the FENIX devices were explanted, usually within 6 months; there were no explants in the sacral nerve stimulation group.

Data were available from 96 out of 99 (97%) participants for analysis of the secondary end point: the Cleveland Clinic Incontinence Score. The results showed that having a device in use led to a statistically significant reduction in the Cleveland Clinic Incontinence Score of 3.04 points, but with no difference observed between the treatment groups. The quality-of-life analysis using the EuroQol-5 Dimensions, five-level version questionnaire showed a statistically significant improvement in quality of life in participants with the device in use (EuroQol-5 Dimensions, five-level version score 0.13 higher; $p = 0.004$), with no difference between the treatment groups. This finding was not replicated in the analysis of the Visual Analogue Scale score, with the randomised treatment producing no benefit and no difference observed between the treatment groups over time. Analysis of faecal incontinence quality of life showed a statistically significant improvement across all four domains when the device was in use, but no significant difference between the treatment groups. Despite obstructed defaecation being the most commonly reported complication following FENIX implantation, no significant difference was observed between the treatment groups, and whether or not a participant had a device in use did not produce a statistically significant difference in the Obstructed Defecation Score. The benefits of having a device in use appeared to be due to improvements in physical component scores rather than mental component scores, as assessed by the Short Form questionnaire-12 items, with no differences between the treatment groups.

At the end of the trial period, slightly higher costs and quality-adjusted life-years (incremental = £305.50 and 0.005, respectively) were observed in the FENIX arm. The sample size available for the analysis limits our ability to draw robust conclusions. The trial results were reversed over the lifetime horizon (incremental = –£1306 and –0.23 for costs and quality-adjusted life-years, respectively) with sacral nerve stimulation being the optimal option (net monetary benefit = –£3283). These analyses were relatively robust to deterministic sensitivity analyses; however, there was significant uncertainty, with sacral nerve stimulation having only a 55% chance of cost-effectiveness over a lifetime. Given the small sample sizes available for parameter value generation, caution is needed in interpreting the results.

Conclusions

Interpretation of the results is limited because of the early termination of the study, which means that the numbers available for analysis are small, with a high proportion of participants in the sacral nerve stimulation group not undergoing permanent sacral nerve stimulation implantation or not completing the 18-month follow-up. With this caveat, the rates of success for both FENIX and sacral nerve stimulation are disappointing and much lower than previously reported in the literature. The complication rates associated with the FENIX device are high, with around one-third of participants undergoing explantation. For those participants for whom a device remained in use, there were benefits in terms of improvement in continence score and quality of life. Based on the cost-effectiveness analysis, if the FENIX device were still available, it is uncertain whether or not it would be recommended for routine use in patients with faecal incontinence given the costs associated with complications.

Trial registration

This trial is registered as ISRCTN16077538.

Funding

This project was funded by the National Institute for Health Research (NIHR) Health Technology Assessment programme and will be published in full in *Health Technology Assessment*; Vol. 25, No. 18. See the NIHR Journals Library website for further project information.

Chapter 1 Treatment options for faecal incontinence and rationale for SaFaRI clinical trial

Much of the text included in this chapter has been taken from the SaFaRI protocol.¹ The research team has previously published the protocol in the *International Journal of Colorectal Disease*. Reproduced with permission from Williams *et al.*¹ This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The text below includes minor additions and formatting changes to the original text.

Introduction

Faecal incontinence (FI) is a distressing condition that affects between 5% and 10% of the adult population. It is more common in female patients and with advancing age, and is the second most common cause of admission to a nursing home. It has an impact on social, physical and mental well-being and is a substantial burden on NHS resources.

Current treatment options

Current treatment strategies for adult FI are summarised in the National Institute for Health and Care Excellence (NICE) 2014 guidance.² All patients should undergo a thorough history and physical examination to determine the nature and severity of the problem, and to identify a probable aetiological cause. Initial management consists of a combination of patient education, dietary modification and antidiarrhoeal medication. If this is unsuccessful, investigation in the form of endoscopic visualisation of the colorectum, anorectal manometry (pudendal nerve testing optional), and endoanal ultrasound is performed to further characterise the underlying disorder and inform treatment options.

Conservative therapies

Conservative therapies include pelvic floor retraining, with or without biofeedback therapy, and irrigation techniques (rectal or antegrade irrigation). Biofeedback therapy aims to increase the patient's awareness of the muscles of continence and rectal sensation. Incontinent symptoms are improved in around 50% of patients, although there appears to be a significant placebo effect, with a marked decrease in efficacy on long-term follow-up.³ Rectal irrigation, for example using the Peristeen® system (Coloplast, Humlebæk, Denmark), aims to clear the rectum and lower colon of faecal residue. In the short term it can have beneficial effects, but as a long-term solution patients frequently find it unacceptably time-consuming and inconvenient. Recently, there has been interest in the use of bulking agents to augment the anal sphincter. Data on the efficacy of these agents is limited, but they may have a role in controlling minor incontinence or 'seepage', or where an isolated sphincter defect is causing incomplete closure of the anal canal.⁴

Surgical interventions

Surgical interventions are indicated for those patients with moderate to severe FI that is resistant to the conservative therapies listed in *Conservative therapies*.

Anterior sphincteroplasty, artificial bowel sphincter and dynamic graciloplasty

Anterior sphincteroplasty may be considered for patients with discrete sphincter defects, which occur typically as a result of obstetric injury. Through a perineal incision, the disrupted sphincter muscle is isolated and an overlapping sutured repair performed. Short-term results are reasonable, with 70% of patients reporting an improvement in continence; however, there is a drop-off in the longer term, with fewer than 50% of patients experiencing a benefit at 5 years.⁵ Patients who do worse following anterior sphincteroplasty include those with coexistent pudendal neuropathy, multiple sphincter defects or sphincter atrophy, and irritable bowel syndrome. Because of the poor long-term results, there has been a move away from sphincter repair, except in well-defined cases, and an increased enthusiasm for sacral nerve stimulation (SNS).

Another surgical intervention which may be considered to treat FI is the artificial bowel sphincter (ABS). The ABS consists of (1) a fluid-filled silicone cuff placed around the anus, (2) a fluid-filled, pressure-regulating balloon positioned in the abdominal wall and (3) a manual pump connecting these components, placed in either the labia majora or the scrotum. When the cuff is inflated, the anal canal is sealed. The fluid is transferred to the balloon by the manual pump, deflating the cuff and opening the anal canal to allow defaecation. A successfully functioning device improves continence and quality of life (QoL); however, it is expensive, with the device alone costing around £4000. The main problem with the ABS is the high complication rate. Revisional surgery is needed in between 12.5% and 50% of cases, with explantation rates between 16.7% and 41.2%.⁶ The majority of revisions are for cuff leaks that are thought to arise from microperforations caused by repeated cycles of inflation and deflation over a number of years. Most explantations are for infective complications. As a consequence, the ABS is not in common usage.

Dynamic graciloplasty involves mobilisation of the gracilis muscle from the inner thigh and wrapping around the anus to augment sphincter function. A neurostimulation device with an impulse generator is implanted to adapt the type II, fast-twitch muscle fibres to type I, slow twitch, fatigue-resistant fibres. The patient uses an external programming device to deactivate the electrical stimulation, relaxing the muscular contraction and enabling defaecation at a voluntary time. The success rate of the operation is between 40% and 60%.⁷ Like the ABS, the main problem is the high complication (infections, 28%; device malfunction, 15%; and leg pain, 13%) and reintervention rates. The use of dynamic graciloplasty in the UK has largely been superseded by SNS.

Sacral nerve stimulation

Sacral nerve stimulation for FI was first described in 1995⁸ and has grown in popularity, gaining NICE recognition as a minimally invasive treatment for moderate to severe FI. SNS works by a combination of anal sphincter augmentation and modulation of spinal/supraspinal pathways. It benefits from a two-stage procedure, which enables the patient to assess acceptability and the clinician to evaluate efficacy prior to commitment to a permanent and expensive implant. An initial percutaneous nerve evaluation, or temporary stimulation, is performed under local, regional or general anaesthetic as a day-case procedure. A fine needle is inserted percutaneously into the sacral foramina (S3 or S4) on both sides to determine the best response in terms of anal sphincter contraction and dorsiflexion of the great toe (S3 stimulation). Once a satisfactory response is obtained, the temporary electrode is inserted, secured to the skin and connected to an external test stimulator, allowing the patient to alter the stimulation voltage. The patient is asked to keep a bowel diary for the 2–3 weeks of stimulation, which allows the clinician to quantify the degree of response. A positive response is defined as a reduction in incontinence episodes or incontinence score of $\geq 50\%$ during the stimulation period.

Around 70% of patients have a good response and proceed to a permanent implant. Of these, 10% never gain any significant improvement and 26% experience loss of efficacy, usually within the first year.^{5,9–11} A further 2–5% suffer irresolvable complications and undergo explantation. Thus, from a decision-to-treat perspective, the long-term efficacy is around 45–50%. Overall, only 50% of patients thought to be eligible for SNS have a functioning device in the long term.

The reasons for loss of efficacy are not clear, but may relate to device malfunction or fibrosis of the stimulating electrode leading to loss of conduction. Pain or discomfort at the stimulator site, down the leg or into the vagina is another commonly reported complication, experienced by 38.1% of patients. Overall, only 58.5% of patients who have a permanent implant have a good or acceptable result in the medium term.⁵

Although SNS is an effective treatment for FI, it is also very costly. The component costs alone (excluding other direct and indirect medical costs) are £200 for the test stimulation and £9393 for the permanent stimulator.¹² A European study has calculated the 5-year cumulative costs for SNS at €22,150 per patient, which compares with €33,996 for a colostomy and €3234 for conservative treatment.¹³ Despite this, SNS has been shown to be cost-effective. The incremental cost-effectiveness ratio (ICER) for SNS is £25,070 per quality-adjusted life-year (QALY) gained, which is within the £30,000 per QALY threshold recommended by NICE as an effective use of NHS resources.

NICE first issued its guidance on SNS for FI in 2004¹⁴ and concluded that current evidence on safety and efficacy appeared to support its use, but that the procedure should only be performed in specialist units by clinicians with a particular interest in the condition. A systematic review at that time included six case series and 266 patients. In patients who had permanent implants, complete continence was achieved in 41 to 75%, while 75 to 100% of patients experienced a decrease of $\geq 50\%$ in the number of incontinent episodes. Improvements were noted in both disease-specific and general QoL scores. The most recent review, including 13 studies and 929 patients, has confirmed the short-term efficacy of SNS.¹⁵ Although the extent of the therapeutic effect varied between studies, a significantly beneficial effect was noted. Functional improvement was observed in 77% with idiopathic FI, 76% in sphincter rupture/episiotomy, 78% after anal repair, and 73% after neurological injury. The benefit was not restricted to improved continence, with several studies showing a significant improvement in QoL.^{9,10,13}

FENIX™ continence restoration system (FENIX™ magnetic sphincter augmentation)

The FENIX™ continence restoration system, or FENIX™ magnetic sphincter augmentation (MSA) (Torax Medical, Minneapolis, MN, USA), is a device that has been designed to reinforce the native sphincter for the treatment of FI that is resistant to conservative therapies. It consists of a ring of 14–20 titanium beads with magnetic cores that are linked together to form a structure to be surgically placed around the anal sphincter complex. To defecate, the patient strains in a normal way and the force generated separates the beads to open the anal canal. Continence is restored by means of passive attraction of the beads. Once implanted, the device does not require patient input to function.

The FENIX MSA costs £4000. Data on efficacy are limited, but they suggest a $\geq 50\%$ improvement in continence in 70% of patients. Complications can occur in around 20% of patients, leading to explantation in around 10%.

Preliminary results are promising, with 70% of patients reporting a benefit; however, studies have been small and a more rigorous evaluation is required prior to its widespread adoption.

The device is manufactured in different lengths to accommodate variations in anal canal circumference, and has been CE (Conformité Européenne) marked since November 2011. FENIX MSA has been used in selected European and US centres to support a feasibility trial and was first used in the NHS in 2013.

The available evidence on safety and efficacy is limited but encouraging. Barussaud *et al.*¹⁶ published data on a series of 24 patients who were implanted with FENIX between 2008 and 2012. All patients were female, with a mean age of 64 years (range 35–78 years) and the mean duration of FI being 8.8 years (range 1–40 years). The mean follow-up was 17.6 months. There was one immediate

postoperative complication: cardiac arrest due to drug intolerance. The patient recovered without further sequelae. Two patients (8.7%) had the device explanted, one for device separation and one for perineal abscess at 6 months post implant. The procedure was considered a failure for five patients (21%) due to a lack of improvement in FI symptoms. Bowel diary results showed a significant improvement in the number of weekly FI episodes, decreasing from 32 to 8 in a 3-week diary. The mean Wexner score [Cleveland Clinic Incontinence Score (CCIS)] was reduced significantly from 16 points at baseline to 7 points, 8 points and 5 points at 12, 24 and 36 months, respectively. All four domains of the faecal incontinence quality-of-life (FIQoL) questionnaire scores significantly improved and remained stable postoperatively compared with the score at baseline.

A retrospective, case-matched comparison of the FENIX MSA with the ABS (Acticon® Neosphincter; American Medical Systems, Minneapolis, MN, USA) in 20 patients with severe FI¹⁷ showed that the FENIX MSA and ABS produced similar significant improvements in FI and QoL. Compared with the ABS, the FENIX MSA was associated with a significantly shorter operating time (FENIX MSA: 62 minutes vs. ABS: 97.5 minutes; $p = 0.0273$) and length of hospitalisation (FENIX MSA: 4.5 days vs. ABS: 10 days; $p = 0.001$). No difference was observed in postoperative complications. The ABS was associated with more explants/revisions (FENIX MSA: 1 vs. ABS: 4; $p = 0.830$), a greater incidence in postoperative constipation, and was more expensive.

Permanent stoma

For patients for whom the above surgical attempts fail to restore normal continence, the options are limited. A permanent stoma (usually colostomy) is often the last resort for patients with intractable FI. It is an effective strategy, but one that carries psychological and physical morbidity. Although most patients adapt to a permanent stoma, there is a continual fear of appliance leakage that can have an impact on social functioning. Around 50% of permanent stomas are complicated by parastomal herniation that may require surgical intervention. Moreover, a stoma is not a cheap intervention, with the 5-year cumulative costs estimated at £28,000.¹³

Rationale for the SaFaRI trial

New technologies have often been introduced into clinical practice without rigorous evaluation of safety, efficacy and cost-effectiveness. Objective assessment has been overlooked because of the intrinsic appeal of new innovation, the need to be a part of a 'pioneering group' or, worse, because of the financial incentives from industry. Once introduced, low-grade observational evidence is often used to keep practices going. As a result, it has often been easier to 'stop them starting' than to 'start them stopping'.¹⁸ Ideally, any new technology introduced into clinical practice should be simultaneously evaluated, and in most cases the best way of doing this is by randomised comparison with an already established technique. The National Institute for Health Research Horizon Scanning Centre (NIHR HSC) was established to 'supply timely information to key health policy and decision-makers within the NHS about emerging health technologies that may have a significant impact on patients or the provision of health services in the near future'.¹⁹ In May 2012, the NIHR HSC reported on the FENIX Continence Restoration System (FENIX MSA) and concluded that 'in order to determine its potential place in the pathway of care for FI larger long term studies of the safety, effectiveness and cost-effectiveness of FENIX in comparison to existing treatments are needed'.²⁰ Therefore, although FENIX MSA may have a role to play in the treatment of FI, the evidence was not robust enough to support its widespread adoption.

The SaFaRI trial was thus designed to undertake a rigorous, prospective assessment of the new FENIX MSA as it was adopted into the NHS. The aim was for reliable data, collected independently from commercial interests, to be made available on the safety and efficacy of the device. This would include information on safety, efficacy, QoL and cost-effectiveness. Important information would be gained on the costs associated with the device, enabling the ICER per QALY to be determined. This would allow health-care providers to make informed decisions about value for money and future provision of the technology.

Sacral nerve stimulation was chosen as the comparator to FENIX MSA as SNS is currently the preferred, and NICE-recommended,² surgical intervention for FI that is resistant to conservative therapies; the NIHR HSC report¹⁹ from May 2012 also identified SNS as the preferred comparator for any randomised comparison with FENIX MSA.

Furthermore, the SaFaRI trial was designed to collect additional, important data about SNS. SNS is a costly yet effective treatment for FI; however, concerns have been expressed about the lack of efficacy when analysed on an intention-to-treat basis and the loss of efficacy on longer-term follow-up. The SaFaRI study provided an additional opportunity to clarify the indications for SNS and the indicators of success.

The opportunity also presented itself to comprehensively document, for the first time, the treatment and associated costs for patients for whom either SNS or FENIX MSA is not successful. In effect, these patients would provide comparative, longitudinal data of the patient pathway where FENIX MSA or SNS is either unsuitable or unavailable.

In addition to the costs detailed above, the health economics would provide data on the short- and long-term cost-effectiveness of FENIX MSA compared with SNS. Within the analyses, use of two measures of health-related QoL to produce QALYs, the Short Form questionnaire-12 items (SF-12) together with the EuroQol-5 Dimensions (EQ-5D), would allow assessment of the sensitivity of the EuroQol-5 Dimensions, five-level version (EQ-5D-5L) to detect changes in FI, which is to date unproven. The disease-specific questionnaire chosen to assess QoL, the FIQoL questionnaire, collects important information on many social and psychological aspects of FI (shame, depression, enjoyment, etc.). These aspects of FI have received little previous recognition in the literature and remain poorly defined.

Methods

Aim and objectives

The overall objectives of the study were to:

- determine the short-term safety and efficacy of FENIX MSA and SNS in adult FI
- assess FENIX MSA and SNS in terms of impact on QoL and cost-effectiveness.

Aim

The aim was to conduct a thorough evaluation of the FENIX MSA device, compared with SNS, for the treatment of adult FI.

Primary outcome measure

The primary outcome measure was success, defined as device in use and $\geq 50\%$ improvement in the participant-reported CCIS at 18 months post randomisation.

Secondary outcome measures

- Length of hospital stay.
- Complications.
- Reinterventions.
- Constipation.
- QoL.
- Cost-effectiveness.

Trial design

SaFaRI was a prospective, UK multisite, parallel-group, randomised clinical study investigating the safety and efficacy of the FENIX MSA for adult FI. The comparator was SNS, a preferred treatment recommended by NICE for the treatment of FI that is resistant to conservative therapies.² Participants were randomised on a 1 : 1 basis to receive either FENIX MSA or SNS.

Prior to randomising participants, all participating surgeons had to have performed a minimum of 10 permanent SNS implantations, observed a minimum of one FENIX MSA procedure and performed two FENIX MSA procedures under proctorship.

A registration phase was incorporated into the study design to enable surgeons without the required FENIX MSA experience prior to study participation to gain the relevant experience within the scope of the study. Within the registration phase, the first two eligible patients providing consent were registered to the study under the training surgeon's name and received FENIX MSA implants (there was no randomisation in the registration phase); these two operations, which were performed under proctorship, were considered study training cases and were not included in the main study/main trial analysis. Once the required FENIX MSA experience had been obtained, the surgeon could progress to the randomisation phase.

The trial received national ethics approval in the UK. The trial conduct was overseen by an independent Trial Steering Committee (TSC) and Data Monitoring and Ethics Committee (DMEC). The trial had public and patient involvement during the trial design phase and throughout the course of the study. The trial was registered on the International Standard Randomised Controlled Trial Number (ISRCTN) register (16077538).

Participants

The inclusion criteria were as follows:

- aged ≥ 18 years
- able to provide written informed consent
- FI for > 6 months
- incontinent episodes of ≥ 2 per week
- suitable candidate for surgery, as judged by the operating surgeon
- suitable for either FENIX MSA or SNS (unless the patient was being registered as a training case, in which event they only needed to be suitable for the FENIX MSA)
- anal sphincter defect $< 180^\circ$ as documented on endoanal ultrasound scan
- able and willing to comply with the terms of the protocol including QoL questionnaires.

The exclusion criteria were as follows:

- previous interventions for FI (i.e. SNS, FENIX MSA or ABS) (unless the patient was registered as a training case, in which event they could have had previous interventions for FI)
- chronic gastrointestinal motility disorders causing incontinence due to diarrhoea
- obstructed defaecation, defined as an inability to satisfactorily evacuate the rectum [it was recommended that the Obstructed Defecation Score (ODS) was calculated and was ≤ 8 for trial inclusion]
- anal sphincter defect $\geq 180^\circ$, as documented on endoanal ultrasound scan
- an electric or metallic implant within 10 cm of anal canal
- co-existent systemic disease (e.g. scleroderma) affecting continence
- active anorectal sepsis
- diagnosis of colorectal or anal cancer within previous 2 years
- external rectal prolapse
- significant scarring of the anorectum that, as judged by the treating surgeon, would prohibit FENIX MSA implantation or put the patient at high risk of implant erosion

- pregnancy (it was the local surgeon's responsibility to assess pregnancy in women of childbearing potential)
- immunocompromised, including haematological abnormalities and treatment with steroids or other immunomodulatory medicines
- congenital spinal abnormalities, preventing SNS implantation
- known requirement for future magnetic resonance imaging surveillance, which would be contraindicated in the presence of metallic implant
- suspected or known allergies to titanium.

Interventions

Preoperative investigation and preparation were as per institutional protocol, which included, as standard practice, visualisation of the colorectum (flexible sigmoidoscopy as a minimum), anorectal manometry (pudendal nerve testing optional) and endoanal ultrasound.

Sacral nerve stimulation

Sacral nerve stimulation implantation was performed in accordance with each research site's usual practice. SNS implantation is a two-stage procedure. As per standard care, a temporary device was implanted during a day-case procedure and the degree of response to the device recorded by the participant over the course of 2 weeks. Response was assessed in accordance with each research site's usual practice. For the purposes of the trial, the CCIS was recorded at this time point regardless of how the response was assessed locally.

As per routine care, if the response was positive (defined as a $\geq 50\%$ improvement in incontinence episodes or $\geq 50\%$ improvement in CCIS), then a second day-case procedure was scheduled and a permanent SNS device was implanted. If the response was negative, the temporary device was removed and the participant did not receive any further study intervention but continued follow-up for the required 18-month period. Further treatment was as per standard practice but participants were not permitted to undergo FENIX MSA implantation during the 18-month post-randomisation follow-up period.

Postoperative care was as per routine care, but participants had to be reviewed at clinic for trial purposes 2 weeks postoperatively for both temporary and permanent device implants, and at 6, 12 and 18 months post randomisation as a minimum. Any further visits were according to local standard clinical practice, but were captured on the follow-up case report forms (CRFs).

FENIX magnetic sphincter augmentation

FENIX MSA implantation was usually performed during an in-patient stay (usually of 1–3 days). Participants for whom FENIX MSA failed were not permitted to undergo SNS during the 18-month follow-up period. No postoperative care was required above routine wound care, but participants had to be reviewed for trial purposes at 2 weeks postoperatively and at 6, 12 and 18 months post randomisation as a minimum. Any further visits were performed according to local standard clinical practice and were recorded on the follow-up CRFs.

Participant-completed questionnaires

Participants completed a number of questionnaires designed to capture FI symptoms prior to randomisation (baseline), at 2 weeks post operation and at 6, 12 and 18 months post randomisation:

- The CCIS.²¹ The CCIS assesses five parameters associated with incontinence – incontinence to solid, incontinence to liquid, incontinence to gas, use of pads, and lifestyle restriction. Each parameter is scored 0–4, with '0' for never and '4' for every day. The five parameters are added to give a total score out of 20.
- The ODS.²² The ODS consists of five items: excessive straining, incomplete rectal evacuation, use of enemas and/or laxatives, vaginal-anal-perineal digitations, and abdominal discomfort and/or pain. Each item is graded from 0 to 4 with a score ranging from 0 (no symptoms) to 20 (very severe symptoms).

- The FIQoL questionnaire.²³ This questionnaire is composed of 29 items that make up four scales: lifestyle (10 items), coping/behaviour (9 items), depression/self-perception (7 items) and embarrassment (3 items). Scoring is derived from a participant-completed questionnaire that assesses the impact of FI on four domains of QoL. Scales range from 1 to 5, with 1 indicating a lower functional QoL. Scale scores are derived by averaging the response to all items in the scale.
- Health and Social Care Resource Use. The questionnaire is composed of questions related to contact with primary, community and social care services. The questionnaire consists primarily of 'tick-box' completion questions.
- SF-12.²⁴ The SF-12 is a 12-item subset of the Short Form questionnaire-36 items version 2 that measures the same eight domains of health. It is a brief, reliable measure of overall health status. It is useful in large population health surveys and has been used extensively as a screening tool.
- EQ-5D-5L.²⁵ This is a well-validated questionnaire used to assess generic QoL; it provides a simple descriptive profile and a single index value for health status.

Participants completed all of the above listed questionnaires at baseline and at 6, 12 and 18 months post randomisation. In addition to these time points, participants completed the CCIS and the Health and Social Care Resource Use questionnaire at 2 weeks postoperatively (for temporary SNS and FENIX MSA only). For the permanent SNS, participants completed the Health and Social Care Resource Use questionnaire at 2 weeks postoperatively.

Summary of protocol changes

A summary of all substantial amendments to the SaFaRI protocol can be found in *Table 1*.

Early trial closure

On 23 March 2017, the SaFaRI trial team received formal notice from Torax Medical (the manufacturer of the FENIX MSA device) that the decision had been made to suspend the commercial sale of the FENIX MSA device in the UK and other European countries for 'strategic and business reasons'. As a result of this, recruitment into the SaFaRI trial, regrettably, had to cease with immediate effect.

Following the withdrawal of the FENIX MSA device and with approval from the Research Ethics Committee, all patients who had consented for the trial up to 23 March 2017, and had been randomised to the FENIX MSA arm but had not yet had surgery, were given the chance to have the FENIX MSA device implanted if they still wished to proceed with the operation as randomised. Any patients not wishing to undergo a FENIX MSA implantation, in the light of the fact that the device had been withdrawn, were offered alternative FI treatment as per local standard practice (this included the option of undergoing the alternative study intervention, SNS).

In total, 99 participants were randomised into the SaFaRI trial and 23 participants were registered as FENIX MSA training cases. A consequence of recruiting only 99 patients out of the target sample size of 350 is that the study is substantially underpowered to detect differences between the treatment arms, in particular with respect to the primary end point. Although the recruitment total was significantly less than the originally planned sample size of 350 participants, it was felt that continuing to follow up all randomised participants until the end of the planned follow-up period (i.e. 18 months post randomisation) would still provide valuable data that, at the very least, could provide some initial evidence; at the time of trial closure, SaFaRI was also the largest randomised trial of SNS to date.

The National Institute for Health Research was amenable to the trial team's proposal to continue with the planned follow-up period, and, as a result, all patients who had been randomised into the SaFaRI trial continued to be followed up until 18 months post randomisation. The last participant's final follow-up took place in September 2018.

TABLE 1 Summary of protocol changes

Version and date	Summary of changes
V1.0, 9 April 2014	N/A: original protocol submitted for ethics review
V2.0, 10 October 2014	<ul style="list-style-type: none"> • Update to contacts • Clarification of primary end point/outcome: 'in situ' amended to 'in use' • Text regarding FENIX MSA device supply arrangements revised • Update to data being collected (confirmation that device is both 'in situ' and 'in use') • Deaths to be reported up to 18 months post randomisation rather than 18 months after the last participant is randomised • CTRU responsibilities for safety reporting to the REC: serious complications removed • Study summary and GP letter for the Bladder and Bowel Foundation website • Addition of text to explain that a thank-you letter will be sent on receipt of a completed QoL questionnaire, or, if not received, a reminder letter will be sent • Inclusion of cover, reminder and thank-you letters for the postal QoL questionnaires that are being sent to participants from CTRU at 6, 12 and 18 months post randomisation
V3.0, 30 April 2015	<ul style="list-style-type: none"> • Inclusion of Scottish sites in the trial • A new consent form was drafted, for participants recruited in Scotland only, to include advance consent to continue to collect follow-up and safety data should capacity be lost during the course of the trial • Process for Scottish participants detailed in a new paragraph in the protocol • The provision to obtain assent from personal or nominated consultees remained; however, these sections were applicable only to participants recruited in England and relevant sections in the protocol were highlighted as such • Clarification to inclusion criterion 6: if the patient is being registered as a training case, they only need to be suitable for the FENIX MSA • Clarification to exclusion criterion 1: if the patient is being registered as a training case, they can have had previous interventions for FI
V4.0, 7 March 2016	<ul style="list-style-type: none"> • Removal of option for participants to complete the baseline and 2-week postoperative QoL packs (temporary SNS, permanent SNS and FENIX MSA) at home and enforce that these questionnaires are completed in clinic only. Wording on the baseline and 2-week postoperative QoL pack coversheets and in the protocol amended to reflect this change • Introduction of a questionnaire guidance sheet to assist with participant-reported CCIS questionnaire completion in clinic at baseline and 2 weeks postoperatively • Clinical CCIS to be collected at all follow-up time points to guard against attrition of the primary end point due to non-completion of the patient-reported outcome. CCIS collected in clinic to be incorporated into methods used to impute missing participant-reported CCIS • Wording in protocol amended to ensure that it is clear that the primary end point would be based on the participant-reported CCIS (and not the clinical score) • Date of completion, specifically for the CCIS, added to the participant-completed QoL questionnaires at all follow-up time points as this fed directly into the primary end point and thus would allow for a more robust primary analysis • Clarification that the 6-month follow-up assessment could be conducted via telephone if the participant was unable to attend the clinic in person • Removal of the word 'endoscopic' before visualisation of colorectum with regard to the preoperative investigations as it had been clinically confirmed that the visualisation of the colorectum did not need to be carried out endoscopically • A semantic alteration to clarify that the primary end point is the 'success' of each device, and not the 'difference in the percentage of successes [...] between the two treatments' • Recommendation that the ODS is calculated and is no more than 8 for inclusion in the trial added for guidance • Minor administrative changes throughout the protocol
V4.0, 7 March 2016	No change to protocol. Substantial amendment submitted to formally notify the REC of the early trial closure owing to the withdrawal from the market of the FENIX MSA device

CTRU, Clinical Trials Research Unit; GP, general practitioner; N/A, not applicable; REC, Research Ethics Committee.

End points

Primary end point

The primary end point was success, defined as device in use and $\geq 50\%$ improvement (between the baseline and 18-month scores) in the participant-reported CCIS, at 18 months post randomisation.

Secondary end points

Secondary end points included:

- the safety of FENIX MSA or SNS, as judged by explant rates, operative (this included those occurring during theatre time and post-surgery hospital stay) and postoperative (up to and including 12 months from the date of the last study surgery) complications
- change from baseline in generic and disease-specific QoL as measured by CCIS, ODS, FIQoL, EQ-5D-5L and SF-12 at 6, 12 and 18 months post randomisation
- cost-effectiveness
- success at 6 and 12 months as defined in the primary end point.

Sample size

A total of 350 participants were required to detect at least a 20% difference in the percentage of successes at 18 months post randomisation (where success was defined as device in use and $\geq 50\%$ CCIS improvement from baseline) between FENIX MSA and SNS at a 5% level of significance, with 90% power, assuming approximately 40% success in the SNS arm and allowing for 20% loss to follow-up. However, the number of patients recruited was 99.

Randomisation

Following confirmation of written informed consent and eligibility, patients were randomised into the trial by authorised members of staff at the trial sites. Randomisation was performed centrally using the Clinical Trials Research Unit (CTRU) automated 24-hour telephone randomisation system. Authorisation codes and personal identification numbers (PINs), provided by the CTRU, were required to access the randomisation system.

Participants were randomised on a 1 : 1 basis to receive either FENIX MSA or SNS, and were allocated a unique study number. A computer-generated minimisation programme that incorporated a random element was used with the following minimisation factors:

- treating surgeon
- participant sex (male or female)
- severity of incontinence (CCIS)
 - mild to moderate: CCIS ≤ 10 points
 - moderate to severe: CCIS > 10 points.
- degree of anal sphincter defect on endoanal ultrasound
 - no anal sphincter defect
 - anal sphincter defect $\leq 90^\circ$
 - $> 90^\circ$ anal sphincter defect $< 180^\circ$.

Blinding

The study was not blinded to participants, medical staff or clinical trial staff because of the difference between the two devices being compared (SNS treatment requires a temporary implant followed by a permanent implant if successful and involves patient input to function).

Statistical methods

Unless otherwise stated, all analyses were prespecified and conducted on the intention-to-treat population (i.e. all randomised participants were categorised into treatment groups based on their randomisation regardless of what treatment they subsequently received). All hypothesis tests were two-sided and conducted at the 5% level of significance. Estimates and their corresponding 95% confidence intervals (CIs) and *p*-values are presented for fixed effects. For all end points, missing outcome data were assumed to be missing at random, and the treatment effect was therefore estimated via maximum likelihood estimation using all participants with non-missing outcome data for non-longitudinal end points (this is referred to as a complete case analysis for the remainder of the report). All models were fitted using SAS® v9.4 (SAS Institute Inc., Cary, NC, USA). (SAS and all other SAS Institute Inc. product or service names are registered trademarks or trademarks of SAS Institute Inc. in the USA and other countries. ® indicates USA registration.)

Primary end point: device in use and $\geq 50\%$ improvement in CCIS at 18 months post randomisation

The primary analysis was a complete case analysis. Multilevel logistic regression was used to estimate the odds ratios between treatment groups for a 'success' in terms of the primary end point, adjusting for all minimisation factors. All minimisation factors were included as fixed effects, except randomising surgeon, which was included as a random effect. A random intercept model was fitted using maximum likelihood via adaptive quadrature, and all modelling was performed using the SAS v9.4 glimmix procedure.

A sensitivity analysis was performed to consider additional covariates in the primary analysis regression model that were thought to be related to patient outcome. These covariates were:

- age (years)
- body mass index (BMI)
- American Society of Anesthesiologists (ASA) grade
- aetiology of incontinence (obstetric trauma, idiopathic, iatrogenic, neurological conditions)
- type of incontinence (urge predominant, passive predominant, mixed urge and passive)
- ODS
- FI medication.

Secondary end point: device in use and $\geq 50\%$ improvement in CCIS at 6 months or 12 months post randomisation

Success at 6 or 12 months was analysed using a multilevel logistic regression model, adjusting for all of the minimisation factors, to estimate the odds ratios. All minimisation factors were included as fixed effects, except randomising surgeon, which was included as a random effect.

Secondary end point: intraoperative complications

Intraoperative complications were modelled using a multilevel logistic model to estimate the odds ratio between the treatment groups for whether or not participants had an intraoperative complication, adjusting for all minimisation factors. All minimisation factors were included as fixed effects, except randomising surgeon, which was included as a random effect with a random intercept.

Secondary end point: postoperative complications and reinterventions

Postoperative complications were modelled using a multilevel logistic model to estimate the odds ratio between the treatment groups for whether participants had a postoperative complication or not, adjusting for all minimisation factors. All minimisation factors were included as fixed effects, except randomising surgeon, which was included as a random effect with a random intercept and slope.

Secondary end point: device explants

The number of explants was analysed using a multilevel logistic regression model, adjusting for all of the minimisation factors, to estimate the odds ratios. All minimisation factors were included as fixed effects, except randomising surgeon, which was included as a random effect.

Quality-of-life end points

All QoL end points were modelled using a three-level multilevel model to account for the hierarchical nature of the repeated measures data and also for the clustering effect of the operating surgeon. All models were adjusted for the minimisation factors, with the minimisation factors included as fixed effects, except for the randomising surgeon, which was included as a random intercept.

Chapter 2 Results

Recruitment

Between 30 October 2014 and 23 March 2017, 322 patients were assessed for eligibility across 18 sites. Ninety-nine of these patients were randomised into the SaFaRI study and 23 participants were registered as FENIX MSA training cases. Recruitment by site can be seen in *Table 2*. A Consolidated Standards of Reporting Trials (CONSORT) flow diagram showing all patients screened for eligibility can be seen in *Figure 1*.

Baseline data

The minimisation factors are summarised by treatment arm across all randomised patients in *Table 3*. Summaries of additional baseline characteristics are given in *Table 4*. All the minimisation factors and baseline characteristics are well balanced between the two treatment arms.

TABLE 2 Recruitment by site

Site number	Site name	Randomised patients, <i>n</i>	Registered patients, <i>n</i>
00050	St James's University Hospital	45	1
00170	University Hospital of North Durham	4	1
00114	Southampton General Hospital	2	0
00232	The Northern General Hospital	14	0
00052	St Peter's Hospital	5	0
00108	Poole Hospital	7	2
00002	Royal Devon and Exeter Hospital	3	0
00153	The Churchill Hospital	0	3
00172	Wythenshawe Hospital	4	0
00099	Good Hope Hospital	2	3
10908	University College London Hospital	0	2
00080	Manchester Royal Infirmary	4	0
00117	Bristol Royal Infirmary	4	2
00072	Royal Victoria Infirmary	2	2
00023	Dewsbury District Hospital	1	2
00317	St Mark's Hospital	1	2
00031	Leicester Royal Infirmary	1	2
00118	Derriford Hospital	0	1




FIGURE 1 The CONSORT flow diagram. MND, motor neuron disease.

The time from randomisation to surgery can be seen in *Figure 2*. The time to implant of the permanent device was very different between the two treatment arms, with some SNS participants not receiving a permanent device until more than 18 months post randomisation. The reason for the delays in the SNS arm were mostly due to surgical capacity.

TABLE 3 Minimisation factors

Stratification factor	FENIX MSA, n (%)	SNS, n (%)	Total, n (%)
Surgeon ID			
2	10 (20.0)	12 (24.5)	22 (22.2)
1	11 (22.0)	8 (16.3)	19 (19.2)
3	6 (12.0)	8 (16.3)	14 (14.1)
31	3 (6.0)	4 (8.2)	7 (7.1)
26	2 (4.0)	3 (6.1)	5 (5.1)
4	3 (6.0)	1 (2.0)	4 (4.0)
12	3 (6.0)	1 (2.0)	4 (4.0)
21	2 (4.0)	2 (4.1)	4 (4.0)
30	2 (4.0)	2 (4.1)	4 (4.0)
36	1 (2.0)	3 (6.1)	4 (4.0)
32	2 (4.0)	1 (2.0)	3 (3.0)
8	1 (2.0)	1 (2.0)	2 (2.0)
15	1 (2.0)	1 (2.0)	2 (2.0)
25	1 (2.0)	1 (2.0)	2 (2.0)
5	0 (0.0)	1 (2.0)	1 (1.0)
23	1 (2.0)	0 (0.0)	1 (1.0)
999	1 (2.0)	0 (0.0)	1 (1.0)
Total	50 (100)	49 (100)	99 (100)
Sex			
Female	49 (98.0)	47 (95.9)	96 (97.0)
Male	1 (2.0)	2 (4.1)	3 (3.0)
Total	50 (100)	49 (100)	99 (100)
CCIS			
≤ 10 points (mild to moderate)	6 (12.0)	5 (10.2)	11 (11.1)
> 10 points (moderate to severe)	44 (88.0)	44 (89.8)	88 (88.9)
Total	50 (100)	49 (100)	99 (100)
Anal sphincter defect			
No anal sphincter defect	27 (54.0)	27 (55.1)	54 (54.5)
≤ 90°	19 (38.0)	18 (36.7)	37 (37.4)
> 90° to < 180°	4 (8.0)	4 (8.2)	8 (8.1)
Total	50 (100)	49 (100)	99 (100)

Primary end point: device in use and ≥ 50% improvement in Cleveland Clinic Incontinence Score at 18 months post randomisation

A total of 80 out of 99 (80.8%) participants were included in the primary analysis as 19 participants had missing primary outcome data. The pathway for participants in the primary analysis can be seen in *Figure 3*.

RESULTS

TABLE 4 Baseline characteristics

Characteristic	FENIX MSA	SNS	Total
Age (years)			
Mean (SD)	60.6 (13.1)	60.8 (14.3)	60.7 (13.7)
Median (range)	61.5 (30.0–82.0)	59.0 (35.0–90.0)	59.0 (30.0–90.0)
IQR	50.0–71.0	52.0–72.0	52.0–71.0
Missing	0	0	0
<i>n</i>	50	49	99
Ethnicity, n (%)			
White	47 (94.0)	49 (100.0)	96 (97.0)
Mixed	1 (2.0)	0 (0.0)	1 (1.0)
Asian (Indian)	1 (2.0)	0 (0.0)	1 (1.0)
Black	1 (2.0)	0 (0.0)	1 (1.0)
Total	50 (100)	49 (100)	99 (100)
Incontinence type, n (%)			
Mixed urge and passive	16 (32.0)	20 (40.8)	36 (36.4)
Urge predominant	13 (26.0)	12 (24.5)	25 (25.3)
Passive predominant	8 (16.0)	4 (8.2)	12 (12.1)
Missing	13 (26.0)	13 (26.5)	26 (26.3)
Total	50 (100)	49 (100)	99 (100)
FI aetiology, n (%)			
Obstetric trauma	29 (58.0)	26 (53.1)	55 (55.6)
Idiopathic	11 (22.0)	15 (30.6)	26 (26.3)
Iatrogenic	5 (10.0)	1 (2.0)	6 (6.1)
Neurological conditions	1 (2.0)	1 (2.0)	2 (2.0)
Other	4 (8.0)	6 (12.2)	10 (10.1)
Total	50 (100)	49 (100)	99 (100)
ASA grade, n (%)			
1	25 (50.0)	23 (46.9)	48 (48.5)
2	24 (48.0)	21 (42.9)	45 (45.5)
3	1 (2.0)	5 (10.2)	6 (6.1)
Total	50 (100)	49 (100)	99 (100)
Length of time suffered FI (months)			
Mean (SD)	71.6 (64.6)	89.9 (90.8)	80.6 (78.6)
Median (range)	60.0 (9.0–384)	60.0 (12.0–480)	60.0 (9.0–480)
IQR	36.0–84.0	27.0–114	36.0–96.0
Missing	0	1	1
<i>n</i>	50	48	98
Average number of episodes per week			
Mean (SD)	7.4 (6.95)	7.0 (7.01)	7.2 (6.94)
Median (range)	5.0 (2.0–28.0)	4.0 (1.5–30.0)	4.3 (1.5–30.0)
IQR	3.0–7.0	2.0–10.0	2.5–8.0
Missing	1	0	1
<i>n</i>	49	49	98

TABLE 4 Baseline characteristics (continued)

Characteristic	FENIX MSA	SNS	Total
Resting pressure (cmH₂O)			
Mean (SD)	52.4 (28.2)	58.4 (29.2)	55.4 (28.7)
Median (range)	48.0 (10.0–120)	53.0 (19.0–133)	51.5 (10.0–133)
IQR	33.0–73.0	35.0–77.5	34.0–75.0
Missing	0	1	1
<i>n</i>	50	48	98
Squeeze pressure (cmH₂O)			
Mean (SD)	77.0 (50.0)	84.8 (42.0)	80.8 (46.2)
Median (range)	69.5 (7.0–269)	85.5 (22.0–199)	71.0 (7.0–269)
IQR	44.0–105	49.0–116	46.0–109
Missing	0	1	1
<i>n</i>	50	48	98
Threshold volume (ml)			
Mean (SD)	57.6 (35.0)	53.1 (39.3)	55.5 (36.9)
Median (range)	52.0 (14.0–180)	46.0 (10.0–200)	47.0 (10.0–200)
IQR	32.5–79.0	30.0–60.0	30.0–74.0
Missing	10	15	25
<i>n</i>	40	34	74
Maximum tolerated volume (ml)			
Mean (SD)	127 (60.6)	137 (69.1)	131 (64.4)
Median (range)	112 (35.0–270)	117 (37.0–292)	114 (35.0–292)
IQR	85.0–160	77.0–195	80.0–180
Missing	10	15	25
<i>n</i>	40	34	74
Internal anal sphincter defect, n (%)			
Yes	8 (16.0)	10 (20.4)	18 (18.2)
No	41 (82.0)	39 (79.6)	80 (80.8)
Missing	1 (2.0)	0 (0.0)	1 (1.0)
Total	50 (100)	49 (100)	99 (100)
External anal sphincter defect, n (%)			
Yes	20 (40.0)	19 (38.8)	39 (39.4)
No	29 (58.0)	30 (61.2)	59 (59.6)
Missing	1 (2.0)	0 (0.0)	1 (1.0)
Total	50 (100)	49 (100)	99 (100)
ODS			
Mean (SD)	7.4 (3.78)	7.1 (3.62)	7.3 (3.68)
Median (range)	7.0 (2.0–18.0)	8.0 (2.0–17.0)	7.0 (2.0–18.0)
Missing	5	6	11
<i>n</i>	45	43	88

continued




FIGURE 3 Diagram of participants in primary analysis. a, Due to other health issues; b, due to complication during procedure.

The characteristics for participants included in the primary analysis are presented in *Table 5*.

The primary end point was not evaluable for 19 out of 99 randomised patients: 9 out of 50 patients randomised to FENIX MSA and 10 out of 49 patients randomised to SNS (see *Figure 3*). For the remaining 80 out of 99 patients with an evaluable primary end point, the rate of 'success' was 10 out of 80 (12.5%) patients overall: 6 out of 41 (14.6%) patients in the FENIX MSA arm and 4 out of 39 (10.3%) patients in the SNS arm (*Table 6*). The unadjusted odds ratio was 1.50 (95% CI 0.39 to 5.78; $p = 0.56$).

Summaries of the two individual components of the primary end point have been provided in *Appendix 1, Tables 30–33*.

RESULTS

TABLE 5 Characteristics for participants included in primary analysis

Variable	FENIX MSA	SNS	Total
CCIS at baseline (points)			
Mean (SD)	14.2 (3.23)	14.6 (3.05)	14.4 (3.13)
Median (range)	14.0 (6.0–20.0)	14.5 (9.0–20.0)	14.0 (6.0–20.0)
Missing, n	2	1	3
n	39	38	77
CCIS at 18 months (points)			
Mean (SD)	11.3 (4.42)	12.0 (4.57)	11.7 (4.48)
Median (range)	11.5 (3.0–17.0)	12.0 (1.0–19.0)	12.0 (1.0–19.0)
Missing, n	13	6	19
n	28	33	61
Sex, n (%)			
Male	1 (2.4)	2 (5.1)	3 (3.8)
Female	40 (97.6)	37 (94.9)	77 (96.3)
Total	41 (100)	39 (100)	80 (100)
CCIS stratification factor, n (%)			
≤ 10 points (mild to moderate)	6 (14.6)	5 (12.8)	11 (13.8)
> 10 points (moderate to severe)	35 (85.4)	34 (87.2)	69 (86.3)
Total	41 (100)	39 (100)	80 (100)
Anal sphincter defect, n (%)			
No anal sphincter defect	22 (53.7)	22 (56.4)	44 (55.0)
≤ 90°	16 (39.0)	13 (33.3)	29 (36.3)
> 90° to < 180°	3 (7.3)	4 (10.3)	7 (8.8)
Total	41 (100)	39 (100)	80 (100)
Randomising surgeon, n (%)			
1	10 (24.4)	6 (15.4)	16 (20.0)
2	10 (24.4)	10 (25.6)	20 (25.0)
3	5 (12.2)	6 (15.4)	11 (13.8)
4	3 (7.3)	1 (2.6)	4 (5.0)
5	0 (0.0)	1 (2.6)	1 (1.3)
8	0 (0.0)	1 (2.6)	1 (1.3)
12	2 (4.9)	1 (2.6)	3 (3.8)
15	1 (2.4)	1 (2.6)	2 (2.5)
21	2 (4.9)	2 (5.1)	4 (5.0)
23	1 (2.4)	0 (0.0)	1 (1.3)
25	0 (0.0)	1 (2.6)	1 (1.3)
26	1 (2.4)	2 (5.1)	3 (3.8)
30	0 (0.0)	1 (2.6)	1 (1.3)
31	3 (7.3)	2 (5.1)	5 (6.3)
32	2 (4.9)	1 (2.6)	3 (3.8)
Other	1 (2.4)	3 (7.7)	4 (5.0)
Total	41 (100)	39 (100)	80 (100)
SD, standard deviation.			

TABLE 6 Number of successes at 18 months post randomisation

Success	FENIX MSA	SNS	Total
18 months post randomisation, n (%)			
Unsuccessful	35 (85.4)	35 (89.7)	70 (87.5)
Successful	6 (14.6)	4 (10.3)	10 (12.5)
Total	41 (100)	39 (100)	80 (100)

The odds ratio adjusting for the minimisation factors was 1.45 (95% CI 0.36 to 5.83; $p = 0.59$).

The adjusted estimates of odds ratios and 95% CIs are presented in *Table 7*. The model shows no statistically significant differences between any of the minimisation factors, although this would be expected owing to the small number of patients recruited, which has led to large standard errors (SEs) of the estimates (i.e. wide CIs and underpowered hypothesis tests).

The estimated random effects with respect to surgeons were equal to 0. The SE was not estimable owing to the low numbers of patients operated on by each surgeon (see *Table 5*).

In *Appendix 1, Figure 30* shows the empirical probability plot for the primary analysis model, which can be used to compare actual Pearson residuals with expected Pearson residuals. The y-axis is the actual Pearson residual value, the x-axis is the empirical median Pearson residual expected under our fitted model assumptions. Each dot represents the actual Pearson residual for an individual patient. If the model fitted perfectly, we would expect all of the dots to lie on the reference line. The band in *Appendix 1, Figure 30*, represents the interval between the empirical 2.5th percentile and 97.5th percentile empirical Pearson residual. There are some areas where the reference line is outside the band, meaning that the model may not fit the data too well, but this is due to the small sample size and the rarity of successful outcomes.

In *Appendix 1, Figure 31* presents the plot of exponentiated delta-betas (y-axis) versus patient identifier. Exponentiated delta-betas further from 1 indicate greater influence of the observation on the estimated treatment effect. Patients with a success for the primary end point are more influential than patients without success for the primary end point, which may be expected given that success was an uncommon occurrence. There does not appear to be any other observations with a large influence on the model.

Sensitivity analysis: additional covariates

Owing to the small number of 'successes', performing the sensitivity analysis with additional covariates that was described in the analysis plan was inappropriate.

TABLE 7 Fixed effects of primary end point model

Effect	Odds ratio	Odds ratio 95% CI	p-value
FENIX MSA (vs. SNS)	1.453	0.362 to 5.827	0.5926
Male (vs. female)	< 0.001	< 0.001 to infinity	0.9957
Baseline CCIS > 10 points (vs. baseline CCIS ≤ 10 points)	0.631	0.110 to 3.614	0.5993
Anal sphincter defect ≤ 90° (vs. no defect)	1.108	0.260 to 4.723	0.8883
Anal sphincter defect > 90° to < 180° (vs. no defect)	1.238	0.115 to 13.298	0.8579

Longitudinal analysis

As the primary end point was measured at 6, 12 and 18 months post randomisation, a longitudinal analysis using the data at each time point was performed. The model did not adjust for the stratification factors because of the added complexity and the number of missing data. In addition, the stratification factors caused the model to fail to converge.

The fixed effects of the model can be seen in *Table 8*. The estimated random effect caused by the operating surgeon was 1.52 (SE 0.99) and the estimated random effect caused by within patient measurements was 1.19. The model does not show a significant difference between the treatment arms ($p = 0.20$) or in change over time ($p = 0.17$), although, again, it is worth noting that the estimates have large SEs due to the small sample size.

The model results can be seen in *Figure 4*.

Secondary end point: device in use and $\geq 50\%$ improvement in Cleveland Clinic Incontinence Score at 12 months post randomisation

Success at 12 months post randomisation was evaluable for 67 out of 99 (67.7%) participants. Thirteen patients (1 FENIX and 12 SNS) were not included in this analysis as they had not had a permanent device fitted within 12 months of randomisation because of surgical capacity in the trial sites. The other 19 patients were not included because of missing CCIS at baseline, at 12 months post randomisation, or at both time points.

TABLE 8 Fixed effects of longitudinal model

Effect	Estimate	Estimate 95% CI	p-value
FENIX MSA (vs. SNS)	1.429	0.281 to 7.261	0.20
Time (months)	1.039	0.914 to 1.180	0.17
Time and treatment interaction	1.278	0.902 to 1.812	0.27




FIGURE 4 Longitudinal model results.

A total of 4 out of 27 (14.8%) patients did not have a temporary SNS device fitted (and therefore did not have a permanent device implanted) because they withdrew from surgery before the temporary SNS operation. A total of 4 out of 40 (10%) patients in the FENIX arm did not have a FENIX device implanted as a result of withdrawing from surgery. Seventeen out of 27 (63%) patients did not have a permanent SNS device implanted because of either withdrawal ($n = 5$) or the lack of success of the temporary device ($n = 12$). Ten out of 40 (25%) patients who were randomised to FENIX had the device explanted; there were no explants in the SNS arm. The number of successes is summarised in *Table 9*. Owing to the small number of successes at 12 months, the models fitted were not meaningful and so are not presented.

Secondary end point: device in use and $\geq 50\%$ improvement in Cleveland Clinic Incontinence Score at 6 months post randomisation

Success at 6 months post randomisation was evaluable for 56 out of 99 (56.6%) patients. Twenty-three patients (3 FENIX and 20 SNS) were not included in this analysis because they had not had a permanent device fitted within 6 months of randomisation owing to surgical capacity at trial sites. The other 20 patients were not included because of missing CCIS at baseline, 6 months post randomisation, or at both time points.

A total of 3 out of 18 (16.7%) patients did not have a temporary SNS device implanted (and therefore no permanent SNS device was implanted) because of withdrawal from surgery. Two out of 38 (5.3%) of patients randomised to FENIX did not have a permanent device implanted because of withdrawal from surgery. Thirteen out of 18 (72.2%) patients did not have a permanent device fitted because of withdrawal ($n = 4$) or the lack of efficacy of the temporary SNS device ($n = 9$). Nine patients in the FENIX arm (76.3%) had the device explanted within 6 months of randomisation and there were no explants in the SNS arm. The number of successes in each arm is summarised in *Table 10*. Owing to the small number of successes at 6 months, the models fitted did not converge and so are not presented.

TABLE 9 Number of successes at 12 months post randomisation

Success	FENIX MSA	SNS	Total
<i>12 months post randomisation, n (%)</i>			
Unsuccessful	35 (87.5)	26 (96.3)	61 (91.0)
Successful	5 (12.5)	1 (3.7)	6 (9.0)
Total	40 (100)	27 (100)	67 (100)

TABLE 10 Number of successes at 6 months post randomisation

Success	FENIX MSA	SNS	Total
<i>6 months post randomisation, n (%)</i>			
Unsuccessful	33 (86.8)	18 (100.0)	51 (91.1)
Successful	5 (13.2)	0 (0.0)	5 (8.9)
Total	38 (100)	18 (100)	56 (100)

Secondary end point: intraoperative complications

In total, 89 out of 99 patients (89.9%) had intraoperative complication data; 45 patients received an operation in the FENIX arm and 44 patients received an operation in the SNS arm.

There were four intraoperative complications in four patients; 3 out of 45 patients (6.7%) who were randomised to FENIX and received the operation had an intraoperative complication, and 1 out of 44 patients (2.3%) who were randomised to SNS and received at least one operation had an intraoperative complication, giving an overall complication rate of 4.5% (4/89) in randomised patients.

The complication in the SNS arm was an unexpected serious complication (USC). The details are as follows:

- USC – anaphylaxis
- Clavien–Dindo grade – IVb
- USC description – patient became tachycardic, hypotensive and flushed following administration of Teicoplanin (Targocid, Sanofi, Paris, France) prior to their anaesthetic for insertion of SNS
- Outcome – recovered.

There were three intraoperative complications in the FENIX arm, none of which was serious. The complications were bleeding, cyst found in recto vaginal septum, and rectal perforation.

Secondary end point: postoperative complications

A total of 85 out of 99 patients (85.9%) were included in the analysis of postoperative complications; 10 patients were not included due to not receiving an operation and four patients were not included due to missing follow-up forms. There were 42 out of 85 (49.4%) patients who experienced at least one postoperative complication: 33 out of 45 patients (73.3%) in the FENIX MSA arm and 9 out of 40 patients (22.5%) in the SNS arm (*Table 11*). The unadjusted odds ratio of having a complication in the FENIX MSA arm was 7.77 (95% CI 3.0 to 20.0; $p < 0.001$).

Figure 2 shows the time from randomisation to operation and *Table 12* shows the time from randomisation to first complication. Complications data were collected at 6, 12 and 18 months' post-randomisation

TABLE 11 Number of patients experiencing at least one postoperative complication

Did patient experience a postoperative complication?, n (%)	FENIX MSA	SNS	Total
Yes	33 (73.3)	9 (22.5)	42 (49.4)
No	12 (26.7)	31 (77.5)	43 (50.6)
Total	45 (100)	40 (100)	99 (100)

TABLE 12 Time from operation to first complication

Parameter	FENIX MSA	SNS	Total
Mean (SD) (days)	80.0 (148.72)	79.3 (123.87)	79.8 (142.33)
Median (range) (days)	12.0 (0.0–540.0)	15.0 (0.0–355.0)	13.0 (0.0–540.0)
Participants, n	33	9	42

SD, standard deviation.

follow-up visits; therefore, patients who had their SNS devices fitted more than 18 months post randomisation will not have any complications recorded. There were five complications from the temporary SNS operation: (1) neurological, (2) haemorrhoid discomfort, (3) intermittent flare of eczema at SNS dressing site, (4) lack or loss of efficiency, and (5) device failure/separation. There were eight complications from the permanent SNS operation: (1) lead migration/fragmentation, (2) pain at battery site, (3) lack or loss of efficiency, (4) transient anal/rectal pain, (5) rectal/anal pain on defaecation, (6) neurological, (7) cardiorespiratory and (8) device reoperation: replacement of SNS wire. There were 88 complications across 33 patients in the FENIX MSA arm (Table 13).

The adjusted model odds ratios can be seen in Table 14. The model results show a statistically significant difference in the odds of a complication between the two treatment arms, with an adjusted odds ratio of 12.91 (95% CI 2.75 to 60.68; $p = 0.004$). The random effect with respect to surgeon (i.e. the 'random intercept') was 0.79 (SE 1.38), and the random effect with respect to the interaction between surgeon and the difference between treatments (i.e. the 'random slope') was < 0.0001 (SE 2.17).

The time from randomisation to operation is presented in Figure 2. There is a large difference between the arms, with four patients in the SNS arm not having a permanent operation within 18 months of randomisation; therefore, the model results may be biased against the FENIX MSA arm as patients in

TABLE 13 Complications in FENIX arm

Complication	Total, n (%)
Worsening constipation/obstructed defecation	18 (20.5)
Other ^a	17 (19.3)
Transient anal/rectal pain	14 (15.9)
Device erosion	8 (9.1)
Wound infection	7 (8.0)
Device explant/reoperation	7 (8.0)
Implant infection	5 (5.7)
Bleeding/wound haematoma	4 (4.5)
Neurological	3 (3.4)
Lack or loss of efficiency	3 (3.4)
Urinary retention	1 (1.1)
Device failure/separation	1 (1.1)
Total	88 (100)

a Other complications:

- increase discharge of wound and breakdown
- small sinus around perianal area
- right upper quadrant pain
- wound dehiscence
- small swelling at edge of wound
- poor evacuation/constipation ($n = 3$)
- pain lasting 4 weeks
- pinching of skin around device
- superficial dehiscence of wound not exposing muscle or the implant
- pain at base of right back
- lack of control of bowels
- wound pain
- covering stoma
- wound still not healed following explant, fistula
- bleeding from prolapse.

TABLE 14 Postoperative complications regression model

Effect	Odds ratio	Odds ratio 95% CI	p-value
FENIX MSA (vs. SNS)	12.91	2.75 to 60.68	0.0042
Male (vs. female)	1.04	0.03 to 36.87	0.98
Baseline CCIS > 10 points (vs. baseline CCIS ≤ 10 points)	0.79	0.11 to 5.59	0.81
Anal sphincter defect ≤ 90° (vs. no defect)	1.43	0.44 to 4.68	0.55
Anal sphincter defect > 90° to ≤ 180° (vs. no defect)	0.41	0.03 to 6.78	0.53

this arm will have had longer postoperative follow-up and, therefore, longer exposure to the risk of postoperative complication within the trial follow-up period. However, *Table 12* shows the number of days from operation to first complication and this shows that most complications occurred within 3 months, with some happening a long time after the operation. As seen in *Figure 2*, most SNS patients were followed up for > 3 months post randomisation.

Appendix 1, Figure 32, shows the empirical probability plot for the postoperative complications model, which can be used to compare actual Pearson residuals with expected Pearson residuals. The y-axis is the actual Pearson residual value, the x-axis is the empirical median Pearson residual expected under our fitted model assumptions. Each dot represents the actual Pearson residual for an individual patient. If the model fitted perfectly we would expect all of the dots to lie on the reference line. The band in *Figure 32* represents the interval between the empirical 2.5th percentile and 97.5th percentile empirical Pearson residual. No values lie outside this region, indicating that we do not have any substantial outliers.

Appendix 1, Figure 33, presents the plot of exponentiated delta-betas (y-axis) versus patient identifier. Exponentiated delta-betas further from 1 indicate greater influence of the observation on the estimated treatment effect. There does not appear to be any unusually influential observations in the delta-beta plot.

Secondary end point: device explants

The number of devices in situ and explanted within 18 months of randomisation can be seen in *Table 15*. The mean number of days from device implant to explant was 164 (SD 168) and the median number of days was 112 (range 0–449).

Figure 5 shows the periods of time (measured from randomisation) for which each patient had a permanent device in situ. The mean number of days from implant to explant was 164 (SD = 168) and the median number of days was 112 (range 0–449). As there were no explants in the SNS arm, the device explants end point has not been modelled because any regression models would not converge.

TABLE 15 Device status at 18 months post randomisation

Device status	FENIX MSA, n (%)	SNS, n (%)	Total, n (%)
Device in situ	30 (60.0)	24 (49.0)	54 (54.5)
Device explanted	15 (30.0)	0 (0.0)	15 (15.2)
No device fitted	5 (10.0)	21 (42.9)	26 (26.3)
Device implanted more than 18 months post randomisation	0 (0.0)	4 (8.2)	4 (4.0)
Total	50 (100)	49 (100)	99 (100)




FIGURE 5 Time to explant: (a) FENIX MSA; and (b) SNS. Dashed line indicates that patient had device explanted.

The reasons for explant are given in *Table 16* and include participant intolerance ($n = 4$), device erosion/migration and infection, implant infected and visible through the skin, device eroded, infection/erosion ($n = 3$), rectal perforation the device was removed/no implant, chronic sinus infection, participant did not have adequate response, and device malfunction.

Secondary end point: Cleveland Clinic Incontinence Score

The CCIS assesses five parameters associated with incontinence: incontinence to solid, incontinence to liquid, incontinence to gas, use of pads and lifestyle restrictions. Each parameter is scored 0–4 with '0' for never and '4' for every day. The five parameters are added to give a total score out of 20, with a lower score indicating a better QoL.

Summary measures of the CCIS split by time point and treatment arm are given in *Table 17*.

The model diagnostics can be seen in *Appendix 1, Figure 34*.

RESULTS

TABLE 16 Reasons for explant

Reason	Days from randomisation to explant
Participant intolerance	257
Device erosion/migration + infection	16
Implant infected and is visible through the skin	154
Participant did not have adequate response	399
Device eroded	350
Infection/erosion of FENIX beads SAE	14
Rectal perforation the device was removed/no implant	0
Infection and erosion, device explanted	7
Infection erosion	69
Chronic sinus infection	169
Participant did not have adequate response	409
Device malfunction	449

SAE, serious adverse event.

TABLE 17 Crude summaries of CCIS at baseline and at 6, 12 and 18 months post randomisation

CCIS (points)	FENIX MSA	SNS	Total
Baseline			
Mean (SD)	14.4 (3.15)	14.7 (2.95)	14.6 (3.04)
Median (range)	14.5 (6.0–20.0)	15.0 (9.0–20.0)	15.0 (6.0–20.0)
Missing	4	3	7
<i>n</i>	46	46	92
6 months post randomisation			
Mean (SD)	11.7 (4.61)	13.8 (3.45)	12.8 (4.16)
Median (range)	12.0 (2.0–20.0)	14.0 (7.0–20.0)	13.0 (2.0–20.0)
Missing	15	10	25
<i>n</i>	35	39	74
12 months post randomisation			
Mean (SD)	11.8 (4.86)	12.9 (3.78)	12.3 (4.37)
Median (range)	12.0 (3.0–20.0)	13.0 (5.0–20.0)	12.0 (3.0–20.0)
Missing	15	15	30
<i>n</i>	35	34	69
18 months post randomisation			
Mean (SD)	11.1 (4.35)	12.1 (4.55)	11.6 (4.45)
Median (range)	11.0 (3.0–17.0)	12.5 (1.0–19.0)	12.0 (1.0–19.0)
Missing	20	15	35
<i>n</i>	30	34	64

SD, standard deviation.

Secondary end point: EuroQol-5 Dimensions, five-level version

The EQ-5D-5L is made up of 5 dimensions (mobility, self-care, usual activities, pain/discomfort and anxiety/depression), each of which has 5 levels (no problems, slight problems, moderate problems, severe problems and extreme problems). These levels can be combined to calculate an index value, which ranges from 1 (best possible health), through 0 (death) to -0.594 (worse than death).

Summary measures of the EQ-5D-5L index values, and the EQ-5D-5L Visual Analogue Scale (VAS) scores, split by time point and treatment arm, are given in *Table 18*.

TABLE 18 Crude summaries of EQ-5D-5L score and EQ-5D-5L VAS at baseline and at 6, 12 and 18 months post randomisation

Variable	FENIX MSA	SNS	Total
EQ-5D-5L score			
<i>Baseline</i>			
Mean (SD)	0.666 (0.256)	0.647 (0.261)	0.657 (0.257)
Median (range)	0.740 (-0.02-1.00)	0.736 (-0.05-1.00)	0.736 (-0.05-1.00)
Missing	4	3	7
<i>n</i>	46	46	92
<i>6 months post randomisation</i>			
Mean (SD)	0.687 (0.225)	0.639 (0.295)	0.662 (0.263)
Median (range)	0.736 (-0.02-1.00)	0.732 (-0.22-1.00)	0.736 (-0.22-1.00)
Missing	12	7	19
<i>n</i>	38	42	80
<i>12 months post randomisation</i>			
Mean (SD)	0.684 (0.258)	0.673 (0.263)	0.679 (0.259)
Median (range)	0.750 (-0.06-1.00)	0.752 (0.023-1.00)	0.750 (-0.06-1.00)
Missing	13	13	26
<i>n</i>	37	36	73
<i>18 months post randomisation</i>			
Mean (SD)	0.660 (0.258)	0.683 (0.255)	0.672 (0.255)
Median (range)	0.732 (0.054-1.00)	0.743 (-0.05-1.00)	0.735 (-0.05-1.00)
Missing	20	17	37
<i>n</i>	30	32	62
EQ-5D-5L VAS score			
<i>Baseline</i>			
Mean (SD)	65.7 (20.3)	66.8 (24.8)	66.2 (22.5)
Median (range)	70.0 (0.00-95.0)	70.0 (5.00-100)	70.0 (0.00-100)
Missing	2	5	7
<i>n</i>	48	45	93

continued

RESULTS

TABLE 18 Crude summaries of EQ-5D-5L score and EQ-5D-5L VAS at baseline and at 6, 12 and 18 months post randomisation (continued)

Variable	FENIX MSA	SNS	Total
<i>6 months post randomisation</i>			
Mean (SD)	66.9 (19.6)	72.7 (19.3)	69.9 (19.5)
Median (range)	70.0 (30.0–95.0)	75.0 (15.0–95.0)	75.0 (15.0–95.0)
Missing	11	7	18
<i>n</i>	39	42	81
<i>12 months post randomisation</i>			
Mean (SD)	65.5 (21.1)	73.2 (19.7)	69.4 (20.6)
Median (range)	70.0 (10.0–96.0)	75.0 (20.0–100)	72.5 (10.0–100)
Missing	14	13	27
<i>n</i>	36	36	72
<i>18 months post randomisation</i>			
Mean (SD)	66.0 (19.6)	72.1 (14.0)	69.2 (17.0)
Median (range)	67.5 (20.0–95.0)	70.0 (33.0–100)	70.0 (20.0–100)
Missing	20	16	36
<i>n</i>	30	33	63
SD, standard deviation.			

Secondary end point: faecal incontinence quality of life

The FIQoL questionnaire is composed of 29 items that make up four scales: lifestyle (10 items), coping/behaviour (9 items), depression/self-perception (7 items) and embarrassment (3 items). Scales range from 1 to 5 and scale scores are derived by averaging the response to all items in the scale. A lower score indicates a lower functional QoL.

Summary measures of the four FIQoL domains split by time point and treatment arm are provided in *Table 19*.

Secondary end point: Obstructed Defecation Score

The ODS consists of five items: excessive straining, incomplete rectal evacuation, use of enemas and/or laxatives, vaginal-anal-perineal digitations, and abdominal discomfort and/or pain. Each item is graded from 0 to 4, with scores ranging from 0 (no symptoms) to 20 (very severe symptoms), meaning that a lower score indicates better QoL.

Summary measures of the total ODS split by time point and treatment arm are presented in *Table 20*.

Secondary end point: Short Form questionnaire-12 items

The SF-12 version 2 is a generic health survey that measures eight health domains: physical functioning, role – physical, bodily pain, general health, vitality, social functioning, role – emotional and mental health. The Physical Component Summary (PCS) and Mental Component Summary (MCS)

TABLE 19 Crude summaries of FIQoL domains at baseline and at 6, 12 and 18 months post randomisation

FIQoL domain	FENIX MSA	SNS	Total
Lifestyle (baseline)			
Mean (SD)	2.17 (0.864)	2.12 (0.887)	2.15 (0.871)
Median (range)	2.00 (1.00–3.80)	2.00 (1.00–3.90)	2.00 (1.00–3.90)
Missing	5	5	10
<i>n</i>	45	44	89
Lifestyle (6 months post randomisation)			
Mean (SD)	2.64 (1.09)	2.39 (0.918)	2.51 (1.01)
Median (range)	2.70 (1.00–4.00)	2.30 (1.00–4.00)	2.50 (1.00–4.00)
Missing	13	11	24
<i>n</i>	37	38	75
Lifestyle (12 months post randomisation)			
Mean (SD)	2.74 (1.08)	2.35 (1.03)	2.56 (1.06)
Median (range)	3.05 (1.00–4.00)	2.30 (1.00–3.90)	2.75 (1.00–4.00)
Missing	14	17	31
<i>n</i>	36	32	68
Lifestyle (18 months post randomisation)			
Mean (SD)	2.73 (1.03)	2.62 (1.02)	2.68 (1.02)
Median (range)	2.90 (1.00–4.00)	2.90 (1.00–4.00)	2.90 (1.00–4.00)
Missing	21	22	43
<i>n</i>	29	27	56
Coping (baseline)			
Mean (SD)	1.44 (0.534)	1.50 (0.616)	1.47 (0.573)
Median (range)	1.33 (1.00–3.67)	1.33 (1.00–3.44)	1.33 (1.00–3.67)
Missing	9	9	18
<i>n</i>	41	40	81
Coping (6 months post randomisation)			
Mean (SD)	2.04 (0.984)	1.63 (0.610)	1.81 (0.818)
Median (range)	1.67 (1.00–4.00)	1.56 (1.00–3.22)	1.56 (1.00–4.00)
Missing	23	16	39
<i>n</i>	27	33	60
Coping (12 months post randomisation)			
Mean (SD)	2.24 (1.02)	1.57 (0.671)	1.91 (0.923)
Median (range)	2.11 (1.00–4.00)	1.44 (1.00–3.22)	1.56 (1.00–4.00)
Missing	23	22	45
<i>n</i>	27	27	54

continued

RESULTS

TABLE 19 Crude summaries of FIQoL domains at baseline and at 6, 12 and 18 months post randomisation (*continued*)

FIQoL domain	FENIX MSA	SNS	Total
Coping (18 months post randomisation)			
Mean (SD)	1.94 (0.895)	1.97 (0.820)	1.96 (0.847)
Median (range)	1.67 (1.00–3.89)	2.11 (1.00–3.44)	1.89 (1.00–3.89)
Missing	30	28	58
<i>n</i>	20	21	41
Depression (baseline)			
Mean (SD)	2.52 (0.956)	2.33 (0.864)	2.43 (0.911)
Median (range)	2.14 (1.14–4.29)	2.29 (1.00–4.29)	2.14 (1.00–4.29)
Missing	11	11	22
<i>n</i>	39	38	77
Depression (6 months post randomisation)			
Mean (SD)	2.74 (1.07)	2.48 (0.891)	2.60 (0.978)
Median (range)	2.36 (1.43–4.43)	2.43 (1.14–4.14)	2.43 (1.14–4.43)
Missing	24	19	43
<i>n</i>	26	30	56
Depression (12 months post randomisation)			
Mean (SD)	2.70 (1.11)	2.57 (0.898)	2.64 (1.01)
Median (range)	2.29 (1.14–4.29)	2.64 (1.29–4.00)	2.43 (1.14–4.29)
Missing	20	21	41
<i>n</i>	30	28	58
Depression (18 months post randomisation)			
Mean (SD)	2.53 (1.01)	2.89 (0.948)	2.72 (0.981)
Median (range)	2.43 (1.29–4.43)	3.00 (1.43–4.43)	2.57 (1.29–4.43)
Missing	29	25	54
<i>n</i>	21	24	45
Embarrassment (baseline)			
Mean (SD)	1.74 (0.699)	1.64 (0.640)	1.69 (0.668)
Median (range)	1.67 (1.00–3.33)	1.33 (1.00–3.33)	1.67 (1.00–3.33)
Missing	4	2	6
<i>n</i>	46	47	93
Embarrassment (6 months post randomisation)			
Mean (SD)	2.12 (0.996)	1.87 (0.789)	1.99 (0.897)
Median (range)	1.67 (1.00–4.00)	1.67 (1.00–4.00)	1.67 (1.00–4.00)
Missing	11	7	18
<i>n</i>	39	42	81

TABLE 19 Crude summaries of FIQoL domains at baseline and at 6, 12 and 18 months post randomisation (continued)

FIQoL domain	FENIX MSA	SNS	Total
Embarrassment (12 months post randomisation)			
Mean (SD)	2.20 (0.970)	1.87 (0.791)	2.04 (0.894)
Median (range)	2.33 (1.00–4.00)	1.67 (1.00–3.67)	1.83 (1.00–4.00)
Missing	13	12	25
<i>n</i>	37	37	74
Embarrassment (18 months post randomisation)			
Mean (SD)	2.24 (1.02)	1.99 (0.827)	2.11 (0.926)
Median (range)	2.33 (1.00–4.00)	1.83 (1.00–4.00)	2.00 (1.00–4.00)
Missing	21	17	38
<i>n</i>	29	32	61
SD, standard deviation.			

TABLE 20 Crude summaries of ODS at baseline and at 6, 12 and 18 months post randomisation

ODS	FENIX MSA	SNS	Total
Baseline			
Mean (SD)	7.44 (3.78)	7.09 (3.62)	7.27 (3.68)
Median (range)	7.00 (2.00–18.0)	8.00 (2.00–17.0)	7.00 (2.00–18.0)
Missing	5	6	11
<i>n</i>	45	43	88
6 months post randomisation			
Mean (SD)	9.10 (4.83)	7.53 (3.48)	8.40 (4.33)
Median (range)	8.00 (0.000–20.0)	7.00 (0.000–16.0)	8.00 (0.000–20.0)
Missing	10	17	27
<i>n</i>	40	32	72
12 months post randomisation			
Mean (SD)	8.34 (4.86)	7.52 (3.75)	7.94 (4.34)
Median (range)	8.00 (1.00–20.0)	8.00 (2.00–15.0)	8.00 (1.00–20.0)
Missing	15	16	31
<i>n</i>	35	33	68
18 months post randomisation			
Mean (SD)	8.58 (4.77)	6.90 (4.34)	7.67 (4.58)
Median (range)	8.00 (0.000–18.0)	6.00 (0.000–16.0)	7.00 (0.000–18.0)
Missing	24	18	42
<i>n</i>	26	31	57
SD, standard deviation.			

RESULTS

measures are calculated using a combination of the eight domains. Scores are calibrated so that the mean score is 50 with a standard deviation (SD) of 10 in the 2009 general US population. All scores range from 0 to 100, with a higher value indicating better functioning and well-being.

Summary measures of the SF-12 PCS and MCS components split by time point and treatment arm are presented in *Table 21*.

TABLE 21 Crude summaries of SF-12 PCS and MCS components at baseline and at 6, 12 and 18 months post randomisation

SF-12 component	FENIX MSA	SNS	Total
PCS (baseline)			
Mean (SD)	44.9 (9.36)	46.9 (10.2)	45.9 (9.78)
Median (range)	46.5 (18.6–61.2)	47.3 (23.6–64.0)	47.3 (18.6–64.0)
Missing	3	1	4
<i>n</i>	47	48	95
PCS (6 months post randomisation)			
Mean (SD)	46.2 (9.96)	47.7 (9.71)	47.0 (9.80)
Median (range)	45.6 (27.6–68.5)	48.7 (23.9–70.0)	47.1 (23.9–70.0)
Missing	11	7	18
<i>n</i>	39	42	81
PCS (12 months post randomisation)			
Mean (SD)	47.1 (9.64)	47.4 (9.49)	47.2 (9.50)
Median (range)	49.0 (26.7–65.3)	49.8 (20.3–64.1)	49.2 (20.3–65.3)
Missing	12	12	24
<i>n</i>	38	37	75
PCS (18 months post randomisation)			
Mean (SD)	43.9 (10.9)	47.0 (10.4)	45.5 (10.7)
Median (range)	43.9 (26.9–59.7)	46.8 (23.7–63.3)	46.6 (23.7–63.3)
Missing	19	16	35
<i>n</i>	31	33	64
MCS (baseline)			
Mean (SD)	41.2 (11.4)	41.2 (11.2)	41.2 (11.2)
Median (range)	42.7 (22.8–61.8)	40.9 (17.0–63.5)	41.3 (17.0–63.5)
Missing	3	1	4
<i>n</i>	47	48	95
MCS (6 months post randomisation)			
Mean (SD)	41.6 (13.2)	42.5 (12.5)	42.0 (12.8)
Median (range)	40.8 (15.8–62.0)	45.2 (15.4–61.2)	43.1 (15.4–62.0)
Missing	10	7	17
<i>n</i>	40	42	82

TABLE 21 Crude summaries of SF-12 PCS and MCS components at baseline and at 6, 12 and 18 months post randomisation (continued)

SF-12 component	FENIX MSA	SNS	Total
MCS (12 months post randomisation)			
Mean (SD)	41.6 (13.3)	43.8 (11.7)	42.7 (12.5)
Median (range)	41.9 (20.5–60.5)	43.2 (17.4–62.4)	43.2 (17.4–62.4)
Missing	12	12	24
<i>n</i>	38	37	75
MCS (18 months post randomisation)			
Mean (SD)	43.5 (12.5)	44.3 (12.2)	43.9 (12.2)
Median (range)	41.2 (15.2–63.1)	46.7 (19.2–69.6)	44.6 (15.2–69.6)
Missing	19	16	35
<i>n</i>	31	33	64

Registered training cases

Prior to randomising patients in the study, all surgeons were required to have experience of a minimum of one observed FENIX MSA procedure and two FENIX MSA procedures under proctorship. Surgeons who did not have this experience before study participation joined the registration phase of the study in which the first two eligible patients providing consent were registered to receive FENIX MSA implants. These two operations, performed under proctorship, were considered training cases and were not included in the main study.

A total of 23 patients were registered as training cases for the FENIX MSA operation. Of these 23 patients, 22 received the FENIX operation and one withdrew, giving the reason as travelling scheduled for after the surgery date; the patient withdrew after discussing with a research nurse advice about travelling post surgery, and decided not to risk discomfort and complications.

There was one operative complication, with the complication described as 'Perf skin posterior. Primary repair'.

Only complications that occurred within 30 days of the operation were reported for the registered training cases. There were 22 postoperative complications across 13 patients:

- wound infection ($n = 7$)
- transient anal/rectal pain ($n = 3$)
- wound dehiscence ($n = 2$)
- worsening constipation/obstructed defecation ($n = 2$)
- urinary retention
- bleeding/urinary retention
- device explant/reoperation
- urinary tract infection
- incomplete emptying of bladder on micturition
- prickly sensation
- incomplete emptying
- superficial wound dehiscence.

Chapter 3 Health economics

This section outlines the approach to the manipulation and analysis of the health economic data from the SaFaRI trial for the purpose of conducting an economic evaluation of the FENIX device.

The economic evaluation followed the NICE reference case²⁶ and, hence, was a cost–utility analysis presenting cost per incremental QALY from the perspective of the health-care and personal social service’s provider. The evaluation used both trial-based and model-based analyses, and adopted a lifetime horizon. A model was required given that the likely costs and benefits of the interventions continue to accrue after the trial time horizon (18 months). Our primary analysis is based on the combined trial and model results, but we also report cost-effectiveness at the trial end.

The evaluation assumes a willingness-to-pay threshold (λ) of £20,000 per QALY gain, and estimates the incremental costs and benefits of the FENIX magnetic anal sphincter (MAS) device versus usual SNS for adult FI.

Methods: trial

Health utility and quality-adjusted life-year calculation

Health-related quality-of-life (HRQoL) data to enable QALY calculations were collected during the trial at baseline and at 6, 12 and 18 months post randomisation. Data were available on the EQ-5D-5L^{27,28} and the SF-12 [Short Form questionnaire-6 Dimensions (SF-6D)].²⁹

The primary analysis is based on the EQ-5D-5L direct valuation.³⁰ The secondary analysis estimated QALYs based on the SF-12 (SF-6D) instrument. A supplementary analysis reported results based on the EQ-5D-5L crosswalk (to the EQ-5D-3L).³¹

The QALY calculation was estimated on an area under the curve approach, adopting the following assumptions:

- All follow-ups were carried out at the exact, prespecified time period (0, 6, 12 and 18 months).
- A patient who died with their last EQ-5D-5L observation as positive had a linear fall in HRQoL from this point until death.
- A patient who died with their last EQ-5D-5L observation as negative had this constant level of HRQoL from this point until death.

Notation: E_0 = baseline utility, E_6 = utility at 6 months, E_{12} = utility at 12 months, E_{18} = utility at 18 months, t = duration in each health state in days. If EQ-5D-5L was present at baseline, 6, 12 and 18 months’ follow-up, QALYs were calculated by:

$$\text{Total QALYs} = (((t/365) \times (E_0 + E_6))/2) + (((t/365) \times (E_6 + E_{12}))/2) + (((t/365) \times (E_{12} + E_{18}))/2). \quad (1)$$

Resource use and costs

The perspective for the costs was the health and social care provider (i.e. wider costs, such as productivity loss or out-of-pocket expenses, were not included).

We used a combination of patient-reported data, CRF data capture and published estimates to derive costs during the evaluation. The general approach, individual unit costs and costing assumptions were verified with and refined following input from the lead clinician on the trial. The cost categories and

overall approach to establishing the cost of each intervention over the trial period and the associated source of resource use/costs is as follows:

1. device costs – taken from publications and online resources
2. procedure costs – taken from national tariffs and online resources
3. medications – based on patient report
4. primary/social care – based on patient report
5. secondary care after the initial procedure – generated by attributing costs to explants, complications and further treatments/surgery.

The strategy for secondary care resource use capture deviates from the strategy that was planned. We had originally aimed to request NHS digital data to capture this aspect of resource; however, as a result of the early stopping of the trial, the associated reduction in research funding and the resultant limited sample size, this was considered to be impractical and was not pursued. This decision was made following consultation with the trial team. Since secondary care costs (hospital visits and stay) were not captured directly elsewhere in the trial, we elected to capture them via the length of stay of the original procedure and by costing subsequent explants, complications and surgeries.

The combined device and procedure costs are included in *Table 22* and include the device-specific cost and a day-case rate for insertion of a generic neurostimulator for treatment of FI costs. These costs are

TABLE 22 Device and procedure costs

Item	Cost (2018/19 prices)	Source and assumptions	Inflated from year
SNS: temporary			
Temporary device cost	£680	Hounsome and Roukas ³²	N/A
Temporary device procedure costs	£1819	Insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³)	N/A
Excess bed-day cost for stays > 1 day	£362	Insertion of neurostimulator for treatment of faecal incontinence, elective short stay excess bed-day (FF47Z) (NHS Reference Costs 2017–2018 ³³)	N/A
SNS: permanent			
Permanent device cost	£7750	Hounsome and Roukas ³²	N/A
Permanent procedure costs	£1819	Insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³)	N/A
Excess bed-day cost for stays > 1 day	£362	Insertion of neurostimulator for treatment of faecal incontinence, elective short stay excess bed-day (FF47Z) (NHS Reference Costs 2017–2018 ³³)	N/A
FENIX: MSA			
Sizing tool	£300	URL: www.io.nihr.ac.uk/wp-content/uploads/migrated/2220.838f0fdf.FENIXforfaecalinalcontinenceFINAL.pdf (accessed 12 February 2021)	N/A
Device cost	£4000	URL: www.io.nihr.ac.uk/wp-content/uploads/migrated/2220.838f0fdf.FENIXforfaecalinalcontinenceFINAL.pdf (accessed 12 February 2021)	N/A
Procedure cost	£1819	Assumed same as SNS: insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³)	N/A
Excess bed-day cost for stays > 1 day	£362	Insertion of neurostimulator for treatment of faecal incontinence, elective short stay excess bed-day (FF47Z) (NHS Reference Costs 2017–2018 ³³)	N/A
N/A, not applicable.			

fixed, except for the cost of the hospital stay. Any nights spent in hospital (including the first night) were costed using the unit cost shown for an excess bed-day. Data collected in the trial (e.g. relating to theatre time during procedure) may have allowed for a more granular approach to costing the procedure; however, given the small sample sizes, it may be prone to undue bias from outliers; thus, we opted to use NHS reference cost tariffs.

An operative form was completed for each patient following the procedure. The form captures operation details, including which procedure the patient received (temporary SNS and/or permanent SNS, or FENIX MSA) and any intraoperative complications that may occur. The form also reports whether patients receive a local, general or spinal anaesthetic, and whether or not the procedure has been completed. Furthermore, a 2-week postoperative review was issued for each patient to gather information on whether or not patients progress to permanent SNS, if they were randomised to receive temporary SNS, whether or not the device they received is still in situ and a reason provided if the device has been explanted. This discharge form was also used to record whether or not patients who have been discharged receive further surgery or suffer from any postoperative complications. Follow-up forms were completed by the local research teams at 6, 12 and 18 months following randomisation and asked for similar patient information. Individual complications and further surgeries were costed using the unit costs given in *Appendix 1, Tables 39 and 40*, respectively. With the exception of explantation, we assumed that the costs of any intraoperative complications were captured in the length of stay of the initial procedure.

At baseline and at 6-month, 12-month and 18-month follow-up, patients were provided with a specially designed form to report on their (primary and social) health-care resource use in the past 3 months: general practitioner (GP), practice nurse, district nurse, physiotherapist, occupational therapist, counsellor in the GP surgery visits, clinic (non-hospital) by telephone/e-mail or at home. They were also asked if they received support from social services, including meals-on-wheels deliveries, laundry services, care workers/help at home or social worker visits, and the frequency of this support. Furthermore, they were asked to provide information on the use of community/residential services, including any visits and overnight stays at convalescent homes, nursing homes and day centres. Patients were also asked to note what medication relating to their FI they were taking on a regular basis. Unit costs for medications are listed in *Appendix 1, Table 41*, and for primary care/social care in *Appendix 1, Tables 42 and 43*. We assumed that any follow-up care costs associated with the interventions are captured in the self-reported primary care use data. The 3-month recall data were multiplied by two to obtain 6-month values.

In a majority of cases, unit costs were based on national resources such as the Personal Social Services Research Unit (PSSRU) report, NHS Reference Costs and the Electronic Market Information Tool (eMIT). Elsewhere, we used other web resources and previous publications as the basis for unit costs for health-care resources. All costs are presented in Great British pounds and the price year is 2018 (costs are inflated to 2018/19 prices where necessary using the Consumer Price Index for health).

Analysis

The main analysis was based on an intention-to-treat principle. The trial analysis produced ICERs over the trial period (18 months). We used seemingly unrelated regression^{34,35} to account for the expected correlation between costs and QALYs while controlling for any baseline imbalances. Although we had planned to use the same covariates as the statistical analysis, analyses were run with no adjustment, and subsequently adjusting for baseline EQ-5D-5L and (primary/social care) costs given the data limitations. The regression models estimated costs and QALYs that will allow the calculation of ICERs.

Our primary analysis incorporated multiple imputation (MI) to impute missing data; however, we also conducted a complete-case analysis. Individuals with missing EQ-5D-5L items were not allocated a utility index score. Where missing values could not be dealt with in the manner described above (see *Health utility and quality-adjusted life-year calculation*), we conducted MI as part of the model estimation procedure. Costs were imputed for each individual at each time point for the following

categories: (1) total primary care and social care costs, (2) total secondary care costs and (3) total medication costs.

We present a range of sensitivity analyses to explore the impact of the assumptions made in the base case analysis on the cost-effectiveness estimates. These include the methods used to assess utility in the trial and model covariates. We also conducted non-parametric bootstrapping to calculate the probability that FENIX was cost-effective at the trial end.

The final cost and QALY estimates at 18 months were added to the beginning of the decision model time horizon, and additional costs and benefits were estimated in the model from 18 months over a lifetime.

Methods: decision model

We generated a de novo decision-analytic model to estimate cost per QALY estimates over a lifetime horizon. In addition to extending the evaluation period to a relevant horizon, the model also allows the exploration of uncertainty in the parameter estimates, individually and jointly, as part of a probabilistic sensitivity analysis (PSA). The latter will allow the production of cost-effectiveness acceptability curves (CEACs)³⁶ and estimates of the value of information (further research). The modelling was conducted in Microsoft Excel® 2016 (Microsoft Corporation, Redmond, WA, USA).

Model type, cycle length, structure and health states

The model is a Markov model describing a simplified patient pathway from the end of the trial period (18 months) over a lifetime in terms of discrete health states. The average age of the patients at the model start reflected that of the trial participants (62 years). Patients move between these health states at the end of every model cycle (in this case, 1 year) according to prespecified transition probabilities.

The structure of the decision model is shown in *Figure 6*. The structure was informed by other models used in this population^{32,37} and was finalised following input from the lead clinician.

In the model, we defined the health states as being related to effectiveness according to the percentage improvement on baseline CCIS (100–76% improvement, 75–51% improvement, $\leq 50\%$ improvement) and whether or not the device is still in situ (device explant). ‘Better’ health states (i.e. higher percentage of improvement) are associated with lower costs and higher QoL. Although we did not a priori expect the interventions to have an impact on survival, because the model is over a lifetime horizon, we also modelled background, all-cause mortality (dead health state).

Each health state is associated with a specific average cost and HRQoL (utility value). A hypothetical cohort of FI patients for each treatment arm was distributed between these health states at model start with the distribution reflecting that of the trial sample and intervention effectiveness/functionality at 18 months. These patients moved between health states at the end of each model cycle, with the likelihood of this movement determined by transition probabilities that reflected treatment effectiveness – more specifically, loss of effectiveness over time – and mortality rates.

We assumed that 18 months after initial implant, effectiveness could only deteriorate, and thus patients would be transited downwards in terms of percentage improvement from baseline. We also assumed that once explantation had occurred, another device (either SNS or FENIX) would not be implanted. Patients receiving FENIX initially are unlikely to receive another FENIX device following explantation owing to clinical viability. While it is possible that initial FENIX implants could be replaced




FIGURE 6 Decision model structure.

by SNS devices and that initial SNS devices could be replaced by either another SNS device or a FENIX device, the patient numbers on these pathways would be relatively small. Incorporating these possibilities would require additional modelling complexity, and the additional burden of data requirements (e.g. effectiveness of second implants) was unlikely to be met, and thus additional assumptions would be required. We also included in the explant health state those in the SNS group who did not have a permanent implant. The explant health state was therefore capturing 'conservative management' for those without a device in situ.

The durability of the FENIX device is unknown and the device will in theory continue to function in the long term until a fault or infection occurs. However, the SNS device has a finite battery life (approximately between 5 and 7 years), at which point explantation, battery replacement and implantation is required. We reflected this in a submodel for SNS. After battery replacement, patients were subject to the same distribution among effectiveness categories as observed in the trial arm at 18 months (rather than immediately post procedure) for those that had the permanent device implanted. We did not include another battery replacement at 14 years in the base-case model because of the added modelling complexity and the small proportion likely to have a device in situ but we tested this in simplified sensitivity analyses. Other modelling assumptions are set out in the parameter tables.

Parameter values: transition probabilities

The decision model required a set of parameter values that defined the movement of the cohort around the model (transition probabilities are shown in *Table 23*). Where possible, the transition rates for the model were estimated from the randomised controlled trial (RCT) data (observed data rather than imputed). We used the end distribution of trial patients per arm across percentage improvement groups as the starting proportions in these groups at model start. We used trial data on transition rates between these percentage improvement groups during the trial to specify a rate beyond 18 months. As a result of the limited sample size, we were unable to do this in a robust way per intervention arm, so we assumed these rates to be the same across arms.

TABLE 23 Parameter values: transition probabilities and event rates

Parameter	Value	SE	Distribution ^a	Source/reference
Common parameters				
Starting age of model cohort (years)	62	N/A	Fixed	Mean of the trial sample
Survival	Age dependent	N/A	Fixed	Taken from Office for National Statistics life tables ³⁸
FI population annual incidence in England (for EVPI)	13,126	N/A	Fixed	The NIHR Horizon Scanning Centre (2012) News Brief: FENIX™ Continence Restoration System for severe chronic faecal incontinence ²⁰ 80 new adults per 100,000 population 30 of these referred to surgical specialist Adult population in England = 43,752,473
Number of years for which the decision problem is relevant (for value of information)	10	N/A	Fixed	Assumed

TABLE 23 Parameter values: transition probabilities and event rates (continued)

Parameter	Value	SE	Distribution ^a	Source/reference
FENIX				
Starting proportions				Trial data at 18 months
100–76% improvement	0.00	N/A	Dirichlet	
75–51% improvement	0.16	N/A	Dirichlet	
≤ 50% improvement	0.39	N/A	Dirichlet	
Device explant	0.42	N/A	Dirichlet	
Dead	0.03	N/A	Dirichlet	
Transition between % improvement group				Trial data
100–76% to 75–51%	0.167	N/A	Dirichlet	
100–76% to ≤ 50%	0.667	N/A	Dirichlet	
75–51% to ≤ 50%	0.800	N/A	Dirichlet	
Probability of explant	0.15	N/A	Dirichlet	
SNS				
Starting proportions				Trial data at 18 months
100–76% improvement	0.026	N/A	Dirichlet	
75–51% improvement	0.105	N/A	Dirichlet	
≤ 50% improvement	0.342	N/A	Dirichlet	
Device explant	0.526	N/A	Dirichlet	
Dead	0.000	N/A	Dirichlet	
Transition between % improvement group				Trial data
100–76% to 75–51%	0.167	N/A	Dirichlet	
100–76% to ≤ 50%	0.667	N/A	Dirichlet	
75–51% to ≤ 50%	0.800	N/A	Dirichlet	Survival curve based on trial data
Probability of explant	0.15	N/A	Dirichlet	Assumed same rate as FENIX
Probability of battery replacement	1	N/A	Fixed	Battery replacement every 7 years based on triangulation of several sources and clinical opinion ^{32,39,40}

EVPI, expected value of perfect information; N/A, not applicable.

^a Fixed indicates the value did not vary in the PSA.

We applied parametric survival curves to the time-to-explant data from the trial and extrapolated this beyond 18 months. The survival analysis indicated no increase in hazard over time, and thus we estimated a constant hazard rate using the exponential distribution in Stata[®] version 15 (StataCorp LP, College Station, TX, USA). There were no explants in the SNS group, but a clinical expert felt that this was an artefact of the sample size; hence, we used the same explant rate for both intervention groups. Following input from the same clinical expert, we assumed that survival in the FI population was not significantly different from survival in the general population and, thus, used Office for National Statistics mortality data³⁸ to model this. We also assumed that neither the interventions nor the percentage improvement had an impact on survival.

Parameter values: health state utilities

Economic evaluations are designed to inform resource allocation decisions, thus the model used the QALY outcome measure as prescribed by NICE. The estimation of QALYs requires the production of utility weights for health states; these data were derived from the RCT (Table 24). We used linear regression predicting EQ-5D-5L utility weights and health state groups as an explanatory categorical variable. The regression used multiple observations per person and robust SEs to reflect this approach.

As the interventions were assumed not to influence survival, it was not considered necessary to adjust utility values for age over the time horizon. There was no utility decrement for battery replacement as this was assumed to be a 'tunnel state' straight to percentage CCIS improvement groups (i.e. patients did not stay in this state for a sufficient period to incur substantive HRQoL loss).

Parameter values: costs

We generated health state costs from the trial data (see Table 24). As we expected, costs were at their highest immediately after the initial procedure and then fell over time (with the exception of the battery replacement for SNS). We used the 12-month and 18-month resource use data to estimate ongoing health-care costs in the model. As it was unclear whether or not the trial forms captured ongoing (conservative management) costs for the no device group, we generated a monthly cost following

TABLE 24 Parameter values: utilities and costs

Parameter	Value	SE	Distribution	Source/reference
Health state utility				
100–76% improvement	0.96	0.0304	Beta	Trial data
75–51% improvement	0.90	0.0429	Beta	Trial data
≤ 50% improvement	0.75	0.0275	Beta	Trial data
Device explant (subsequently, conservative management)	0.64	0.0801	Beta	Trial data
Health state costs (per year)				
100–76% improvement	£90.67	79.39	Gamma	Trial data
75–51% improvement	£248.99	154.45	Gamma	Trial data
≤ 50% improvement	£384.07	131.16	Gamma	Trial data
Device explant	£631.91	451.92	Gamma	Trial data
Procedure and device costs				
Device explant: procedure cost	£1819	N/A	Fixed	Insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³) (assumed same for FENIX and SNS)
Battery replacement: procedure cost	£1819	N/A	Fixed	Insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³) (assumed same for FENIX and SNS)
Battery replacement: SNS device cost	£7750	N/A	Fixed	Hounsome and Roukas ³²
N/A, not applicable.				

discussion with the lead clinician (detailed in *Table 24*) and tested this in a sensitivity analysis. We used the full set of cost data to estimate costs immediately post battery replacement, and this was anticipated to capture any complication and additional surgery costs. The explanation of a device or replacement of a battery led to the procedure costs outlined in *Table 24*.

Relevant costs for the evaluation include the FENIX device/SNS acquisition and procedure (including length of hospital stay), costs of treating adverse events/complications, costs of device explants and other surgery, and (for SNS) the cost of battery replacement. These are in addition to the health state costs that were obtained from the trial data. All costs relating to other surgeries, explants and complications within 18 months were observed/captured in the trial data, and thus were not modelled here. After 18 months, complication costs were assumed to be captured within the other health states.

Unit costs were obtained from national sources including the PSSRU,⁴¹ the *British National Formulary*⁴² (BNF) and the NHS Reference Cost³³ database. Internet searches were conducted for other cost elements.

Validation

We validated the model by checking the face validity of the model structure and parameter values with other health economists and with the lead clinician. We tested the internal validity of the model by checking that model outputs followed expectations when changes were made to certain parameter values (e.g. that the cost-effectiveness of FENIX improved following an increase in the SNS device cost or reduced following an increase in the probability of explant in the FENIX arm).

Analysis

The average costs and QALYs accrued during the trial period were inputted at the start of the model, with further costs and benefits added to these over the model horizon. We generated a deterministic ICER per QALY gained and a net monetary benefit (NMB) estimate at the model end. We tested these estimates using prespecified sensitivity analyses to identify drivers of cost-effectiveness. We specified distributions for certain parameter values (denoted in the parameter tables) and ran a PSA where random parameters were selected from these distributions and inputted into the model over 10,000 Monte Carlo simulations. The 10,000 costs and QALY estimates generated from the PSA were plotted on a cost-effectiveness plane and the average of these was used to calculate the probabilistic ICER. We used estimates of NMB to determine the probability that FENIX was cost-effective compared with SNS over a range of willingness-to-pay values on the CEAC. Since the CEAC indicates the probability of cost-effectiveness, we can calculate the probability that the optimal treatment will not be cost-effective. Using this information and the NMB provided by the alternative treatment (the net benefit loss), we used the value of information framework⁴³ to estimate the total population expected value of perfect information (EVPI). The population EVPI is estimated using the expected net benefit loss per person from the decision, the number of people with FI that would be eligible for the treatments per year (annual incidence is estimated to be 13,126) and the period for which the FENIX versus SNS decision is still relevant (here, we assumed this to be 10 years). The higher the EVPI value, the greater the potential cost of uncertainty and the greater the incentive to conduct additional research (e.g. further trials) to reduce that uncertainty prior to making a decision.

We used a NICE willingness-to-pay threshold of £20,000 per QALY to determine cost-effectiveness (or NMB value of > 0). We used a half-cycle correction to reflect the expectation that health state transitions can happen at any time during a model cycle. In accordance with current NICE recommendations, costs and outcomes post year 1 were discounted at 3.5% per annum. The perspective remained that of the health and personal social services provider.

Results

Trial analysis

Descriptive results for the costs (and cost category) are shown by arm in *Table 25*. There is a relatively modest difference in total cost between arms. However, there are notable differences in the individual cost categories, with the FENIX procedure being much cheaper than the SNS procedure, but being associated with significantly higher secondary care costs (for other complications and surgeries). Descriptive statistics on costs (and by cost category) and utility by arm, time-point and sample are included in *Appendix 1, Table 44*.

The length of stay varied between intervention arms (SNS: range 0 to 59 days, median 0 days, mean 1.43 days; FENIX: range 0 to 15 days, median 1 day, mean 1.39 days).

The economic evaluation results for the 18-month trial period are included in *Table 26*. Results are presented for complete case and following MI using EQ-5D-5L valuation, EQ-5D-5L to 3L cross-walk and SF-6D utility scores for alternative adjusted models.

For the EQ-5D-5L and SF-6D analyses, there were a total of 47 cases across both arms that met the complete-case criteria (of which 42 cases were common across both the EQ-5D-5L and the SF-6D complete cases), whereas the MI analysis sample total was 94 cases. Five of the original 99 cases were dropped because they had no baseline data or only baseline data. In all analyses, FENIX was more expensive in the trial period than SNS, although this was reduced in the imputed analyses. In all of the EQ-5D-5L analyses, except for the complete-case, 3L cross-walk with adjustment (for age and baseline utility), FENIX led to a QALY benefit compared with SNS. We present the results for adjustment for operating surgeon, but because of the small sample sizes these figures are highly variable and cannot be considered robust.

The ICER for the primary analysis (MI, direct EQ-5D-5L valuation with adjustment for age and baseline utility) is £44,785.62 per QALY gained and is well above the £20,000 threshold. In this analysis, FENIX was associated with modest additional costs and negligible incremental QALYs compared with SNS. Uncertainty in the results was represented in the CEAC (*Figure 7*). At a willingness-to-pay threshold of £20,000 per QALY gain, FENIX has a 41.3% chance of being cost-effective.

The supplementary analysis using MI and a cross-walk to derive EQ-5D-3L values yields a much higher QALY gain and lower ICER for FENIX. The analyses based on the SF-6D indicated a QALY loss associated with FENIX, which was therefore dominated.

TABLE 25 Cost breakdown

Cost category ^a	FENIX (£)		SNS (£)	
	Mean	SE	Mean	SE
Procedure	6190.55	274.96	8176.49	926.80
Medication	65.48	20.02	31.47	7.91
Implants	541.83	122.65	503.13	119.97
Primary care	701.30	115.38	549.32	102.40
Secondary care	2663.85	1119.02	597.11	283.86
Total	10,163.01	1266.37	9857.51	888.44

^a These values are based on the MI (n = 94), EQ-5D-5L, unadjusted analysis.

TABLE 26 Cost-effectiveness at 18 months

Analysis	Costs (£)					QALYs					ICER		
	n	FENIX (95% CIs)	n	SNS (95% CIs)	Incremental (95% CIs)	n	FENIX (95% CIs)	n	SNS (95% CIs)	Incremental (95% CIs)	Unadjusted	Adjusted for baseline utility + age	Adjusted for baseline utility + age + surgeon
Complete-case analysis													
EQ-5D-5L	24	11,360.25 (6506.76 to 16,213.74)	23	10,913.08 (7870.44 to 13,955.73)	447.17 (-4912.45 to 5806.79)	24	1.157 (1.042 to 1.273)	23	1.148 (1.018 to 1.277)	0.010 (-0.150 to 0.170)	44,051.88 (n = 47)	117,153.71 (n = 47)	440.57 (n = 46)
EQ-5D-3L cross-walk	24	11,360.25 (6506.76 to 16,213.74)	23	10,913.08 (7870.44 to 13,955.73)	447.17 (-4912.45 to 5806.79)	24	1.030 (0.895 to 1.166)	23	1.023 (0.888 to 1.159)	0.007 (-0.170 to 0.185)	62,121.75 (n = 47)	Dominated (n = 47)	303.05 (n = 46)
SF-6D	24	11,546.51 (6685.34 to 16,407.68)	23	9597.22 (6342.64 to 12,851.81)	1949.28 (-3517.45 to 7416.02)	24	0.975 (0.890 to 1.051)	23	1.038 (0.950 to 1.125)	-0.062 (-0.169 to 0.045)	Dominated (n = 47)	Dominated (n = 47)	Dominated (n = 45)
Imputed data set													
EQ-5D-5L	47	10,163.01 (7610.86 to 12,715.17)	47	9857.51 (8067.04 to 11,647.99)	305.50 (-2694.59 to 3305.58)	47	1.129 (1.050 to 1.207)	47	1.124 (1.039 to 1.209)	0.005 (-0.106 to 0.115)	67,697.50 (n = 94)	44,785.62 ^a (n = 94)	124,387.65 (n = 87)
EQ-5D-3L cross-walk	47	10,164.88 (7613.59 to 12,716.16)	47	9869.06 (8077.85 to 11,660.27)	295.82 (-2702.91 to 3294.54)	47	1.019 (0.934 to 1.103)	47	0.986 (0.890 to 1.082)	0.032 (-0.091 to 0.156)	9193.58 (n = 94)	12,804.46 (n = 94)	Dominated (n = 87)
SF-6D	47	10,152.95 (7607.32 to 12,698.58)	47	9853.67 (8062.22 to 11,645.12)	299.28 (-2696.15 to 3294.72)	47	0.994 (0.940 to 1.049)	47	0.997 (0.944 to 1.050)	-0.003 (-0.076 to 0.070)	Dominated (n = 94)	344,418.94 (n = 94)	100,461.45 (n = 87)
a Primary analysis.													




FIGURE 7 Cost-effectiveness acceptability curve.

The proportion of patients distributed between the CCIS percentage improvement groups is shown in *Table 23* (parameter table). For both arms, a majority of patients were either in the < 50% improvement group or did not have a device in situ. Where the device was in situ, the results appear to indicate a slight benefit for the SNS group, with more patients in the 50–75% and > 75% improvement groups. One patient in the FENIX arm had died by 18 months and, although this is unlikely to be attributable to randomisation arm, we retained this patient in the analysis.

The costs and benefits estimated during the trial period were added to the start of the model and assumed to be fixed in the base case deterministic analysis. In each simulation of the probabilistic analysis, we used a bootstrapped cost and QALY generated from the trial data.

Decision modelling

The model Markov trace is included in *Figure 8*. This gives an indication of the predicted intervention effectiveness and survival over time. There are very few noticeable differences in the curves over the lifetime horizon. The kinks in the SNS figure show where those patients with a device in situ receive a battery replacement at 7 years and the redistribution among CCIS percentage improvement groups.

The deterministic model results, including lifetime costs, QALYs and ICER, are presented in *Table 27*. Estimated lifetime costs are higher in the SNS group, as are the QALYs. The ICER and NMB values indicate that FENIX is not cost-effective over this time horizon (SNS is the optimal strategy). These results are mirrored by the probabilistic results, where the ICER for SNS versus FENIX drops.

Figure 9 plots the 10,000 costs and QALYs from the Monte Carlo simulations. The results highlight significant uncertainty in the model outputs. The ICER of the mean value falls just below the willingness-to-pay threshold (for SNS minus FENIX). The uncertainty in the results is further represented in the CEAC (*Figure 10*). At a willingness-to-pay threshold of £20,000 per QALY gain, SNS has a 55% chance of being the optimal strategy (and, therefore, FENIX has a 45% chance of being the optimal strategy).

Table 28 includes a range of deterministic sensitivity analyses. The analyses include using complete-case costs and QALYs from the trial, changing the year at which battery replacement is conducted in the SNS group, changes to health state costs and utilities, and assuming equivalent effectiveness across interventions at 18 months. In most analyses, the decision is unchanged (i.e. SNS remains the optimal strategy). A significant drop in the utility estimates for those without a device in situ substantially reduces the QALY gain from SNS over time and changes the decision such that FENIX would be the optimal strategy.




FIGURE 8 Markov model traces: (a) SNS; and (b) FENIX.

TABLE 27 Deterministic and probabilistic base-case cost-effectiveness results

Intervention	Costs (discounted)	QALYs (discounted)	Incremental cost (discounted)	Incremental QALYs (discounted)	ICER	NMB
Deterministic						
FENIX	£18,657	10.12				£183,806
SNS	£19,972	10.33	£1315	0.20	£6508	£186,531
Probabilistic						
FENIX	£18,668	10.09				£183,130
SNS	£19,975	10.32	£1306	0.23	£5694	£186,413

In all cases, we have reversed the incremental value calculation for ease of interpretation. In the alternative format, FENIX leads to negative costs, QALYs and ICERs, which would denote the money saved per QALY loss.




FIGURE 9 Cost-effectiveness plane: SNS vs. FENIX. The orange dot represents the ICER using the mean from the cost and QALY simulations.




FIGURE 10 Cost-effectiveness acceptability curve.

TABLE 28 Deterministic one-way and scenario analyses

Intervention	Costs	QALYs	Incremental cost	Incremental QALYs	ICER	NMB
Base case scenario						
FENIX	£18,657	10.12				£183,806
SNS	£19,972	10.33	£1315	0.20	£6508	£186,531
Complete-case scenario						
FENIX	£19,855	10.15				£183,189
SNS	£21,027	10.35	£1173	0.20	£5973	£185,943
Battery replacement at year 5						
FENIX	£18,657	10.12				£183,806
SNS	£20,681	10.33	£2023	0.21	£9830	£185,899

TABLE 28 Deterministic one-way and scenario analyses (continued)

Intervention	Costs	QALYs	Incremental cost	Incremental QALYs	ICER	NMB
Additional battery replacement at year 14 (only cost implications)						
FENIX	£18,657	10.12				£183,806
SNS	£20,300	10.33	£1643	0.20	£8134	£186,203
50% increase in FI cost after explant (£1264)						
FENIX	£25,486	10.12				£176,977
SNS	£27,555	10.33	£2069	0.20	£10,245	£178,948
Cost of FI assuming 100% use pads, 7.5% use irrigation, 2.5% use anal plug and 7.5% have a stoma (£1895 on average per year; base case £632)						
FENIX	£32,303	10.12				£170,160
SNS	£34,798	10.33	£2495	0.20	£12,353	£171,705
25% reduction in the utility for patients after device explanted						
FENIX	£18,657	8.32				£147,792
SNS	£20,300	8.43	£1643	0.10	£15,784	£148,230
50% reduction in the utility for patients after device explanted						
FENIX	£18,657	6.52				£111,777
SNS	£20,300	6.53	£1643	0.01	£265,313	£110,258
FENIX proportion of patients at end of trial equal to SNS						
FENIX	£18,904	10.32				£187,542
SNS	£20,300	10.33	£1397	0.00	£489,114	£186,203
In all cases, we have reversed the incremental value calculation for ease of interpretation. In the alternative format, FENIX leads to negative costs, QALYs and ICERs, which would denote the money saved per QALY loss.						

FENIX also becomes the optimal strategy (cost saving but marginally less effective) when the proportions of patients across CCIS percentage improvement and explant groups are assumed to be equivalent at the trial end.

Table 29 includes the per patient and population EVPI. These values reflect a monetary cost (attaching a value of £20,000 to a QALY) of the uncertainty in the current decision problem. At the NICE willingness-to-pay per QALY threshold of £20,000, there was a 55% and a 45% chance that SNS and FENIX, respectively, were the optimal intervention and, hence, there was significant uncertainty remaining in the decision. Given the large number of potential intervention recipients, the cost of this uncertainty over 1 year and 10 years is considerable. The EVPI estimates indicate that further research in this area is warranted to reduce the level of decision uncertainty.

TABLE 29 Expected value of perfect information ($\lambda = £20,000$)

Parameter	Value
Per person EVPI	£9630
FI annual incidence	13,126
Total population EVPI at 1 year	£126,403,677
Total population EVPI at 10 years	£1,264,036,774

Chapter 4 Discussion

Although the SaFaRI trial failed to recruit the planned 350 participants, it is nevertheless the largest reported evaluation of the MAS (FENIX) device to our knowledge, providing important information about circumanal implants to augment anal sphincter function and treat FI. Despite initial encouraging results in small, single-centre studies, which provided the rationale for this larger randomised clinical trial, the FENIX device was taken off the market in 2017, when the manufacturing company (Torax Medical, Minneapolis, MN, USA) was purchased by Johnson and Johnson (New Brunswick, NJ, USA). The decision to discontinue marketing the FENIX device was said to be strategic, with Johnson and Johnson continuing to market a related device (LINX®) for the treatment of gastro-oesophageal reflux.

The SaFaRI trial involved 18 pelvic floor centres, with colorectal surgeons from across the UK who were members of the Association of Coloproctology of Great Britain and Ireland. The majority of cases were recruited from a single site (St James's University Hospital, Leeds), which may have an impact on the generalisability of the results. In mitigation, strenuous attempts were made to standardise the technique for MAS implantation with preceptorship of training cases performed to manufacturer guidelines. The technique for SNS implantation was left to individual surgeons to reflect normal practice.

The study population was typical of that suffering from moderate to severe FI, with the majority of participants being female and a median age of 59.0 years. In accordance with the known aetiology of FI, previous obstetric injury accounted for the majority of cases, with mixed passive and urge incontinence being the predominant symptomatology.

Previous studies of SNS have reported outcomes at time points relative to the patient's operation (either temporary or permanent SNS), which makes comparison with the results of the SaFaRI trial difficult. In the SaFaRI trial, the patient outcomes were measured at time points relative to randomisation (i.e. from the key decision-making event, rather than from operation). This approach evaluates the effect of the decision of which intervention to give the patient. It provides data that is more informative to clinicians and patients when weighing up the risks and benefits of surgical treatment for FI, and is preferred over the previously reported per-protocol analyses.

The 18-month follow-up post randomisation was chosen with the expectation that definitive device operations (FENIX and permanent SNS) would occur within around 6 months post randomisation, giving patients around 12 months' follow-up post operation. The median time of 57.0 days from randomisation to implantation of a FENIX device probably conforms to normal NHS practice. The median time of 371 days from randomisation to implantation of a permanent SNS device, although longer than our a priori expectation, might not be unusual in the NHS, where there are surgical capacity issues and benign conditions tend to be given low priority. This delay in undertaking permanent SNS implantation had a bearing on the number of patients who had a device in situ and had completed 18 months' follow-up at the time of stopping the study.

In total, five (10%) participants withdrew from surgery in the FENIX group and did not undergo device implantation. This compared with five (10.2%) participants not receiving a temporary SNS device, four (8.2%) participants not undergoing implantation, and a further 12 (24.5%) participants not proceeding from a temporary device to a permanent device owing to lack of efficacy. This gives an intention-to-treat figure of 57.1% and a per-protocol figure of 62.2% of patients progressing to a permanent implant, which is generally in keeping with the literature. However, the success rate at 18 months, as judged by the intention-to-treat analysis, was very low at 8.2%. Even on a per-protocol basis, of the 28 participants who received a permanent SNS implant, only four (14.3%) reported a successful outcome with the device in situ and sustained benefit at 18 months' follow-up. This contrasts with the literature, which generally reports success rates between 50 and 70%. In a long-term analysis of 407 patients

undergoing SNS, Altomare *et al.*⁴⁴ reported progression from temporary to permanent implant in 66.8% of patients. At a median of 84 months' follow-up, success was maintained in 71.3% of patients in per-protocol analysis and 47.7% on intention-to-treat analysis.⁴⁴ Similar results were reported in the systematic review published by Thin *et al.*,⁴⁵ with the median success rate of SNS, on intention to treat, being 63%, 58% and 54% on short-, medium-, and long-term follow-up, respectively.

The reason for the large discrepancy between the success rates observed in SaFaRI and those reported in the literature is not clear. One possible explanation is the rigour of data collection within the context of a RCT, giving a more accurate reflection of true success. It is possible that the large number of sites participating in SaFaRI, compared with the single-centre studies reported in the literature, may have captured data from widely different practices, although all participating surgeons were experienced with the SNS technique prior to participation in the study. The proportion of patients who progressed from the temporary SNS device to the permanent SNS device is similar to the proportions seen in the literature. The discrepancy is seen in the number of successes in the patients who received the permanent SNS implant. Four patients did not receive the permanent SNS device before 18 months post randomisation, and so were classed as 'failures' according to the definition of the primary end point, which may explain some of the discrepancy. Most other studies have used a 50% reduction in the number of FI episodes per week as their end point rather than 50% improvement in CCIS, which was used in SaFaRI. This could also explain some of the discrepancy between the results of SaFaRI and those reported in the literature.

A similar low success rate was observed with the FENIX device, with only six (13.3%) successes out of 45 participants who underwent device implantation. Again, this is much lower than in the reported literature. Pakravan *et al.*⁴⁶ reported a success rate of 76% in 18 patients undergoing FENIX implantation, defined as a $\geq 50\%$ reduction in FI rates per week, with no explants. In a longer-term study, Sugrue *et al.*⁴⁷ reported on 37 FENIX patients with a median follow-up of 5 years. Therapeutic success rates at 1, 3 and 5 years were 63%, 66% and 53%, respectively. The lower success rate of the FENIX device in SaFaRI appears to have been because of a higher proportion of patients not experiencing an improvement in CCIS (36.6% of the 41 patients with complete data) and a higher explant rate (33% of the 45 implants).

Both the FENIX and the SNS interventions proved to be relatively safe, with few intraoperative complications. However, the postoperative complications were high, particularly in the FENIX group (73.3%), with participants receiving the FENIX intervention being 11 times more likely to suffer a postoperative complication, a difference that did not appear to be related to the effect of the operating surgeon. Notably, the most commonly reported complication in the FENIX group was worsening symptoms of obstructed defaecation (20.5%). Recommendation was made that patients should have an ODS of < 8 for entry to SaFaRI. The baseline characteristics for the whole group showed a median ODS of 7.0 (range 2.0–18.0), and perhaps with more stringent inclusion criteria this complication would have been reduced. However, there was no difference in ODS between the two treatment arms and having a device in situ did not have a significant effect on ODS. It is possible that patients in the FENIX group were more likely to attribute any obstructed defaecation symptoms to the implanted device. The high complication rate with the FENIX device is mirrored in the literature. Sugrue *et al.*⁴⁷ reported 30 adverse events in 35 patients undergoing FENIX implantation, with the most common being defaecatory dysfunction (20%), followed by pain (14%), erosion (11%) and infection (11%).⁴⁷ In Pakravan *et al.*'s⁴⁶ smaller study of 18 patients, the most common complication was postoperative pain (29%). The FENIX explant rate observed in SaFaRI was much higher than previously reported, with 15 out of the 45 implants (33.3%) being explanted by the 18-month time point, with the majority explanted within 6 months. Similarly, the complication rate in the SNS group (22.5%) was higher than might be expected from the literature. This might be a result of rigorous data collection within a randomised trial and the fact that all possible complications were recorded, which is not always the case in single-centre cohort studies.

For those patients who retained a device, there was a statistically significant benefit in terms of reduction in CCIS, but there was no difference between the treatment groups, with the effect sustained throughout follow-up. This benefit was independent of baseline CCIS. Likewise, for patients who retained a device there was a sustained improvement in QoL as calculated by the EQ-5D-5L score, but not the VAS, with no difference between the treatment groups. The FIQoL showed a similar pattern, with improvement in all QoL domains if the device remained in situ. Both Pakravan *et al.*⁴⁶ and Sugrue *et al.*⁴⁷ reported improvement in FIQoL scores across all domains following FENIX implantation, although several studies attest to the benefit of SNS in improving FIQoL.⁴⁸⁻⁵⁰ The benefit from a functioning FENIX or SNS device appears to be more on physical than mental abilities, as judged by the SF-12 v2 analysis.

An economic evaluation of FENIX compared with SNS was undertaken from the perspective of the social and health-care provider. Estimates of cost-effectiveness were produced over the trial period and over a lifetime horizon using a de novo decision model.

At the end of the trial, slightly lower costs were observed in the SNS group despite the fact that intervention costs were significantly higher in this group. There were notably higher secondary care costs in the FENIX arm, which appears to be driven by the complication and other surgery rates. There were also higher (self-reported) primary care costs in the FENIX arm.

There was a small QALY intervention differential at 18 months in favour of FENIX; however, at 18 months, the ICERs for FENIX compared with SNS were > £20,000, except for analyses using cross-walking to EQ-5D-3L. The patient proportions across health state groups at the end of the trial (feeding into the decision model) slightly favoured SNS, despite this group having a higher proportion of patients without a device in situ. The trial results were highly sensitive to choice of estimation model and utility derivation strategy.

The lifetime expected costs and QALYs generated by the decision model were a reversal of those observed at 18 months; that is, in the long term, FENIX was less costly but also less effective. The base case analysis still indicated that FENIX was not cost-effective compared with SNS and that the latter strategy was optimal. This finding held in most sensitivity analyses, except where there were significant reductions in the explanted group QoL and where the effectiveness at trial end was assumed to be equivalent between arms.

Our model estimated lifetime costs similar to the 5 years' costs (€22,150) estimated elsewhere for SNS,¹³ and it is unclear what drives this disparity. Another evaluation of SNS also used a 5-year time horizon³² yielding similar 5-year SNS costs (£13,829–19,153). There are few other published economic evaluations in this population^{37,51} and none comparing SNS with FENIX. One abstract reports this direct comparison over 1 year and finds similar results to ours (FENIX being more effective and more costly) over this period, but finding an ICER of < £20,000.⁵²

Limitations

The SaFaRI study was limited by the small number of patients recruited as a consequence of the FENIX device being taken off the market less than one-third of the way through the study. Any conclusions therefore have to be taken with caution, given that the sample size was inadequate to show a difference between the two treatments. Despite the small numbers, important information has been gained, particularly relating to the patient pathways, delays in treatment and the complication profiles of the two treatments.

As expected, given the small sample and loss to follow-up, there is significant uncertainty in the cost-effectiveness results. The 18 months' results were influenced by the approach to dealing with missing data and utility assessment. It is possible that the difference in post-procedure follow-up timing across

arms may have affected the relative costs and QALYs captured, increasing uncertainty. The longer-term results were heavily reliant on the effectiveness data from the trial, which are relatively weak. A number of assumptions were necessary in the decision modelling (not all of which were incorporated in sensitivity analyses) and these may have a significant impact on the model outputs. A larger sample size would have permitted greater confidence in the model parameter values generated from the trial data. More nuanced approaches may also have been enabled (e.g. having differing effectiveness decay rates for the two interventions). Our analyses were limited to the health-care perspective; taking a broader view (e.g. including patient out-of-pocket costs) may have affected results, especially if those without a device in situ covered a proportion of conservative management costs themselves.

Given these factors and the level of uncertainty in the results, it is difficult to commend one intervention over another on the basis of the evidence presented. The results from the value of information analysis suggest that further research is warranted.

Future research

Future research might include longer follow-up of SaFaRI participants to ensure accurate capture of efficacy and explant rates in the longer term. There would be scope for further economic modelling to include standard (non-operative) care and additional data from other studies, and new devices for FI. Because SaFaRI was terminated early, as a result of a buyout of the manufacturer, the sample size is greatly reduced from that originally planned. There is a possible option of undertaking a meta-analysis of a similar study undertaken in France⁵² to gain a better understanding of the efficacy and safety of the FENIX and SNS devices.

Chapter 5 Conclusions

The SaFaRI study suggests that neither FENIX nor SNS is as effective in treating FI of moderate to severe severity as previously reported in the literature. The FENIX device was associated with a high postoperative morbidity, including the risk of explantation within the first year. The FENIX device was withdrawn from the market in 2017 and the results from SaFaRI would not support its reintroduction in its current form. Similar high rates of complications and explantation have been observed with other circumanal sphincter augmentation devices, such as the Acticon Neosphincter® (American Medical Systems, Minneapolis, MN, USA). New strategies are needed for anal sphincter augmentation that avoid the use of prosthetic implants that can become infected and erode.

Although the results for SNS, as observed in SaFaRI, were disappointing, it remains a recommended treatment by NICE for patients with moderate to severe FI that is resistant to medical management. It is unlikely, given its established place in surgical practice, that the use of SNS will decline as a consequence of SaFaRI. However, more research is required to unravel why the pathways for SNS are so inefficient, with patients suffering long waiting times for treatment of a disabling condition. The technique for SNS continues to evolve and new devices are coming to the market. The NIHR SUBSoNIC study is currently recruiting and investigating the efficacy and mechanism of subsensory (optimised) neuromodulation in adults with FI.⁵³ It will be interesting to see how the outcomes of this study will compare with those of SaFaRI.

Use of our health economics model to incorporate synthesised evidence from other studies and explore other scenarios (e.g. new devices with longer battery lives) may be informative in the future. Previous research indicated that SNS was only marginally cost-effective compared with conservative management¹² and perhaps the latter strategy could be included in a wider evaluation.

Acknowledgements

We are indebted to the patients who participated in this trial. We would also like to thank the TSC (Charles Knowles, Doreen McClurg, Graeme MacLennan and Tara Willson), the DMEC (Paul-Antoine LeHur, Natalie Rowland and Lynn Shaw), additional members of the Trial Management Group (Gregory Taylor, Sushil Maslekar, David Meads, Armando Vargas-Palacios, Alison Smith, Debbie Beirne, Adam Douglas, Jacqueline Emkes, Catherine Moriarty, David Protheroe, Jen Lodge and Karen Nugent) and the CTRU Project Team for their important contributions. Sarah Abraham provided assistance during the economic evaluation.

Patient and public involvement

The SaFaRI study had patient and public involvement (PPI) throughout its lifetime, from the initial stages of protocol development through to trial completion. Adam Douglas and Jacqueline Emkes served on the Trial Management Group and Tara Willson was a member of the TSC.

All three PPI representatives regularly attended trial oversight committee meetings and were actively involved in trial discussions, providing valued opinions and ideas from a patient and public perspective. PPI input also fed into the protocol design and the drafting/review of participant information resources.

The following institutions and surgeons participated in the trial:

Bristol Royal Infirmary: Jonathan Randall; The Churchill Hospital: Helen Jones, Ian Lindsey and Kim Gorissen; Derriford Hospital: Chris Oppong; Dewsbury and District Hospital: Adeshina Fawole; Good Hope Hospital: Haney Youssef and Mark Chapman; Leicester Royal Infirmary: Andrew Miller; Manchester Royal Infirmary: Finlay Curran; Northern General Hospital: Steve Brown; Poole Hospital: Andrew Clarke; Royal Devon and Exeter Hospital: Patricia Boorman; Royal Victoria Infirmary: Stefan Plusa; Southampton General Hospital: Sophie Pilkington; St James's University Hospital: David Jayne, Julian Hance and Sushil Maslekar; St Mark's Hospital: Carolynne Vaizey; St Peter's Hospital: Pasha Nisar; University College London Hospital: Richard Cohen and James Crosbie; University Hospital of North Durham: Susan Green; Wythenshawe Hospital: Karen Telford.

Contributions of authors

David G Jayne (<https://orcid.org/0000-0002-8725-3283>) (Consultant Colorectal Surgeon), **Neil Corrigan** (<https://orcid.org/0000-0002-1424-9830>) (Statistician), **Julie Croft** (<https://orcid.org/0000-0001-7586-3394>) (Clinical Triallist), **Vicky Napp** (<https://orcid.org/0000-0001-6726-2222>) (Clinical Triallist) and **Julia M Brown** (<https://orcid.org/0000-0002-2719-7064>) (Statistician) designed the study and were involved in study co-ordination.

Annabelle E Williams (<https://orcid.org/0000-0001-8832-6686>) (ST7 Specialist Trainee in Colorectal and General Surgery) was the Clinical Research Fellow for the study, and provided invaluable clinical input during site set-up and the initial stages of recruitment.

Neil Corrigan, Alison Pullan (<https://orcid.org/0000-0001-6060-9123>) (Statistician) and **Julia M Brown** were responsible for the statistical analysis.

Rachel Kelly (<https://orcid.org/0000-0002-2912-2859>) (Clinical Triallist) was responsible for data management and was involved in trial co-ordination.

ACKNOWLEDGEMENTS

David Meads (<https://orcid.org/0000-0003-1369-2483>) (Health Economist), **Armando Vargas-Palacios** (<https://orcid.org/0000-0002-6503-0134>) (Health Economist) and **Adam Martin** (<https://orcid.org/0000-0002-2559-6483>) (Health Economist) conducted the cost analysis and reporting.

Claire Hulme (<https://orcid.org/0000-0003-2077-0419>) (Health Economist) was involved in the design of the health economics component of the study.

David G Jayne, **Steven R Brown** (<https://orcid.org/0000-0002-0980-2793>) (Consultant Colorectal Surgeon), **Sushil Maslekar** (<https://orcid.org/0000-0002-8233-8480>) (Consultant Colorectal Surgeon), **Andrew Clarke** (<https://orcid.org/0000-0003-3585-8704>) (Consultant Colorectal Surgeon) and **Pasha Nisar** (<https://orcid.org/0000-0001-8374-6761>) (Consultant Colorectal Surgeon) made significant contributions to participant recruitment.

Karen Nugent (<https://orcid.org/0000-0002-0408-2950>) (Consultant Colorectal Surgeon), **Jen Lodge** (<https://orcid.org/0000-0002-5814-6630>) (Senior Community Incontinence Advisor), **David Protheroe** (<https://orcid.org/0000-0001-9139-0726>) (Consultant Liaison Psychiatrist) contributed to the trial design and sat on the SaFaRI Trial Management Group.

Steven R Brown also contributed to the trial design.

All authors contributed to data interpretation and the writing and review of the manuscript.

Publication

William AE, Croft J, Napp V, Corrigan N, Brown JM, Hulme C, *et al.* SaFaRI: sacral nerve stimulation versus the FENIX magnetic sphincter augmentation for adult faecal incontinence: a randomised investigation. *Int J Colorectal Dis* 2016;**31**:465–72.

Data-sharing statement

All data requests should be submitted to the corresponding author for consideration. Access to anonymised data may be granted following review.

Patient data

This work uses data provided by patients and collected by the NHS as part of their care and support. Using patient data is vital to improve health and care for everyone. There is huge potential to make better use of information from people's patient records, to understand more about disease, develop new treatments, monitor safety, and plan NHS services. Patient data should be kept safe and secure, to protect everyone's privacy, and it's important that there are safeguards to make sure that it is stored and used responsibly. Everyone should be able to find out about how patient data are used. #datasaveslives You can find out more about the background to this citation here: <https://understandingpatientdata.org.uk/data-citation>.

References

1. Williams AE, Croft J, Napp V, Corrigan N, Brown JM, Hulme C, *et al.* SaFaRI: sacral nerve stimulation versus the FENIX magnetic sphincter augmentation for adult faecal incontinence: a randomised investigation. *Int J Colorectal Dis* 2016;**31**:465–72 <https://doi.org/10.1007/s00384-015-2492-3>
2. National Institute for Health and Care Excellence (NICE). *Faecal Incontinence in Adults [QS54]*. 2014. URL: www.nice.org.uk/guidance/qs54 (accessed 11 May 2019).
3. Ferrara A, De Jesus S, Gallagher JT, Williamson PR, Larach SW, Pappas D, *et al.* Time-related decay of the benefits of biofeedback therapy. *Tech Coloproctol* 2001;**5**:131–5. <https://doi.org/10.1007/s101510100014>
4. Maeda Y, Laurberg S, Norton C. Perianal injectable bulking agents as treatment for faecal incontinence in adults. *Cochrane Database Syst Rev* 2010;**5**:CD007959. <https://doi.org/10.1002/14651858.CD007959.pub2>
5. Maeda Y, Lundby L, Buntzen S, Laurberg S. Suboptimal outcome following sacral nerve stimulation for faecal incontinence. *Br J Surg* 2011;**98**:140–7. <https://doi.org/10.1002/bjs.7302>
6. Mundy L, Merlin TL, Maddern GJ, Hiller JE. Systematic review of safety and effectiveness of an artificial bowel sphincter for faecal incontinence. *Br J Surg* 2004;**91**:665–72. <https://doi.org/10.1002/bjs.4587>
7. Chapman AE, Geerdes B, Hewett P, Young J, Eyers T, Kiroff G, Maddern GJ. Systematic review of dynamic graciloplasty in the treatment of faecal incontinence. *Br J Surg* 2002;**89**:138–53. <https://doi.org/10.1046/j.1365-2168.2002.02018.x>
8. Matzel KE, Stadelmaier U, Hohenfellner M, Gall FP. Electrical stimulation of sacral spinal nerves for treatment of faecal incontinence. *Lancet* 1995;**346**:1124–7. [https://doi.org/10.1016/S0140-6736\(95\)91799-3](https://doi.org/10.1016/S0140-6736(95)91799-3)
9. Melenhorst J, Koch SM, Uludag O, van Gemert WG, Baeten CG. Sacral neuromodulation in patients with faecal incontinence: results of the first 100 permanent implantations. *Colorectal Dis* 2007;**9**:725–30. <https://doi.org/10.1111/j.1463-1318.2007.01241.x>
10. Tjandra JJ, Lim JF, Matzel K. Sacral nerve stimulation: an emerging treatment for faecal incontinence. *ANZ J Surg* 2004;**74**:1098–106. <https://doi.org/10.1111/j.1445-1433.2004.03259.x>
11. Altomare DF, Ratto C, Ganio E, Lolli P, Masin A, Villani RD. Long-term outcome of sacral nerve stimulation for fecal incontinence. *Dis Colon Rectum* 2009;**52**:11–17. <https://doi.org/10.1007/DCR.0b013e3181974444>
12. Dudding TC, Meng Lee E, Faiz O, Parés D, Vaizey CJ, McGuire A, Kamm MA. Economic evaluation of sacral nerve stimulation for faecal incontinence. *Br J Surg* 2008;**95**:1155–63. <https://doi.org/10.1002/bjs.6237>
13. Hetzer FH, Bieler A, Hahnloser D, Löhlein F, Clavien PA, Demartines N. Outcome and cost analysis of sacral nerve stimulation for faecal incontinence. *Br J Surg* 2006;**93**:1411–17. <https://doi.org/10.1002/bjs.5491>
14. National Institute for Health and Care Excellence (NICE). *Interventional Procedure Guidance 99: Sacral Nerve Stimulation for Faecal Incontinence*. 2004. URL: www.nice.org.uk/guidance/ipg99 (accessed 7 February 2021).
15. Matzel KE. Sacral nerve stimulation for faecal incontinence: its role in the treatment algorithm. *Colorectal Dis* 2011;**13**(Suppl. 2):10–14. <https://doi.org/10.1111/j.1463-1318.2010.02519.x>

REFERENCES

16. Barussaud ML, Mantoo S, Wyart V, Meurette G, Lehur PA. The magnetic anal sphincter in faecal incontinence: is initial success sustained over time? *Colorectal Dis* 2013;**15**:1499–503. <https://doi.org/10.1111/codi.12423>
17. Wong MT, Meurette G, Stangherlin P, Lehur PA. The magnetic anal sphincter versus the artificial bowel sphincter: a comparison of 2 treatments for fecal incontinence. *Dis Colon Rectum* 2011;**54**:773–9. <https://doi.org/10.1007/DCR.0b013e3182182689>
18. Gray M. *How to Get Better Value Health Care*. Witney: Alden Press; 2007. <https://doi.org/10.1177/014107680710001019>
19. National Institute for Health Research Horizon Scanning Centre. *FENIX® Contience Restoration System for Severe Chronic Faecal Incontinence*. URL: www.hsc.nihr.ac.uk/topics/fenix-contience-restoration-system-for-severe-chr/
20. NIHR Horizon Scanning Centre. *FENIX® Contience Restoration System for Severe Chronic Faecal Incontinence*. 2012. URL: <http://io.nihr.ac.uk/wp-content/uploads/migrated/2220.838f0fdf.FENIXforfaecalincontinenceFINAL.pdf> (accessed 3 February 2020).
21. Jorge JM, Wexner SD. Etiology and management of fecal incontinence. *Dis Colon Rectum* 1993;**36**:77–97. <https://doi.org/10.1007/BF02050307>
22. Renzi A, Brillantino A, Di Sarno G, d'Aniello F. Five-item score for obstructed defecation syndrome: study of validation. *Surg Innov* 2013;**20**:119–25. <https://doi.org/10.1177/1553350612446354>
23. Rockwood TH, Church JM, Fleshman JW, Kane RL, Mavrantonis C, Thorson AG, et al. Fecal incontinence quality of life scale: quality of life instrument for patients with fecal incontinence. *Dis Colon Rectum* 2000;**43**:9–16. <https://doi.org/10.1007/BF02237236>
24. Ware J, Kosinski M, Keller SD. A 12-item short-form health survey: construction of scales and preliminary tests of reliability and validity. *Med Care* 1996;**34**:220–33. <https://doi.org/10.1097/00005650-199603000-00003>
25. The EuroQol Group. EuroQol – a new facility for the measurement of health-related quality of life. *Health Policy* 1990;**16**:199–208. [https://doi.org/10.1016/0168-8510\(90\)90421-9](https://doi.org/10.1016/0168-8510(90)90421-9)
26. National Institute for Health and Care Excellence. *Guide to the Methods of Technology Appraisal 2013: Process and Methods (PMG9)*. 2013. URL: <https://nice.org.uk/process/pmg9/chapter/foreword> (accessed 3 April 2019).
27. Brooks RG, Jendteg S, Lindgren B, Persson U, Björk S. EuroQol: health-related quality of life measurement. Results of the Swedish questionnaire exercise. *Health Policy* 1991;**18**:37–48. [https://doi.org/10.1016/0168-8510\(91\)90142-K](https://doi.org/10.1016/0168-8510(91)90142-K)
28. Kind P, Dolan P. The effect of past and present illness experience on the valuations of health states. *Med Care* 1995;**33**(Suppl. 4):A255–63.
29. Brazier JE, Roberts J, Deverill M. The estimation of a preference based measure of health from the SF-36. *J Health Econ* 2002;**21**:271–92. [https://doi.org/10.1016/S0167-6296\(01\)00130-8](https://doi.org/10.1016/S0167-6296(01)00130-8)
30. Devlin NJ, Shah KK, Feng Y, Mulhern B, van Hout B. Valuing health-related quality of life: an EQ-5D-5L value set for England. *Health Econ* 2018;**27**:7–22. <https://doi.org/10.1002/hec.3564>
31. van Hout B, Janssen MF, Feng YS, Kohlmann T, Busschbach J, Golicki D, et al. Interim scoring for the EQ-5D-5L: mapping the EQ-5D-5L to EQ-5D-3L value sets. *Value Health* 2012;**15**:708–15. <https://doi.org/10.1016/j.jval.2012.02.008>
32. Hounsome N, Roukas C. Cost-effectiveness of sacral nerve stimulation and percutaneous tibial nerve stimulation for faecal incontinence. *Therap Adv Gastroenterol* 2018;**11**:1756284818802562. <https://doi.org/10.1177/1756284818802562>

33. NHS Improvement. *NHS Reference Costs 2017–2018*. URL: <https://improvement.nhs.uk/resources/reference-costs/#rc1718> (accessed 31 May 2019).
34. Fiebig DG. Seemingly Unrelated Regression. In Baltagi B, editor. *A Companion to Theoretical Econometrics*. Malden, MA: Blackwell Publishing Ltd; 2001. pp. 101–21. <https://doi.org/10.1002/9780470996249.ch6>
35. Willan AR, Briggs AH, Hoch JS. Regression methods for covariate adjustment and subgroup analysis for non-censored cost-effectiveness data. *Health Econ* 2004;**13**:461–75. <https://doi.org/10.1002/hec.843>
36. Fenwick E, O'Brien BJ, Briggs A. Cost-effectiveness acceptability curves – facts, fallacies and frequently asked questions. *Health Econ* 2004;**13**:405–15. <https://doi.org/10.1002/hec.903>
37. van Wunnik BP, Visschers RG, van Asselt AD, Baeten CG. Cost-effectiveness analysis of sacral neuromodulation for faecal incontinence in The Netherlands. *Colorectal Dis* 2012;**14**:e807–14. <https://doi.org/10.1111/codi.12002>
38. Office for National Statistics (ONS). *National Life Tables: UK*. 2020. URL: <https://ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/lifeexpectancies/datasets/nationallifetablesunitedkingdomreferencetables> (accessed 3 February 2020).
39. NHS England. *Clinical Commissioning Policy: Sacral Nerve Stimulation (SNS) for Faecal Incontinence (Adult)*. June 2013. Reference NHSE A08/P/b. URL: www.england.nhs.uk/wp-content/uploads/2013/08/a08-p-b.pdf (accessed 26 April 2019).
40. National Institute for Health and Care Excellence. *Axonics Sacral Neuromodulation System for Overactive Bladder and Faecal Incontinence. Medtech Innovation Briefing [MIB164]*. December 2018. URL: www.nice.org.uk/advice/mib164/chapter/The-technology (accessed 26 April 2019).
41. Curtis LA and Burns A. *Unit Costs of Health and Social Care 2018*. Canterbury: PSSRU, University of Kent; 2018. URL: <https://doi.org/10.22024/UniKent/01.02.70995> (accessed 29 May 2019).
42. Joint Formulary Committee. *British National Formulary* (online). London: BMJ Group and Pharmaceutical Press. URL: www.medicinescomplete.com (accessed 31 May 2019).
43. Wilson EC. A practical guide to value of information analysis. *Pharmacoeconomics* 2015;**33**:105–21. <https://doi.org/10.1007/s40273-014-0219-x>
44. Altomare DF, Giuratrabocchetta S, Knowles CH, Muñoz Duyos A, Robert-Yap J, Matzel KE, European SNS Outcome Study Group. Long-term outcomes of sacral nerve stimulation for faecal incontinence. *Br J Surg* 2015;**102**:407–15. <https://doi.org/10.1002/bjs.9740>
45. Thin NN, Horrocks EJ, Hotouras A, Palit S, Thaha MA, Chan CL, *et al*. Systematic review of the clinical effectiveness of neuromodulation in the treatment of faecal incontinence. *Br J Surg* 2013;**100**:1430–47. <https://doi.org/10.1002/bjs.9226>
46. Pakravan F, Helmes C. Magnetic anal sphincter augmentation in patients with severe fecal incontinence. *Dis Colon Rectum* 2015;**58**:109–14. <https://doi.org/10.1097/DCR.000000000000263>
47. Sugrue J, Lehur PA, Madoff RD, McNevin S, Buntzen S, Laurberg S, Mellgren A. Long-term experience of magnetic anal sphincter augmentation in patients with fecal incontinence. *Dis Colon Rectum* 2017;**60**:87–95. <https://doi.org/10.1097/DCR.0000000000000709>
48. Duelund-Jakobsen J, Lehur PA, Lundby L, Wyart V, Laurberg S, Buntzen S. Sacral nerve stimulation for faecal incontinence – efficacy confirmed from a two-centre prospectively maintained database. *Int J Colorectal Dis* 2016;**31**:421–8. <https://doi.org/10.1007/s00384-015-2411-7>
49. Irwin GW, Dasari BV, Irwin R, Johnston D, Khosraviani K. Outcomes of sacral nerve stimulation for faecal incontinence in Northern Ireland. *Ulster Med J* 2017;**86**:20–4.

REFERENCES

50. Moya P, Arroyo A, Lacueva J, Candela F, Soriano-Irigaray L, López A, *et al.* Sacral nerve stimulation in the treatment of severe faecal incontinence: long-term clinical, manometric and quality of life results. *Tech Coloproctol* 2014;**18**:179–85. <https://doi.org/10.1007/s10151-013-1022-y>
51. Indinnimeo M, Ratto C, Moschella CM, Fiore A, Brosa M, Giardina S. Sacral neuromodulation for the treatment of fecal incontinence: analysis of cost-effectiveness. *Dis Colon Rectum* 2010;**53**:1661–9. <https://doi.org/10.1007/DCR.0b013e3181f46309>
52. Bulsei J, Lehur P, Durand-Zaleski I and the MOS STIC Study Group, Health Economics and Health Policy Research Unit, Assistance Publique – Hôpitaux de Paris, Paris, France, University Hospital of Nantes, Nantes, France, Colorectal Units, France. *A Comparison of Magnetic Anal Sphincter and Sacral Neuromodulation for Fecal Incontinence: the MOS-STIC Study Cost-effectiveness Results*. European Society of Coloproctology. September 2018. URL: <https://app.oxfordabstracts.com/events/472/program-app/submission/51805> (accessed 13 November 2019).
53. National Institute for Health Research. *SUBSoNIC Clinical Trial*. URL: www.qmul.ac.uk/pctu/about-us/clinical-strengths-and-studies/our-studies/subsonic/ (accessed 19 May 2019).
54. National Institute for Health and Care Excellence (NICE). *Peristeen Transanal Irrigation System for Managing Bowel Dysfunction*. Medical Technologies Guidance [MTG36]. London: NICE; 2018. URL: www.nice.org.uk/guidance/mtg36 (accessed 10 February 2021).
55. National Institute for Health and Care Excellence (NICE). *Infliximab, Adalimumab and Golimumab for Treating Moderately to Severely Active Ulcerative Colitis After the Failure of Conventional Therapy*. Technology Appraisal Guidance [TA329]. London: NICE; 2015. URL: www.nice.org.uk/guidance/ta329/resources/ (accessed 10 February 2021).
56. Curtis LA, Burns A. *Unit Costs of Health and Social Care 2017*. Canterbury: PSSRU, University of Kent; 2017. URL: www.pssru.ac.uk/project-pages/unit-costs/unit-costs-2017/ (accessed 29 May 2019).
57. Curtis LA, Burns A. *Unit Costs of Health and Social Care 2015*. Canterbury: PSSRU, University of Kent; 2015. URL: www.pssru.ac.uk/project-pages/unit-costs/unit-costs-2015/ (accessed 29 May 2019).

Appendix 1

Summaries of components of the 'success' end point at 6, 12 and 18 months post randomisation

TABLE 30 Success defined by device in use and $\geq 50\%$ improvement in CCIS

Success	FENIX MSA, n (%)	SNS, n (%)	Total, n (%)
6 months post randomisation			
Unsuccessful	33 (86.8)	18 (100.0)	51 (91.1)
Successful	5 (13.2)	0 (0.0)	5 (8.9)
Total	38 (100)	18 (100)	56 (100)
12 months post randomisation			
Unsuccessful	35 (87.5)	26 (96.3)	61 (91.0)
Successful	5 (12.5)	1 (3.7)	6 (9.0)
Total	40 (100)	27 (100)	67 (100)
18 months post randomisation			
Unsuccessful	35 (85.4)	35 (89.7)	70 (87.5)
Successful	6 (14.6)	4 (10.3)	10 (12.5)
Total	41 (100)	39 (100)	80 (100)

TABLE 31 Success defined by $\geq 50\%$ improvement in CCIS

$\geq 50\%$ improvement in CCIS?	FENIX MSA, n (%)	SNS, n (%)	Total, n (%)
6 months post randomisation			
No	33 (86.8)	18 (100.0)	51 (91.1)
Yes	5 (13.2)	0 (0.0)	5 (8.9)
Total	38 (100)	18 (100)	56 (100)
12 months post randomisation			
No	35 (87.5)	25 (92.6)	60 (89.6)
Yes	5 (12.5)	2 (7.4)	7 (10.4)
Total	40 (100)	27 (100)	67 (100)
18 months post randomisation			
No	34 (82.9)	34 (87.2)	68 (85.0)
Yes	7 (17.1)	5 (12.8)	12 (15.0)
Total	41 (100)	39 (100)	80 (100)

TABLE 32 Success defined by device in use

Device in use?	FENIX MSA, n (%)	SNS, n (%)	Total, n (%)
6 months post randomisation			
Yes	27 (71.1)	5 (27.8)	32 (57.1)
No	11 (28.9)	13 (72.2)	24 (42.9)
Total	38 (100)	18 (100)	56 (100)
12 months post randomisation			
Yes	25 (62.5)	11 (40.7)	36 (53.7)
No	15 (37.5)	16 (59.3)	31 (46.3)
Total	40 (100)	27 (100)	67 (100)
18 months post randomisation			
Yes	21 (51.2)	16 (41.0)	37 (46.3)
No	20 (48.8)	23 (59.0)	43 (53.8)
Total	41 (100)	39 (100)	80 (100)

TABLE 33 Cross-tabulation of the components

Improvement in continence	Device in use?		Total, n (%)
	No, n (%)	Yes, n (%)	
FENIX MSA (6 months post randomisation)			
≥ 50% improvement in CCIS?			
No	11 (100.0)	22 (81.5)	33 (86.8)
Yes	0 (0.0)	5 (18.5)	5 (13.2)
Total	11 (100)	27 (100)	38 (100)
SNS (6 months post randomisation)			
≥ 50% improvement in CCIS?			
No	13 (100.0)	5 (100.0)	18 (100.0)
Yes	0 (0.0)	0 (0.0)	0 (0.0)
Total	13 (100)	5 (100)	18 (100)
FENIX MSA (12 months post randomisation)			
≥ 50% improvement in CCIS?			
No	15 (100.0)	20 (80.0)	35 (87.5)
Yes	0 (0.0)	5 (20.0)	5 (12.5)
Total	15 (100)	25 (100)	40 (100)
SNS (12 months post randomisation)			
≥ 50% improvement in CCIS?			
No	15 (93.8)	10 (90.9)	25 (92.6)
Yes	1 (6.3)	1 (9.1)	2 (7.4)
Total	16 (100)	11 (100)	27 (100)

TABLE 33 Cross-tabulation of the components (continued)

Improvement in continence	Device in use?		
	No, n (%)	Yes, n (%)	Total, n (%)
FENIX MSA (18 months post randomisation)			
CCIS improvement?			
No	19 (95.0)	15 (71.4)	34 (82.9)
Yes	1 (5.0)	6 (28.6)	7 (17.1)
Total	20 (100)	21 (100)	41 (100)
SNS (18 months post randomisation)			
CCIS improvement?			
No	22 (95.7)	12 (75.0)	34 (87.2)
Yes	1 (4.3)	4 (25.0)	5 (12.8)
Total	23 (100)	16 (100)	39 (100)

Exploratory analyses: additional covariates

Cleveland Clinic Incontinence Score

There were 96 out of 99 (97%) patients included in the analysis, meaning that they each provided their CCIS at least once at one of the baseline, 6-, 12- or 18-month time points.

The CCIS data were modelled using a three-level multilevel model, with time points clustered within patients and patients clustered within surgeons. The estimates of the fixed effects can be seen in Table 34. The random effect caused by surgeon was 0 and the random effect caused by within-patient measurements was 5.93 (SE 1.26). The residual of the random effects was 7.13 (SE 0.71).

TABLE 34 Fixed effects

Covariate	Estimate	SE	p-value	95% CI
Intercept	11.1227	1.1579	< 0.0001	8.6546 to 13.5907
FENIX MSA vs. SNS	0.1052	0.6643	0.8744	-1.2048 to 1.4152
Male vs. female	1.0012	1.6859	0.5533	-2.3234 to 4.3258
Baseline CCIS > 10 points (moderate to severe) vs. ≤ 10 points (mild to moderate)	3.4638	1.1137	0.0021	1.2677 to 5.6600
Anal sphincter defect ≤ 90° vs. no anal sphincter defect	0.5030	0.6384	0.4316	-0.7547 to 1.7624
Anal sphincter defect > 90° to < 180° vs. no anal sphincter defect	0.7415	1.1433	0.5174	-0.7558 to 1.7619
Time (months)	-0.07833	0.07518	0.2987	-1.5131 to 2.9961
Device in use	-3.0386	0.7209	< 0.0001	-4.4601 to -1.6171
Time and baseline CCIS interaction (CCIS > 10 points vs. CCIS ≤ 10 points)	0.03473	0.07793	0.6564	-0.1189 to 0.1884
Device in use and treatment interaction	-0.1304	0.8309	0.8755	-1.7688 to 1.5080

Patients who were randomised to the FENIX MSA arm have a baseline CCIS 0.11 points lower than patients randomised to SNS; however, this result is not statistically significant ($p = 0.9$). Patients having a device in use reduces the CCIS by 3.04 points, a result that is statistically significant ($p < 0.0001$); however, the interaction term is not statistically significant ($p = 0.9$). The estimated change in CCIS over time is small; there is a reduction of 0.078 points per month on average, although the CI around this estimate is large owing to the small number of patients. The baseline CCIS (≤ 10 points or > 10 points) has a significant effect on the CCIS ($p = 0.005$), as would be expected due to them using the same measure; however, the interaction between baseline CCIS and time was not significant ($p = 0.8$).

The results of the model can be seen in *Figure 11*. This figure shows the difference between two identical patients, with the only difference being the randomised treatment, and assuming they received the device after 6 months and it remained in use at 12 and 18 months post randomisation.

EuroQol-5 Dimensions, five-level version

There were 96 out of 99 (97%) patients included in the analysis, meaning that 96 patients provided a score at least once at one of the baseline, 6-, 12- or 18-month time points.

The EQ-5D-5L score was modelled using a three-level multilevel model, with time points clustered within patients and patients clustered within surgeons. The fixed effects of the model are presented in *Table 35*. The random effect caused by surgeon was 0.00 (SE 0.003) and the random effect caused by within-patient measurements was 0.04 (SE 0.007). The residual of the random effects was 0.03 (SE 0.003). The model did not show a significant difference in the score between treatment arms ($p = 0.76$) or over time ($p = 0.18$). The device being in use has a statistically significant difference of 0.13 ($p = 0.004$), meaning that patients with their device in use have an EQ-5D-5L score 0.13 points higher than patients without the device in use, and this difference is the same for patients randomised to both treatment arms, as can be seen by the non-significance of the device in use and treatment interaction ($p = 0.2$). There is no indication that there is a clustering effect caused by the randomising surgeon.

The results of the model can be seen in *Figure 12*, which presents estimates and 95% CIs for the EQ-5D-5L score at 6, 12 and 18 months post randomisation for two patients who are identical except for their randomised treatment, and assuming that they received the device after 6 months and it remained in use at 12 and 18 months post randomisation.




FIGURE 11 Cleveland Clinic Incontinence Score model results.

TABLE 35 Fixed effects of EQ-5D-5L model

Covariate	Estimate	SE	p-value	95% CI
EQ-5D-5L score model				
Intercept	0.7408	0.07531	< 0.0001	0.5803 to 0.9013
FENIX MSA vs. SNS	0.01453	0.04771	0.7610	-0.07953 to 0.1086
Male vs. female	0.03300	0.1249	0.7918	-0.2132 to 0.2792
Baseline CCIS > 10 points (moderate to severe) vs. ≤ 10 points (mild to moderate)	-0.1180	0.06974	0.0922	-0.2555 to 0.01950
Anal sphincter defect ≤ 90° vs. no anal sphincter defect	0.01844	0.04755	0.6986	-0.07530 to 0.1122
Anal sphincter defect > 90° to < 180° vs. no anal sphincter defect	-0.02196	0.08345	0.7927	-0.1865 to 0.1426
Time (months)	-0.00250	0.001854	0.1785	-0.00616 to 0.001152
Device in use	0.1300	0.04494	0.0042	0.04144 to 0.2186
Device in use and treatment interaction	-0.06466	0.05166	0.2121	-0.1665 to 0.03718
EQ-5D-5L VAS model				
Intercept	81.1491	6.4029	< 0.0001	67.5016 to 94.7967
FENIX MSA vs. SNS	-4.3211	3.7654	0.2525	-11.7444 to 3.1022
Male vs. female	-4.9907	9.9532	0.6166	-24.6127 to 14.6313
CCIS > 10 points (moderate to severe) vs. ≤ 10 points (mild to moderate)	-14.1538	6.1549	0.0225	-26.2878 to -2.0199
Anal sphincter defect ≤ 90° vs. no anal sphincter defect	1.4308	3.7453	0.7028	-5.9529 to 8.8144
Anal sphincter defect > 90° to < 180° vs. no anal sphincter defect	2.7676	6.7168	0.6807	-10.4741 to 16.0093
Time (months)	-0.02979	0.3702	0.9359	-0.7596 to 0.7001
Device in use	5.8162	3.5563	0.1035	-1.1947 to 12.8272
Time and CCIS interaction (CCIS ≤ 10 points vs. CCIS > 10 points)	0.04212	0.3831	0.9126	-0.7131 to 0.7973
Device in use and treatment interaction	-4.6053	4.0771	0.2600	-12.6431 to 3.4324




FIGURE 12 EQ-5D score model results.

The EQ-5D-5L questionnaire also records the respondent's self-rated health on a vertical VAS ranging from 0–100, in which higher numbers indicate better health.

A total of 97 out of 99 (98%) patients were included in the VAS analysis, meaning that they provided a VAS score at least at one time point.

The VAS was modelled using a three-level multilevel model with time points clustered within patients and patients clustered within surgeons. The fixed effects of the model can be seen in *Table 35*. The random effect caused by surgeon was 0 and the random effect caused by within-patient measurements was 233.05 (SE 44.06). The residual of the random effects was 169.71 (SE 16.62). The randomised treatment has no significant effect on the VAS score ($p = 0.25$) and there is no significant difference over time between the two treatment arms ($p = 0.9$). The baseline CCIS minimisation factor has a statistically significant effect ($p = 0.023$); however, the effect over time is not significant, meaning that the baseline VAS score is higher for patients who have a baseline CCIS ≤ 10 points, but the VAS score changes at the same rate for patients with a baseline CCIS ≤ 10 points and > 10 points. The device being in use does not have a significant effect on the VAS score ($p = 0.1$) and the effect is not significantly different between the treatment arms ($p = 0.3$). There is no indication of a clustering effect caused by the randomising surgeon.

The results of the model can be seen in *Figure 13*, which presents estimates and 95% CIs for the EQ-5D-5L VAS score at 6, 12 and 18 months post randomisation for two patients who are identical except for their randomised treatment, and assuming that they received the device after 6 months and it remained in use at 12 and 18 months post randomisation.

The model diagnostics can be seen in *Figures 14* and *15*.

Faecal incontinence quality of life

The results of the models fitted are summarised in *Table 36*. None of the domains shows any significant difference between the treatment arms. All of the domains show a statistically significant improvement in FIQoL when the device is in use and there is no significant difference in the improvement between the two treatment arms.




FIGURE 13 EQ-5D VAS model results.




FIGURE 14 Model residuals for EQ-5D-5L score. (a) Raw residuals vs. linear predictor value; (b) histogram of the raw residuals with normal density overlay for reference; (c) normal Q-Q plot of the raw residuals; and (d) box plot of the raw residuals.




FIGURE 15 Model residuals for EQ-5D VAS. (a) Raw residuals vs. linear predictor value; (b) histogram of the raw residuals with normal density overlay for reference; (c) normal Q-Q plot of the raw residuals; and (d) box plot of the raw residuals.

TABLE 36 Lifestyle domain model results: fixed effects

Covariate	Estimate	SE	p-value	95% CI
Lifestyle domain				
Intercept	2.9449	0.3039	< 0.0001	2.2971 to 3.5927
FENIX MSA vs. SNS	0.01600	0.1824	0.9302	-0.3437 to 0.3757
Male vs. female	-0.4174	0.4905	0.3958	-1.3850 to 0.5501
CCIS > 10 points (moderate to severe) vs. ≤ 10 points (mild to moderate)	-0.8378	0.2893	0.0042	-1.4085 to -0.2672
Anal sphincter defect ≤ 90° vs. no anal sphincter defect	-0.03691	0.1849	0.8420	-0.4017 to 0.3278
Anal sphincter defect > 90° to < 180° vs. no anal sphincter defect	-0.01219	0.3427	0.9717	-0.6882 to 0.6639
Time (months)	0.000148	0.000462	0.7491	-0.00076 to 0.001059
Device in use	0.4124	0.1526	0.0075	0.1113 to 0.7135
Time and CCIS interaction (CCIS ≤ 10 points vs. CCIS > 10 points)	0.000119	0.000482	0.8051	-0.00083 to 0.001070
Device in use and treatment interaction	0.06514	0.1701	0.7022	-0.2705 to 0.4007
Coping domain				
Intercept	2.0434	0.2566	< 0.0001	1.4931 to 2.5937
FENIX MSA vs. SNS	0.004001	0.1468	0.9783	-0.2862 to 0.2942
Male vs. female	0.1544	0.3750	0.6811	-0.5869 to 0.8958
CCIS > 10 points (moderate to severe) vs. ≤ 10 points (mild to moderate)	-0.6468	0.2415	0.0083	-1.1242 to -0.1694
Anal sphincter defect ≤ 90° vs. no anal sphincter defect	0.03141	0.1497	0.8341	-0.2645 to 0.3273
Anal sphincter defect > 90° to < 180° vs. no anal sphincter defect	-0.03069	0.2533	0.9037	-0.5314 to 0.4700
Time (months)	0.000235	0.000544	0.6666	-0.00084 to 0.001311
Device in use	0.5340	0.1441	0.0003	0.2493 to 0.8188
Time and CCIS interaction (CCIS ≤ 10 points vs. CCIS > 10 points)	0.000168	0.000568	0.7685	-0.00096 to 0.001291
Device in use and treatment interaction	0.2217	0.1693	0.1926	-0.1131 to 0.5564
Depression domain				
Intercept	3.2273	0.3340	< 0.0001	2.5109 to 3.9436
FENIX MSA vs. SNS	0.05262	0.1948	0.7874	-0.3323 to 0.4376
Male vs. female	0.01222	0.4939	0.9803	-0.9638 to 0.9882
CCIS > 10 points (moderate to severe) vs. ≤ 10 points (mild to moderate)	-0.8544	0.3116	0.0069	-1.4701 to -0.2387
Anal sphincter defect ≤ 90° vs. no anal sphincter defect	-0.1388	0.2016	0.4924	-0.5372 to 0.2597
Anal sphincter defect > 90° to < 180° vs. no anal sphincter defect	-0.1428	0.3481	0.6823	-0.8307 to 0.5452

TABLE 36 Lifestyle domain model results: fixed effects (continued)

Covariate	Estimate	SE	p-value	95% CI
Time (months)	-6.32E-6	0.000557	0.9910	-0.00111 to 0.001095
Device in use	0.4244	0.1617	0.0096	0.1048 to 0.7439
Time and CCIS interaction (CCIS ≤ 10 points vs. CCIS > 10 points)	0.000017	0.000581	0.9772	-0.00113 to 0.001165
Device in use and treatment interaction	-0.04992	0.1872	0.7901	-0.4199 to 0.3201
Embarrassment domain				
Intercept	2.3581	0.2489	< 0.0001	1.8276 to 2.8886
FENIX MSA vs. SNS	-0.01209	0.1465	0.9343	-0.3008 to 0.2766
Male vs. female	0.3032	0.3921	0.4402	-0.4697 to 1.0761
CCIS > 10 points (moderate to severe) vs. ≤ 10 points (mild to moderate)	-0.7196	0.2382	0.0067	-1.1891 to -0.2500
Anal sphincter defect ≤ 90° vs. no anal sphincter defect	0.003584	0.1475	0.9806	-0.2873 to 0.2945
Anal sphincter defect > 90° to < 180° vs. no anal sphincter defect	-0.2464	0.2634	0.3505	-0.7656 to 0.2727
Time (months)	0.000492	0.000441	0.2662	-0.00038 to 0.001361
Device in use	0.3192	0.1252	0.0115	0.07244 to 0.5659
Time and CCIS interaction (CCIS ≤ 10 points vs. CCIS > 10 points)	-0.00038	0.000457	0.4107	-0.00128 to 0.000525
Device in use and treatment interaction	0.2677	0.1433	0.0632	-0.01482 to 0.5503

In the lifestyle domain, the random effect caused by surgeon was 0 and the random effect caused by within-patient measurements was 0.61 (SE 0.10). The residual of the random effects was 0.25 (SE 0.03).

In the coping domain, the random effect caused by surgeon was 0 and the random effect caused by within-patient measurements was 0.32 (SE 0.06). The residual of the random effects was 0.22 (SE 0.03).

In the depression domain, the random effect caused by surgeon was 0 and the random effect caused by within-patient measurements was 0.60 (SE 0.11). The residual of the random effects was 0.27 (SE 0.03).

In the embarrassment domain, the random effect caused by surgeon was 0 and the random effect caused by within-patient measurements was 0.38 (SE 0.07). The residual of the random effects was 0.21 (SE 0.02).

The results of the models can be seen in *Figures 16–19*, which present estimates and 95% CIs for each of the FIQoL domains at 6, 12 and 18 months post randomisation for two patients who are identical except for their randomised treatment, and assuming that they received the device after 6 months and it remained in use at 12 and 18 months post randomisation.

The model diagnostics corresponding to models of the lifestyle, coping, depression and embarrassment domains of the FIQoL can be seen in *Figures 20–23*, respectively.




FIGURE 16 The FIQoL lifestyle model results.




FIGURE 17 The FIQoL coping model results.




FIGURE 18 The FIQoL depression model results.




FIGURE 19 The FIQoL embarrassment model results.




FIGURE 20 Model residuals for FIQoL: lifestyle. (a) Raw residuals vs. linear predictor value; (b) histogram of the raw residuals with normal density overlay for reference; (c) normal Q-Q plot of the raw residuals; and (d) box plot of the raw residuals.




FIGURE 21 Model residuals for FIQoL: coping. (a) Raw residuals vs. linear predictor value; (b) histogram of the raw residuals with normal density overlay for reference; (c) normal Q-Q plot of the raw residuals; and (d) box plot of the raw residuals.




FIGURE 22 Model residuals for FIQoL: depression. (a) Raw residuals vs. linear predictor value; (b) histogram of the raw residuals with normal density overlay for reference; (c) normal Q-Q plot of the raw residuals; and (d) box plot of the raw residuals.




FIGURE 23 Model residuals for FIQoL: embarrassment. (a) Raw residuals vs. linear predictor value; (b) histogram of the raw residuals with normal density overlay for reference; (c) normal Q-Q plot of the raw residuals; and (d) box plot of the raw residuals.

Obstructed Defecation Score

The results of the model fitted to the ODS data can be seen in *Table 37*. There is no significant difference in the ODS between the two treatment arms ($p = 0.64$). Whether or not the patient has the device in use does not have a statistically significant difference on the ODS ($p = 0.58$).

The random effect caused by surgeon was 0 and the random effect caused by within patient measurements was 8.63 (SE 1.73). The residual of the random effects was 8.08 (SE 0.83).

TABLE 37 Obstructed Defecation Score model results: fixed effects

Covariate	Estimate	SE	p-value	95% CI
Intercept	5.2302	1.2865	0.0010	2.4882 to 7.9723
FENIX MSA vs. SNS	1.1086	0.7205	0.1256	-0.3130 to 2.5301
Male vs. female	0.3308	2.0760	0.8736	-3.7648 to 4.4264
CCIS > 10 points (moderate to severe) vs. ≤ 10 points (mild to moderate)	1.8101	1.2449	0.1476	-0.6458 to 4.2661
Anal sphincter defect ≤ 90° vs. no anal sphincter defect	0.5190	0.7554	0.4929	-0.9713 to 2.0092
Anal sphincter defect > 90° to < 180° vs. no anal sphincter defect	-1.0357	1.3198	0.4336	-3.6394 to 1.5680
Time (months)	-0.00207	0.002504	0.4098	-0.00701 to 0.002871
Device in use	0.3027	0.5458	0.5798	-0.7740 to 1.3795
Time and CCIS interaction (CCIS ≤ 10 points vs. CCIS > 10 points)	0.002587	0.002631	0.3267	-0.00260 to 0.007777

The results of the model can be seen in *Figure 24*, which presents estimates and 95% CIs for each of the ODSs at 6, 12 and 18 months post randomisation for two patients who are identical except for their randomised treatment, and assuming that they received the device after 6 months and it remained in use at 12 and 18 months post randomisation.

The model diagnostics can be seen in *Figure 25*.




FIGURE 24 Obstructed Defecation Score model results.




FIGURE 25 Model residuals for ODS. (a) Raw residuals vs. linear predictor value; (b) histogram of the raw residuals with normal density overlay for reference; (c) normal Q-Q plot of the raw residuals; and (d) box plot of the raw residuals.

Short Form questionnaire-12 items version 2

The fixed effects of the multilevel model fitted to the data can be seen in *Table 38*. Both models show that there is no difference between the PCS/MCS for patients randomised to the two treatment arms. The PCS model shows a statistically significant increase in score of 3.07 if the patient has the device in use ($p = 0.04$), and this difference is not significantly different for the treatment arms ($p = 0.6$). The MCS model does not show any statistically significant difference in the score if the patient has the device in use. The MCS model shows a significant difference between the scores for patients who have mild to moderate CCIS versus moderate to severe CCIS ($p = 0.004$); however, this effect is not significantly different over time ($p = 0.48$).

For the PCS model, the random effect caused by surgeon was 2.45 (SE 5.84) and the random effect caused by within-patient measurements was 61.14 (SE 11.45). The residual of the random effects was 30.52 (SE 2.93).

TABLE 38 Fixed effect estimates of PCS model

Covariate	Estimate	SE	p-value	95% CI
PCS model				
Intercept	51.5532	3.0005	< 0.0001	45.1578 to 57.9486
FENIX MSA vs. SNS	-1.8281	1.8383	0.3211	-5.4514 to 1.7953
Male vs. female	1.6534	5.0070	0.7416	-8.2158 to 11.5225
CCIS > 10 points (moderate to severe) vs. ≤ 10 points (mild to moderate)	-4.8303	2.7803	0.0838	-10.3104 to 0.6498
Anal sphincter defect ≤ 90° vs. no anal sphincter defect	-0.7844	1.8763	0.6763	-4.4827 to 2.9139
Anal sphincter defect > 90° to < 180° vs. no anal sphincter defect	-1.4812	3.3644	0.6602	-8.1126 to 5.1502
Time (months)	-0.1045	0.05933	0.0797	-0.2214 to 0.01247
Device in Use	3.0712	1.4691	0.0377	0.1756 to 5.9668
Device in use and treatment interaction	-0.9068	1.7147	0.5975	-4.2866 to 2.4729
MCS model				
Intercept	51.4858	3.7708	< 0.0001	43.4485 to 59.5231
FENIX MSA vs. SNS	-1.6192	2.1049	0.4426	-5.7680 to 2.5295
Male vs. female	-1.5167	5.8981	0.7973	-13.1419 to 10.1085
Baseline CCIS > 10 points (moderate to severe) vs. ≤ 10 points (mild to moderate)	-10.7417	3.6346	0.0035	-17.9056 to -3.5778
Anal sphincter defect ≤ 90° vs. no anal sphincter defect	-0.4134	2.2128	0.8520	-4.7749 to 3.9481
Anal sphincter defect > 90° to < 180° vs. no anal sphincter defect	2.1652	3.9816	0.5871	-5.6826 to 10.0130
Time (months)	-0.1289	0.2084	0.5368	-0.5397 to 0.2818
Device in use	2.4745	1.3824	0.0749	-0.2503 to 5.1993
Time and baseline CCIS interaction	0.1546	0.2168	0.4764	-0.2726 to 0.5819

For the PCS model, the random effect caused by surgeon was 0 and the random effect caused by within-patient measurements was 82.30 (SE 15.16). The residual of the random effects was 58.73 (SE 5.62).

The results of the models can be seen in *Figures 26 and 27*, which present estimates and 95% CIs for each of the PCS and MCS components of the SF-12 at 6, 12 and 18 months post randomisation for two patients who are identical except for their randomised treatment, and assuming that they received the device after 6 months and it remained in use at 12 and 18 months post randomisation.

The model diagnostics can be seen in *Figures 28 and 29*.




FIGURE 26 Physical Component Summary model results.




FIGURE 27 Mental Component Summary model results.




FIGURE 28 Model residuals for SF-12: PCS. (a) Raw residuals vs. linear predictor value; (b) histogram of the raw residuals with normal density overlay for reference; (c) normal Q-Q plot of the raw residuals; and (d) box plot of the raw residuals.




FIGURE 29 Model residuals for SF-12: MCS. (a) Raw residuals vs. linear predictor value; (b) histogram of the raw residuals with normal density overlay for reference; (c) normal Q-Q plot of the raw residuals; and (d) box plot of the raw residuals.

Model diagnostics




FIGURE 30 Empirical probability plot: primary end point.

The observations that lie furthest from the line are those patients who have a higher probability of a successful outcome owing to the stratification factors entered at randomisation.




FIGURE 31 Delta-betas: primary end point.




FIGURE 32 Empirical probability plot: postoperative complications.




FIGURE 33 Delta-betas: postoperative complications.

The observations with the greatest delta-betas are FENIX patients who did not have a complication, where the operation was performed by a surgeon that had a slightly higher probability of a complication occurring.




FIGURE 34 Model residuals for CCIS end point. (a) Raw residuals vs. linear predictor value; (b) histogram of the raw residuals with normal density overlay for reference; (c) normal Q-Q plot of the raw residuals; and (d) box plot of the raw residuals.

Health economics

TABLE 39 Complication costs (secondary care)

Item	Cost (2018/19 prices)	Source and assumptions	Inflated from year
Postoperative complications: listed			
Cardiorespiratory: serious	£1056	Non-elective short stay (DZ19 K): other respiratory disorders with single intervention, with CC score 0–4 (NHS Reference Costs 2017–2018 ³³)	N/A
Urinary retention: serious	£359	Non-elective short stay (LB16 K): urinary incontinence or other urinary problems, without interventions, with CC score 0–1 (NHS Reference Costs 2017–2018 ³³)	N/A
Neurological pain: serious	£358	Non-elective short stay (WH08B): unspecified pain with CC score 0 (NHS Reference Costs 2017–2018 ³³)	N/A
Bleeding/wound haematoma: serious	£1069	Non-elective short stay (FD03E): gastrointestinal bleed with single intervention, with CC score 0–4 (NHS Reference Costs 2017–2018 ³³)	N/A
Wound infection: serious	£967	Non-elective short stay (WH07D): infection or other complications of procedures, with single interventions, with CC score 0–1 (NHS Reference Costs 2017–2018 ³³)	N/A

TABLE 39 Complication costs (secondary care) (continued)

Item	Cost (2018/19 prices)	Source and assumptions	Inflated from year
Implant infection: serious	£967	Non-elective short stay (WH07D): infection or other complications of procedures, with single interventions, with CC score 0–1 (NHS Reference Costs 2017–2018 ³³)	N/A
Lead migration/fragmentation: serious or not serious	£1100	Minor anal procedures, ≥ 19 years, day case (FF42 A) (NHS Reference Costs 2017–2018 ³³)	N/A
Pain at battery site (permanent SNS) due to non-infective cause (e.g. battery rotation): serious or not serious	£1100	Minor anal procedures, ≥ 19 years, day case (FF42 A) (NHS Reference Costs 2017–2018 ³³)	N/A
Lack or loss of efficiency: serious or not serious	N/A	Assumed cost to be explant, which is included separately	N/A
Transient anal/rectal pain: serious	£585	Minimal anal procedure, day case (FF43Z) (NHS Reference Costs 2017–2018 ³³)	N/A
Device failure/separation: serious or not serious	£1100	Minor Anal Procedures, ≥ 19 years, day case (FF42 A) (NHS Reference Costs 2017–2018 ³³)	N/A
Device erosion: serious or not serious	£1819	Assume explant occurs and same cost as insertion: insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³)	N/A
Device explant/reoperation: serious or not serious	£1819	Assume explant occurs and same cost as insertion: insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³)	N/A
Worsening constipation/obstructive defaecation: serious	£585	Minimal anal procedure, day case (FF43Z) (NHS Reference Costs 2017–2018 ³³)	N/A
Other serious complications			
Wound still not healed following explant, fistula	£967	Non-elective short stay (WH07D): infection or other complications of procedures, with single interventions, with CC score 0–1 (NHS Reference Costs 2017–2018 ³³)	N/A
Wound dehiscence	£967	Non-elective short stay (WH07D): infection or other complications of procedures, with single interventions, with CC score 0–1 (NHS Reference Costs 2017–2018 ³³)	N/A
Covering stoma	£1093	Minor anal procedures, ≥ 19 years, non-elective short stay (FF42 A) (NHS Reference Costs 2017–2018 ³³)	N/A
Presented with constipation	£585	Minimal anal procedure, day case (FF43Z) (NHS Reference Costs 2017–2018 ³³)	N/A
All non-serious complications (listed and other)			
All other non-serious complications	£105	Colorectal surgery, consultant-led, non-admitted face-to-face attendance, follow-up (WF01 A) (NHS Reference Costs 2017–2018 ³³)	N/A
CC, complication and comorbidity; N/A, not applicable.			

TABLE 40 Further treatment/surgery costs (secondary care)

Item	Cost (2018/19 prices)	Source and assumptions	Inflated from year
Explants			
Explant of device: temporary or permanent – SNS	£1819	Insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³)	N/A
Explant of device: temporary or permanent – FENIX	£1819	Insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³) (Assumed same as SNS)	N/A
Further surgery and treatments			
Biofeedback therapy	£345	Initial consultation (£75) and three subsequent 30-minute sessions (£90 each) (BrainTrain UK; URL: www.braintrainuk.com/about-us/prices/ ; last accessed 20 February 2021)	
Colostomy formed	£13,851	Colectomy £9218 complex large intestine procedures, ≥ 19 years, with CC score 3–5 (FF31C) (NHS Reference Costs 2017–2018 ³³) Ileoanal pouch formation/reversal of ileostomy £4633 major small intestine procedures, ≥ 19 years, with CC score 0–1 (FF22D) (NHS Reference Costs 2017–2018 ³³)	
EUA of rectum and manual evacuation	£1100	'Minor anal procedure'. (NHS Reference Costs 2017–2018 ³³)	
EUA of rectum and evacuation of haematoma	£1100	'Minor anal procedure'. (NHS Reference Costs 2017–2018 ³³)	
Insertion of temporary SNS	£3692	£680 temporary device cost (Hounsome and Roukas ³²) £1819 insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³) £1193 insertion of neurostimulator electrodes for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³)	
Insertion of permanent SNS	£10,762	£7750 permanent device cost (Hounsome and Roukas ³²) £1819 insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³) £1193 Insertion of neurostimulator electrodes for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³)	
Laparoscopic insertion of pudendal nerve stimulator	£10,762	£7750 permanent device cost (Hounsome and Roukas ³²) £1819 insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³) £1193 insertion of neurostimulator electrodes for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³)	

TABLE 40 Further treatment/surgery costs (secondary care) (continued)

Item	Cost (2018/19 prices)	Source and assumptions	Inflated from year
Laparoscopic sigmoid colectomy	£13,851	Colectomy £9218 complex large intestine procedures, ≥ 19 years, with CC score 3–5 (FF31C) (NHS Reference Costs 2017–2018 ³³) Ileoanal pouch formation/reversal of ileostomy £4633 major small intestine procedures, ≥ 19 years, with CC score 0–1 (FF22D) (NHS Reference Costs 2017–2018 ³³)	
Laparoscopic ventral mesh rectopexy	£1300	Major anal procedures, ≥ 19 years, with CC score 0 (FF40B) (NHS Reference Costs 2017–2018 ³³)	
Physiotherapy	£57	Physiotherapist, adult, one to one, allied health professionals, community health services (A08A1) (NHS Reference Costs 2017–2018 ³³)	
Posterior tibial nerve stimulation	£2579	Hounscome and Roukas ³²	
Rectal irrigation	£76.28	NICE ⁵⁴	
Removal of FENIX	£1819	Insertion of neurostimulator for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³) (Assumed same as SNS)	
Replacement of SNS wire	£1193	Insertion of neurostimulator electrodes for treatment of faecal incontinence, day case (FF47Z) (NHS Reference Costs 2017–2018 ³³)	N/A
Sigmoidoscopy	£402	Diagnostic flexible sigmoidoscopy, ≥ 19 years, day case (FE35Z) (NHS Reference Costs 2017–2018 ³³)	
Stoma	£13,851	Colectomy £9218 complex large intestine procedures, ≥ 19 years, with CC score 3–5 (FF31C) (NHS Reference Costs 2017–2018 ³³) Ileoanal pouch formation/reversal of ileostomy £4633 major small intestine procedures, ≥ 19 years, with CC score 0–1 (FF22D) (NHS Reference Costs 2017–2018 ³³)	
Annual ongoing stoma care	£2444	NICE ⁵⁵	2014–15
Wound closed up	£967	Non-elective short stay (WH07D): infection or other complications of procedures, with single interventions, with CC score 0–1 (NHS Reference Costs 2017–2018 ³³)	
Excluded as unrelated to FI and the interventions			
Bilateral foot surgery			
Bilateral left and right knee repair			
Intramedullary nailing surgery			
Left shoulder surgery			
Right submandibular gland excision			
Right total knee replacement			
CC, complexity and comorbidity; EUA, examination under anaesthetic; N/A, not applicable.			

TABLE 41 Medication costs

Item	Cost (2018/19 prices)	Source and assumptions
Abena pads	£10.40	Pack of 30, premium Abena Abri-San-7-premium. URL: www.allaboutincontinence.co.uk/incontinence-brands/abena/abena-abri-san (accessed 12 February 2021)
Adcal	£2.95	Adcal-D3 750-mg/200-unit caplets (Kyowa Kirin Ltd, Galashiels, UK), 112 tablets (BNF ⁴²)
Alendronic acid	£1.59	Alendronic acid 10-mg tablets (AAH Pharmaceuticals Ltd, Coventry, UK). 28 tablets (BNF ⁴²)
Alverine citrate	£3.60	Alverine 60-mg capsule (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Amitriptyline (10 mg)	£0.91	Amitriptyline 10-mg tablets (AAH Pharmaceuticals Ltd) 28 tablets (BNF ⁴²)
Amitriptyline (25 mg)	£0.72	Amitriptyline 25-mg tablets (AAH Pharmaceuticals Ltd) 28 tablets (BNF ⁴²)
Anal plug	£66.17	Peristeen anal plug 1450 small: 12–37 mm. URL: www.clearchemist.co.uk/peristeen-anal-pl-1450-s-12-37mm.html?gclid=EAlaQobChMIq6v-5ene4AIVE4XVCh0gXw1LEAQYASABEgKAePD_BwE (accessed 1 February 2019)
Andrews Original Salts	£5.49	250 mg. URL: www.boots.com/andrews-original-salts-250g-10007017 (accessed 1 February 2019)
Aqua flush	£33.95	URL: www.bcapharmulary.nhs.uk/2118-trans-anal-irrigation-systems (accessed 1 February 2019)
Aspirin (75 mg)	£1.13	Aspirin 75-mg tablets (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Atorvastatin (20 mg)	£0.78	Atorvastatin 20 mg (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Braltus®	£25.80	Braltus 10-mg inhalation powder capsules with Zonda inhaler (Teva UK Ltd, Castleford, UK). 30 capsules (BNF ⁴²)
Buscopan (10 mg)	£3.00	Buscopan 10-mg tablets (Sanofi, Reading, UK). Hyoscine butylbromide 10 mg. 56 tablets (BNF ⁴²)
Colpermin	£3.77	Colpermin gastro-resistant modified-release capsules (McNeil Products Ltd, High Wycombe, UK). Peppermint oil 200 µl, 20 capsules (BNF ⁴²)
Candesartan (2 mg)	£2.02	Candesartan 2-mg tablets (AAH Pharmaceuticals Ltd). Candesartan cilexetil 2 mg. 7 tablets (BNF ⁴²)
Cefalexin	£1.57	Cefalexin 250-mg tablets (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Celluvisic® dry eye drops	£4.80	Celluvisic 0.5% eye drops 0.4-ml unit dose (Allergan Ltd, Marlow, UK). Carmellose sodium 5 mg per 1 ml. 30 unit dose (BNF ⁴²)
Cetirizine	£0.81	Cetirizine 10-mg tablets. Cetirizine hydrochloride 10 mg (BNF ⁴²)
Cholestyramine	£10.76	Questran 4-g oral powder sachets (Bristol-Myers Squibb Pharmaceuticals Ltd, Uxbridge, UK). Colestyramine anhydrous 4 g (BNF ⁴²)
Cinnarizine	£5.06	Cinnarizine 15-mg tablets (AAH Pharmaceuticals Ltd). Cinnarizine 15 mg. 84 tablets (BNF ⁴²)
Ciprofloxacin	£2.11	Ciprofloxacin 100-mg tablets [Alliance Healthcare (Distribution) Ltd, Chessington, UK]. Ciprofloxacin hydrochloride 100 g. 6 tablets (BNF ⁴²)
Citalopram (20 mg)	£1.00	Citalopram 20-mg tablets (AAH Pharmaceuticals Ltd). Citalopram hydrochloride. 28 tablets (BNF ⁴²)
Clindamycin	£3.45	Clindamycin 150-mg capsules (AAH Pharmaceuticals Ltd). Clindamycin hydrochloride 150 mg. 24 capsules (BNF ⁴²)
Clonidine (25 mg)	£4.50	Clonidine 25-mg tablets (AAH Pharmaceuticals Ltd). Clonidine hydrochloride 25 mg. 112 tablets (BNF ⁴²)

TABLE 41 Medication costs (continued)

Item	Cost (2018/19 prices)	Source and assumptions
Co-amoxiclav	£1.77	Co-amoxiclav 250-mg/125-mg tablets (AAH Pharmaceuticals Ltd). Amoxicillin (as amoxicillin trihydrate) 250 mg, Clavulanic acid (as potassium clavulanate) 125 mg. 21 tablets (BNF ⁴²). Co-codamol
Co-codamol (30 mg)	£3.90	Co-codamol 30-mg/500-mg caplets (AAH Pharmaceuticals Ltd). 30 tablets (BNF ⁴²)
Codeine	£0.74	Codeine 15-mg tablets (AAH Pharmaceuticals Ltd). Codeine phosphate 15 mg. 28 tablets (BNF ⁴²)
Co-dydramol (10 mg/500 mg)	£0.71	Co-dydramol 10-mg/500-mg tablets (AAH Pharmaceuticals Ltd). 30 tablets (BNF ⁴²)
Colecalciferol (1.5 mg)	£3.65	Adcal-D3 lemon chewable tablets (Kyowa Kirin Ltd). Calcium carbonate 1.5 g, colecalciferol 400 unit. 56 tablets (BNF ⁴²)
Colestyramine	£10.76	Questran 4-g oral powder sachets (Bristol-Myers Squibb Pharmaceuticals Ltd). 50 sachets (BNF ⁴²)
Colpermin	£3.77	Colpermin gastro-resistant modified release capsules (McNeil Products Ltd). Peppermint oil 200 µl. 20 capsules (BNF ⁴²)
Coloplast anal catheters	£76.28	NICE ⁵⁴
Cream for piles	£2.49	Anusol® HC ointment (Church & Dwight UK Ltd, Folkestone, UK). 30 g (BNF ⁴²)
Peristeen irrigation	£76.28	NICE ⁵⁴
Diocetyl	£2.09	Diocetyl 100-mg capsules (UCB Pharma Ltd, Slough, UK). Docusate sodium 100 mg. 30 capsules (BNF ⁴²)
Doxycycline	£17.30	Periostat® 20-mg tablets (Alliance Pharmaceuticals Ltd). 56 tablets (BNF ⁴²)
Dulcolax®	£2.35	Dulcolax 10-mg suppositories (Sanofi). 12 suppository (BNF ⁴²)
Entrolax	£2.35	Dulcolax 10-mg suppositories (Sanofi). 12 suppository (BNF ⁴²)
Estradiol pessaries (10 mg)	£16.72	Vagifem® 10-mg vaginal tablets (Novo Nordisk Ltd, Gatwick, UK). 24 pessary (BNF ⁴²)
Femigel	£10.99	URL: www.amazon.co.uk/Australian-Bodycare-Femigel-Vaginal-Moisturiser/dp/B01LXQPO79/ref=sr_1_1?keywords=femigel%26qid=1552575258%26s=drugstore%26sr=1-1-catcorr (accessed 12 February 2021)
Fittleworth sense catheters	£1.72	Brand cost not found: assumed same as Lofric Sense. URL: www.supplychain.nhs.uk/savings/price-ranking/~/_media/Files/Price%20Ranking/Price%20Ranking%20Urology%20March%202018.ashx (accessed 1 February 2019)
Flucloxacillin	£1.00	Flucloxacillin 250-mg capsules (AAH Pharmaceuticals Ltd) 28 capsules (BNF ⁴²)
Fluoxetine (40 g)	£1.80	Fluoxetine 40-mg capsules [Alliance Healthcare (Distribution) Ltd]. 30 capsules (BNF ⁴²)
Folic acid	£2.93	Folic acid 400-µg tablets (Phoenix Healthcare Distribution Ltd, Runcorn, UK). 90 tablets (BNF ⁴²)
Furosemide (20 mg)	£2.10	Furosemide 20-mg tablets (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Fybogel	£2.73	Fybogel 3.5-g effervescent granules sachets plain SF [Reckitt Benckiser Healthcare (UK) Ltd, Slough, UK]. 30 sachets (BNF ⁴²)

continued

TABLE 41 Medication costs (continued)

Item	Cost (2018/19 prices)	Source and assumptions
GAVISCON®	£4.46	GAVISCON Advance Mint chewable tablets [Reckitt Benckiser Healthcare (UK) Ltd]. 24 tablets (BNF ⁴²)
Glycerin suppositories	£1.04	Glycerol 1-g suppositories (DE Pharmaceuticals, Prudhoe, UK). 12 suppository (BNF ⁴²)
Ibuprofen	£3.60	Ibuprofen 600-mg tablets (AAH Pharmaceuticals Ltd). 84 packets (BNF ⁴²)
ICaps®	£13.49	30 tablets URL: www.boots.com/icaps-extra-lutein-tablets-30s-10114811 (accessed 1 February 2019)
IMODIUM®	£4.21	IMODIUM plus caplets (McNeil Products Ltd). 12 tablets (BNF ⁴²)
IMODIUM	£1.17	IMODIUM 1-mg/5-ml oral solution (Janssen-Cilag Ltd, High Wycombe, UK). 100 ml (BNF ⁴²)
Infliximab	£377.00	FLIXABI® 100-mg powder for concentrate for solution for infusion vials (Biogen Idec Ltd, Maidenhead, UK). Infliximab 100 mg. 1 vial (BNF ⁴²)
Infliximab + B ₁₂ injection	£14.50	Cytamen 1000-µg/1-ml solution for injection ampoules (RPH Pharmaceuticals AB, Ashton-under-Lyne, UK). Five ampoules (BNF ⁴²)
Irrigation	£76.28	NICE ⁵⁴
Kira® Menopause relief (6.5 mg)	£10.29	URL: www.hollandandbarrett.com/shop/product/kira-menopause-relief-tablets-6-5mg-60032265 (accessed 1 February 2019)
Lactulose	£2.28	Lactulose 3.1-3.7-g/5-ml oral solution (Phoenix Healthcare Distribution Ltd). 500 ml. (BNF ⁴²)
Lansoprazole	£0.65	Lansoprazole 15-mg gastro-resistant capsules (AAH Pharmaceuticals Ltd). 28 capsules (BNF ⁴²)
Lascido®	£2.63	Macrogol compound oral powder sachets sugar free [Alliance Healthcare (Distribution) Ltd]. 20 sachets (BNF ⁴²)
Laxicol®	£2.63	Macrogol compound oral powder sachets sugar free [Alliance Healthcare (Distribution) Ltd]. 20 sachets (BNF ⁴²)
Laxido®	£2.63	Macrogol compound oral powder sachets sugar free [Alliance Healthcare (Distribution) Ltd]. 20 sachets (BNF ⁴²)
Laxiolo®	£2.63	Macrogol compound oral powder sachets sugar free [Alliance Healthcare (Distribution) Ltd]. 20 sachets (BNF ⁴²)
Laxulo®	£2.63	Macrogol compound oral powder sachets sugar free [Alliance Healthcare (Distribution) Ltd]. 20 sachets (BNF ⁴²)
Lercanidipine (10 mg)	£5.34	Lercanidipine 10-mg tablets (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Levothyroxine (100 mg)	£0.99	Levothyroxine sodium 100-µg tablets (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Lisinopril	£0.82	Lisinopril 2.5-mg tablets (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Loperamide	£2.93	Loperamide 2-mg tablets (AAH Pharmaceuticals Ltd). 30 tablets. 30 tablets (BNF ⁴²)
Lactulose	£2.28	Lactulose 3.1-3.7-g/5-ml oral solution (Phoenix Healthcare Distribution Ltd). 500 ml. (BNF ⁴²)
Macrogol	£2.63	Macrogol compound oral powder sachets sugar free [Alliance Healthcare (Distribution) Ltd]. 20 sachets (BNF ⁴²)
Magnesium hydroxide	£5.31	Magnesium hydroxide 7.45-8.35% oral suspension BP (AAH Pharmaceuticals Ltd). 500 ml (BNF ⁴²)

TABLE 41 Medication costs (continued)

Item	Cost (2018/19 prices)	Source and assumptions
MOVICOL®	£5.41	MOVICOL oral powder 13.8-g sachets lemon and lime (Forum Health Products Ltd, Redhill, UK). 20 sachets (BNF ⁴²)
Mebeverine	£6.00	Mebeverine 135-mg tablets (AAH Pharmaceuticals Ltd). 100 tablets (BNF ⁴²)
Methocarbamol	£12.65	Methocarbamol 750-mg tablets (AAH Pharmaceuticals Ltd). 100 tablets (BNF ⁴²)
Metronidazole (400 mg)	£18.00	Metronidazole (AAH Pharmaceuticals Ltd). 21 tablets (BNF ⁴²)
Mini irrigation system	£76.28	Assumed: NICE ⁵⁴
Mirabegron	£29.00	Betmiga 25-mg modified-release tablets (Astellas Pharma Ltd, Woking, UK). Mirabegron 25 mg. 30 tablets (BNF ⁴²)
Movelat	£8.39	Gel: 80 g URL: www.boots.com/movelat-relief-gel-80g-10023902 (accessed 1 February 2019)
MoviCell	£25.55	Converted from Euros. URL: www.iafstore.com/uk/promopharma/movicell-drena-plus-codp33303 (accessed 1 February 2019)
MOVICOL	£8.11	MOVICOL chocolate oral powder 13.9-g sachets (Forum Health Products Ltd). 30 sachets (BNF ⁴²)
Naproxen (500 mg)	£12.00	Naproxen 500-mg tablets (AAH Pharmaceuticals Ltd)/28 tablets (BNF ⁴²)
Nitrofurantoin (500 mg)	£15.42	Nitrofurantoin 50-mg capsules (AAH Pharmaceuticals Ltd). 30 capsules (BNF ⁴²)
Nortriptyline	£8.55	Nortriptyline 10-mg tablets (AAH Pharmaceuticals Ltd). 30 tablets (BNF ⁴²)
Octasa (400 mg)	£16.58	Octasa 400-mg MR gastro-resistant tablets (Tillotts Pharma UK Ltd, Wellingore, UK). 90 tablets (BNF ⁴²)
Omacor (1000 mg)	£6.00	Omega 3-acid-ethyl esters 1000-mg capsules (Glenmark Pharmaceuticals Europe Ltd, Harrow, UK) 28 tablets (BNF ⁴²)
Omeprazole (10–40 mg)	£5.79	Omeprazole 20-mg gastro-resistant (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
OTC laxatives	£1.00	Senna 7.5-mg tablets (AAH Pharmaceuticals Ltd). 20 tablets (BNF ⁴²)
Oxybutynin hydrochloride	£3.05	Oxybutynin 2.5-mg tablets (AAH Pharmaceuticals Ltd). 56 tablets (BNF ⁴²)
Zapain® (500 mg)	£3.85	Zapain 30-mg/500-mg capsules (Advanz Pharma, London, UK). 100 capsules (BNF ⁴²)
Pantoprazole	£0.80	Pantoprazole 20-mg gastro-resistant tablets (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Paracetamol	£1.53	Paracetamol 500-mg caplets (AAH Pharmaceuticals Ltd). 100 tablets (BNF ⁴²)
Peristeen bowel wash	£76.28	NICE ⁵⁴
Pregabalin	£3.43	Pregabalin 25-mg capsules (AAH Pharmaceuticals Ltd). 56 capsules (BNF ⁴²)
Proctosedyl Ointment	£10.34	Proctosedyl Ointment (Sanofi). 30 g (BNF ⁴²)
Procyclidine	£3.47	Procyclidine 5-mg tablets (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Questran® powder	£10.76	Questran 4-g oral powder sachets (Bristol-Myers Squibb Pharmaceuticals Ltd). 50 sachets (BNF ⁴²)
Questran	£10.76	Questran 4-g oral powder sachets (Bristol-Myers Squibb Pharmaceuticals Ltd). 50 sachets (BNF ⁴²)

continued

TABLE 41 Medication costs (continued)

Item	Cost (2018/19 prices)	Source and assumptions
Quinine	£2.04	Quinine sulfate 200-mg tablets (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Qufora® IrriSedo Mini system	£59.00	URL: www.stomacarehandbook.com/product/2470/qufora_irrisedo_mini_system (accessed 1 February 2019) (includes 15 cones)
Ranitidine	£1.50	Ranitidine 150-mg tablets (AAH Pharmaceuticals Ltd). 60 tablets (BNF ⁴²)
Replens MD™ Vaginal Moisturiser	£11.49	Replens MD Vaginal Moisturiser: 35 g. URL: www.boots.com/replens-md-vaginal-moisturiser-35g-10025232 (accessed 1 February 2019)
Senna	£1.00	Senna 7.5-mg tablets (AAH Pharmaceuticals Ltd). 20 tablets (BNF ⁴²)
Senokot	£3.23	Senokot Max Strength 15-mg tablets [Reckitt Benckiser Healthcare (UK) Ltd] (BNF ⁴²)
Sertraline	£0.76	Sertraline 50-mg tablets (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Simvastatin	£0.55	Simvastatin 10-mg tablets (AAH Pharmaceuticals Ltd) 28 tablets (BNF ⁴²)
Suppositories (generic)	£3.49	URL: www.boots.com/boots-constipation-relief-12-suppositories-10006837 (accessed 12 February 2021) (12 tablets)
Telmisartan	£6.00	Telmisartan 20-mg tablets (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Thyroxine	£1.20	Levothyroxine sodium 12.5-µg tablets (AAH Pharmaceuticals Ltd). 28 tablets (BNF ⁴²)
Trajenta	£33.26	Trajenta 5-mg tablets (Boehringer Ingelheim Ltd, Bracknell, UK). 28 tablets (BNF ⁴²)
Tramadol	£0.76	Tramadol 50-mg capsules (AAH Pharmaceuticals Ltd). 30 capsules (BNF ⁴²)
TRUSOPT® eye drops	£6.33	TRUSOPT 20-mg/-ml eye drops (Santen UK Ltd, St Albans, UK). 5 ml (BNF ⁴²)
Vitamineral Green	£98.22	500 g. URL: www.amazon.co.uk/Healthforce-Vitamineral-Green-Powder-500-Grams/dp/B001H0T4TA/ref=sr_1_fmnull_1?keywords=vitamineral±green%26qid=1552578134%26s=gateway%26sr=8-1-fknull (accessed 1 February 2019)

TABLE 42 Primary care costs

Item	Cost (2018/19 prices)	Source and assumptions	Inflated from year
GP, home visit (face to face)	£95.47	Curtis and Burns ⁵⁶ estimates for costs/minute and Curtis and Burns ⁵⁷ estimates for duration	2017
GP, surgery visit (face to face)	£37.00	Curtis and Burns ⁵⁶ estimates for a consultation duration of 9.22 minutes	2017
GP telephone/e-mail	£28.40	Curtis and Burns ⁵⁶ estimates for costs/minutes and Curtis and Burns ⁵⁷ lasting 7.1 minutes at a per patient contact time of £4. Curtis and Burns ⁵⁷ estimates for duration. The same cost is assumed for any values entered for telephone contacts in the telephone contracts in the final column for all of 'GP surgery visit', 'GP home visit', 'GP out-of-hours home visit'	2017
District nurse (face to face)	£38.45	'District nurse, adult, face to face' (<i>NHS Reference Costs 2017-2018</i> ³³) under community health services	
District nurse (telephone/e-mail)	£18.88	'District nurse, adult, none face to face' (<i>NHS Reference Costs 2017-2018</i> ³³) under community health services	
GP out of hours (face to face)	£124.06	Curtis and Burns ⁵⁶ estimates for travelling time costs/minute and National Audit Office (2014) estimates for out-of-hours consultations	2017
Practice nurse (face to face)	£14.11	Curtis and Burns ⁴¹ estimate per hour £54.60 (£42 × 1.30 ratio for direct contact time). An average consultation of 15.5 minutes (PSSRU, 2014/2015)	
Practice nurse (telephone/e-mail)	£4.26	Curtis and Burns ⁴¹ page 182 under 'Time use of community care professionals' states that GP practice nurses dedicate 5.3% of their time to telephone consultations	
Occupational physiotherapist (face to face)	£57	'Physiotherapist, adult, one to one' (<i>NHS Reference Costs 2017-2018</i> ³³)	
Occupational physiotherapist, telephone/e-mail	£33.06	'Physiotherapy, non-admitted non-face-to-face attendance, follow-up', using volume-weighted average of consultant-led and non-consultant-led costs (<i>NHS Reference Costs 2017-2018</i> ³³)	2016
Occupational therapist, face to face	£81.66	'Occupational therapist, adult, one to one' (<i>NHS Reference Costs 2017-2018</i> ³³)	2016
Occupational therapist, telephone/e-mail	£43.64	'Occupational therapy, non-admitted none face-to-face attendance, follow-up', (consultant-led and non-consultant-led reference cost identical) (<i>NHS Reference Costs 2017-2018</i> ³³)	2016
Counsellor (face to face)	£44	Curtis and Burns ⁴¹ counsellor (band 6) under 'scientific and professional staff'	2018
Counsellor (telephone/e-mail)	£23.32	Assumed same ratio face to face: telephone as occupational therapist (53%)	

TABLE 43 Other service use

Item	Cost (£, 2018/19 prices)	Source and assumptions	Inflated from year
Other support from personal social services			
Meals on wheels (frozen)	£4.65	URL: www.leeds.gov.uk/adult-social-care/help-at-home (accessed 3 April 2019). Assumed cost of £3.65 for a main course and £1.00 for dessert (uppermost values)	2018
Meals on wheels (hot)	£8.80	URL: www.leeds.gov.uk/adult-social-care/help-at-home (accessed 3 April 2019). £6.00 for the hot main meal and dessert and a cost of £2.80 for the tea	2018
Laundry services	£14.00	URL: www.laundryheap.co.uk/ (accessed 3 April 2019). £14 per 6 kg wash	2018
Home help, face to face (assumed same cost as that of cleaner, carer, home care and health and social care)	£58.54	'Health visitor, other clinical interventions' (NHS Reference Costs 2017–2018 ³³)	2017
Community and residential based services			
Nursing home/hospice stay, per day	£468.26	'Inpatient day in hospice care' (Public Health England, 2017)	2012
Convalescent care	£158.00	Per day permanent resident week 'Local authority own-provision residential care for older people (age 65+)' from Curtis and Burns ⁴¹	2017–18

TABLE 44 Descriptive statistics

Group	Number and mean	All data collected as part of the trial				Complete-case analysis			
		Baseline ^a	6 months	12 months	18 months	Baseline ^a	6 months	12 months	18 months
Utilities: EQ-5D-5L									
Total	<i>n</i>	89	80	73	63	47	47	47	47
	Mean	0.762	0.754	0.769	0.750	0.798	0.756	0.793	0.770
FENIX	<i>n</i>	44	38	37	31	24	24	24	24
	Mean	0.761	0.771	0.777	0.731	0.802	0.772	0.792	0.757
SNS	<i>n</i>	45	42	36	32	23	23	23	23
	Mean	0.763	0.738	0.760	0.768	0.794	0.740	0.794	0.783
Costs: health and social care									
Total	<i>n</i>		81	76	67	47	47	47	47
	Mean		£116.72	£126.34	£121.02		£103.95	£121.71	£142.50
FENIX	<i>n</i>		39	38	32		24	24	24
	Mean		£107.70	£125.26	£176.15		£110.45	£128.64	£203.39
SNS	<i>n</i>		42	38	35		23	23	23
	Mean		£125.10	£127.41	£70.62		£97.17	£114.48	£78.97

TABLE 44 Descriptive statistics (continued)

Group	Number and mean	All data collected as part of the trial				Complete-case analysis			
		Baseline ^a	6 months	12 months	18 months	Baseline ^a	6 months	12 months	18 months
Costs: secondary care									
Total	<i>n</i>		94	92	91		47	47	47
	Mean		£416.83	£501.89	£131.52		£716.18	£464.17	£69.36
FENIX	<i>n</i>		47	46	45		24	24	24
	Mean		£603.07	£847.81	£180.45		£954.11	£820.70	£86.13
SNS	<i>n</i>		47	46	46		23	23	23
	Mean		£230.60	£155.97	£83.66		£467.91	£92.14	£51.87
Costs: medications									
Total	<i>n</i>		81	76	67		47	47	47
	Mean		£9.75	£7.52	£10.84		£8.60	£9.46	£12.87
FENIX	<i>n</i>		39	38	32		24	24	24
	Mean		£10.75	£9.15	£20.22		£11.87	£12.77	£23.14
SNS	<i>n</i>		42	38	35		23	23	23
	Mean		£8.81	£5.89	£2.27		£5.19	£6.00	£2.15

^a Baseline costs were not used in this study.

EME
HS&DR
HTA
PGfAR
PHR

Part of the NIHR Journals Library
www.journalslibrary.nihr.ac.uk

*This report presents independent research funded by the National Institute for Health Research (NIHR).
The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the
Department of Health and Social Care*

Published by the NIHR Journals Library

