



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/182165/>

Version: Accepted Version

Conference abstract:

Teece, A, Baker, J and Smith, H (2021) Understanding How Critical Care Nurses Make The Decision To Restrain A Delirious Patient: A "Think Aloud" Study. [Conference abstract]

<https://doi.org/10.1111/nicc.12693>

© 2021 The Authors. Nursing in Critical Care © 2021 British Association of Critical Care Nurses. This is the peer reviewed version of the following article: Teece, A, Baker, J and Smith, H (2021) Understanding How Critical Care Nurses Make The Decision To Restrain A Delirious Patient: A "Think Aloud" Study. In: Nursing in Critical Care. BACCN Conference 2021, 13-14 Sep 2021 Wiley , pp. 16-17., which has been published in final form at <https://doi.org/10.1111/nicc.12693>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Oral, Primary Research

Angela Teece, University of Leeds

John Baker, University of Leeds

Helen Smith, University of Leeds

UNDERSTANDING HOW CRITICAL CARE NURSES MAKE THE DECISION TO RESTRAIN A DELIRIOUS PATIENT: A "THINK ALOUD" STUDY

Background: Patients with psychomotor agitation due to hyperactive delirium are at risk of disrupting life-sustaining therapies. Chemical or physical restraint are often cited by staff as the main method of preserving patient safety, however its efficacy is unproven (Ai et al., 2018) and seems rooted in culture and routine rather than a clear evidence base (Teece et al., 2020).

Aim: To explore the process through which critical care nurses make the decision to restrain their patient.

Method: A "Think Aloud" approach was taken. 30 critical care practitioners were invited to view audio-visual vignettes featuring simulated patient scenarios with a variety of risk inferences and levels of agitation. The participant was invited to "Think Aloud" their responses to the scenario. Reflexive thematic analysis was undertaken following transcription of the interviews. HRA and University ethics approval received.

Results: Five themes were identified: 1. Intrinsic beliefs and aptitudes and their influence on the decision to restrain; 2. The influence of handover and sharing of labels on the decision to restrain; 3. Failure to maintain a consistent approach to restraint; 4. Reduced staffing ratios can lead to restraint; 5. The tyranny of the now.

Implications for practice: Decision-making varied with participants' level of clinical experience, with established critical care nurses being most likely to apply physical or chemical restraint. Caring for patients with psychomotor agitation is an emotional and physical challenge. Newly-qualified and Advanced Practitioners were less likely to use restraint, perhaps due to reduced continued bedside exposure to agitation. Critical care nurses hold preconceptions about delirious patients which may influence their decision to apply restraint.

Decision making around restraint is complex. The ability to cope with delirium varies between nurses and it is difficult to objectively quantify the point at which restraint becomes clinically appropriate.

References

1. Ai Z-P, Gao X-L, Zhao X-L. Factors associated with unplanned extubation in the Intensive Care Unit for adult patients: a systematic review and meta-analysis. *Intensive Crit Care Nurs.* 2018;47:62-68.
2. Teece A, Baker J, Smith H. Identifying determinants for the application of physical or chemical restraint in the management of psychomotor agitation on the critical care unit. *J Clin Nurs.* 2020;29(1-2):5-19.