

This is a repository copy of *Metabarcoding reveals selective dietary responses to environmental availability in the diet of a nocturnal, aerial insectivore, the European Nightjar (Caprimulgus europaeus)*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/182479/>

Version: Accepted Version

Article:

Mitchell, Lucy J., Horsburgh, Gavin J., Dawson, Deborah A. et al. (2 more authors) (2022) Metabarcoding reveals selective dietary responses to environmental availability in the diet of a nocturnal, aerial insectivore, the European Nightjar (*Caprimulgus europaeus*). *Ibis*. pp. 60-73. ISSN 0019-1019

<https://doi.org/10.1111/ibi.13010>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **Metabarcoding reveals selective dietary responses to**
2 **environmental availability in the diet of a nocturnal, aerial**
3 **insectivore, the European Nightjar (*Caprimulgus europaeus*)**
4

5 LUCY J. MITCHELL,^{1,2*} GAVIN J. HORSBURGH,² DEBORAH A. DAWSON,²

6 KATHRYN H. MAHER² & KATHRYN E. ARNOLD¹

7 ¹*Department of Environment and Geography, University of York, Wentworth Way,*
8 *Heslington, York, YO10 5NG, UK*

9 ²*NERC Biomolecular Analysis Facility, Department of Animal and Plant Sciences,*
10 *University of Sheffield, Alfred Denny Building, Western Bank, Sheffield, S10 2TN, UK*

11 * Corresponding author: Lucy Mitchell lucyjayneryan@gmail.com; [ORCID ID: 0000-0003-](https://orcid.org/0000-0003-2162-013X)
12 [2162-013X](https://orcid.org/0000-0003-2162-013X); [Current address:](#) Department of Biological and Marine Sciences, University
13 of Hull, Cottingham Road, Hull, HU6 7RX, UK

14
15
16
17
18
19
20
21
22 Many bird species are vulnerable to environmental change, so knowledge of their diet
23 and its variation can help to understand population status and flexibility to respond to
24 change. Insectivorous species are predicted to have a flexible diet within and between
25 individuals, which can respond to naturally fluctuating prey abundance, thus allowing

26 opportunistic exploitation of available resources. We analysed the diet of a nocturnal,
27 aerial insectivore, the European Nightjar *Caprimulgus europaeus*, using high-throughput
28 metabarcoding. We quantified diet diversity and composition of 130 faecal samples from
29 nests and roosts on a northern breeding site in the UK from 2015 – 2018, and compared
30 differences among individuals and years. Although dominated by moths, diet varied
31 significantly between individual faecal samples and between years and months. Prey
32 species composition varied between years, and was more variable between samples in
33 2017, compared to other years. Faecal samples were significantly more likely to contain
34 moth species with a wingspan of >60mm and less likely to contain moth species of
35 <25mm wingspan. This indicated size-selective foraging, which also varied between
36 months and years. Diet was driven by inter-individual variation, indicating population-
37 level flexibility in prey choice. Metabarcoding provided a valuable tool in the exploration
38 of insectivorous diets, but efforts are needed to build comprehensive reference libraries,
39 in order to compile full prey species lists.

40

41 *Keywords:* avian ecology, diet, metabarcoding, faecal sampling, 16S rRNA gene, moth,
42 Lepidoptera

43 Understanding variation in the composition and diversity of diet within a bird
44 population can assist understanding of how a population or species is likely to fare in
45 response to anthropogenic change (Howells *et al.* 2017). Environmental change is
46 likely to affect certain groups of species more acutely than others; for example,
47 specialist predators reliant on ephemeral insect prey may be more sensitive (Nocera
48 *et al.* 2012, Stanton *et al.* 2017). Species that take flying insect prey on the wing
49 ('aerial insectivores') are identified globally as being at risk (Nebel *et al.* 2010,
50 Nocera *et al.* 2012) due in part to the significant decline in insect populations
51 worldwide (Hallmann *et al.* 2017, van Strien *et al.* 2019). The diet composition of
52 aerially-feeding, insectivorous birds may reflect localised spatial and temporal
53 variation in prey abundance related to habitat type (Mills *et al.* 2020) or weather
54 conditions (Imlay *et al.* 2017). Their diet may also reflect the ease of catching
55 particular prey items, their nutritional content or habitat quality (Garlapow 2007,
56 English 2009, Sharps 2013, Razeng & Watson 2015).

57 Difficulty observing feeding, and in finding and dissecting faecal matter
58 (Mumma *et al.* 2016, Nielsen *et al.* 2018) mean that there is little information on the
59 diet of adult, insectivorous birds, especially nocturnal species. Moreover, soft-bodied
60 moths and flies that are primarily taken by this feeding guild are not readily
61 identifiable in faecal matter (Razgour *et al.* 2011, Trevelline *et al.* 2016). Recently,
62 rapid development of molecular techniques such as metabarcoding, has allowed
63 researchers to acquire valuable information relevant to species conservation through
64 diet analysis (Gillet *et al.* 2015, Kress *et al.* 2015, Gerwing *et al.* 2016).
65 Metabarcoding facilitates diet analysis using faecal samples, by identifying short
66 sequences from specific genes within the DNA of prey items that remain in the faecal
67 matter. Commonly-used barcodes include the cytochrome c oxidase subunit I (COI)

68 and the 16S rRNA gene, and these can effectively distinguish between species
69 (Metzker 2010, Alberdi *et al.* 2012, da Silva *et al.* 2019) where the barcodes have
70 high interspecific but low intraspecific diversity. As well as allowing samples to be
71 processed in bulk (Taberlet *et al.* 2012), these methods are non-invasive and can be
72 more comprehensive than visual identification (Ji *et al.* 2013, De Barba *et al.* 2014,
73 Krehenwinkel *et al.* 2017), aspects that are particularly beneficial for vulnerable or
74 secretive species (Thalinger *et al.* 2018).

75 The European Nightjar *Caprimulgus europaeus* (henceforth, 'Nightjar') is a
76 nocturnal insectivore of conservation concern in the UK (Amber listed - Eaton *et al.*
77 2015, IUCN classification is least concern, BirdLife International 2016). This species
78 completes an annual migration from central Africa to the UK to breed from May until
79 August, and feeds on arthropods (Cramp 1985, Sharps 2013). Limited diet data have
80 been obtained through stomach content analysis and physical dissection of faeces
81 from European Nightjars and the Common Nighthawk *Chordeiles minor* (Sierro *et al.*
82 2001, Sharps 2013, Knight *et al.* 2018). However, these methods result in a bias
83 towards remaining hard parts of consumed arthropods such as beetle elytra, legs
84 and antennae, which means that methods such as metabarcoding that are capable
85 of identifying partially or fully digested remains are needed. Extracting sufficient DNA
86 from avian faeces is challenging, especially after degradation by exposure to sun,
87 rain and soil microbes (Deagle *et al.* 2006, Oehm *et al.* 2011, Kamenova *et al.* 2018).
88 The high concentration of uric acid (Eriksson *et al.* 2017) in conjunction with
89 environmental exposure inhibits extraction and amplification of DNA, making the
90 extraction and clean-up stages of metabarcoding, along with choice of primer
91 sequences, critical to success (Alberdi *et al.* 2017, Carneiro *et al.* 2020).

92 Here we have analysed the diet of a sample of Nightjar individuals at a key
93 breeding site in the UK. To understand variation in Nightjar diet within this population
94 we used metabarcoding techniques, methods which have had limited application to
95 the diet of this species prior to now (Evens *et al.* 2020). We used metabarcoding to
96 test the following hypotheses:

97 1) Diet composition and diversity varies among individuals (as in Alberdi *et al.*
98 2020), as individual Nightjars at this site have been shown to vary significantly in
99 their habitat preferences (Mitchell *et al.* 2020).

100 2) Females have lower diet diversity than males (da Silva *et al.* 2020), as they
101 only spend a short time foraging due to higher investment in parental care.

102 3) Diet diversity and composition varies between years. Nightjars are
103 constrained in their breeding schedule due to migration, so their short window of
104 opportunity should coincide with peaks of preferred moth prey. However, weather,
105 and thus emergence and activity of moths, can vary substantially throughout the
106 breeding season and between years (Clare *et al.* 2011, 2014).

107 4) Nightjars select large moths more often than their relative abundance in the
108 environment, due to both nutritional value (Razeng & Watson 2015), and greater
109 ease of detecting large moths in low light (Bayne & Brigham 1995).

110

111 **METHODS**

112 This study took place at the Humberhead levels National Nature Reserve (NNR;
113 South Yorkshire, UK, comprising Thorne Moor (53.636N, 0.898W) and Hatfield Moor
114 (53.545N, -0.938W). The site is classified as a degraded raised bog, with 28% of the
115 site covered by bog or marsh, 10% by standing water, and 15% mixed woodland

116 (JNCC 2015). It is also a European Special Protection Area (SPA) for its breeding
117 Nightjar population (1.9% of the UK population; JNCC 2015). The NNR has
118 undergone substantial habitat management since 2014, primarily the removal of 562
119 km² of birch woodland and an increase in wet habitat through the damming of
120 drainage channels. Both techniques will influence vegetation structure and insect
121 prey availability; selective and patchy clearance of woodland has been shown to
122 increase species richness of Lepidoptera (Summerville & Crist 2002), and should
123 also provide an increased amount of breeding space for Nightjars (Sharps 2013).
124 Use of wetland habitat (e.g. bogs, mires) by nightjars is little considered in the
125 literature explicitly as a breeding habitat, but there is some indication that they act as
126 suitable foraging areas, containing a high number of invertebrates (English *et al.*
127 2017, Evens *et al.* 2017).

128

129 **Faecal sample collection**

130 Nightjar faecal samples were collected from June to August, 2015 to 2018 (2015: $n =$
131 9, 2016: $n = 20$, 2017: $n = 35$, 2018: $n = 36$) and were obtained from previously
132 located roosts and nests, during the day. Upon approaching the nest or roost, we
133 used binoculars to view the bird in situ. If the bird was not visible, we slowly
134 approached the location until the bird flew away, whereupon the sex of the bird was
135 visible (male Nightjars possess bright white spots on primaries 3,4 and 5 whilst
136 females have smaller, beige spots). Nests and roosts of Nightjars and the size and
137 shape of their faeces are distinctive and identifiable, as they are discrete curled
138 pellets of between 3 – 8mm in width, found individually or in small clusters (see
139 image in supplementary information). Faeces were placed into sterile, 30ml screw-
140 top tubes (Elkay labs, UK) with wooden toothpicks, then labelled with sample

141 number, date, location, and the sex of the bird. Because samples were taken from
142 roosts and nests, some samples at nests may have been from juveniles, which
143 cannot be sexed. Samples from nests were therefore excluded from comparative
144 analyses between years, months and sexes. All samples were included in analyses
145 of prey size. Birds were not individually identifiable in the field, so it is possible that
146 some samples are repeats from the same bird. Roost and nest details were
147 documented and these locations are held faithfully by individuals throughout the
148 season, so we consider it unlikely that there is much duplication of individuals within
149 the dataset. Where we have repeatedly sampled the same roost, or nest, we
150 document these as likely to be from the same bird. Although defecations at roost and
151 nest sites could potentially be from more than one bird (e.g. the male and the female
152 at the nest), the male only visits the nest for short periods of time, and where we find
153 fresh defecations during the day when the female is on the nest, we associate these
154 with the female. Additionally, males were found by Berry (1979) not to roost with the
155 female at the nest, but separately, often up to 100 metres away. This ties in with
156 other information presented by Berry where strict fidelity to roost sites by individual
157 male Nightjars was found within and between seasons.

158 Fresh faecal samples, identified by their soft consistency, were targeted.
159 Where there were multiple faecal pellets at a location, we took softer, fresher pellets
160 over dry samples. Where there were only drier, older pellets, we took a sample, but
161 visited the site again in the next 2-3 days in order to find a fresher sample, both in
162 order to improve the amount of DNA extracted and to better pinpoint deposition date.
163 Samples were transferred to a -20°C freezer within four hours of collection, and then
164 to a -80°C freezer in batches every seven days, and stored there until processing. All

165 laboratory work took place at the NERC Biomolecular Analysis Facility (NBAF) within
166 the Department of Animal and Plant Sciences, at the University of Sheffield, UK.

167

168 **DNA extraction**

169 We extracted DNA from faecal samples using the QIAmp DNA Fast Mini Stool Kit
170 (Qiagen, Germany), following the human DNA extraction protocol (Supplementary
171 Material, Appendix I), adapted to suit more highly degraded DNA from bird faeces by
172 a) increasing the amount of sample (300mg vs standard 220mg), b) using the
173 inhibitex buffer provided with the kit at a higher volume than recommended (1.4ml vs
174 standard 1ml), and c) incubating samples for 10 minutes longer than recommended
175 after the lysis buffer was added to increase the number of cells broken down from
176 the sample and increase the DNA yield. We quantified DNA concentration using a
177 Fluostar Optima (BMG Labtech, Germany) and kept the DNA extracts in a -20°C
178 freezer until PCR.

179

180 **Polymerase Chain Reaction (PCR)**

181 Primer sets were selected based on their specificity and their ability to amplify short
182 fragments of degraded DNA (fragments of between 100 – 180 base pairs (bp)),
183 based on data from previous studies (Zeale *et al.* 2011, Clarke *et al.* 2014, Alberdi *et al.*
184 2018). Two commonly-used primer sets targeting the mitochondrial cytochrome
185 oxidase COI gene: the 'Uniminibar' primer set (Meusnier *et al.* 2008) and the 'ZBJ'
186 primer set (Zeale *et al.* 2011) were tested, but not used because they produced
187 inconsistent amplification across different groups of insects (also highlighted in
188 Brandon-Mong *et al.* 2015), and amplified primer dimer (Brownie *et al.* 1997), when

189 specific products failed to amplify. We then tested the Ins16S_1Short F and R primer
190 set that amplifies a 156bp fragment of the mitochondrial 16S region (Clarke *et al.*
191 2014), identified in the literature as providing wider taxonomic coverage across
192 invertebrate orders. The use of these 16S rRNA primers resulted in more consistent
193 amplification levels across all samples. A temperature-gradient PCR from 55 – 70°C
194 (Supplementary Material, Appendix I) with a random subset of faecal samples ($n =$
195 4), moth ($n = 4$), Nightjar control samples (DNA from egg albumen collected at nests,
196 $n = 4$) and negative water samples ($n = 2$). The optimal PCR annealing temperature
197 was identified as 62°C. This temperature amplified moth tissue positive control
198 samples but not Nightjar DNA. PCR products were visualised on a 1% agarose gel
199 with a 100bp ladder (Invitrogen, Thermo Fischer Scientific, UK) to verify amplification
200 success.

201

202 **Reference DNA library**

203 Although the INS16S 1Short primers provided good taxonomic coverage, there are
204 unfortunately many fewer reference sequences available with which to compare
205 sequence reads (Clarke *et al.* 2014, Tournayre *et al.* 2020). To improve the ability to
206 identify sequences from faecal samples, we created a local reference 16S sequence
207 library, representative of some of the assemblage present on the site. We extracted
208 DNA from 81 specimens of 80 species of moths and beetles collected from the moth
209 traps, to supplement those already available in the BLAST global database (Altschul
210 *et al.* 1990). Specimens were identified to species based on visual and microscopic
211 examination by LJM, using the criteria of Townsend and Waring (2014). Extractions
212 followed an ammonium acetate precipitation method (Bruford *et al.* 1998), where a
213 digestion solution, Proteinase K and ammonium acetate were added to each sample

214 of moth legs and incubated overnight, to penetrate the invertebrates' solid chitin-
215 based exterior. Extracted insect DNA was Sanger-sequenced using di-dedoxy dye
216 terminators (see Supplementary Material, Appendix I for protocol) using both the *ZBJ*
217 COI primers (Zeale *et al.* 2011) and the INS16S 1Short primers (Clarke *et al.* 2014).
218 Sequences were visualised using an ABI3730 DNA Sequencer (Applied Biosystems,
219 ThermoFischer Scientific, USA). Due to the lack of British 16S moth sequences in
220 the NCBI /EMBL /DDJB databases, we verified species identification by comparing
221 the sequenced moth 16S genes against COI sequences, which are more frequently
222 present in the BLAST database. Finally, 16S sequences from 78 invertebrate
223 species were submitted to the NCBI GenBank database (sequence accession
224 numbers: MK620910 – MK620988).

225

226 **Sequencing and bioinformatics**

227 Detailed sequencing preparation and procedures are provided in the Appendices.
228 Briefly, samples were cleaned, individually labelled, quantified and then pooled for
229 sequencing. We then used the high-powered computer clusters '*Iceberg*' and '*Sharc*'
230 at the University of Sheffield, to process and quantify the raw reads generated by the
231 Illumina Miseq Benchtop Sequencer.

232 Sequences were processed, trimmed and clustered into 'Molecular
233 Operational Taxonomic Units' (MOTUs). The threshold at which sequences are
234 clustered – i.e. their level of similarity – strongly influences how many MOTUs are
235 produced. MOTUs are taxonomic proxies (Hemprich-Bennett *et al.* 2018), and are
236 frequently interpreted as equating to species (da Silva *et al.* 2020). Depending on the
237 level of clustering used however (most commonly between 96% and 99%; Clare *et*

238 *al.* 2016), sequences may be equated to family or order (Clare *et al.* 2016,
239 Hemprich-Bennett *et al.* 2018, Gordon 2019). We clustered our MOTUs at 97%,
240 based on our knowledge of the close-relatedness of Lepidoptera species (Rytönen
241 *et al.* 2019), as well as an understanding that if degradation has taken place, as is
242 common with DNA within faecal matter, it may be better to take a more conservative
243 approach (Clare *et al.* 2016).

244 Sequences were then compared to those stored in the NCBI GenBank
245 nucleotide database (including those of the 78 potential prey species that we
246 submitted) using a Nucleotide BLAST (blastn; Altschul *et al.* 1990). Because of the
247 lack of 16S reference sequences of British Lepidoptera, some MOTUs were
248 automatically assigned to species not present in the UK or to multiple species with
249 an identical level of certainty. We therefore manually assigned such species to
250 genus or family, using the NCBI taxonomy browser. For all diversity analyses we
251 therefore used a matrix of MOTUs, rather than species and/or genera because of the
252 described discrepancy in assigning units (as in Razgour *et al.* 2011). Classifying
253 sequence reads this way allowed us to include all units identified through the
254 clustering process, whether or not we were able to assign them to a known taxa
255 (Clare *et al.* 2011, Hawlitschek *et al.* 2018, da Silva *et al.* 2020). MOTUs that had
256 been assigned to specific non-target species (*Homo sapiens*, *Sus scrofa*, *Columba*
257 *palumbus* and *Akkermansia sp*), were removed (da Silva *et al.* 2020).

258 Finally, we created matrices of final read numbers and presence/absence
259 matrices in R (v. 3.6), which were used to filter out low numbers of reads (Galan
260 2018), that are likely to represent contamination rather than true sequences.
261 Following De Barba *et al.* (2014), we used our negative control samples to set
262 thresholds for each PCR amplification, sequences with numbers below these

263 thresholds were considered as noise, and removed (see Supplementary material,
264 Appendix II for more detail). False positives can occur at multiple stages of the
265 extraction and PCR procedures (Ficedola *et al.* 2015, Zepeda-Mendoza *et al.* 2016,
266 Alberdi *et al.* 2018) and caution over such sequences is important (Froslev *et al.*
267 2017).

268 Sequencing data will be available online upon publication through the
269 European Nucleotide Archive (ENA), part of the European Molecular Biology
270 Laboratory (EMBL), as fastq files.

271

272 **Statistical analyses**

273 To examine the diet of this population of Nightjars as a whole, we calculated the
274 frequency of occurrence of sequences (FOO) – i.e. the proportion of samples in
275 which each MOTU was found. Conversion of sequence data into a more quantitative
276 estimate of diet, known as relative read abundance (RRA), has flaws (see discussion
277 in Deagle *et al.* 2019). However, the use of FOO alone can be conservative, as it
278 essentially equates one sequence that registers 10,000 reads, with another that only
279 registers 10 reads, because the only criterion is to be present (Deagle *et al.* 2019).
280 Thus, we use RRA to calculate all metrics, bearing in mind that numbers of
281 sequence reads do not perfectly translate to percentage biomass (Rytönen *et al.*
282 2019).

283 To examine variation in composition between samples, we used multivariate
284 PERMANOVAs (using ‘*adonis*’ in ‘*vegan*’). These tested for differences in the means
285 of Chao dissimilarity indices (R package ‘*vegan*’, function ‘*vegdist*’; Oksanen *et al.*
286 2019) between sex, month and year (McClenaghan *et al.* 2019a). Because it was not

287 possible to sex chicks, these were removed from the data for calculation of the
288 PERMANOVAs, as were within-sample repeats. Sex, month and year in which the
289 sample was collected, were modelled separately as they could not be included in the
290 same PERMANOVA, because there were missing data from both the sex and month
291 categories. We used Chao dissimilarity indices within the PERMANOVAs in order to
292 account for unknown species, and because they were stable in our unbalanced
293 sample size scenario (Chao *et al.* 2005, Chao & Chiu 2016).

294 PERMANOVAs are a non-parametric version of a MANOVA and are stable
295 when faced with heterogeneity within groups (Anderson 2017). We used package
296 'pairwise Adonis' (Martinez Arbizu 2020) to conduct among-year pairwise
297 comparisons. As we had some unbalanced sample sizes, we calculated
298 homogeneity of group variances (function '*betadisper*' in *vegan* (Oksanen *et al.*
299 2019)), which calculates the distance from each sample to the group centroid
300 (Anderson, 2006) to estimate compositional variance within groups, using the Chao
301 dissimilarity indices again. We calculated 95% confidence intervals for all tests using
302 Tukey's HSD.

303 For a random subset of the samples, we undertook repeat extraction, PCR
304 and sequencing. Although most variation is attributed to between sample variation
305 (Mata *et al.* 2019), it is possible that due to a large number of variables such as size
306 of the dropping, digestion time and size of the prey items (Oehm *et al.* 2011), there
307 may be variation within a faecal sample as well (Ando 2020). We compared the
308 within-sample repeats using Chao dissimilarity indices, calculated from RRA as
309 above. The repeat samples are included in the supplementary information file,
310 marked with the suffix '_r'.

311 **Prey size variation and selection by Nightjars**

312 To understand if size-selective foraging took place, we used moth traps to obtain an
313 indication of flying insect prey availability on the breeding site. We placed 15W
314 actinic light traps (made in the electronics department at the University of York –
315 details and photos within the supplementary information), at eight locations within the
316 boundary of Thorne and Hatfield Moors ($n = 16$). Moths were trapped at four sites for
317 one night in each alternate week (i.e. sites 1-4 for one night (e.g. overnight Thursday
318 - Friday) in weeks 1, 3, 5 etc., followed by sites 5-9 for the same night in weeks 2, 4,
319 6 etc.) starting at the beginning of June until the middle of August. Traps were placed
320 at each site within two hours prior to dusk and emptied the following morning within
321 one hour of dawn. Trap locations were identified as grid squares on a map of the
322 site, and the traps placed in a randomly chosen but consistent location on the ground
323 within this square. Although the same grid squares were being sampled (each
324 square was sampled $n = 4$ times), the two-week interval between sampling
325 occasions should remove some concern about repeat sampling (Truxa & Fiedler
326 2012). The sites chosen for moth trapping included all habitat types present on the
327 nature reserve. Habitat types were classified as woodland (trees with canopy >2
328 metres high), scrub (trees with canopy <2 metres high), heather, wetland (rush-
329 dominated, <50% standing water), bog (>50% standing water, *Eriophorum*
330 dominated, bare-ground (bare peat areas with minimal (< 10% vegetation cover) and
331 'off site' – all areas outwith the boundary of the NNR). Habitat structure was also
332 incorporated, so as to include information about habitat manipulation (there were
333 three categories here: unmanaged, machine-manipulated (mass management using
334 flail/digger) or hand-manipulated (selective chainsawing, volunteers with handtools).
335 We recorded numbers of species and individuals, calculated diversity (Inverse

336 Simpson's dominance; Morris *et al.* (2014)) and allocated species to a size category
337 (in mm; <25, 25-30, 30-40, 40-50, 50-60, or >60) based on Townsend and Waring
338 (2014).

339 To explore size composition for the subset of MOTUs identified to species
340 level within each sample, including those fed to chicks, we allocated each species
341 identified within the faecal samples to a size category according to mean wingspan,
342 using Townsend and Waring (2014). To test for significant differences in the
343 frequency of occurrence of different size classes between years we used Kruskal-
344 Wallis rank sum tests. To test for size selection of moth prey by Nightjars, we used
345 proportional Z-tests to compare the frequency of occurrence of each size class found
346 in the faecal samples, to those found in the traps, between years and months.

347

348 **RESULTS**

349 We retrieved a total of 13,900,616 sequence reads from 141 Nightjar faecal samples
350 using the Illumina MiSeq Benchtop sequencer, which comprised 1631 MOTUs after
351 trimming and clustering. After processing and filtering (i.e. removing 'impossible'
352 reads likely to be a result of contamination and degradation: De Barba *et al.* 2014),
353 we retained 625 unique MOTUs from 130 samples. Some samples ($n = 11$),
354 contained no reads once filtered, meaning that the amount of non-arthropod
355 contamination was high, as has been found in other studies (Regnaut *et al.* 2006).
356 65% of MOTUs were identified to species, 5% to genus, 12% to family and 18% only
357 to order (all Lepidoptera). Lepidoptera were found in 99% of samples, followed by
358 Diptera (27%), Coleoptera (9%), Neuroptera (7%) and <1% of both Hemiptera and
359 Hymenoptera. Of the sequences identified to species level, the most common moth

360 species present were Poplar Hawk-moth *Lathoe populi* (in 43% of samples), Silver Y
361 *Autographa gamma* (45%), Yellow-tail *Euproctis similis* (45%), True Lover's Knot
362 *Lycophotia porphyrea* (49%), Smoky Wainscot *Mythimna impura* (50.4%), and Large
363 Yellow Underwing *Noctua pronuba* (52%). Of the Coleoptera we were able to
364 identify, we found *Harpalus* sp. (Carabidae; 2% of samples), *Melanotus villosus* and
365 *Stenagostus rhombeus* (both Elateridae and in 6% of samples). Diptera sequences
366 identified belonged to the families Chironomidae, Sciaridae, Culicidae and Tipulidae,
367 but were not identifiable to any more detailed level than that of family. All identified
368 species are listed in the Supplementary Information (Appendix III).

369 *Variation in composition and diversity among samples*

370 Samples contained an average of 15.6 MOTUs (+/- SD 9.7). Mean Chao dissimilarity
371 of adult samples (2015: $n = 8$, 2016: $n = 17$, 2017: $n = 27$, 2018: $n = 27$; no chicks
372 and no repeat samples, and a reduced total number of 429 MOTUs) was 0.91 (+/-
373 SD 0.17), which shows relatively high individual variation. Mean Chao dissimilarity of
374 the repeat samples was 0.85 (+/- SD 0.19), therefore making related samples slightly
375 more similar.

376 No significant differences were found in sample composition between sexes
377 (Female: $n = 26$, Male: $n = 43$; $F_{1,68} = 0.36$, $P = 0.99$), nor between the months in
378 which the samples were collected (June: $n = 21$; July: $n = 32$; August: $n = 15$; $F_{2,70} =$
379 1.21 , $P = 0.16$). Between-year differences in sample composition were significant
380 (2015: $n = 8$, 2016: $n = 17$, 2017: $n = 27$, 2018: $n = 27$; $F_{3,75} = 3.3$, $P = 0.001$; $r^2 =$
381 0.12), supported by a non-significant difference in the distribution of samples around
382 the centroid ($F_{3,75} = 1.37$, $P = 0.26$; Figure 1), although some caution must be taken
383 due to uneven sample sizes. Specific significant among-year differences in variance
384 were shown between 2016 – 2017 and 2017 -2018 (Table 1).

385

386 **Moth species sizes within faecal samples**

387 96% of faecal samples contained moths with a wingspan of between 30 and 40mm
388 and 46% of samples contained the largest moths (>60 mm wingspan). The five most
389 frequent species in the samples all had a mean wingspan of more than 30mm; two of
390 these had a wingspan of more than 50mm (*N. pronuba*, *L. populi*). The largest moths
391 (>60mm) were present in significantly more samples in 2018 (66%), than all other
392 years (2015: 0%; 2016: 14%; 2017: 36%; $X^2 = 27.08$, $df = 3$, $P < 0.0001$).

393

394 **Size selection in diet samples compared to prey availability**

395 Faecal samples were more likely than moth traps to contain the largest moths
396 (>60mm). This was significant in 2017 (diet samples: 0.4, moth traps: 0.18; $X^2 = 4.61$,
397 $df = 1$, $P = 0.03$) and 2018 (diet samples: 0.67, moth traps: 0.34; $X^2 = 9.5$, $df = 1$, $P =$
398 0.002 ; Figure 2). In contrast, moth traps were more likely than faecal samples to
399 contain smaller moths (<25mm), but the difference was only significant in 2018 (diet
400 samples: 0.18, moth traps: 0.49; $X^2 = 9.55$, $df = 1$, $P = 0.002$; Figure 3).

401 Across all years, less than 18% of diet samples contained moths with the
402 smallest wingspan (<25mm). In June and August, faecal samples were more likely
403 than moth traps to contain the largest moths (June: 50-60mm: diet samples: 22%;
404 moth traps: 4%, $X^2: 11.66$, $df = 1$, $P = < 0.001$. August: >60mm: diet samples: 15%,
405 moth traps: 3%, $X^2: 5.73$, $df = 1$, $P = 0.02$).

406

407 **DISCUSSION**

408 We have demonstrated the use of high-throughput sequencing in investigating the
409 diet of an insectivorous, avian migrant to the UK, joining a growing bank of literature
410 exploring the use of metabarcoding to reduce the bias shown by physical dissection
411 of faeces towards retained, undigested fragments of prey items (Evens *et al.* 2020,
412 Mills *et al.* 2020). Our analyses showed strong intra- and inter-annual variation in diet
413 richness and size-based prey selection, which varied annually and seasonally.
414 Differences among years in Nightjar diet potentially reflected a response by Nightjars
415 to environmental fluctuation influencing the annual life cycle of moth species.

416 There was a dominance of Lepidoptera over other arthropods in Nightjar diet,
417 which has been reported previously (Sierro *et al.* 2001, Sharps 2013, Evens *et al.*
418 2020), although this does not hold true for all species of caprimulgid (Knight *et al.*
419 2018). The prevalence of Lepidoptera in the diet of this population may reflect local
420 availability of this prey type (exceedingly high numbers of larger moths, or very few
421 Coleoptera) and an adjustment of the population to being almost entirely Lepidoptera
422 specialists. It is also possible that due to our molecular optimisation procedures,
423 there has been a lack of amplification of groups other than Lepidoptera. However,
424 previous tests using the primers in our study showed very high taxonomic coverage
425 (>96%) for 11 invertebrate orders, and showed higher success for Coleoptera than
426 for Lepidoptera (Clarke *et al.* 2014).

427 Moths present in the faecal samples were diverse in terms of their habitat
428 preferences and population trends. The most common species found in faecal
429 samples were all habitat generalists (e.g. *N. pronuba*, *E. similis*). True Lover's Knot,
430 however, is a specialist of heathlands, its larvae feeding exclusively on heather
431 (Townsend & Waring 2007, Thomsen *et al.* 2015). A number of heathland and wet

432 woodland specialist species (including for example, Oak Eggar *Lasiocampa quercus*,
433 Map-winged swift *Hepialus fusconebulosa* and Gold Spot *Plusia festucae* were
434 caught in the light traps, but most species found in both locations were generalists,
435 found in such widespread habitats as hedgerows, woodlands and gardens. Although
436 some such generalist species have more positive population trends than habitat
437 specialists, large moths such as those preferentially selected by Nightjars, and
438 another migrant insectivore the Common Cuckoo *Cuculus canorus* are, overall, in
439 decline (Coulthard *et al.* 2019, Mills *et al.* 2020, Fox *et al.* 2021). The most
440 commonly identified moth species within Nightjar faeces, the Large Yellow
441 Underwing, has increased over the past 50 years by 72%, clearly thriving in
442 generalist habitats (Harrower *et al.* 2020). Conversely, True Lover's Knot, Silver Y,
443 Yellow-Tail, Poplar Hawkmoth and Smoky Wainscot, the other five most common
444 components of Nightjar faecal matter, have all declined since 1968 by between 1.5
445 and 83% (the latter being the heathland specialist, True Lover's Knot).

446 Samples differed significantly from each other. Although some samples may
447 have been taken from the same individual, intraspecific variation in diet is common
448 and can reflect both resource availability and the avoidance of competition through
449 resource partitioning (Kotler & Brown 1988, Maldonado *et al.* 2019, Alberdi *et al.*
450 2020), or differences in habitat selection. We tracked a number of individuals in this
451 population over the same four years of this study, and this tracking work highlighted
452 a lack of overall population preference for any particular habitat type (Mitchell *et al.*
453 2020). The lack of similarity in diet among samples potentially also indicates a lack of
454 association with a particular habitat for foraging. For species exploiting patchy,
455 ephemeral resources as the Nightjars are here, it is beneficial to be flexible to be

456 able to adjust to prey availability and abundance (Maldonado *et al.* 2019, Szigeti *et*
457 *al.* 2019).

458 Contrary to expectation, no significant difference in sample composition or
459 diversity was found between male and female Nightjars (Mata *et al.* 2016, Knight *et*
460 *al.* 2018). We expected to find that females had less diverse diets than males,
461 because of their limited foraging opportunity (Houston 1997). However, not all
462 females were incubating during the period when the faecal samples were acquired,
463 so perhaps were not restricted in their foraging duration. Differences between years
464 in sample variance and prey species richness may indicate that the population could
465 respond to variation in prey availability, by taking advantage of emergences of
466 particular species, perhaps including influxes of migratory moths, several species of
467 which (e.g. Silver Y) were identified within our Nightjar samples (Lee & McCracken
468 2005) . Annual and seasonal variation in moth community diversity and overall
469 abundance have been shown to be related to variation in climate, as well as habitat
470 management (Summerville & Crist 2004, Summerville *et al.* 2007).

471 By comparing diet and moth trap samples, we examined consumption
472 compared to local availability, bearing in mind that not all the MOTUs identified in the
473 diet samples could be translated into species and that moth trap catches do not
474 sample moths representatively (see Truxa & Fiedler 2012, Jonason *et al.* 2014 for
475 information on bias in trapping and Evens *et al.* 2020 for use of the same type of
476 trap). It is also pertinent to note that some Nightjars make trips away from their
477 nesting areas to separate foraging locations, in this and other studies (Alexander &
478 Cresswell 1992, Evens *et al.* 2020, Mitchell *et al.* 2020), which means that
479 depending on habitat type at any distant foraging locations, we will not have
480 completed accounted for all prey available to Nightjars across their individual

481 foraging ranges. Again, their preference for large moths, which should provide higher
482 energy gain per item than small moths is a strategy also recognised in other
483 nocturnal insectivores (Clare *et al.* 2009, Vesterinen *et al.* 2016). Twilight foraging
484 limits the time available for capturing prey, which should drive size bias upwards in
485 line with the energy maximization – time minimization rule (Pyke 1984, Schoener
486 2003), and our findings suggest that availability of larger moths might be a key
487 aspect of Nightjar foraging strategy and requirements (Araújo *et al.* 2011, Schrimpf
488 *et al.* 2012, Vesterinen *et al.* 2016, Evens *et al.* 2020).

489 Understanding the demographic consequences of variation in prey availability
490 is important in order to make practical decisions about future management. The
491 habitat management that occurred on our study area has resulted in a reduction in
492 birch woodland, a rich invertebrate prey source. However, the outcome of this is not
493 necessarily negative, if regenerating, herbaceous, understorey vegetation that
494 results from the clearance sustains sufficient numbers of moths (Summerville & Crist
495 2004) and continuing low-intensity management supports large moth species (Mills
496 *et al.* 2020). Whilst Nightjars may be resilient to variation in prey availability on an
497 annual basis, it remains especially important to encourage habitat types that provide
498 larval food for moths during the Nightjar chick raising period to allow offspring to
499 fledge.

500

501 **Use of metabarcoding for insectivorous diets**

502 Metabarcoding has made it possible to identify physically unrecognisable remains of
503 prey specimens for many taxa (e.g. Kartzinel & Pringle 2015, Hawlitschek *et al.*
504 2018, McClenaghan *et al.* 2019a). Despite considerable progress, such methods are

505 still hindered by a few key challenges. Firstly, contamination of samples with non-
506 target DNA means that caution must be taken when conducting laboratory work and
507 the post-sequencing bioinformatics (Zepeda-Mendoza *et al.* 2016). Although one of
508 the benefits of metabarcoding is that less common prey species can be identified,
509 the post-sequencing filtering thresholds that must be established to produce
510 confident results, can sometimes remove rare sequences.

511 Additionally, some studies have found within-sample variation in MOTU
512 composition and diversity (Alberdi *et al.* 2018, Mata *et al.* 2018), although between-
513 sample variation was always stronger. We did find lower within-sample variation than
514 between-sample variation, but both measures were higher than in these other
515 referenced studies. We do not know a sufficient amount about Nightjar gut transit
516 time so we are unable to quantify how many specimens or prey capture events were
517 contained in each faecal sample, but high within-sample variation could relate to a
518 high number of specimens per defaecation, or may relate more to the methods used
519 to extract the DNA. This does indicate that there are several steps that may be
520 advised in order to ensure maximum information and variation is obtained, such as
521 multiple extracts and PCR replicates of the same sample, as well as repeated
522 sampling from the same individuals across the season, where known (Alberdi *et al.*
523 2018). The latter would ensure that individual specialisation in diet could be recorded
524 through longitudinal data collection as opposed to a 'snapshot' of an individual's diet
525 (Araujo *et al.* 2011). Although there is concurrence across general metabarcoding
526 protocols, the specific details such as number of replicates and use of controls are
527 not standardised, which can limit utility and cross-study comparison (Deagle *et al.*
528 2019, Ando *et al.* 2020).

529 For our study, by far the biggest of these challenges was a lack of reference
530 sequences for particular groups of species in global databases (such as NCBI
531 Genbank used for BLAST processing). This lack of reference sequences meant that
532 species could not be identified for around 35% of our MOTUs, even after the addition
533 of 78 species of moth to the 16S reference database (compared to 90% of COI
534 sequences identified to species in Evens *et al.* 2020). This means that despite being
535 able to assess sample richness, variation and change between months and years,
536 understanding the ecological information that comes with a species-level
537 assignment, which we are lacking, is invaluable (Hebert *et al.* 2003). Although we did
538 contribute several dozen 16S sequences to GenBank, we know from a
539 comprehensive compilation of Lepidoptera records on Thorne Moors alone (not both
540 Moors that constitute the Humberhead Peatlands NNR), that there have been 717
541 moth species recorded (382 macro-, 335 micromoths; Moat 2014). This indicates
542 that much more structured, systematic barcoding of the 16S rRNA gene of British
543 moth species needs to take place, to further this type of work.

544

545 **Conclusions**

546 Nightjars showed within-population and annual diet variation, as well as size
547 selection. The population appears able to adjust its diet in response to variation in
548 prey availability, reflecting ecological theory. This has consequences for Nightjars'
549 ability to withstand habitat change, with populations and species exhibiting diverse
550 diets often able to adapt better. For cryptic, insectivorous species such as the
551 Nightjar, metabarcoding provides a very useful tool to identify diet components,
552 despite continuing challenges related to extraction through to sequencing.

553

554 We thank all staff and volunteers involved in sample collection, especially Paul Shawcroft,
555 Colin Neale, Vivian Hartwell, Stephen Mosely, Gracie Adams, Tim Jones and George Day.
556 Thank you to Tim Kohler at Natural England for facilitating sample collection and storage.
557 Many thanks to the three anonymous reviewers, and to the Associate Editor, who provided
558 excellent feedback and thoughtful comments on two submissions of this paper, helping to
559 improve the manuscript.

560

561 Data availability statement

562 Raw Fastq sequencing files will be openly-available upon publication through the
563 ENA, part of the EMBL under the primary accession number PRJEB44974.
564 Summary files for statistical analysis are available as an excel file in SOM 4.

565 **REFERENCES**

- 566 Alberdi, A., Garin, I., Aizpurua, O. and Aihartza, J. 2012. The foraging ecology of the
567 mountain long-eared bat *Plecotus macrobullaris* revealed with DNA mini-
568 barcodes. - PLoS One 7: e35692.
- 569 Alberdi, A., Aizpurua, O., Gilbert, M. T. P. and Bohmann, K. 2018. Scrutinizing key
570 steps for reliable metabarcoding of environmental samples. - Methods Ecol.
571 Evol. 9: 134–147.
- 572 Aldridge, H. D. J. N. and Brigham, R. M. 2008. Factors influencing foraging time in
573 two aerial insectivores: the bird *Chordeiles minor* and the bat *Eptesicus fuscus*,
574 Can. J. Zool., 69 (1), pp. 62–69.
- 575 Alexander, I. and Cresswell, B. 1990. Foraging by Nightjars *Caprimulgus europaeus*
576 away from their nesting areas. *Ibis*, 132 (4), pp.568-574.
- 577 Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. 1990. Basic
578 Local Alignment Search Tool (BLAST). - J. Mol. Biol. 215: 403–410.
- 579 Anderson, M. J. 2006. Distance-based tests for homogeneity of multivariate
580 dispersions. - Biometrics 62: 245–253.
- 581 Araújo, M. S., Bolnick, D. I. and Layman, C. A. 2011. The ecological causes of
582 individual specialisation. - Ecol. Lett. 14: 948–958.
- 583 Bayne, E. M. and Brigham, R. M. 1995. Prey selection and foraging constraints in
584 common poorwills (*Phalaenoptilus nuttallii*: Aves: Caprimulgidae). - J. Zool. 235:
585 1–8.
- 586 Berry, R. 1979. Nightjar habitats and breeding in East Anglia. *Br. Birds*, 72 (11).

587 Bolnick, D.I., Svanbäck, R., Fordyce, J.A., Yang, L.H., Davis, J.M., Hulseley, C.D. and
588 Forister, M.L. 2002. The ecology of individuals: Incidence and implications of
589 individual specialization, *Am. Nat.* 161 (1), pp. 1–28.

590 Bolnick, D. I., Yang, L. H., Fordyce, J. A., Davis, J. M. and Svanback, R. 2011.
591 Measuring Individual-Level Resource Specialization. - *Ecology* 83: 2936–2941.

592 Both, C., Van Turnhout, C.A., Bijlsma, R.G., Siepel, H., Van Strien, A.J. and Foppen,
593 R.P. 2010. Avian population consequences of climate change are most severe
594 for long-distance migrants in seasonal habitats, *Proc. R. Soc. B, Biol. Sci.* 277
595 (1685), pp. 1259–1266.

596 Bowles, E., Schulte, P. M., Tollit, D. J., Deagle, B. E. and Trites, A. W. 2011.
597 Proportion of prey consumed can be determined from faecal DNA using real-
598 time PCR. - *Mol. Ecol. Res.* 11: 530–540.

599 Brandon-Mong, G. J., Gan, H. M., Sing, K. W., Lee, P. S., Lim, P. E. and Wilson, J.
600 J. 2015. DNA metabarcoding of insects and allies: an evaluation of primers and
601 pipelines. - *Bull. Entomol. Res.* 105: 717–727.

602 Brownie, J., Shawcross, S., Theaker, J., Whitcombe, D., Ferrie, R., Newton, C. and
603 Little, S. 1997. The elimination of primer-dimer accumulation in PCR. *Nucl.*
604 *Acids Res.* 25 (16), pp. 3235 - 3241.

605 Chao, A., Chazdon, R. L. and Shen, T. J. 2005. A new statistical approach for
606 assessing similarity of species composition with incidence and abundance data.
607 *Ecol. Lett.* 8: 148–159.

608 Chao, A. and Chiu, C.H. 2016. *Species Richness: Estimation and Comparison*. Wiley
609 *StatsRef Stat. Ref. Online.* 1–26.

610 Charmantier, A., McCleery, R.H., Cole, L.R., Perrins, C., Kruuk, L.E. and Sheldon,
611 B.C. (2019) 'Adaptive Phenotypic Plasticity in Response to Climate Change in a
612 Wild Bird Population', *Science*, 320 (5877), pp. 800–803.

613 Clare, E. L., Fraser, E. E., Braid, H. E., Fenton, M. B. and Hebert, P. D. 2009.
614 Species on the menu of a generalist predator, the eastern red bat (*Lasiurus*
615 *borealis*): using a molecular approach to detect arthropod prey. *Mol. Ecol.* 18:
616 2532–2542.

617 Clare, E. L., Barber, B. R., Sweeney, B. W., Hebert, P. D. and Fenton, M. B. 2011.
618 Eating local: influences of habitat on the diet of little brown bats (*Myotis*
619 *lucifugus*).*Mol. Ecol.* 20: 1772–1780.

620 Clare, E. L., Symondson, W. O. C., Broders, H., Fabianek, F., Fraser, E. E.,
621 MacKenzie, A., Boughen, A., Hamilton, R., Willis, C. K. R., Martinez-Nunez, F.,
622 Menzies, A. K., Norquay, K. J. O., Brigham, M., Poissant, J., Rintoul, J., Barclay,
623 R. M. R. and Reimer, J. P. 2014. The diet of *Myotis lucifugus* across Canada:
624 assessing foraging quality and diet variability. *Mol. Ecol.* 23: 3618–3632.

625 Clarke, L. J., Soubrier, J., Weyrich, L. S. and Cooper, A. 2014. Environmental
626 metabarcodes for insects: In silico PCR reveals potential for taxonomic bias.
627 *Mol. Ecol. Res.* 14: 1160 - 1170.

628 Coulthard, E., Norrey, J., Shortall, C. and Harris, W.E. (2019) 'Ecological traits
629 predict population changes in moths', *Biol.Conserv.* 233: 213–219.

630 Cramp, S. 1985. Handbook of the Birds of Europe, the Middle East and North Africa:
631 The Birds of the Western Palearctic; Volume IV. Terns to Woodpeckers. -
632 Oxford University Press.

633 Cucco, M. and Malacarne, G. 1996. Reproduction of the pallid swift (*Apus pallidus*)
634 in relation to weather and aerial insect abundance. *Ital. J. Zool.* 63: 247–253.

635 De Barba, M., Miquel, C., Boyer, F., Mercier, C., Rioux, D., Coissac, E. and Taberlet,
636 P. 2014. DNA metabarcoding multiplexing and validation of data accuracy for
637 diet assessment: application to omnivorous diet. *Mol. Ecol. Res.* 14: 306–323.

638 Deagle, B. E., Thomas, A. C., McInnes, J. C., Clarke, L. J., Vesterinen, E. J., Clare,
639 E. L., Kartzinel, T. R. and Eveson, J. P. 2018. Counting with DNA in
640 metabarcoding studies: How should we convert sequence reads to dietary data?
641 *Mol. Ecol.* 28: 391 - 406.

642 Downs, N. C., Cresswell, W. J., Reason, P., Sutton, G., Wells, D. and Wray, S. 2016.
643 Sex-Specific Habitat Preferences of Foraging and Commuting Lesser
644 Horseshoe Bats *Rhinolophus hipposideros* (Borkhausen, 1797) in Lowland
645 England . *Acta Chiropt.* 18: 451–465.

646 Dunn, P.O., Winkler, D.W., Whittingham, L.A., Hannon, S.J. and Robertson, R.J.
647 2011. A test of the mismatch hypothesis: how is timing of reproduction related to
648 food abundance in an aerial insectivore? *Ecology*, 92 (2), pp.450-461.

649 Durst, S. L., Theimer, T. C., Paxton, E. H. and Sogge, M. K. 2008. Age, Habitat, and
650 Yearly Variation in the Diet of a Generalist Insectivore, the Southwestern Willow
651 Flycatcher. *Condor* 110: 514–525.

652 Eaton, M. A., Aebischer, N. J., Brown, A. F., Hearn R.D., Lock, L., Musgrove, A. J.,
653 Noble, D. G., Stroud, D. A. and Gregory, R. D. 2015. Birds of Conservation
654 Concern 4: the population status of birds in the United Kingdom, Channel Islands
655 and Isle of Man. *Br. Birds* 108: 708–746.

656 Emlen, J. M. 1966. The role of time and energy in food preference. *Am. Nat.* 100
657 (916), pp. 611-617.

658 English, P. A. 2009. A role for insect availability in limiting populations of a
659 threatened Nightjar *Antrostomus vociferous*. - PhD Thesis, Simon Fraser
660 University.

661 English, P.A., Nocera, J.J., Pond, B.A. and Green, D.J. 2017. Habitat and food
662 supply across multiple spatial scales influence the distribution and abundance of
663 a nocturnal aerial insectivore, *Landsc. Ecol.* 32 (2), pp. 343–359.

664 Eriksson, P., Mourkas, E., Gonzalez-Acuna, D., Olsen, B. and Ellstrom, P. 2017.
665 Evaluation and optimization of microbial DNA extraction from fecal samples of
666 wild Antarctic bird species. *Infect. Ecol. Epidemiol.* 7: 1386536.

667 Evens, R., Beenaerts, N., Witters, N. and Artois, T., 2017. Study on the foraging behaviour
668 of the European Nightjar *Caprimulgus europaeus* reveals the need for a change in
669 conservation strategy in Belgium. *J. Avian Biol.* 48 (9), pp.1238-1245.

670 Evens, R., Conway, G., Franklin, K., Henderson, I., Stockdale, J., Beenaerts, N., Smeets, K.,
671 Neyens, T., Ulenaers, E. and Artois, T., 2020. DNA diet profiles with high-resolution
672 animal tracking data reveal levels of prey selection relative to habitat choice in a
673 crepuscular insectivorous bird. *Ecol. Evol.* 10 (23), pp.13044-13056.

674 Fox, R., Oliver, T.H., Harrower, C., Parsons, M.S., Thomas, C.D. and Roy, D.B.
675 2014. Long-term changes to the frequency of occurrence of British moths are
676 consistent with opposing and synergistic effects of climate and land-use
677 changes', *J. Appl. Ecol.* 51 (4), pp. 949–957.

678 Fox, R., Dennis, E.B., Harrower, C.A., Blumgart, D., Bell, J.R.; Cook, P., Davis, A.M., Evans-
679 Hill, L.J., Haynes, F., Hill, D.; Isaac, N.J.B., Parsons, M.S., Pocock, M.J.O., Prescott,

680 T., Randle, Z., Shortall, C.R., Tordoff, G.M., Tuson, D., Bourn, N.A.D.. 2021 The state
681 of Britain's larger moths 2021. Wareham, Butterfly Conservation, Rothamsted Research
682 and UK Centre for Ecology & Hydrology, 44pp.Fuentes-Montemayor, E., Goulson,
683 D., Cavin, L., Wallace, J.M. and Park, K.J. (2012) 'Factors influencing moth
684 assemblages in woodland fragments on farmland: Implications for woodland
685 management and creation schemes', *Biol. Conserv.* 153, pp. 265–275.

686 Garlapow, R. M. 2007. Whip-poor-will prey availability and foraging habitat:
687 implications for management in pitch pine/scrub oak barrens habitat. Masters
688 Thesis, University of Massachusetts Amherst.

689 Hallmann, C. A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan, H.,
690 Stenmans, W., Muller, A., Sumser, H., Horren, T., Goulson, D. and de Kroon, H.
691 2017. More than 75 percent decline over 27 years in total flying insect biomass
692 in protected areas. *PLoS One* 12: e0185809.

693 Harrower, C.A., Bell, J.R., Blumgart, D., Botham, M.S., Fox, R., Isaac, N.J.B., Roy, D.B.,
694 Shortall, C.R. (2020) Moth trends for Britain and Ireland from the Rothamsted Insect
695 Survey light-trap network (1968 to 2016). NERC Environmental Information Data
696 Centre. <https://doi.org/10.5285/0a7d65e8-8bc8-46e5-ab72-ee64ed851583>.

697 Hawlitschek, O., Fernández-González, A., Balmori-de la Puente, A. and Castresana, J.
698 2018. A pipeline for metabarcoding and diet analysis from fecal samples developed for
699 a small semi-aquatic mammal. *PLoS One* 13: 1–19.

700 Hebert, P. D. and Gregory, T. R. 2005. The promise of DNA barcoding for taxonomy. *Syst.*
701 *Biol.* 54: 852–859.

702 Hebert, P. D., Cywinska, A., Ball, S. L. and deWaard, J. R. 2003. Biological identifications
703 through DNA barcodes. *Proc. Biol. Sci.* 270: 313–321.

704 Henderson, I., Hunter, D. and Conway, G. 2018. Comparing moth abundance between the
705 breeding and foraging locations of the European Nightjar *Caprimulgus europaeus*, in
706 Thetford Forest. *Ibis*: 2016–2019.

707 Howells, R. J., Burthe, S. J., Green, J. A., Harris, M. P., Newell, M. A., Butler, A., Johns, D.
708 G., Carnell, E. J., Wanless, S. and Daunt, F. 2017. From days to decades: short- and
709 long-term variation in environmental conditions affect offspring diet composition of a
710 marine top predator. *Mar. Ecol. Prog. Ser.* 583: 227–242.

711 Imlay, T. L., Mann, H. A. R. and Leonard, M. L. 2017. No effect of insect abundance on
712 nestling survival or mass for three aerial insectivores. *Avian Conserv. Ecol.* 12: 4–13.

713 Jackson, H. D. 2000. Food of the Nightjars in Zimbabwe. *Ostrich* 71: 404–407.

714 Jetz, W., Steffen, J. and Linsenmair, K. E. 2003. Effects of light and prey availability on
715 nocturnal, lunar and seasonal activity of tropical Nightjars, *Oikos*, 103 (3), pp. 627–639.

716 Ji, Y., Ashton, L., Pedley, S. M., Edwards, D. P., Tang, Y., Nakamura, A., Kitching, R.,
717 Dolman, P. M., Woodcock, P., Edwards, F. A., Larsen, T. H., Hsu, W. W., Benedick, S.,
718 Hamer, K. C., Wilcove, D. S., Bruce, C., Wang, X., Levi, T., Lott, M., Emerson, B. C.
719 and Yu, D. W. 2013. Reliable, verifiable and efficient monitoring of biodiversity via
720 metabarcoding. *Ecol. Lett.* 16: 1245–1257.

721 Jonason, D., Franzen, M. and Ranius, T. 2014. Surveying Moths Using Light Traps: Effects
722 of Weather and Time of Year. *PLoS One* 9: e92453.

723 Kartzinel, T. R. and Pringle, R. M. 2015. Molecular detection of invertebrate prey in
724 vertebrate diets: Trophic ecology of Caribbean island lizards. *Mol. Ecol. Res.* 15: 903–
725 914.

726 Knight, E. C., Ng, J. W., Mader, C. E., Brigham, R. M. and Bayne, E. M. 2018. “An inordinate
727 fondness for beetles”: first description of Common Nighthawk (*Chordeiles minor*) diet in
728 the boreal biome. *Wilson J. Ornithol.* 130: 525–531.

729 Kohn, M. H. and Wayne, R. K. 1997. Facts from feces revisited. *Trends Ecol. Evol.* 12: 223–
730 227.

731 Kotler, B.P. and Brown, J.S. 1988. Environmental heterogeneity and the coexistence of
732 desert rodents, *Annu. Rev. Ecol. Syst.* 19 (1), pp.281-307.

733 Krehenwinkel, H., Kennedy, S., Pekár, S. and Gillespie, R. G. 2017. A cost-efficient and
734 simple protocol to enrich prey DNA from extractions of predatory arthropods for large-
735 scale gut content analysis by Illumina sequencing. *Methods Ecol. Evol.* 8: 126–134.

736 Lazaridis, E. 2014. *lunar: Lunar phrase & distance, seasons and other environmental*
737 *factors.*: v 0.1-04.

738 Lee, Y. F. and McCracken, G. F., 2005. Dietary variation of Brazilian free-tailed bats links to
739 migratory populations of pest insects. *J. Mammalogy*, 86 (1), pp.67-76.

740 Magoč, T. and Salzberg, S. L. 2011. FLASH: Fast length adjustment of short reads to
741 improve genome assemblies. *Bioinformatics* 27: 2957–2963.

742 Maldonado, K., Bozinovic, F., Newsome, S. D. and Sabat, P. 2017. Testing the niche
743 variation hypothesis in a community of passerine birds. *Ecology* 98: 903–908.

744 Maldonado, K., Newsome, S. D., Razeto-Barry, P., Ríos, J. M., Piriz, G. and Sabat, P. 2019.
745 Individual diet specialisation in sparrows is driven by phenotypic plasticity in traits
746 related to trade-offs in animal performance. *Ecol. Lett.* 22: 128–137.

747 Mata, V. A., Amorim, F., Corley, M. F., McCracken, G. F., Rebelo, H. and Beja, P. 2016.
748 Female dietary bias towards large migratory moths in the European free-tailed bat
749 (*Tadarida teniotis*). *Biol. Lett.* 12 (3): 20150988.

750 McClenaghan, B., Kerr, K.C. and Nol, E., 2019a. Does prey availability affect the
751 reproductive performance of Barn Swallows (*Hirundo rustica*) breeding in Ontario,
752 Canada?. *Can. J. Zool.*, 97 (11), pp.979-987.

753 McClenaghan, B., Nol, E. and Kerr, K.C., 2019b. DNA metabarcoding reveals the broad and
754 flexible diet of a declining aerial insectivore. *The Auk: Ornithological Advances*, 136 (1).

755 Merckx, T., Slade, E. M., Basset, Y. and Christie, F. 2014. Macro-moth families differ in their
756 attraction to light: implications for light-trap monitoring programmes. *Insect Conserv.*
757 *Divers.* 7: 453–461.

758 Met Office. 2012. Met Office Integrated Data Archive System (MIDAS), LAnd and Marine
759 Surface Stations Data (1853 - current). NCAS Br. Atmos. Data Cent.

760 Metzker, M. L. 2010. Applications of Next Generation Sequencing technologies - the next
761 generation. *Nat. Rev. Genet.* 11: 31–46.

762 Meusnier, I., Singer, G.A., Landry, J.F., Hickey, D.A., Hebert, P.D. and Hajibabaei, M. 2008.
763 A universal DNA mini-barcode for biodiversity analysis, *BMC Genomics*, 9 (1), p. 214.

764 Mitchell, L.J., Kohler, T., White, P.C. and Arnold, K.E. 2020. High interindividual variability in
765 habitat selection and functional habitat relationships in European Nightjars over a
766 period of habitat change. *Ecol. Evol.* 10 (12), 5932 -5945.

767 Moat, R. 2014. Checklist of the Lepidoptera of Thorne Moors 1837 - 2014, Thorne and
768 Hatfield Moors Conservation Forum Technical Report No. 20 (Ed. Buckland, P. C.).
769 Thorne and Hatfield Moors Conservation Forum, Doncaster, UK.

770 Møller, A. 2013 Long-term trends in wind speed, insect abundance and ecology of an
771 insectivorous bird study area, *Ecosphere*, 4 (1), pp. 1–11.

772 Morris, E. K., Caruso, T., Buscot, F., Fischer, M., Hancock, C., Maier, T. S., Meiners, T.,
773 Müller, C., Obermaier, E., Prati, D., Socher, S. A., Sonnemann, I., Wäschke, N., Wubet,
774 T., Wurst, S. and Rillig, M. C. 2014. Choosing and using diversity indices: insights for
775 ecological applications from the German Biodiversity Exploratories. *Ecol. Evol.* 4: 3514–
776 24.

777 Mumma, M. A., Adams, J. R., Zieminski, C., Fuller, T. K., Mahoney, S. P. and Waits, L. P.

778 2016. A comparison of morphological and molecular diet analyses of predator scats. *J.*
779 *Mammalogy* 97: 112–120.

780 Nebel, S., Mills, A., Mccracken, J. D., Taylor, P. D., Nebel, S., Mills, A., Mccracken, J. D. and
781 Taylor, P. D. 2010. Declines of Aerial Insectivores in North America Follow a
782 Geographic Gradient. *Avian Conserv. Ecol.* 5: 1.

783 Nielsen, J. M., Clare, E. L., Hayden, B., Brett, M. T. and Kratina, P. 2018. Diet tracing in
784 ecology: Method comparison and selection. *Methods Ecol. Evol.* 9: 278–291.

785 Nocera, J. J., Blais, J. M., Beresford, D. V, Finity, L. K., Grooms, C., Kimpe, L. E., Kyser, K.,
786 Michelutti, N., Reudink, M. W. and Smol, J. P. 2012. Historical pesticide applications
787 coincided with an altered diet of aerially foraging insectivorous chimney swifts. *Proc.*
788 *Biol. Sci.* 279: 3114–3120.

789 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P.
790 R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E. and
791 Wagner, H. 2019. *vegan: Community Ecology Package.*: version 2.5-4.

792 Orłowski, G., Karg, J. and Karg, G. 2014. Functional invertebrate prey groups reflect dietary
793 responses to phenology and farming activity and pest control services in three
794 sympatric species of aerially foraging insectivorous birds. *PloS one*, 9 (12), e114906.

795 Paiva, V.H., Geraldles, P., Ramirez, I., Meirinho, A., Garthe, S. and Ramos, J.A. 2010.
796 Foraging plasticity in a pelagic seabird species along a marine productivity gradient,
797 *Mar. Ecol. Progr. Ser.* 398, pp: 259 – 274.

798 Pyke, G. H. 1984. Optimal Foraging Theory: A Critical Review. *Annu. Rev. Ecol. Syst.* 15:
799 523–575.

800 Razeng, E. and Watson, D. M. 2015. Nutritional composition of the preferred prey of
801 insectivorous birds: Popularity reflects quality. *J. Avian Biol.* 46: 89–96.

802 Razgour, O., Clare, E. L., Zeale, M. R. K., Hanmer, J., Schnell, I. B., Rasmussen, M.,

803 Gilbert, T. P. and Jones, G. 2011. High-throughput sequencing offers insight into
804 mechanisms of resource partitioning in cryptic bat species. *Ecol. Evol.* 1 (4), pp.556-
805 570.

806 Regnaut, S., Lucas, F. S. and Fumagalli, L. (2006) 'DNA degradation in avian faecal
807 samples and feasibility of non-invasive genetic studies of threatened capercaillie
808 populations', *Cons. Gen.*, 7 (3), pp. 449–453.

809 Roughgarden, J. 1974. Niche width: biogeographic patterns among *Anolis* lizard
810 populations. *Am. Nat.* 108 (962), pp.429-442..

811 Saino, N., Ambrosini, R., Rubolini, D., von Hardenberg, J., Provenzale, A., Hüppop, K.,
812 Hüppop, O., Lehikoinen, A., Lehikoinen, E., Rainio, K. and Romano, M. 2011. Climate
813 warming, ecological mismatch at arrival and population decline in migratory birds. *Proc.*
814 *R. Soc. B, Biol. Sci.* 278 (1707), pp.835-842.

815 Schoener, T. W. 2003. Theory of Feeding Strategies. - *Annu. Rev. Ecol. Syst.* 2: 369–404.

816 Schrimpf, M. B., Parrish, J. K. and Pearson, S. F. 2012. Trade-offs in prey quality and
817 quantity revealed through the behavioral compensation of breeding seabirds. *Mar. Ecol.*
818 *Prog. Ser.* 460: 247–259.

819 Sharps, K. 2013. The conservation ecology of the European Nightjar (*Caprimulgus*
820 *europaeus*) in a complex heathland-plantation landscape. PhD Thesis, University of
821 East Anglia, UK.

822 Sharps, K., Henderson, I., Conway, G., Armour-Chelu, N. and Dolman, P. M. 2015. Home-
823 range size and habitat use of European Nightjars *Caprimulgus europaeus* nesting in a
824 complex plantation-forest landscape, *Ibis* 157: 260–272.

825 Siervo, A., Arlettaz, R., Naef-Daenzer, B., Strebel, S. and Zbinden, N. 2001. Habitat use and
826 foraging ecology of the Nightjar (*Caprimulgus europaeus*) in the Swiss Alps: Towards a
827 conservation scheme. *Biol. Conserv.* 98: 325–331.

828 Stahl, J., Tolsma, P.H., Loonen, M.J.J.E. and Drent, R.H. 2001. Subordinates explore but
829 dominants profit: resource competition in high Arctic barnacle goose flocks. *Anim.*
830 *Behav.* 61: 257-264.

831 Stanton, R., Clark, R. G. and Morrissey, C. A. 2017. Intensive agriculture and insect prey
832 availability influence oxidative status and return rates of an aerial insectivore.
833 *Ecosphere* 8: e01746.

834 Sullins, D. S., Haukos, D. A., Craine, J. M., Lautenbach, J. M., Robinson, S. G., Lautenbach,
835 J. D., Kraft, J. D., Plumb, R. T., Reitz, J. H., Sandercock, B. K. and Fierer, N. 2018.
836 Identifying the diet of a declining prairie grouse using DNA metabarcoding. *Auk* 135:
837 583–608.

838 Summerville, K.S. and Crist, T.O. 2002. Effects of timber harvest on forest lepidoptera:
839 Community, guild, and species responses. *Ecol. Appl.* 12: 820 - 835.

840 Summerville, K.S. and Crist, T.O. 2004. Contrasting effects of habitat quantity and quality on
841 moth communities in fragmented landscapes. *Ecography* 27: 3 -12.

842 Summerville, K.S., Bonte, A.C. and Fox, L.C. 2007. Short-term temporal effects on
843 community structure of lepidoptera in restored and remnant tallgrass prairies. *Restor.*
844 *Ecol.* 15: 179 - 188.

845 Szigeti, V., Korosi, A., Harnos, A. and Kis, J. 2019. Lifelong foraging and individual
846 specialisation are influenced by temporal changes of resource availability. *Oikos* 128:
847 649–658.

848 Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C. and Willerslev, E. 2012. Towards
849 next-generation biodiversity assessment using DNA metabarcoding. *Mol. Ecol.* 21:
850 2045–2050.

851 Thomas, D. W., Brigham, R. M. and Lapierre, H. 1996. Field metabolic rates and body mass
852 changes in common poorwills (*Phalaenoptilus nuttallii*: Caprimulgidae). *Ecoscience* 3:

853 70–74.

854 Thomas, A. C., Deagle, B. E., Eveson, J. P., Harsch, C. H. and Trites, A. W. 2016.
855 Quantitative DNA metabarcoding: improved estimates of species proportional biomass
856 using correction factors derived from control material. *Mol. Ecol. Res.* 16: 714–726.

857 Thomsen P. F., Jørgensen, P. S., Bruun, H. H., Pedersen, J., Riis-Nielsen, T., Jonko, K.,
858 Słowińska, I., Rahbek, C., Karsholt, O. 2016. Resource specialists lead local insect
859 community turnover associated with temperature—analysis of an 18-year full-seasonal
860 record of moths and beetles. *J. Anim. Ecol.* 85 (1), pp. 251-61.

861 Townsend, M. and Waring, P. 2014. Concise guide to the moths of Great Britain and
862 Ireland. British Wildlife Publishing.

863 Trevelline, B. K., Latta, S. C., Marshall, L. C., Nuttle, T. and Porter, B. A. 2016. Molecular
864 analysis of nestling diet in a long-distance Neotropical migrant, the Louisiana
865 Waterthrush (*Parkesia motacilla*). *Auk* 133: 415–428.

866 Truxa, C. and Fiedler, K., 2012. Attraction to light—from how far do moths (Lepidoptera)
867 return to weak artificial sources of light? *Eur. J. Entomol.* 109 (1), pp. 77- 84.

868 van Strien, A. J., van Swaay, C. A. M., van Strien-van Liempt, W. T. F. H., Poot, M. J. M. and
869 WallisDeVries, M. F. 2019. Over a century of data reveal more than 80% decline in
870 butterflies in the Netherlands. *Biol. Conserv.* 234: 116–122.

871 Vesterinen, E. J., Ruokolainen, L., Wahlberg, N., Pena, C., Roslin, T., Laine, V. N., Vasko,
872 V., Saaksjarvi, I. E., Norrdahl, K. and Lilley, T. M. 2016. What you need is what you
873 eat? Prey selection by the bat *Myotis daubentonii*. *Mol. Ecol.* 25: 1581–1594.


874 Wilson, J. J., Rougerie, R., Schonfeld, J., Janzen, D. H., Hallwachs, W., Hajibabaei, M.,
875 Kitching, I. J., Haxaire, J. and Hebert, P. D. N. 2011. When species matches are
876 unavailable are DNA barcodes correctly assigned to higher taxa? An assessment using
877 sphingid moths. *BMC Ecol.* 11: 18.

878 Winiger, N., Korner, P., Arlettaz, R. and Jacot, A. 2018. Vegetation structure and decreased
879 moth abundance limit the recolonisation of restored habitat by the European Nightjar.
880 Rethink. Ecol. 3: 25–39.

881 Zeale, M. R., Butlin, R. K., Barker, G. L., Lees, D. C. and Jones, G. 2011. Taxon-specific
882 PCR for DNA barcoding arthropod prey in bat faeces. Mol. Ecol. Res. 11: 236–244.

883

884




885

886 **Figure 1.** Mean and variance of faecal sample distances to group centroids for each year samples
887 were collected, produced by the *'betadisper'* function in r. Sample sizes vary between groups and are
888 presented in the text.

889

890



891

892 **Figure 2.** Proportion of each moth size class present in diet samples and moth trap catches, for each
 893 year. Differences between faecal samples and moth trap samples were calculated using proportional
 894 Z-tests and significance levels are denoted using: ** = <0.05, *** = <0.01. Note: No moth trapping took
 895 place in 2015.

896

897 **Table 1.** Pairwise comparisons of among-year differences in mean sample composition using the
 898 function pairwise.Adonis in R which takes the Chao dissimilarity values as input. Significant adjusted
 899 *P* values (< 0.05) in bold.

Pairs	df	Sums of Sq	<i>F</i>	<i>R</i>²	<i>P</i>	<i>P</i> adjusted
2015 vs 2016	1	0.663	1.587	0.065	0.034	0.204
2015 vs 2017	1	0.623	1.452	0.042	0.089	0.534
2015 vs 2018	1	0.872	2.301	0.065	0.006	0.036
2016 vs 2017	1	1.656	4.064	0.088	0.001	0.006
2016 vs 2018	1	2.495	6.775	0.139	0.001	0.006
2017 vs 2018	1	1.061	2.758	0.050	0.006	0.036

900

901