UNIVERSITY of York

This is a repository copy of A Multi-Proxy Holocene Palaeoenvironmental Record of Climate Change and Prehistoric Human Activity from Lough Cullin, southeast Ireland.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/182949/</u>

Version: Accepted Version

Article:

Kearney, Kevin, Gearey, Benjamin, Hegarty, Susan et al. (11 more authors) (2022) A Multi-Proxy Holocene Palaeoenvironmental Record of Climate Change and Prehistoric Human Activity from Lough Cullin, southeast Ireland. The Holocene. pp. 262-279. ISSN 0959-6836

https://doi.org/10.1177/09596836211066593

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

The Holocene

A Multi-Proxy Holocene Palaeoenvironmental Record of Climate Change and Prehistoric Human Activity from Lough Cullin, southeast Ireland

Journal:	The Holocene
Manuscript ID	HOL-21-0083.R1
Manuscript Type:	Paper
Date Submitted by the Author:	08-Nov-2021
Complete List of Authors:	Kearney, Kevin; University College Cork, Department of Archaeology Gearey, Benjamin; University College Cork, Department of Archaeology Hegarty, Susan; Dublin City University, School of History and Geography Richer, Suzi; Richer Environmental Ferreira, Carla; Queen's University Belfast, Archaeology and Palaeoecology O'Carrol, Ellen; University College Dublin, School of Archaeology Becker, Katharina; University College Cork, Department of Archaeology Hamilton, Derek; Scottish Universities Environmental Research Centre Eogan, James; Transport Infrastructure Ireland McClatchie, Meriel; University College Dublin, School of Archaeology Armit, Ian; University of York, Department of Archaeology Nagle, Caitlin ; The University of Sheffield Taylor, Kate; TVAS (Ireland) Ltd Hull, Graham; TVAS (Ireland) Ltd.
Keywords:	Palaeoecology, Pollen Analysis, Chronology, Archaeology, Ireland, Human Impact, Climate
Abstract:	A multiproxy (pollen, microcharcoal, loss-on-ignition, magnetic susceptibility and geochemistry) sequence from Lough Cullin, southeast Ireland, supported by a high-resolution radiocarbon chronology, modelled using Bayesian approaches, provides a record of environmental change for much of the Holocene. Following the establishment of mixed deciduous woodland, climatic deterioration was likely responsible for pronounced vegetation change and erosion, 7615-6500 cal. BC to 6245- 5575 cal. BC, evidence for the '8.2 Kyr' BP climate event. The so-called 'elm decline' is dated to 4220-3980 cal. BC and whilst there are possible indications of an anthropogenic cause, clear evidence of woodland clearance with cereal pollen is recorded at 3900-3700 cal. BC, 3790- 3580 cal. BC and 3760-3650 cal. BC, during a period of clearance and farming of 320-450 years duration. A reduction in farming/settlement and woodland regeneration during the Middle Neolithic parallels the archaeological record, with low levels of activity during the Late Neolithic/Chalcolithic after 2960-2525 cal. BC, prior to increases during the Bronze Age then woodland clearance and agriculture between 1500- 1410 and 1275-1000 cal. BC, corresponding with the archaeological evidence. A subsequent 'step-wise' reduction in human activity follows,

http://mc.manuscriptcentral.com/holocene

	from the latter date to 815-685 cal. BC, and a brief but pronounced
	cessation at 690-535 cal. BC. Renewed woodland clearance and agriculture commenced until 415-250 cal. BC. From the latter date unti cal. AD 390-540, the Late Iron Age/Early Medieval period, a phase of woodland recovery is attested, followed by renewed landscape disturbance and arable agriculture in particular, continuing to the close of the record at cal. AD 780-1035.
L	
	SCHOLARONE [™]
	Manuscripts
	http://mc.manuscriptcentral.com/holocene

1

2		
3	1	A Multi-Proxy Holocene Palaeoenvironmental Record of Climate Change and
4 5	2	Prehistoric Human Activity from Lough Cullin, southeast Ireland.
6	2	Tremstorie fruman Activity from Lough Cumit, southeast fretand.
7 8	3	Ву
9 10	4	Kevin Kearney, Benjamin Gearey, Susan Hegarty, Suzi Richer,
11 12	5	Carla Ferreira, Ellen O' Carroll, Katharina Becker, Derek Hamilton, James Eogan, Meriel
13 14	6	McClatchie, Ian Armit, Caitlin Nagle, Kate Taylor and Graham Hull.
15 16 17	7	
18 19	8	Kevin Kearney
20 21	9	Department of Archaeology, University College Cork.
22 23	10	kevin.kearney@ucc.ie
24 25	11	Benjamin Gearey
26 27 28	12	Department of Archaeology, University College Cork.
29 30	13	b.gearey@ucc.ie
31 32	14	Susan Hegarty
33 34	15	School of History and Geography, Dublin City University
35 36 37	16	susan.hegarty@dcu.ie
38 39	17	Suzi Richer
40 41	18	Richer Environmental
42 43	19	suzi@richerenvironmental.com
44 45 46	20	Carla Ferreira
47 48	21	Archaeology & Palaeoecology, School of Natural and Built Environment, Queen's University
49 50	22	Belfast, Northern Ireland.
51 52	23	cferreira01@qub.ac.uk
53 54	24	Ellen O' Carroll
55 56	25	School of Archaeology, University College Dublin
57 58 50	26	ellen.ocarrol@ucd.ie
59 60	27	Katharina Becker

28	Department of Archaeology, University College Cork.
29	katharina.becker@ucc.ie
30	Derek Hamilton
31	Scottish Universities Environmental Research Centre
32	derek.hamilton.2@glasgow.ac.uk
33	James Eogan
34	Transport Infrastructure Ireland
35	james.eogan@tii.ie
36	Meriel McClatchie
37	School of Archaeology, University College Dublin
38	meriel.mcclatchie@ucd.ie
39	Ian Armit
40	Department of Archaeology, University of York
41	ian.armit@york.ac.uk
42	Caitlin Nagle
43	Department of Archaeology, University of Sheffield
44	cnagle1@sheffield.ac.uk
45	cnagle1@sheffield.ac.uk Kate Taylor
46	TVAS (Ireland) Ltd.
47	kate@tvasireland.ie
48	Graham Hull
49	TVAS (Ireland) Ltd.
50	graham@tvasireland.ie
51	
52	
53	
	 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52

54 Abstract.

A multiproxy (pollen, microcharcoal, loss-on-ignition, magnetic susceptibility and geochemistry) sequence from Lough Cullin, southeast Ireland, supported by a high-resolution radiocarbon chronology, modelled using Bayesian approaches, provides a record of environmental change for much of the Holocene. Following the establishment of mixed deciduous woodland, climatic deterioration was likely responsible for pronounced vegetation change and erosion, 7615-6500 cal. BC to 6245-5575 cal. BC, evidence for the '8.2 Kyr' BP climate event. The so-called 'elm decline' is dated to 4220-3980 cal. BC and whilst there are possible indications of an anthropogenic cause, clear evidence of woodland clearance with cereal pollen is recorded at 3900-3700 cal. BC, 3790-3580 cal. BC and 3760-3650 cal. BC, during a period of clearance and farming of 320-450 years duration. A reduction in farming/settlement and woodland regeneration during the Middle Neolithic parallels the archaeological record, with low levels of activity during the Late Neolithic/Chalcolithic after 2960-2525 cal. BC, prior to increases during the Bronze Age then woodland clearance and agriculture between 1500-1410 and 1275-1000 cal. BC, corresponding with the archaeological evidence. A subsequent 'step-wise' reduction in human activity follows, from the latter date to 815-685 cal. BC, and a brief but pronounced cessation at 690-535 cal. BC. Renewed woodland clearance and agriculture commenced until 415-250 cal. BC. From the latter date until cal. AD 390-540, the Late Iron Age/Early Medieval period, a phase of woodland recovery is attested, followed by renewed landscape disturbance and arable agriculture in particular, continuing to the close of the record at cal. AD 780-1035.

76 Keywords: Palaeoecology, Pollen Analysis, Chronology, Archaeology, Ireland.

1. Introduction

Understanding the character and chronology of Holocene vegetation development and the respective roles of climatic and anthropogenic impact on patterns and process of landscape change is a key focus of palaeoecological research. There is a long history of such research in Ireland, where the wide distribution of peatland and lake deposits provides the required sampling sites, and investigations continue to provide new insights into environmental change. Various questions concern the timing and character of human activity, climatic change and its relationship to archaeology, continue to be a subject of ongoing debate (see Kelly and O'Carragain 2021 for a summary). A particular methodological issue concerns the production

Page 5 of 49

HOLOCENE

of tightly constrained chronologies that are now available for the archaeological record (Cooney et al. 2011; McClatchie et al. 2014; McLaughlin et al. 2016) and the utility of formal modelling procedures to generate robust estimates of the tempo of environmental changes such as those around the mid-Holocene 'elm decline' (e.g. Whitehouse et al. 2014; Kearney and Gearey, 2020) and the Bronze Age to Iron Age transition (e.g. Gearey et al., 2021). Whilst it has long been largely common practice to use 'single point' estimates to describe palaeoenvironmental chronologies, largely for reasons of perceived clarity of data presentation, we utilise posterior density estimates, as these reflect the inherent probabilistic uncertainty associated with radiocarbon dates. It is now usual in much archaeological research to present chronological information in this way (Whittle et al. 2011). In order to allow meaningful, robust comparisons between archaeological and palaeoenvironmental datasets, it is essential we standardise chronological 'currencies' and hence develop a 'best practice' approach.

98 Recent detailed palaeoenvironmental studies have focussed on the west and north of the island 99 (e.g Chique *et al.* 2017; Spencer *et al.* 2020; Stolze and Monecke 2020) but there has been 100 relatively little research in the southeast (see Eogan *et al.* 2015). In this paper, we describe and 101 discuss the results of a multi-proxy palaeoenvironmental analysis of a 7.75m core from Lough 102 Cullin, Co. Kilkenny, southeast Ireland, which is supported by a high-resolution radiocarbon 103 chronology. Compilation of archaeological evidence from the close vicinity of the sampling 104 site provides an independent record of the relative intensity of human activity through time.

2. Background

Lough Cullin (52° 19' N, -7° 06' W). is a small lake (c.5 hectares) located in the catchment of the river Suir, in the south of County Kilkenny, southeast Ireland. The lake is situated in a lowlying landscape (11m OD), has a maximum depth of 3.4m and is fed by a series of small streams from the north (with a stream flowing through a channel incised into the Lower Palaeozoic and Devonian rocks at Catsrock), east and south. The outflow is via a small stream to the south-west. The lake is nestled in the centre of a syncline, with the underlying geology of the lake itself being well-bedded pale grey lower Dinantian limestones of the Ballysteen Formation (Tietzsch-Tyler and Sleeman 1994, the Hook Head formation of Keeley 1983). The slope increases significantly 2.5 km to the east and north of the lake basin, as the geology changes from limestone to Devonian sandstones. The recession of the lake due to nineteenth century drainage can be seen on the various editions of Ordnance Survey maps produced during the

nineteenth and early twentieth centuries. John O'Donovan, who was one of the civilians employed by the Ordnance Survey of Ireland in the 1830s and compiled the name books for the Ordnance Survey, remarked: 'I remember this lake when it was of considerable extent. It is now nearly drained, and the bog nearly all cut out' (OSI, n.d.). There is also sedimentological evidence the lake was once larger, with lacustrine sediments filling an area of 2 km² around the lake. The presence of a glaciofluvial sand and gravel deposit to the north of the lake between it and the channel at Catsrock, suggests the lake may, partly at least, owe its origin to glacial meltwater.

3. Methods and Materials

3.1. Core sampling and lithostratigraphy.

In June 2012, a 7.75 metre sediment core was extracted from the deepest part of the lake using a Livingstone piston corer from a platform secured in the centre of the lake. The core was extracted and wrapped on site, and later transferred to the lab where it was cut in half along its length. One half was used for geochemical analysis, and both halves were subsampled at 1 cm intervals down to 6.33 metres (the top of the clays).

135 3.2. Palynological analysis

Sub-samples of 1 cm³ were prepared for pollen analyses following standard techniques including KOH digestion, HF treatment and acetylation (Moore et al., 1991). Pollen concentrations were established by adding a known concentration of Lycopodium clavatum spore (batch number 177745) to the samples before treatment (Stockmarr 1971). Pollen counts were made using a Leica DM 1000 LED microscope at x400, x800 and x1000 magnification under oil immersion for critical examination of pollen sculpture and measurement of pollen grains. A minimum pollen sum of 400 Total Land Pollen (TLP) grains, excluding spores and aquatics, was employed. Pollen grains were identified mainly using the key from Moore et al. (1991) and Bennett (1995), with reference to Fægri et al. (1989). Pollen and spore identification was made to the lowest taxonomic level possible using the available references and following the nomenclature of Stace (1997), with suggestions from Bennett et al. (1994). The programmes TILIA and TILIA - GRAPH (Grimm 2013) were used to construct spreadsheets and pollen diagrams. Local pollen assemblage zones (LPAZ) were defined based on visible changes in the pollen record.

3.3. Loss on Ignition (LoI)

Sediment samples ($c.5 \text{ cm}^3$) were taken at 2 cm intervals to calculate loss-on-ignition (LoI %). The samples were dried for 24 hours at 105°C and subsequently transferred to porcelain crucibles and ashed for 4 hours at 550 °C (after Bengtsson and Enell, 1986; Heiri *et al.*, 2001).

154 3.4. Geochemical analysis

Geochemical analysis was carried out through non-destructive X-ray fluorescence (XRF) core scanning of the split cores using a Cox Analytical Systems ITRAX XRF Core Scanner located in the School of Geography, University College Dublin. 28 trace elements were selected for analysis (Mg, Al, Si, P, S, Cl, Ar, K, Ca, Ti, V, Cr, Mn, Fe, Ni, Cu, Zn, Ge, As, Br, Rb, Sr, Y, Zr, Ba, Nd, Pb, U), of which 10 key elements (Fe, Si, Ti, Ca, K, Sr, Rb, Mn, Cu, and Br) and the ratio of inc:coh were chosen for interpretation. Cores were scanned at 600 µm step intervals with a count time of 15 seconds, using a Mo tube operating at 30 kV and 30 mA.

The elemental data obtained are counts only, and so should be considered as semi-quantitative in nature (Croudace *et al.* 2006; Weltje and Tjallingii 2008). The data was normalized using kilo counts per second (kcps) to account for the variation in XRF intensities (Jouve *et al.* 2013; Turner *et al.* 2015; Gregory *et al.* 2019). Relative elemental concentrations within a core can provide valuable information on the sedimentology and palaeoclimatic history of an area (Guyard *et al.* 2007), particularly when coupled with data from other methodologies.

Along with the geochemical data, the ITRAX scanner returns a profile of the magnetic susceptibility (MS) measurement of the core at 4 mm resolution. MS is the amount of magnetisation acquired by the sediments of the core due to the application of a weak magnetic field. MS is related to the amount of dia- para- and ferrimagnetic minerals within the core (Støren *et al.* 2016), and changes in MS can be indicative of sediment flux and erosion in lake catchments (Evans and Slaymaker 2004).

- **3.5.** Chronology
- 176 3.5.1. Radiocarbon Dating

Thirty-three samples (52 measurements in total) were dated by Accelerator Mass Spectrometry at the Scottish Universities Environmental Research Centre (SUERC-), Queen's University Belfast (UBA-) and Beta Analytic (Beta-). All measurements were obtained from organic sediment samples except for *UBA-22422*, which was obtained from a branch fragment of *Salix*

sp. (willow). Where possible, duplicate measurements on the humin (the acid and alkaliinsoluble) and humic (alkali-soluble, acid insoluble) fractions (Cook *et al.* 1998, 21) were
obtained to test the reliability of sediment derived radiocarbon determinations (cf. Brock *et al.*2011) and the statistical consistency (Ward and Wilson 1978) of each paired date assessed
using the OxCal (Bronk Ramsey 2009a) programme.

The resulting radiocarbon determinations are presented as conventional radiocarbon ages (Stuiver and Polach 1977) quoted in accordance with the international standard established by the Trondheim Convention (Stuiver and Kra 1986). The calibrations have been calculated using the published datasets, IntCal20, (Reimer et al. 2020) and the computer program OxCal v4.3 (Bronk Ramsey 2009a) and are cited with the end points rounded outward to 10 years. The calibrated date ranges have been calculated using the maximum intercept method (Stuiver and Reimer 1986) and the graphical distribution of the calibrated result derived from the probability method (Stuiver and Reimer 1993). All radiocarbon dates are cited at 95.4% confidence, unless stated otherwise.

3.5.2. Bayesian Modelling.

In order to refine the chronology of palynological features and extend the chronological framework to changes which fall between dated horizons, a Bayesian approach was employed to the modelling of the radiocarbon dates (Buck et al. 1992). The Bayesian approach has been widely discussed in the literature (e.g. Blaauw and Christensen 2005; Blaauw et al. 2007a; Blaauw et al. 2007b; Bronk Ramsey 2008). The approach taken here uses the software OxCal v4.3 program (Bronk Ramsey 2009a), and the *P* Sequence function (Bronk Ramsey 2008) was chosen to allow for a Poisson process, or a potentially random rate, of sediment formation. The prior information for the deposition rate was defined as log $10(k/k \ 0)$ where k 0 = 1. This allows k to take any value from 0.01-100. This value used was estimated from the radiocarbon dates following the approach outlined in Blockley et al. (2007; 2008)

The command Outlier() was used in OxCal (Bronk Ramsey 2009b) to identify any measurements that were statistically defined as outliers at a 0.05 probability (1 in 20 chance). The model has been constrained using Boundaries, the depths of which have been defined based on stratigraphic changes (see below). The ranges quoted in italics are *posterior density* estimates, derived from this Bayesian modelling of the radiocarbon chronology (Bayliss et al. 2007) and are cited at 95.4% probability, unless stated otherwise.

2	
3 4	213
5 6	214
7 8	215
9 10	216
11	217
12 13	218
14 15	219
16 17	220
18	
19 20	221
21 22	222
23	223
24 25	224
26 27	225
28	226
29 30	227
31 32	228
33 34	229
35 36	230
37 38	231
39 40	232
41 42	233
43 44	234
45	235
46 47	236
48 49	237
50	238
51 52	
53 54	239
55	240
56 57	241
58	242
59 60	243

14 4.1. Stratigraphy.

4. Results.

The core lithology was (from top to bottom) light grey, organic-rich sediment (0 - 1.75 m); dark brown gyttja (1.75 m - 6.26 m), with a layer of sandier gyttja between 5.55 m and 5.74 m. The gyttja lies above 6 cm of sand (6.26 m - 6.32 m). At the base of the sequence was 1.65 metres of laminated grey-brown silty clay, with an angular pebble within the sediment at 7 metres.

221 4.2. Palynological analysis.

A total of forty-nine different taxa, including spores, were recorded across the Lough Cullin pollen profile. The data is presented in Table 1 and the percentage pollen diagram LC (Figure 2) and pollen concentration diagram (Figure 3) have been divided in 13 local pollen assemblage zones (LPAZ). LPAZ boundaries are given as *posterior density estimates* derived from the agedepth model shown in Figure 5. In addition, a curve for 'anthropogenic indicator' (*sensu* Behre 1981) taxa has been calculated (comprising *Cerealia*-type, *Artemisia*-type, Asteraceae undif., Caryophyllaceae, *Plantago lanceolata, Plantago major/media, Rumex, Trifolium*-type, and *Urtica dioica*).

4.3. Loss on Ignition (LoI), Magnetic Susceptibility (MS) and Geochemistry

The results of loss on ignition are presented in Table 1 and, with the results of geochemistry and magnetic susceptibility analyses, shown in Figure 4. Zonation for LoI data follows the LPAZ used in the pollen diagram. LoI results are relatively stable throughout the sequence, with some generally minor fluctuations. However, there is a marked decrease in LoI values at 5.09 m, with a significant and more sudden decrease at 4.90 m, possibly indicating accelerated minerogenic sedimentation into the lake. LoI begins to increase at 3.14 m to a peak at 2.15 m, from where it decreases to the top of the core.

The geochemical data, given the high resolution, is 'noisy' and must be interpreted with some caution. These data do not always track the changes in LoI, but there are observable patterns which may be hypothesised as reflecting weathering and erosion of different areas of the catchment: in particular, increased values for Calcium (Ca) may derive predominantly from the limestone bedrock areas (see above, Section 2), with Iron (Fe) from the sandstone areas. The latter element does not consistently track the MS curve, possibly because of the dissolution ofmagnetic minerals.

247 4.4. Chronology

4.4.1. Radiocarbon dating and Bayesian modelling.

The measurement on the humic fraction GU43402 from 2.61m failed, therefore the humin measurement (SUERC-72571) from this depth was used in the model. Four pairs of measurements from 2.65m (SUERC-70823 and SUERC-70824), 2.71m (SUERC-70825 and SUERC-70826), 3.26m (SUERC-76326 and SUERC-76666) and 5.34m (SUERC-70843 and SUERC-70844) are not statistically consistent (see Table 2). In all instances the humic fraction is younger than the humin fraction; this is likely caused by the downwards penetration of humic acids from above, therefore the humin fractions (SUERC-70824, SUERC-70826, SUERC-76666 and SUERC-70844) have been used in the modelling process for these depths. Measurement UBA-22422, a fragment of Salix (willow) twig, was identified as a possible outlier and has been excluded from the model; this determination was considerably older than those below it, suggesting that reworked material had been incorporated into the profile.

An age-depth model (Bronk Ramsey 2008) for the profile was constructed to provide a chronology for the sequence. The P Sequence model (Figure 5) has good agreement (A_{model}=99) and has been used to provide estimates for the boundaries between the local pollen assemblage zones (Table 1). Except for UBA-22422 (the Salix twig) the model shows good agreement, suggesting the sequence is intact below 2.19m.

The pollen sequence can be estimated to cover a period of 8830-9560 years (95.4% probability, distribution not shown), probably 9060-9330 years (68.2% probability, distribution not shown). Estimates derived from the age-depth model suggest an average deposition rate of 4-5 cm/100yr (distribution not shown) between the upper and basal radiocarbon measurements; it can be estimated that 21-23 years (distribution not shown) are represented in every centimetre.

The effectiveness of the model can be seen by considering the radiocarbon determination UBA-22423: this provided a calibrated (unmodelled) date of 750-400 cal. BC. The output on the model (Figure 5) constrained this to 595-445 cal. BC (Figure 5), a significant reduction in the bandwidth, from 350 to 150 years. This reduction was partly achieved through obtaining two determinations that fell on the 'steep' part of the radiocarbon calibration curve (SUERC-70824

HOLOCENE

and Beta-426775) thus helping to constrain the determinations between them, in particular
those on the 'Hallstatt plateau'.

5. Interpretation

280 LC-1: 6.25-6.03m, 9050-8090 cal. BC to 7615-6500 cal. BC

The pollen diagram covers much of the Holocene from 9030-8065 cal. BC to cal. AD 780-1035, the Early Mesolithic to the Medieval period. *Quercus* (oak) and *Ulmus* (elm) are well established from the base of the sequence and constitute the main tall canopy trees in the landscape around Lough Cullin. Corvlus avellana-type (probably largely hazel) is abundant, comprising a major component of the vegetation, probably as an under-storey shrub, while Betula (birch) and Salix (willow) had minor presences in the landscape. Low values for non-arboreal pollen (NAP),<5%, suggest the woodland had a rather closed structure. The presence of Hedera helix (ivy) and Polypodium vulgare (common polypody) probably indicate the presence of epiphytes (Molloy and O'Connell 1991).

291 LC-2: 6.03-5.88m, 7615-6500 cal. BC to 6245-5575 cal. BC

Zone LC-2 is marked by a decrease in hazel (c.50% to 30%) and oak (c.30% to 20%) and corresponding increases in *Pinus sylvestris* (Scots' pine) (c.4% to 20%) and birch (c.4% to 15%) between 5.97 to 5.93m. Hazel falls to its lowest value for the zone at 5.89m, corresponding to a peak in birch; whilst oak reaches its lowest percentage at 5.93m, at which point pine demonstrates a clear peak, attaining its highest values for the entire sequence before falling again. There are also reductions in elm and ivy apparent across these levels. Total NAP values do not show any fluctuations, suggesting the vegetation changes involved the woodland canopy taxa. Total pollen concentrations display a marked reduction across these levels, from 232x10³ to 155x10³ grains cm⁻³, which may reflect fluctuations in relative pollen productivity.

In terms of geochemistry, this zone sees the highest values for Ti, Ca and Mn for the entire diagram, tracking increasing values for MS, although LoI percentages remain fairly steady (c.30%). This pattern may indicate a change in the source of inorganic material eroded from the catchment, with weathering of the soils on the limestone (Ca) as well as the sandstone bedrock (Fe) implied. These fluctuations are closely associated with the changes in woodland composition outlined above. Taken collectively, the data may well indicate the impact of

climatic deterioration, with destabilised soils across the catchment associated with pronounced declines in thermophilous arboreal taxa (oak, hazel, elm and ivy), and related expansion in trees/shrubs (pine and birch) tolerant of colder conditions.

Towards the top of LC-2, Alnus glutinosa (alder) percentages begin to rise, implying the establishment of this tree, probably as alder carr at the lake edge. The date for the establishment of alder (5.89m, 6350-5910 cal. BC, the 'rational limit', sensu Smith and Pilcher 1978) appears to be slightly earlier than elsewhere across Ireland. This spread and expansion of alder post-7000 BP appears to have been erratic in space and time (Bennett and Birks 1990): for example, alder became established after 5490-5010 cal. BC (6315±110, D-115) at Belle Lake (Craig 1978), after 5480-5070 cal. BC (6330±80, Beta-65095) at Clara Bog (Connolly 1999; Crushell et al. 2008), whilst at Lough Kinale (Ballywillin Crannog core) this tree expanded after 5720-5560 cal. BC (6720±40, Beta-173320; Brown et al. 2005; Selby et al. 2005).

LC-3: 5.88-5.19m, 6245-5575 cal. BC to 4320-4020 cal. BC

By the close of LC-2, oak and hazel recovered as the dominant woodland taxa. Arboreal taxa remain very well represented in LC-3 (minimum 95%) with few other marked changes. However, *Ilex aquifolium* (holly) is recorded from midway through the zone (5.30m): this shrub tends to become established in more open conditions (Molloy and O'Connell 1987), but there is no evidence of significant changes to the woodland structure that might suggest disturbance similar to the previous zone. Indicators of open environments such as *Plantago lanceolata* (ribwort plantain), are present only sporadically prior to the establishment of holly, but cease during the initial expansion of the latter. The low levels of NAP (<5%) suggest open environments were limited, although the extent of openness will have been greater than implied by the percentages.

The LoI values do not show any marked changes, falling slightly towards the top of the zone. The geochemical data show reductions for Ca and Mn, although there are increases in Mn midway through the zone. Fluctuating values for Fe with pronounced decreases at the top and base of the zone are apparent. These fluctuations are slightly enigmatic but might imply some destablisation of the local soils, albeit not on a similar scale to LC-2.

338 LC-4a: 5.90-5.01m, 4320-4020 cal. BC to 4050-3960 cal. BC

Zone LC-4a is defined by a pronounced reduction in elm from 4220-3980 cal. BC, accompanied by a temporary fall in hazel, although other arboreal taxa remain steady. NAP increases (c.10%), primarily Poaceae (grass), but with sporadic grains of other herbs including ribwort plantain, Asteraceae (daisies) and Caryophyllaceae (pink family). There are decreases in LoI percentages across the zone coincident with increasing anthropogenic indicators, indicating increased mineral inwash into the lake and/or changes in accumulation rates associated with the reduction in woodland and expansion of open environments. This is supported by the rather abrupt fluctuations in Ti, Fe and Ca, indicating weathering and erosion in the lake catchment, again potentially involving soils on both the limestone and sandstone geologies.

350 LC-4b: 5.01-4.82m, 4050-3960 cal. BC to 3860-3650 cal. BC and LC4c: 4.82-4.71m, 3860351 3650 cal. BC to 3685-3530 cal. BC

Further reductions in arboreal pollen (AP), c. 80%, specifically oak and hazel are recorded from 4040-3910 cal. BC, with rises in NAP (c. 20%) and a greater diversity of ruderal herbaceous taxa including ribwort plantain, Ranunculaceae (buttercups), Rumex (docks), daisies, Artemisia-type (mugwort) and Urtica dioica (nettles). This indicates the expansion of open, pastoral environments, at the expense of deciduous woodland, especially oak but with hazel affected to a lesser degree. The identification of *Cerealia*-type (cereal) grains at: 3890-3700 cal. BC, 3790-3580 cal. BC and 3760-3560 cal. BC demonstrate arable farming in the pollen catchment. The LoI percentages show an abrupt drop to their lowest values for the entire sequence (4.90m) corresponding to a peak in total anthropogenic indicators, followed by an equally abrupt recovery, indicating inwash of minerogenic material. There are no clear corresponding changes in the geochemistry, other than some fluctuations in Ti, nor does the MS show any response.

This phase lasted until *3860-3650 cal. BC*, at which point AP began to recover (LC-4c). However, although there are reductions in NAP across this subzone, anthropogenic indicators including ribwort plantain and cereal-type are still recorded. The impression is of clearance of woodland and some destabilisation of soils during LC-4b, followed by a less intense period of human activity with some recovery of trees and shrubs in LC-4c. The initial phase lasted for

70-320 years, and the subsequent phase 10-270 years, with the pollen data implying both pastoral and arable land-use.

LC-5: 4.71-4.35m, 3685-3530 cal. BC to 3355-3120 cal. BC and LC-6; 4.38-4.08m, 3355-3120 *cal.* BC to 2960-2525 *cal.* BC

The overall AP values increase, especially elm, and the ribwort plantain curve is interrupted, reflecting woodland recovery. Elm populations regenerated, although not to their previous extent, while Fraxinus (ash) played an increased role in the arboreal vegetation dynamics, presumably expanding into the gaps created in the woodland in LC-4. This recovery of elm and associated woodland regeneration was sustained until 3310-3090 cal. BC, when a second reduction in elm is apparent, primarily concomitant with increased representation of alder, a slight increase in ash and marginally increased levels of NAP, primarily in wet-loving taxa such as Cyperaceae (sedges) and *Filipendula* (meadowsweet). This suggests the expansion of wetland environments, probably in the form of alder carr at the lake's edge, perhaps as processes of hydroseral succession created favourable habitats. The muted response in NAP aside from the wetland taxa, raises questions concerning the cause and process of the decline in elm (see below). LoI values are stable (c.36%) across LC-5 and LC-6 but decline towards the top of the latter zone (c.30%). Some fluctuations in Fe are recorded, possibly indicating changes in the source of the minerogenic input into the lake.

LC-7a: 4.08-3.90m, 2960-2525 cal. BC to 2620-2240 cal. BC

A small but pronounced reduction in AP and re-establishment of a continuous curve for ribwort plantain and other light demanding ruderal/grassland taxa (including buttercups, daisies, mugwort, Lactuceae (dandelions etc.) are apparent from 2960-2525 cal. BC, indicating an expansion of open, meadow environments. The woodland regeneration phase in the previous zones (LC-5 and 6) lasted between 660-1120 years. The appearance of cereal-type pollen at 2860-2420 cal. BC suggests arable farming around Lough Cullin. After the latter date, NAP continues to increase at the expense of primarily hazel and oak. Charcoal shows a slight increase and loss on ignition values are generally stable (c.30-34%); MS values increase coincident with a spike in Fe at c.4m, the highest values for anthropogenic indicators for this zone. These data collectively indicate catchment scale disturbance, related to the evidence for woodland reduction and the expansion of open environments.

HOLOCENE

401 LC-7b: 3.90-3.56m, 2620-2240 to 1880-1740 cal. BC and LC-7c: 3.56-3.34m, 1880-1740 to
402 1500-1410 cal. BC

Zone LC-7b is primarily defined on the basis of steadily increasing NAP, primarily grasses, with continuing low values of other herbs, suggesting open areas persisted but were of limited extent. Low values of microcharcoal are recorded, indicating restricted burning during this zone. This picture changes at the transition from LC-7b to LC-7c, around the Middle Bronze Age (1880-1740 cal. BC), with increases in ribwort plantain and the appearance of other anthropogenic indicators including dandelions and docks towards the close of the zone, reflecting an expansion in open, pastoral environments. However, percentages of arboreal pollen remain relatively steady, although there are fluctuations in hazel values. Charcoal values do not increase. The overall impression is of some expansion in open grassland environments, but only small decreases in woodland cover. The LoI values are largely stable, rising slightly across 7b (from c. 30 to 40%) before falling slightly across 7c. There are no clear changes in the MS or geochemical data other than a rise in Ca, possibly indicating disturbance on the limestone areas.

417 LC-8: 3.34-3.14m, *1500-1410* to *1275-1000 cal. BC*

The opening of LC-8 marks the beginning of a period of intensive human impact on the environment, the most pronounced recorded for the entire sequence. This seems to follow a brief period of reduced impact at the opening of the zone with reduced total anthropogenic indicators and increases in hazel, followed by a very pronounced increase in NAP (c. 45%), especially grasses, and associated decline in AP demonstrating a significant opening up of the woodland cover. The rising values for ribwort plantain (peaking c.7%) alongside grasses and to a lesser extent sedges, demonstrate the expansion of open, pastoral environments to form the greater proportion of the landscape around the lake. The occurrence of a single grain of cereal-type pollen at 3.20m (1380-1220 cal. BC) indicates arable farming. There are also reductions in alder, suggesting the alder carr on the wetter soils, probably close to the lake edges, were affected by clearance at this time.

However, a relatively restricted range of herbs is recorded: Chenopodiaceae (fat fen family),
 buttercups and *Succisa* (devil's bit scabious) appear alongside the peak of ribwort plantain, all
 of which are typical of open grassland/meadow environments. The concomitant increase in
 Pteridium aquilinum (bracken) across this zone is further evidence for the opening up of

the landscape on drier soils, whilst a spike in charcoal demonstrates increased burning. The
LoI percentages fall, generally tracking the increase in NAP, whilst in the geochemical data,
Fe also shows pronounced peaks at the opening and close of the zone, coincident with small
rises in Mn and Ti, and increases in the MS curves.

The combined data seem to indicate a short episode of reduced human activity and the recovery of shrubs, prior to intensive clearance and farming leading to soil disturbance across the catchment. Overall, the period between *1500-1410* and *1275-1000 cal. BC* was one of woodland reduction. Most parts of the landscape were affected, from the fertile dryland soils through to the 'marginal' wetland contexts. There is evidence for farming, predominantly of a pastoral character, and since cultivation is significantly underrepresented palynologically, arable plots were also present locally.

445 LC-9: 3.14-2.925m, *1275-1000 cal. BC* to *815-685 cal. BC* and LC-10: 2.925-2.82m, *815-685*446 *cal. BC* to *690-535 cal. BC*

The previous period of woodland clearance and farming was followed by the recovery of woodland taxa (50-60%), especially hazel, oak and alder, accompanied by reductions in grasses and ribwort plantain (LC-9). However, increased diversity in herbs is recorded, specifically the anthropogenic indicators fat hen family, nettles and dandelions. Micro-charcoal values also increase, whilst a single cereal grain coincides with the peak in ribwort plantain at *1020-915 cal. BC* indicating the presence of arable and pastoral habitats.

However, the subsequent zone, LC-10, is notable for a marked fall in ribwort plantain and recovery of AP, especially hazel and oak, but with elm values also forming a low but consistent curve for much of the zone. Elm tends to thrive on fertile, better- drained soils that are often suitable for arable agriculture (as observed in the previous zones), implying that trees and shrubs re-colonised much of the landscape. The almost complete absence of microcharcoal also contributes to the impression of little anthropogenic activity, until the close of the zone. There is a small but sustained increase in LoI across the zone, with Fe values reduced relative to the previous zone, suggesting some stabilisation of the catchment soils related to the recovery of the woodland.

HOLOCENE

464 LC-11: 2.82-2.64m, 690-535 cal. BC to 415-250 cal. BC and LC-12: 2.64-2.36m, 415-250 cal.
465 BC to 390-540 cal. AD

The landscape changed again at the opening of LC-11, with a fairly abrupt increase in ribwort plantain defining the zone transition, suggesting opening of the landscape. The response from other herbs, grasses included, is slightly muted, although levels of dandelions rise towards the close of the zone and there are occasional records of other grains such as mugwort, fat hen family, meadowsweet and buttercups. There is a steady reduction in hazel across the zone and oak also displays a fall, following a peak at the opening of the zone, contributing to falling AP from c.90% to 70% across the zone. Total anthropogenic indicators reach their highest percentage at the top of LC-11, at which point cereal-type pollen is recorded at 430-360 cal. BC. However, microcharcoal values show only modest increases. The record therefore indicates a resurgence in human activity, which led initially to an expansion in grassland and pasture and the clearance of areas of woodland, with evidence for arable cultivation towards the end of the zone.

Almost immediately following the peak in pastoral and arable farming, there is evidence for what can only be described, on the basis of the pollen record, as the apparent possibly complete cessation of human activity. This is manifested through an abrupt drop in ribwort plantain at the opening of LC-12, and the disappearance of this species for the entire zone. Grasses also display a slight reduction, whilst few other herbs are recorded. Microcharcoal values are minimal and absent from the middle section of the zone, closely paralleling the behaviour of total anthropogenic indicators. AP percentages increase, mainly hazel and oak, but with some recovery in elm and birch. Overall, from around 415-250 cal. BC to 390-540 cal. AD, woodland dominated the landscape around Lough Cullin.

488 LC-13: 2.36-2.15m, *cal. AD 390-540* to *cal. AD 780-1035*

The final zone, LC-13, opens with the relatively abrupt resumption of the ribwort plantain curve, a steady rise in grasses and reductions in hazel and subsequently oak. Elm effectively disappears from the record and microcharcoal values also increase. Other herbs reappear including docks, mugworts and daisies etc., all of which form continuous curves. Total anthropogenic indicators demonstrate a marked increase whilst cereal-type pollen is recorded in almost all levels, indicating a significant expansion and persistence of arable cultivation in the pollen source area. Despite this, the maintenance of AP (*c*.80%) until the close of the

diagram, suggests sizable areas of woodland remained locally. The diagram closes at *cal. AD 780-1035* with an indication of renewed falls in arboreal taxa suggesting further anthropogenic
pressure on the woodland. There is a hint that by this time, clearance of woodland was starting
to impact all parts of the landscape again, with alder falling to its lowest values since LPAZ
LC-2. The LoI values increase across the zone, with the rising values for Fe that began in LC12 also continuing.

503 6. Discussion

504 Overview

The Lough Cullin sequence provides a record of vegetation change from the Early Mesolithic (9050-8090 cal. BC) through to the Medieval period (cal. AD 780-1035), a period of 10020-9025 years. A series of phases of expansions and contractions of AP and NAP are apparent: with one exception (see below), these appear to reflect human disturbance, characterized by increases in taxa indicative of open, pastoral environments and disturbed soils (especially ribwort plantain, grasses, docks, dandelions and occasional grains of cereal pollen suggesting arable agriculture) and falls in values of tree/shrub pollen showing contraction in wood/scrubland.

It should be noted total tree and shrub pollen percentages rarely drop below 75% of total land pollen for the entire record, implying that although herbaceous taxa are probably underrepresented palynologically (Caseldine and Fyfe 2006), wood/scrub cover remained of greater extent than open ground, even during episodes of human impact and woodland clearance (see below). Episodes of intensified human activity seem to have impacted most on the populations of hazel and oak, with elm affected at some points, especially at the start of the Neolithic (4040-3900 cal. BC). The behaviour of Fraxinus (ash) suggests this fast growing and light demanding tree was favoured by the opening up of the woodland canopy during episodes of increased human activity (Caseldine and Hatton 1996).

The values for microcharcoal, an indicator of the frequency of burning in the landscape around the site (probably mainly from domestic fires, but potentially from natural burns), closely track these phases, with increased values associated with rises in pollen taxa indicating human activity. Microcharcoal falls noticeably during episodes of woodland recovery. The fluctuations in the LoI, MS and geochemical data provide additional information concerning the patterns of inwash and provenance of inorganic material into the Lough during the episodes

Page 19 of 49

HOLOCENE

of landscape disturbance. On the basis of these data, the most pronounced episodes of weathering at the catchment scale by far occurred during LC2, 4a, 7a and 7c, with the first presenting the clearest signal for the entire record.

532 The 8.2ka Event: Climatic forcing and vegetation change (LC-2)

The earlier Holocene landscape was dominated by deciduous woodland, consisting predominantly of hazel, oak and elm, with some birch and willow probably growing on the damper soils. Scots' pine was probably growing locally, despite the possibility of long-distance transport of this species' pollen (Pilcher *et al.* 1995, cf. Lageard *et al.* 1999). This limited representation is comparable with records from the east and midlands (e.g. Caseldine and Hatton 1996; Selby *et al.* 2005), although Scots' pine had become established locally at Kelly's Lough, Co. Wicklow (Leira *et al.* 2007) by 7450-7070 cal. BC (8220±50 BP, β -173463).

The abrupt reduction in hazel and oak in LC-2 indicates significant changes in this woodland composition, also reflected by a pronounced fall in pollen concentrations. Hazel pollen production can be adversely affected by climatic shifts such as air frosts, drought or excessive precipitation levels (Tallantire 2002), while oak is also sensitive to early spring and autumn frosts (Giesecke et al. 2008). On the other hand, pine and birch are tolerant of low summer temperatures and can be rapid colonisers, given suitable conditions (Atkinson 1992; Richardson and Rundel 1998; Paus 2010). The expansion of pine and birch alongside the small increases in NAP, and the fluctuations in the geochemical, LoI and MS data, can be interpreted as the impact of climatic deterioration, possibly a shift to a more continental climate, with colder winters but also lower summer temperatures and a considerably lower annual thermal sum (cf. Huntley 2012).

The date range at Lough Cullin (7120-6230 cal. BC to 6350-5910 cal. BC) correlates with climate anomalies in the Greenland ice-core records (Rasmussen et al. 2007; Thomas et al. 2007) between 8.5 and 7.9 kyr BP (Rohling and Pälike 2005). This climatic deterioration might have resulted from changes in the strength of the Atlantic meridional overturning circulation (cf. Alley et al. 1997; Alley and Ágústsdóttir 2005; Ellison et al. 2006; Hede et al. 2010), leading to colder and drier conditions in the North Atlantic region (Alley et al. 1997; Barber et al. 1999; Alley and Ágústsdóttir 2005; Wiersma and Renssen 2006). The clear palaeoenvironmental signature at Lough Cullin provides persuasive evidence for the impact of this climatic 'event' on the eastern as well as western seaboards of Ireland, with the chronology

of the increase in Scots' pine overlapping with dendrochronological evidence for expansion in
this tree after 6250 BC (Torbenson *et al.*, 2015).

Similar reductions in thermophilus taxa coincident with increases in cool-tolerant trees are exhibited in other Irish pollen records: Cooney Lough, County Sligo showed a marked reduction in pollen from hazel and oak, synchronous with expansion of pine and birch (Ghilardi and O'Connell 2012). Fluctuations in the frequency of juniper at An Loch Mór, Inis Oírr (Molloy and O'Connell 2004) and increased representations of pine at the expense of hazel are recorded during a period of erosion at Lough Maumeen, Connemara (Huang 2002).

569 The 'Elm Decline': Neolithic woodland clearance and agriculture

Mesolithic impacts on the environment have been identified in pollen records from the UK and Ireland (e.g. Smith 1970; Simmons and Innes 1996; Innes et al. 2003; Warren et al. 2014; Warren 2020), but there is little evidence for such 'disturbance events' in the Lough Cullin record, although the archaeological record indicates Mesolithic activity (Gleeson and Breen 2006a; 2006b; Wren 2006a; 2006b; Russell 2010). A continuous ribwort plantain curve and increased values for NAP (especially ruderal taxa) are evident from the Early Neolithic around 4040-3900 cal. BC, and some 10-260 years after the initial decline in elm at 4220-3980 cal. BC. This may indicate the 'elm decline' was unrelated to human activity, although rising values for grass and trace values for anthropogenic indicators are coeval with reductions in this tree. However, this might equally reflect open environments resulting from the mortality of elm trees due to disease (Casedline and Fyfe 2006). The fluctuations in the geochemical data, LoI and MS are distinct, if not as pronounced as in LC-2, indicating the decline in elm was coeval with disturbance to local soils and inwash of minerogenic matter into the Lough. Other Irish records (O'Connell and Molloy 2001; Ghilardi and O'Connell 2013; Molloy et al. 2014; Kearney and Gearey 2020; McClatchie and Potito 2020), also indicate a 'gap' between the 'Elm Decline' and clear palynological evidence for human disturbance. Overall, this may refute an anthropogenic cause for the reduction in elm, although the chronological disparity could also reflect increased stimulation of arboreal pollen production resulting from openings to the woodland canopy (cf. Aaby 1986).

The Early Neolithic (LC-4a) is marked by the continuous presence of ribwort plantain from
 4040-3900 cal. BC until 3720-3540 cal. BC, a duration of 220-450 years. During this period,
 there was a substantial opening up of the woodland and expansion in open habitats, presumably

relatively close to the lake, although this is considerably earlier than the estimated date (3720-3680 cal. BC) for the Neolithic 'house horizon' and the first cereal cultivation in Ireland (McSparron 2008; Cooney et al. 2011; McClatchie et al. 2014; Whitehouse et al. 2014; McLaughlin et al. 2016). Overall, the impact of Early Neolithic agriculture in Irish pollen records is rather variable (Connolly 1999; Crushell et al. 2008; O'Carroll 2012), perhaps reflecting the fact that Early Neolithic agriculture was spatially heterogeneous (Whitehouse et al. 2014) with varying levels of intensity, and the relative proximity of activity to sampling sites would further attenuate the signal.

Cereal-type pollen is recorded at 3900-3700 cal. BC, an estimated 50-290 years after the start of clearance and again from 3790-3580 cal. BC. Palynological evidence from Newrath, south of Lough Cullin (Timpany 2009) also strongly suggests Neolithic arable agriculture at this time. The most reliable evidence is the presence of charred cereal grain in Neolithic archaeological contexts (e.g Hughes 2006; Monteith 2009; 2011; McKinstry 2010). The chronological model estimates farming continued for 220-450 years, implying Early Neolithic farmers did not engage in shifting cultivation contra Iversen's (1941) Landnam model, but rather longer-term, fixed-plot agriculture (Whitehouse et al. 2014; McClatchie and Potito, 2020).

This phase of agriculture was followed by woodland regeneration and a decline in anthropogenic indicators from 3720-3540 cal. BC. A similar pattern of re-afforestation is implied during the Middle Neolithic in other parts of Ireland (e.g. Ghilardi and O'Connell 2013; Molloy et al. 2014; Chique et al. 2017; Stolze and Monecke 2020). It has been suggested the Middle Neolithic saw a reduction in human activity overall (Whitehouse et al. 2014), rather than a shift in location, as reflected by the archaeological record: the number of confirmed Middle Neolithic archaeological sites in the region is substantially lower than in the preceding Early Neolithic, also in keeping with the archaeological evidence from across Ireland. This has also been linked to climatic change, dubbed the '5.2 ka event' (Roland et al. 2015), although the precise nature and timing of this remains unclear (Plunkett et al., 2020).

Evidence for sustained human activity is subsequently absent until 2960-2525 cal. BC, a period of woodland regeneration of estimated 660-1120 years, followed by a renewal of arable and pastoral activity. The reappearance of agricultural activity during the Late Neolithic is indicated in other pollen records (Molloy and O'Connell 2004; Molloy et al. 2014; Stolze and Monecke 2020) whilst the archaeological record also suggests a limited Late Neolithic presence in south-east Ireland (Tierney 2005; Wren 2006b; Monteith 2011).

The reduction in Scots pine from 2620-2240 cal BC, probably coincides with the beginnings of the 'Pine Decline' in the region. The decline of Scots pine has been well documented across much of Ireland, northern Scotland and England (e.g. Bennett 1984; Bridge et al. 1990; Gear and Huntley 1991; Pilcher et al. 1995; Lageard et al. 1999; Mighall et al. 2004), although recent studies have suggested the persistence of the taxa in certain regions (McGeever and Mitchell 2016; Sassoon et al. 2021). Numerous competing hypotheses have been proposed for the cause of this reduction, from climate (Bradshaw and Browne 1987; Mighall et al. 2004) to anthropogenic activity (Tipping et al. 2008), however Lough Cullin provides no conclusive evidence for the cause of this at the site.

635 Human activity from The Bronze Age to Early Medieval Period.

Relatively low levels of human activity are indicated during the Late Neolithic (LC-7a) through to the Chalcolithic/Early Bronze Age (LC-7b), followed by resurgence into the Middle Bronze Age (LC-7c), peaking in the Middle to Late Bronze Age (from 1500-1410 cal. BC to 1275-1000 cal. BC). Excursions in the geochemical data, especially Ca during LC-7c, followed by increases in Fe and associated peaks in MS towards the top of LC-8, support the picture that the Middle Bronze Age saw significant scrub/woodland clearance, farming and associated destabilisation of soils. Following this, human activity then seems to have decreased in two 'steps': an initial reduction, which may be described as a contraction, during the Late Bronze Age from 1275-1000 cal. BC to 815-685 cal. BC (LC-9) followed by a pronounced 'collapse' from 815-685 cal. BC until 690-535 cal. BC (LC-10), the Bronze Age to Iron Age transition. Human impact on the landscape seems to have almost completely ceased, quite abruptly in this latter period.

This pattern is paralleled by the changes in archaeological site distributions across this entire period (cf. Armit et al. 2013.; Becker et al. 2017); extensive landscape utilisation until the beginning of the Hallstatt plateau, followed by a marked contraction. This characteristic pattern can also be traced in the archaeology in the southeast (Eogan et al. 2015). Comparison with data from recently published Irish pollen records indicates a broadly coherent pattern during the Early to Middle Bronze Age of consistent and increasing woodland clearance and agriculture, followed by fluctuating human activity during the Late Bronze Age (Chique et al. 2017; Spencer et al. 2020; Stolze & Monecke, T. 2020). tt (2009) has previously drawn attention to a similar pattern, suggesting a link between increased archaeological visibility during the Late Bronze Age and palynological evidence for intensified pastoral activity, and

HOLOCENE

indicating that the end of this period may have been characterised by consolidation ofsettlement and agriculture.

Climate change and human activity during Later Prehistory

It has long been suggested there is an apparent coincidence between the 'end' of the Bronze Age (ca. 1150-800/600 cal. BC) and evidence for climatic deterioration (wetter/colder conditions) across northwest Europe; the period known as the 'Sub-Boreal to Sub-Atlantic transition' or the '2.8 Ka event' (see Gearey et al., 2020 for a recent review). Some scholars have suggested a direct causal link, with the onset of wetter/colder conditions, impacting directly on Bronze Age settlement and agriculture. However, Armit et al. (2014) have presented combined palaeodemographic (Armit et al. 2013) and hydroclimatic data (peatland Bog Surface Wetness/BSW records; Swindles et al. 2013) from Ireland, arguing a 'peak' in human activity c.1050 to 900 BC, was followed by a reduction to c.800 BC and a rapid 'collapse' to c.750 BC. The combined BSW data indicate climatic deterioration c.750-550 BC, which may suggest that this 'event' can be causally 'disconnected' from the 'end' of the Bronze Age, at least as indicated by the proxy demographic data.

There are few comparative palaeotemperature datasets for this period; although a recent chironomid-based reconstruction (expressed as mean July air temperature, referred to as Chironomid-inferred temperatures: C-ITs) is available from Lough Meenachrinna, Co. Donegal (Taylor et al. 2018). These data indicate C-ITs of 11.5°C during the Early Bronze Age (c.1390 BC), rising into the Middle Bronze Age (c.1310-1100 BC) to peak at 13.5 °C (c.1260 BC), then falling to c.12 °C at the beginning of the Late Bronze Age (c.1150-960 BC). Slight increases in C-ITs to 12.8 °C are recorded from the end of the Bronze Age into the Iron Age (c.920-620 BC), later dropping to 11.4 °C (c.660-480 BC). The range of palaeotemperature fluctuations in this record are relatively small and their potential significance for human activity hard to assess (K. Taylor, pers.comm.)

However, 'events' in the Lough Cullin pollen record can be described as chronologically 'imbricated' with the palaeoclimatic 'events': a clear peak in human activity during the Middle Bronze Age, followed by contraction or consolidation during the Late Bronze Age and a pronounced fall into the Iron Age. The initial reduction (LC-9) also overlaps with the beginning of the palaeodemographic drop (Armit et al. 2014) and the subsequent 'collapse' (LC-10) with the postulated climatic deterioration in the BSW records (see also Plunkett et al., 2020 who

point to a gap in germination in the bog oak record between 751-713 BC and 709-668 BC). Correlations are hampered by the variable chronological precision of the different records; even with the improved chronological precision of the Cullin record, the ranges for the palynological 'events' are still 'smeared' over several centuries. Nevertheless, the chronological correspondence between the postulated climatic downturn (c.750 BC) in the BSW records and the contraction of human activity during the Bronze Age-Iron Age transition (690-535 cal. BC) at Lough Cullin can be noted; at the very least leaving open the possibility of a link between the two. Further analysis is also hampered by a general lack of detailed knowledge of the precise nature or severity of climatic changes, and an associated lack of clarity concerning pattern and process of social-cultural changes from Bronze to Iron Age (see Gearey et al., 2020; Coyle-McClung and Plunkett, 2020 for further discussion).

Whatever the cause/s of the decline in anthropogenic activity during the transitional period between Bronze and Iron Ages, two 'steps' can be defined: a rapid recovery following the previous 'collapse' and marked expansion in woodland clearance and farming very soon afterwards, between 690-535 cal. BC and 415-250 cal. BC (LC-11). This was followed by another phase of woodland recovery from the latter date until 390-540 cal. AD (LC-12). A marked period of woodland regeneration is a feature of many Irish pollen records during later prehistory, sometimes referred to as the 'Late Iron Age Lull' (Mitchell and Ryan 1997) although the reduction in activity dates to the Developed Iron Age at Lough Cullin. The timing of woodland regeneration and reduced human activity generally varies between records, from the final centuries BC to the early centuries AD (Chique *et al.* 2017). In some locations, human activity continued, albeit at fairly low levels, across the Late Iron Age (Figure 6). There are indications of drier/warmer conditions in Ireland from c.300 BC-400 AD (Coyle-McClung and Plunkett, 2020), corresponding broadly with the woodland regeneration phase (LC-11) in the Cullin record.

At Lough Cullin, human activity seems to have ceased almost completely until cal. AD 390-540, when there was a relatively abrupt resumption of the ribwort plantain curve, a steady rise in grass and reductions in hazel and oak (LC-13). Anthropogenic indicators demonstrate a marked increase and cereal pollen is recorded in almost all the samples across the zone, indicating a significant expansion of arable cultivation locally. In comparison to the sporadic record of cereal pollen from the rest of the sequence, the Early Medieval period saw the most extensive period of arable cultivation in the Lough pollen catchment. A steady rise in Fe and Ca towards the close of the zone, a peak in MS track rising NAP; the increasing LoI values is

Page 25 of 49

HOLOCENE

contrary to the evidence for increased mineral input, but may reflect increased eutrophication
of the lake due to the input of organic material from intensified arable and pastoral farming;
the record terminates at *cal. AD* 780-1035 with increasing anthropogenic indicators and NAP
and associated reductions in AP.

7. Conclusions

The Lough Cullin sequence provides a nuanced and detailed multi-proxy (pollen, loss-on-ignition, geochemistry and magnetic susceptibility) reconstruction of Holocene environmental and landscape change from 9050-8090 cal. BC to 780-1035 cal. AD, the Mesolithic to the Early Medieval Period; the provision of a high resolution radiocarbon chronology has allowed a comparatively high level of chronological precision for the later prehistoric period in particular, using Bayesian approaches permitting formal, robust estimates of the timing and duration of particular events, avoiding potentially misleading precision resulting from other approaches to age-depth modelling of palaeoenvironmental records. The data provide clear evidence for the impact of the '8.2 Kyr' BP climate event on the environment of the southeastern seaboard of Ireland. Certainly, on the basis of the geochemistry and magnetic susceptibility datasets, the 'signature' of the 8.2 'event' is very pronounced, which is interpreted as reflecting the impact of this colder/drier period across the entire catchment. Later alleged prehistoric climatic 'events' are largely obscured or overlain by human impact, reflecting woodland clearance, settlement and agriculture which began during the Early Neolithic, slackened during the Middle Neolithic before intensifying in the Middle-Late Bronze Age. The compilation of archaeological data has allowed a qualitative comparison of the palaeoenvironmental record with the spatially delimited evidence of human activity and taking account of the uncertainties associated with the interpretation of both records, broad correspondence in terms of implied intensities of human activity are indicated. A steady 'demise' of human activity during the Late Bronze Age-Iron Age is apparent, although there is no clear evidence to illuminate the cause/s of this, there is chronological correspondence with alleged national climatic deterioration around the mid-8th century BC. There is also unambiguous evidence for a reduction in human activity during the Iron Age until the Early Medieval period, providing further indication of the spatial and chronological variability of settlement and agriculture in Ireland across these periods. Despite ongoing archaeological and palaeoenvironmental investigation of the later prehistoric period in particular, uncertainties concerning data interpretation and variations in the chronological resolution, precision and robustness of different proxy climate records,

⁷⁵⁶ hamper the formulation of mechanistic models linking environmental and cultural processes.

The development of new methods and the generation of further, high resolution
palaeoenvironmental records are one key to future progress, but this must include further
research into archaeological data on regionally specific scales.

References.

- 762 Ordnance Survey of Ireland (no date) Name book, Dunkitt parish.
- Aaby, B. 1986. Trees as anthropogenic indicators in regional pollen diagrams from eastern
 Denmark. In Behre, K. E. (ed.) *Anthropogenic indicators in pollen diagrams*. Balkema,
 Rotterdam, p.73-93.
- Alley, R. B. & Ágústsdóttir, A. M. 2005. The 8k event: cause and consequences of a major
 Holocene abrupt climate change. *Quaternary Science Reviews*, 24, p.1123-1149.
- Alley, R. B., Mayewski, P. A., Sowers, T., Stuiver, M., Taylor, K. C. & Clark, P. U. 1997.
 Holocene climatic instability: A prominent, widespread event 8200 yr ago. *Geology*, 25, p.483770 486.
- Armit, I., Swindles, G. and Becker, K. 2013 From dates to demography in later prehistoric Ireland? Experimental approaches to the meta-analysis of large 14 C data-sets. *Journal of Archaeological Science* 40, p.433-438.
- Armit, I., Swindles, G. T., Becker, K., Plunkett, G. & Blaauw, M. 2014. Rapid climate change
 did not cause population collapse at the end of the European Bronze Age. *Proceedings of the National Academy of Sciences*, 111, p.17045-17049.
- 777 Atkinson, M. D. 1992. Betula pendula Roth (*B. verrucosa Ehrh.*) and B. pubescens Ehrh.
 778 *Journal of Ecology*, 80, p.837-870.
- ⁸ 779 Barber, D. C., Dyke, A., Hillaire-Marcel, C., Jennings, A. E., Andrews, J. T., Kerwin, M. W.,
- Bilodeau, G., McNeely, R., Southon, J., Morehead, M. D. & Gagnon, J. M. 1999. Forcing of
 the cold event of 8,200 years ago by catastrophic drainage of Laurentide lakes. *Nature*, 400,
 p.344-348.
- Bayliss, A., Bronk Ramsey, C., van der Plicht, J. & Whittle, A. 2007. Bradshaw and Bayes:
 Towards a timetable for the Neolithic. *Cambridge Archaeological Journal*, 17, p.1-28.

HOLOCENE

Becker, K., Armit, I. and Swindles, G. 2017. New perspectives on the Irish Iron Age: The
impact of NRA development on our understanding of Later Prehistory. In: Stanley, M (eds). *Stories of our past*. Bray: Wordwell Books
Behre, K. E. 1981. The interpretation of anthropogenic indicators in pollen diagrams. *Pollen et Spores*, 23, p.225-245.

- Bengtsson, L. & Enell, M. 1986. Chemical analysis. In Berglund, B. E. (ed.) *Handbook of Holocene Palaeoecology and Palaeohydrology*. Whiley, Chichester, p.423-445.
- 7 792 Bennett, K. D. 1984. The post-Glacial history of *Pinus sylvestris* in the British Isles. *Quaternary* 8 793 Science Reviews 3, 133–155.
- 1 794 Bennett, K. D. 1995. *Pollen Catalogue of the British Isles* [Online]. Available: 2 795 <u>http://chrono.qub.ac.uk/pollen/pc-intro.html</u> [Accessed 11 February 2019].
- 796 Bennett, K. D. & Birks, H. J. B. 1990. Postglacial history of alder (*Alnus glutinosa (L.) Gaertn.*)
 797 in the British Isles. *Journal of Quaternary Science*, 5, p.123-133.
- ⁹ 798 Bennett, K. D., Whittington, G. and Edwards, K. J. 1994 Recent plant nomenclature changes
 ⁰ and pollen morphology in the British Isles. *Quaternary Newsletter* 73, p.1-6.
- ² 800 Blaauw, M., Bakker, R., Christen, J. A., Hall, V. A. & van der Plicht, J. 2007a. A Bayesian
 ⁴ 801 framework for age modeling of radiocarbon-dated peat deposits: Case studies from the
 ⁶ 802 Netherlands. *Radiocarbon*, 49, p.357-367.
- ⁸ 803 Blaauw, M. & Christen, J. A. 2005. Radiocarbon peat chronologies and environmental change.
 ⁹ 804 *Journal of the Royal Statistical Society Series C-Applied Statistics*, 54, p.805-816.
- ¹² 805 Blaauw, M., Christen, J. A., Mauquoy, D., van der Plicht, J. & Bennett, K. D. 2007b. Testing
 ¹³ 806 the timing of radiocarbon-dated events between proxy archives. *The Holocene*, 17, p.283-288.
- ⁴⁶ 807 Blockley, S. P. E., Blaauw, M., Bronk Ramsey, C. & van der Plicht, J. 2007. Building and
 ⁴⁸ 808 testing age models for radiocarbon dates in Lateglacial and Early Holocene sediments.
 ⁴⁹ 809 *Quaternary Science Reviews*, 26, p.1915-1926.
- Blockley, S. P. E., Ramsey, C. B., Lane, C. S. & Lotter, A. F. 2008. Improved age modelling
 approaches as exemplified by the revised chronology for the Central European varved lake
 Soppensee. *Quaternary Science Reviews*, 27, p.61-71.
- ⁵⁷ 813 Bradshaw, R. H. W. & Browne, P. 1987. Changing patterns in the post-Glacial distribution of
- Pinus sylvestris in Ireland. Journal of Biogeography 14, 237–248.

1 2		
2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21	815	Bridge, M. C., Haggart, B. A. & Lowe, J. J. 1990. The history and palaeoclimatic significance
	816	of subfossil remains of Pinus Sylvestris in blanket peats from Scotland. Journal of Ecology 78,
	817	77–99.
	818	Brock, F., Lee, S., Housley, R. A. & Bronk Ramsey, C. 2011. Variation in the radiocarbon age
	819	of different fractions of peat: A case study from Ahrenshöft, northern Germany. Quaternary
	820	Geochronology, 6, p.550-555.
	821	Bronk Ramsey, C. 2008. Deposition models for chronological records. Quaternary Science
	822	<i>Reviews</i> , 27, p.42-60.
	823	Bronk Ramsey, C. 2009a. Bayesian analysis of radiocarbon dates. Radiocarbon, 51, p.337-360.
	824	Bronk Ramsey, C. 2009b. Dealing with Outliers and Offsets in Radiocarbon Dating.
22 23	825	<i>Radiocarbon</i> , 51, p.1023-1045.
24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40	826	Brown, A. G., Hatton, J., O'Brien, C. E., Selby, K. A., Langdon, P. G., Stuijts, I. & Caseldine,
	827	C. J. 2005. Vegetation, landscape and human activity in midland Ireland: Mire and lake records
	828	from the Lough Kinale-Derragh Lough area, central Ireland. Vegetation History and
	829	Archaeobotany, 14, p.81-98.
	830	Buck, C. E., Litton, C. D. & Smith, A. F. M. 1992. Calibration of radiocarbon results related
	831	to archaeological events. Journal of Archaeological Science, 19, p.497-512.
	832	Caseldine, C. J. & Fyfe, R. M. 2006. A modelling approach to locating and characterising elm
	833	decline/landnam landscapes. Quaternary Science Reviews, 25, p.632-644.
	834	Caseldine, C. J. & Hatton, J. 1996. Early land clearance and wooden trackway construction in
41 42	835	the third and fourth Millennia BC at Corlea, Co. Longford. Proceedings of the Royal Irish
43 44	836	Academy. Section B: Biological, Geological, and Chemical Science, 96, p.11-19.
45 46	837	Chique, C., Molloy, K. & Potito, A. 2017. Mid-Late Holocene Vegetational History and Land-
47 48	838	Use Dynamics in County Monaghan, Northeastern Ireland-The Palynological Record of Lough
49 50	839	Muckno. Journal of the North Atlantic, 32, p.1-24.
51 52	840	Connolly, A. 1999. The Palaeoecology of Clara Bog, County Offaly. Unpublished PhD thesis,
52 53 54 55 56	841	Trinity College, Dublin.
	842	Cook, G. T., Dugmore, A. J. & Shore, J. S. 1998. The influence of pretreatment on humic acid
57 58	843	yield and 14C age of Carex peat. Radiocarbon, 40, p.21-27.
59 60		

HOLOCENE

2		
3 4	844	Cooney, G., Bayliss, A., Healy, F., Whittle, A., Danaher, E., Cagney, L., Mallory, J., Smyth,
5	845	J., Kador, T. & O'Sullivan, M. 2011. Ireland. In Whittle, A., Healy, F. & Bayliss, A. (eds.)
6 7	846	Gathering Time: Dating the Early Neolithic Enclosures of Southern Britain and Ireland.
8 9 10 11 12 13	847	Oxbow, Oxford, p.562-669.
	848	Coyle-McClung, L. & Plunkett, G. 2020. Cultural change and the climate record in final
	849	prehistoric and early medieval Ireland. Proceedings of the Royal Irish Academy: Archaeology,
14 15	850	Culture, History, Literature, 120C, p.129-158.
16 17	851	Craig, A. J. 1978. Pollen Percentage and Influx Analyses in South-East Ireland: A Contribution
17 18 19	852	to the Ecological History of the Late-Glacial Period. Journal of Ecology, 66, p.297-324.
20 21 22	853	Croudace, I.W., Rindby, A. and Rothwell, R.G., 2006. ITRAX: description and evaluation of
	854	a new multi-function X-ray core scanner. Geological Society, London, Special
23 24	855	<i>Publications</i> , 267, p.51-63.
25 26 27 28 29 30 31 32	856	Crushell, P., Connolly, A., Schouten, M. & Mitchell, F. J. G. 2008. The changing landscape of
	857	Clara Bog: the history of an Irish raised bog. Irish Geography, 41, p.89-111.
	858	Eogan, J., Becker, K., McClatchie, M., Armit, I., Nagle, C. & Gearey, B. 2015. The Prehistory
	859	of the Southeast. In McGlynn, G. & Stefanini, B. (eds.) The Quaternary of South East Ireland:
33 34	860	A filed guide. Quaternary Research Association and the Irish Quaternary Association, Dublin,
35 36 37 38	861	p.35-43.
	862	Ellison, C. R. W., Chapman, M. R. & Hall, I. R. 2006. Surface and Deep Ocean Interactions
39 40	863	During the Cold Climate Event 8200 Years Ago. Science, 312, p.1929-1932.
40 41 42	864	Evans, M. and Slaymaker, O., 2004. Spatial and temporal variability of sediment delivery from
43	865	alpine lake basins, Cathedral Provincial Park, southern British Columbia. Geomorphology, 61,
44 45	866	p.209-224.
46 47	867	Fægri, K., Iversen, J., Kaland, P. & Krywinski, K. 1989. Textbook of pollen analysis, Wiley,
48 49 50 51 52 53	868	Chichester.
	869	Gear, A. J. & Huntley, B. 1991. Rapid changes in the range limits of scots pine 4000 years
	870	ago. Science 251, 544–547.
54 55	871	Gearey, B. R., Becker, K., Everett, R. & Griffiths, S. 2020. On the brink of Armageddon?
56 57 58	872	Climate change, the archaeological record and human activity across the Bronze Age-Iron Age
	873	Transition in Ireland. Proceedings of the Royal Irish Academy: Archaeology, Culture, History,
59 60	874	<i>Literature</i> , 120C, p.105-128.

Ghilardi, B. & O'Connell, M. 2012. Early Holocene vegetation and climate dynamics with particular reference to the 8.2 ka event: pollen and macrofossil evidence from a small lake in western Ireland. Vegetation History and Archaeobotany, 22, p.99-114. Ghilardi, B. & O'Connell, M. 2013. Fine-resolution pollen-analytical study of Holocene woodland dynamics and land use in north Sligo, Ireland. Boreas, 42, p.623-649. Giesecke, T., Bjune, A. E., Chiverrell, R. C., Seppä, H., Ojala, A. E. K. & Birks, H. J. B. 2008. Exploring Holocene continentality changes in Fennoscandia using present and past tree distributions. Quaternary Science Reviews, 27, p.1296-1308. Gleeson, C. & Breen, G. 2006a. Final Report on archaeological investigations at Sites 1 and 2, in the townland of Granny, Co. Kilkenny. (04E0226). Unpublished excavation report. Headland Archaeology Ltd. on behalf of Kilkenny County Council and the National Roads Authority. Gleeson, C. & Breen, G. 2006b. Final Report on archaeological investigations at Sites 21-23, in the townland of Granny, Co. Kilkenny. (04E0200). Unpublished excavation report. Headland Archaeology Ltd. on behalf of Kilkenny County Council and the National Roads Authority. Gregory, B.R., Patterson, R.T., Reinhardt, E.G., Galloway, J.M. and Roe, H.M., 2019. An evaluation of methodologies for calibrating Itrax X-ray fluorescence counts with ICP-MS concentration data for discrete sediment samples. Chemical Geology, 521, p.12-27. Grimm, E. 2013. TILIA and TGView software, Springfield Museum, Illinois. Hede, M. U., Rasmussen, P., Noe-Nygaard, N., Clarke, A. L., Vinebrooke, R. D. & Olsen, J. 2010. Multiproxy evidence for terrestrial and aquatic ecosystem responses during the 8.2 ka cold event as recorded at Højby Sø, Denmark. Quaternary Research, 73, p.485-496. Guyard, H., Chapron, E., St-Onge, G., Anselmetti, F.S., Arnaud, F., Magand, O., Francus, P. and Mélières, M.A., 2007. High-altitude varve records of abrupt environmental changes and mining activity over the last 4000 years in the Western French Alps (Lake Bramant, Grandes Rousses Massif). Quaternary Science Reviews, 26, p.2644-2660. Heiri, O., Lotter, A. F. & Lemcke, G. 2001. Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. Journal of Paleolimnology, 25, p.101-110.

Huang, C. C. 2002. Holocene landscape development and human impact in the Connemara
uplands, western Ireland. *Journal of Biogeography*, 29, p.153-165.

907 Hughes, J. 2006. N25 Waterford bypass, contract 3. Final report on archaeological
908 investigations at sites 24-30 in the townland of Granny, County Kilkenny. (04E0548).
909 Unpublished excavation report. Headland Archaeology Ltd. on behalf of Kilkenny County
910 Council and the National Roads Authority.

911 Huntley, B. 2012. Reconstructing palaeoclimates from biological proxies: Some often
 912 overlooked sources of uncertainty. *Quaternary Science Reviews*, 31, p.1-16.

913 Innes, J. B., Blackford, J. J. & Rowley-Conwy, P. 2003. The start of the Mesolithic-Neolithic
 914 transition in north-west Europe: The palynological contribution. *Antiquity*, 77(297).

915 Iversen, J. 1941. Landnam i Danmarks Stenalder: en pollenanalytisk undersøgelse over det
 916 første landbrugs indvirkning paa vegetationsudviklingen (Land occupation in Denmark's stone
 917 age: a pollen-analytical study of the influence of farmer culture on the vegetational
 918 development). Danmarks geologiske undersøgelse, II, p.1-68.

Jouve, G., Francus, P., Lamoureux, S., Provencher-Nolet, L., Hahn, A., Haberzettl, T., Fortin,
 D., Nuttin, L. and Team, T.P.S., 2013. Microsedimentological characterization using image
 analysis and μ-XRF as indicators of sedimentary processes and climate changes during
 Lateglacial at Laguna Potrok Aike, Santa Cruz, Argentina. *Quaternary Science Reviews*, 71,
 p.191-204.

9 924 Kearney, K. & Gearey, B. R. 2020. The Elm Decline is dead! Long live declines in elm:
 925 revisiting the chronology of the Elm Decline in Ireland and its association with the
 926 Mesolithic/Neolithic transition. *Environmental Archaeology*.
 927 10.1080/14614103.2020.1721694

928 Keeley M.L. (1983) The Stratigraphy of the Carrick-on-Suir Syncline, Southern Ireland.
 929 *Journal of Earth Science Royal Dublin Society*, 5, p.107-120.

930 Kelly, J. & O'Carragain, T. (eds.) 2021. *Climate and society in Ireland: From prehistory to* 931 *present.* The Royal Irish Academy, Dublin.

Lageard, J. G. A., Chambers, F. M. & Thomas, P. A. 1999. Climatic significance of the
marginalization of scots pine (*Pinus sylvestris L.*) c.2508 BC at White Moss, south Cheshire,
UK. *The Holocene*, 9, p.321-331.

Leira, M., Cole, E. E. & Mitchell, F. J. G. 2007. Long term impacts of atmospheric deposition

 and peat erosion on an oligotrophic lake in eastern Ireland. Journal of Paleolimnology, 38, p.49-71. McClatchie, M., Bogaard, A., Colledge, S., Whitehouse, N. J., Schulting, R. J., Barratt, P. & McLaughlin, T. R. 2014. Neolithic farming in north-western Europe: Archaeobotanical evidence from Ireland. Journal of Archaeological Science, 51, p.206-215. McClatchie, M. & Potito, A. 2020. Tracing environmental, climatic and social change in Neolithic Ireland. Proceedings of the Royal Irish Academy: Archaeology, Culture, History, Literature, 120C, p.23-50. McGeever A.H. & Mitchell F.J. 2016. Redefining the natural range of Scots Pine (Pinus sylvestris L.): a newly discovered microrefugium in western Ireland. Journal of Biogeography 43, 2199-2208. McKinstry, L. 2010. N9/N10 Kilcullen to Waterford Scheme: Waterford to Knocktopher -Phase 2 Archaeological Resolution, Dunkitt to Sheepstown Co. Kilkenny Final Report A032/000, E3005 Site AR031, Earlsrath, Co. Kilkenny. Unpublished excavation report. Valerie J Keeley Ltd. on behalf of Kilkenny County Council and the National Roads Authority. McLaughlin, R. T., Whitehouse, N. J., Schulting, R. J., McClatchie, M., Barratt, P. & Bogaard, A. 2016. The changing face of Neolithic and Bronze Age Ireland: A big data approach to the settlement and burial Records. Journal of World Prehistory, 29, p.117-153. McSparron, C. 2008. Have you no homes to go to? Neolithic housing. Archaeology Ireland, 22, p.18-21. Mighall, T. M., Lageard, J. G. A., Chambers, F. M., Field, M. H. & Mahi, P. 2004. Mineral deficiency and the presence of *Pinus sylvestris* on mires during the mid to late Holocene: Palaeoecological data from Cadogan's Bog, Mizen Peninsula, Co. Cork, southwest Ireland. The Holocene 14, 95-109. Mitchell, F. & Ryan, M. 1997. Reading the Irish Landscape. Tower House, Dublin. Molloy, K., Feeser, I. & O'Connell, M. 2014. A pollen profile from Ballinphuill bog: Vegetation and land-use history. In McKeon, J. & O'Sullivan, J. (eds.) The quiet landscape: Archaeological and palaeoecological investigations on the M6 Galway to Ballinasloe motorway scheme. NRA Scheme Monographs 15 (on CD Rom). The National Roads Authority, Dublin.

Page 33 of 49

HOLOCENE

Molloy, K. & O'Connell, M. 1987. The nature of the vegetational changes at about 5000 B.P. with particular reference to the Elm Decline: Fresh evidence from Connemara, western Ireland. New Phytologist, 107, p.203-220. Molloy, K. & O'Connell, M. 1991. Palaeoecological investigations towards the reconstruction of woodland and land-use history at Lough Sheeauns, Connemara, western Ireland. Review of Palaeobotany and Palynology, 67, p.75-113. Molloy, K. & O'Connell, M. 2004. Holocene vegetation and land-use dynamics in the karstic environment of Inis Oirr, Aran Islands, western Ireland: Pollen analytical evidence evaluated in light of the archaeological record. Quaternary International, 113, p.41-64. Monteith, J. 2009. Kilkeasy. In Bennett, I. (ed.) Excavations 2006: summary accounts of archaeological excavations in Ireland. Wordwell, Bray, p.269-270. Monteith, J. 2011. N9/N10 Kilcullen to Waterford Scheme: Waterford to Knocktopher - Phase 2. Archaeological Resolution, Dunkitt to Sheepstown Co. Kilkenny. Final Report Scart, Co. Kilkenny. (E3001). Unpublished excavation report. Valerie J. Keeley Ltd. on behalf of Kilkenny County Council and the National Roads Authority. Moore, P. D., Webb, J. A. & Collinson, M. E. 1991. Pollen analysis. Blackwell, Oxford. O'Carroll, E. 2012. Quantifying woodland resource usage in the Irish midlands using archaeological and palaeoecological techniques. Unpublished PhD thesis, University of Dublin Trinity College. O'Connell, M. & Molloy, K. 2001. Farming and woodland dynamics in Ireland during the Neolithic. Proceedings of the Royal Irish Academy. Section B: Biological, Geological, and *Chemical Science*, 101, p.99-128. Paus, A. 2010. Vegetation and environment of the Rødalen alpine area, Central Norway, with emphasis on the early Holocene. Vegetation History & Archaeobotany, 19, p.29-51. Pilcher, J. R., Baillie, M. G. L., Brown, D. M., McCormac, F. G., Macsweeney, P. B. & McLawrence, A. S. 1995. Dendrochronology of subfossil pine in the north of Ireland. Journal of Ecology, 83, p.665-671.

Plunkett, G. 2009. Land-use patterns and cultural change in the Middle to Late Bronze Age in Ireland: inference from pollen records. Vegetation History & Archaeobotany, 18, p.273-295.

995 Plunkett, G., Brown, D. M. & Swindles, G. T. 2020. Siccitas magna ultra modum: examining

- the occurrence and societal impact of droughts in prehistoric Ireland. *Proceedings of the Royal Irish Academy: Archaeology, Culture, History, Literature,* 120C, p.83-104.
- Rasmussen, S. O., Vinther, B. M., Clausen, H. B. & Andersen, K. K. 2007. Early Holocene climate oscillations recorded in three Greenland ice cores. *Quaternary Science Reviews*, 26, p.1907-1914.
- Reimer, P. J., Austin, W. E. N., Bard, E., Bayliss, A., Blackwell, P. G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kromer, B., Manning, S. W., Muscheler, R., Palmer, J. G., Pearson, C., van der Plicht, J., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Turney, C. S. M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S. M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A. & Talamo, S. 2020. The INTCAL20 Northern Hemisphere radiocarbon age calibration curve (0-55 cal kBP). Radiocarbon, 62, p.725-757.
- Richardson, D. M. & Rundel, P. W. 1998. Ecology and biogeography of Pinus: an introduction.
 In Richardson, D. M. (ed.) *Ecology and biogeography of Pinus*. Cambridge University Press,
 Cambridge, p.3-46.
- Rohling, E. J. & Pälike, H. 2005. Centennial-scale climate cooling with a sudden cold event around 8,200 years ago. *Nature*, 434, p.975-979.
- Roland, T. P., Daley, T. J., Caseldine, C. J., Charman, D. J., Turney, C. S. M., Amesbury, M.
 J., Thompson, G. J. & Woodley, E. J. 2015. The 5.2 ka climate event: Evidence from stable
 isotope and multi-proxy palaeoecological peatland records in Ireland. *Quaternary Science Reviews*, 124, p.209-223.
- Russell, I. 2010. N25 Waterford Bypass Archaeological Investigation contract 1- Final report on the Archaeological excavation of Killoteran 9. (03E0406). Unpublished excavation report.
 Archaeological Consultancy Services Ltd. on behalf of Waterford County Council and the National Roads Authority.
- Sassoon, D., Fletcher, W. J., Hotchkiss A., Owen, F. & Feng, L. 2021 Scots pine (*Pinus sylvestris*) dynamics in the Welsh Marches during the mid to late-Holocene. *The Holocene*, 31, 1024 1033-1046.

Selby, K. A., O'Brien, C. E., Brown, A. G. & Stuijts, I. 2005. A multi-proxy study of Holocene
lake development, lake settlement and vegetation history in central Ireland. *Journal of Quaternary Science*, 20, p.147-168.

- Simmons, I. G. & Innes, J. B. 1996. Disturbance phases in the mid-Holocene vegetation at
 North Gill, North York Moors: Form and Process. *Journal of Archaeological Science*, 23,
 p.183-191.
- 1031 Smith, A. G. 1970. The influence of Mesolithic and Neolithic man on British vegetation. In
 1032 Walker, D. & West, R. G. (eds.) *Studies in the vegetational history of the British Isles*.
 1033 Cambridge University Press, Cambridge, p.81-96.
- Smith, A. G. & Pilcher, J. R. 1973. Radiocarbon dates and vegetational history of the British
 Isles. *The New Phytologist*, 72, p.903-914.
- I036 Spencer, D., Molloy, K., Potito, A. P. & Jones, C. 2020. New insights into Late Bronze Age
 I037 settlement and farming activity in the southern Burren, western Ireland. *Vegetation History and* I038 Archaeobotany, 29, p.339-356.
- 1039 Stace, C. 1997. New Flora of the British Isles, 2nd Edition. Cambridge University Press,
 1040 Cambridge.
- Stockmarr, J. 1971. Tablets with spores used in absolute pollen analysis. *Pollen et Spores*, 13,
 p.615-621.
- Stolze, S. & Monecke, T. 2020. Neolithic land-use dynamics in northwest Ireland: multi-proxy
 evidence from Lough Arrow, County Sligo. *Vegetation History and Archaeobotany*, p.1 21.
- 1045 Støren, E. W., Paasche, Ø., Hirt, A. M. & Kumari, M. 2016. Magnetic and geochemical
 1046 signatures of flood layers in a lake system. *Geochemistry, Geophysics, Geosystems*, 17, p.4236 1047 4253.
- ⁴⁷₄₈ 1048 Stuiver, M. & Kra, R. S. 1986. Editorial comment. *Radiocarbon*, 28, p.ii.
- ⁴⁹ ₅₀ ¹⁰⁴⁹ Stuiver, M. & Polach, H. A. 1977. Reporting of 14C data. *Radiocarbon*, 19, p.355-363.
- Stuiver, M. & Reimer, P. J. 1986. A computer program for radiocarbon age calculation.
 Radiocarbon, 28, p.1022-1030.
- Stuiver, M. & Reimer, P. J. 1993. Extended 14C database and revised CALIB 3.0 14C age
 calibration program. *Radiocarbon*, 35, p.215-230.

1054 Swindles, G.T.; Lawson, I.T.; Matthews, I.P.; Blaauw, M.; Daley, T.J; Charman, D.J.; Roland,

- 1055 T.P.; Plunkett, G.; Schettler, G.; Gearey, B.R.; Turner, E.; Rea, H.A.; Roe, H.M.; Amesbury,
- 1056 M.J.; Chambers, F.M.; Holmes, J; Mitchell, F.J.G.; Blackford, J.; Blundell, A.; Branch, N.;
- ⁸ 1057 Holmes, J.; Langdon, P.; McCarroll, J.; McDermott, F.; Oksane, P.O.; Pritchard, O.; Stastney,
- 10 1058 P.; Stefanini, B.; Young, D.; Wheeler, J.; Becker, K.; Armit, I. 2013. Centennial-scale climate
- 12 1059 change in Ireland during the Holocene. *Earth Science Review* 126, p.300-320.

1 2 3

4 5

6

7

13

17

- 14 1060 Tallantire, P. A. 2002. The early-Holocene spread of hazel (Corylus avellana L.) in Europe
 15 1061 north and west of the Alps: an ecological hypothesis. *The Holocene*, 12, p.81-96.
- 18 1062 Taylor, K. J., McGinley, S., Potito, A. P., Molloy, K. & Beilman, D. W. 2018. A mid to late
 1063 Holocene chironomid-inferred temperature record from northwest Ireland. *Palaeogeography*,
 1064 *Palaeoclimatology*, *Palaeoecology*, 505, p.274-286.
- Thomas, E. R., Wolff, E. W., Mulvaney, R., Steffensen, J. P., Johnsen, S. J., Arrowsmith, C.,
 White, J. W. C., Vaughn, B. & Popp, T. 2007. The 8.2ka event from Greenland ice cores. *Quaternary Science Reviews*, 26, p.70-81.
- 29 Tierney, J. 2005. N25 Road Realingmant Scheme, Kilmacthomas, C. Waterford. Final 1068 30 31 archaeological testing report on archaeology found in Ahanaglogh and Graigueshoneen 1069 32 townlands, Kilmacthomas, Co. Waterford. (98E0575). Unpublished excavation report. Eachtra 33 1070 34 Archaeological Projects on behalf of Waterford County Council and the National Roads 1071 35 36 Authority. 1072 37
- Tietzsch-Tyler, D., Sleeman, A.G., Boland, M.A., Daly, E.P., Flegg, A.M., O'Connor, P.J.,
 Warren, W.P. (1994) *Geology of South Wexford. A geological description of South Wexford*and adjoining parts of Waterford, Kilkenny and Carlow to accompany the bedrock geology
 1076 1:100,000 scale map series, sheet 23, South Wexford. Geological Survey of Ireland.
- Timpany, S. 2009. Appendix 9: Palaeoenvironmental Analyses Report, Site 34, Newrath 46 1077 47 Townland, Co. Kilkenny. N25 Waterford Bypass, Contract 3. Final Report on archaeological 1078 48 49 Investigations at Site 34 in the townland of Newrath, Co Kilkenny. Unpublished excavation 1079 50 51 report, Headland Archaeology on behalf of Kilkenny County Council and the National Roads 1080 52 Authority. 1081 53 54
- ⁵⁵ 1082 Tipping, R., Ashmore, P., Davies, A. L., Haggart, B. A., Moir, A., Newton, A. J., Sands, R., ⁵⁶
- 57 1083 Skinner, T. & Tisdall, E. W. 2008. Prehistoric *Pinus* woodland dynamics in an upland
- ⁵⁸
 ⁵⁹ 1084 landscape in northern Scotland: The roles of climate change and human impact. *Vegetation*
- ⁶⁰ 1085 *History and Archaeobotany* 17, 251–267.

Torbenson, M. C. A., Plunkett, G., Brown, D. M., Pilcher, J. R. & Leuschner, H. H. 2015.
Asynchrony in key Holocene chronologies: Evidence from Irish bog pines. *Geology*, 43, p.799802.

Turner, J.N., Holmes, N., Davis, S.R., Leng, M.J., Langdon, C. and Scaife, R.G., 2015. A
multiproxy (micro-XRF, pollen, chironomid and stable isotope) lake sediment record for the
Lateglacial to Holocene transition from Thomastown Bog, Ireland. *Journal of Quaternary Science*, 30, p.514-528.

⁶ 1093 Ward, G. K. & Wilson, S. R. 1978. Procedures for comparing and combining radiocarbon age
⁸ 1094 determinations: A critique. *Archaeometry*, 20, p.19-31.

Warren, G. 2020. Climate change and hunter gatherers in Ireland: problems, potentials and
 pressing research questions. *Proceedings of the Royal Irish Academy: Archaeology, Culture,* History, Literature, 120C, p.1-22.

Warren, G., Davis, S., McClatchie, M. & Sands, R. 2014. The potential role of humans in structuring the wooded landscapes of Mesolithic Ireland: A review of data and discussion of approaches. *Vegetation History & Archaeobotany*, 23, p.629-646.

- Weltje, G.J. and Tjallingii, R., 2008. Calibration of XRF core scanners for quantitative
 geochemical logging of sediment cores: Theory and application. *Earth and Planetary Science Letters*, 274, p.423-438.
- Whitehouse, N. J., Schulting, R. J., McClatchie, M., Barratt, P., McLaughlin, R. T., Bogaard, A., Colledge, S., Marchant, R., Gaffrey, J. & Bunting, M. J. 2014. Neolithic agriculture on the European western frontier: The boom and bust of early farming in Ireland. *Journal of Archaeological Science*, 51, p.181-205.
- Whittle, A., Healy, F. & Bayliss, A. 2011. Gathering Time. Dating the Early Neolithic
 Enclosures of Southern Britain and Ireland, Oxbow Books, Oxford.
- Wiersma, A. P. & Renssen, H. 2006. Model-data comparison for the 8.2kaBP event:
 confirmation of a forcing mechanism by catastrophic drainage of Laurentide Lakes.
 Quaternary Science Reviews, 25, p.63-88.

Wren, J. 2006a. N25 Waterford Bypass, Contract 3. Final Report on archaeological
investigations at Site 4 in the townland of Mullinabro, Co. Kilkenny. (04E332). Unpublished
excavation report. Headland Archaeology Ltd. on behalf of Kilkenny County Council and the
National Roads Authority.

Wren, J. 2006b. N25 Waterford Bypass, Contract 3. Final Report on archaeological
Investigations at Site 34 in the townland of Newrath, Co Kilkenny. 04E0289. Unpublished
excavation report. Headland Archaeology Ltd. on behalf of Kilkenny County Council and the
National Roads Authority.

to per perieu

HOLOCENE

LPAZ/Top Depth (m) Depth (m) Depth (m) Posterior density estimate (95.4% probability)		Posterior density estimate (68.2% probability) period		Main pollen features	LoI	
LC13 2.15	780-1035 cal. AD	810-940 cal. AD	Early Medieval	Reductions in AP (c.55%) across zone, especially <i>Corylus</i> (c.20%), <i>Quercus</i> (c.10%) and <i>Alnus</i> (c.10%) towards top of zone. Increase in NAP (c.45%), especially Poaceae (c.20%), Cyperaceae (c.15%), <i>P. lanceolata</i> (c.4%) and <i>Rumex</i> (2%). <i>Cerealia</i> -type recorded across zone. Rise in <i>Pteridium</i> (c.10% TLP+S), Charcoal increases to peak midway through zone.	OM increasing <i>c</i> .40% at base to <i>c</i> .47% at top LPAZ. Average <i>c</i> .43%	
LC12 2.36	390-540 cal. AD	395-470 cal. AD	Late Iron Age	Pronounced increase in AP (<i>c</i> .95%), especially <i>Corylus</i> (<i>c</i> .50%) and <i>Quercus</i> (<i>c</i> .30%), other arboreal taxa remaining steady. Falls in NAP predominantly Poaceae and Cyperaceae, <i>P. lanceolata</i> low and sporadic. Increase in charcoal towards top of zone.	OM increasing from <i>c</i> .36% to <i>c</i> .41% at top LPAZ. Average <i>c</i> .39%	
LC11 2.64	415-250 cal. BC	405-355 cal. BC	Iron Age	Steady reduction in AP across zone (c.70%), primarily <i>Corylus</i> and following an initial increase, <i>Quercus</i> . NAP <i>increase</i> , Poaceae rises (c.15%), also <i>P. lanceolata</i> (c.8%), Lactuceae (2%) and sporadic <i>Filipendula</i> . <i>Cerealia</i> -type recorded 2.65m. Charcoal low.	OM ranges <i>c</i> .34% at base to <i>c</i> .29% across LPAZ. Average <i>c</i> .31%	
LC10 2.82	690-535 cal. BC	630-555 cal. BC	Iron Age transition	Increase in AP ($c.90\%$), mainly <i>Corylus</i> ($c.45\%$), other arboreal taxa remaining steady. Reductions in NAP, primarily Poaceae, <i>P. lanceolata</i> consistently present, at certain points higher values than the former. Charcoal low and sporadic.	OM increasing from <i>c</i> .33 to 37% across LPAZ. Average <i>c</i> .34%	
LC9 2.925	815-685 cal. BC	795-740 cal. BC	Late Bronze Age	Both AP (between 70-85%) and NAP (30-15%) fluctuate cross the zone; former mainly due to changes in <i>Corylus</i> but <i>Quercus</i> also involved and the latter Poaceae; <i>P. lanceolata</i> remaining steady. Two <i>Cerealia</i> -type grains recorded at 3.08m, <i>Pteridium</i> peaks at <i>c</i> .15% and charcoal also shows increase above (3.04m).	OM ranges c.28% at base to c.35% at top of LPAZ. Average c.30%	
LC8 3.14	1275-1000 cal. BC	1225-1080 cal. BC	Middle Bronze Age	Pronounced fall in AP (<i>c</i> .55%), especially <i>Corylus</i> (<i>c</i> .15%) and <i>Quercus</i> (<i>c</i> .10%) towards close of zone. Sharp increase in Poaceae and <i>P. lanceolata</i> , reaching highest values for diagram (<i>c</i> .28% and 7% respectively) also Cyperaceae (7%) and <i>Succisa</i> -type (2%), low values for Chenopodiaceae and <i>Filipendula</i> . Increase in <i>Pteridium aquilinum</i> (11% TLP+S) and Pteropsida monolete (indet.) (9% TLP+S) towards close of zone. Charcoal values very low.	OM increases from <i>c</i> .34 % to <i>c</i> .37% at 3.20m, decreasing to <i>c</i> .29% by top of LPAZ. Average <i>c</i> .32%	
LC7c 3.34	1500-1470 cal. BC (28.1%) 1465-1410 cal. BC (67.3%)	1495-1480 cal. BC (17.5%) 1455-1420 cal. BC (50.7%)		AP remaining steady ($c.80\%$), <i>Corylus</i> increasing ($c.45\%$) towards top of zone. NAP increases initially ($c.20\%$), mainly Poaceae and <i>P. lanceolata</i> , followed by reduction in former ($c.10\%$). Charcoal increases slightly to peak at top of zone.	OM ranges <i>c</i> .34 to <i>c</i> .36% across LPAZ. Average <i>c</i> .35%	
LC7b 3.56	1880-1740 cal. BC	1875-1840 cal. BC (30.0%) 1820-1800 cal. BC (14.1%) 1780-1750 cal. BC (24.2%)	Chalcolithic & Early Bronze Age	NAP remaining steady (15%); the latter mainly represented by Poaceae, low values for <i>P. lanceolata</i> . Charcoal remains low.	OM ranges from <i>c</i> .30% at base to <i>c</i> .40% at top LPAZ. Average <i>c</i> .33%	

HOLOCENE

LC7a 3.90	2620-2240 cal. BC	2505-2320 cal. BC	Late Neolithic	<i>Ulmus</i> remains low, <i>Corylus</i> values fall mid-way to <i>c</i> .25%. Other trees at similar values to previous zone. Increased NAP (<i>c</i> .15%), a second continuous <i>P. lanceolata</i> curve (<i>c</i> .3%) from start of zone, Poaceae also increases, Asteraceae, Apiaceae, <i>Filipendula</i> and Ranunculaceae present. Single grain <i>Cerealia</i> -type (4.02m) alongside <i>Artemisia</i> -type. Pteropsida (monolete) indet. increases to 8% TLP+S. Charcoal values show very slight increase.	OM ranges <i>c</i> .33 to <i>c</i> .35% across LPAZ. Average <i>c</i> .33%
LC6 4.08	2960-2525 cal. BC	2845-2630 cal. BC	Middle Neolithic II	Steady reduction in <i>Ulmus</i> (<i>c</i> .2-3%), <i>Alnus</i> increases (<i>c</i> .25%) and <i>Salix</i> also present in slightly greater values (<i>c</i> .3%). NAP steady at <i>c</i> .7%, predominantly Cyperaceae, continuous Ranunculaceae, sporadic <i>Filipendula</i> , <i>P. lanceolata</i> , <i>Succisa</i> -type. Charcoal remains steady.	OM ranges <i>c</i> .35 to <i>c</i> .37% across LPAZ, dropping to <i>c</i> .30% at top. Average <i>c</i> .34%
LC5 4.38	3355-3120 cal. BC	3275-3150 cal. BC	Middle Neolithic I	AP values increase overall (<i>c</i> .95%), primarily <i>Corylus</i> (<i>c</i> .30%), <i>Quercus</i> (<i>c</i> .25%) and <i>Ulmus</i> (<i>c</i> .8%) and <i>Alnus</i> (<i>c</i> .20%). NAP values decline, only sporadic records of <i>P. lanceolata</i> . Slight increase in <i>Polypodium</i> (6% TLP+S). Charcoal remains steady.	OM ranges from <i>c</i> .35 to <i>c</i> .37% across LPAZ. Average <i>c</i> .36%
LC4c 4.71	3685-3530 cal. BC	3650-3555 cal. BC	Early Neolithic II	<i>Quercus</i> (<i>c</i> .25%) and <i>Ulmus</i> (<i>c</i> .4%) values recover, <i>Corylus</i> remaining <i>c</i> .30%. NAP values are reduced (<i>c</i> .5%) although a continuous but reduced curve for <i>P. lanceolata</i> (<i>c</i> .1% max) is exhibited, <i>Cerealia</i> -type pollen recorded (2 grains 4.77 and 1 grain 4.75m); other herbs similar to previous zone. Charcoal also remaining at similar levels.	OM ranges from $c.31\%$ at base to $c.36\%$ at the top of LPAZ. Average $c.34\%$
LC4b 4.82	3860-3650 cal. BC	3815-3710 cal. BC	Early Neolithic I	Quercus (25 to 15%) and Corylus (40 to 30%) decreasing across zone; NAP increases to c.20%, predominantly Poaceae (10%), continuous presence of <i>P. lanceolata</i> (c.5%) and Ranunculaceae (c.1-2%), <i>Artemisia</i> - type, Asteraceae, <i>Rumex acetosella/acetosa, Filipendula</i> and <i>Urtica dioica</i> recorded at lower values. Single grain of <i>Cerealia</i> -type (4.85m). Charcoal increases at top of the zone.	OM ranges <i>c</i> .10% at 4.90m to <i>c</i> .31% at top of LPAZ. Average <i>c</i> .29%
LC4a 5.01	4050-3960 cal. BC	4040-4010 cal. BC (25.8%) 4005-3975 cal. BC (42.4%)	Late Mesolithic	Ulmus values decrease at the start of zone (c. 10% to c. 3%) and thereafter remain very low; a temporary reduction in <i>Corylus</i> (c.25%), followed by recovery (c.40%). NAP values increase to c. 10%, consisting of Poaceae (c.6%) and other herbs recorded include Asteraceae, Caryophyllaceae and <i>Plantago lanceolata</i> . Pteropsida (monolete) indet. increases to 9% TLP+S. Charcoal remains low.	OM ranges <i>c</i> .30 to <i>c</i> .25% at top of LPAZ. Average <i>c</i> .27%
LC3 5.09	4320-4020 cal. BC	4225-4070 cal. BC		<i>Corylus</i> recovers (<i>c</i> .40%), while <i>Alnus</i> increases steadily to a peak (<i>c</i> .25%), <i>Quercus</i> and <i>Ulmus</i> stable (20% and 10% respectively). The initiation of low but continuous <i>Ilex</i> curve is recorded at top of zone. NAP values remaining low. Slight if sporadic increases in charcoal.	OM ranges from <i>c</i> .29 to <i>c</i> .38%. Average <i>c</i> .34%
LC2 5.84	6245-5575 cal. BC	6120-5780 cal. BC	Early Mesolithic	Temporary reduction in both <i>Corylus</i> ($c.30\%$) and <i>Quercus</i> ($c.20\%$) corresponding with spikes in <i>Betula</i> ($c.15\%$) and <i>Pinus</i> ($c.20\%$), followed by recoveries in the former. <i>Alnus</i> increases abruptly at top of zone to $c.15\%$. NAP remaining low ($c.4\%$). Charcoal remaining very low.	OM ranges <i>c</i> .35 to <i>c</i> .33%. across LPAZ. Average <i>c</i> .34%

_	LC1 6.03	7615-6500 cal. BC	7320-6745 cal. BC	Dominated by AP, high values of <i>Corylus</i> (<i>c</i> .50%) and <i>Quercus</i> (<i>c</i> .30%), <i>Ulmus</i> (<i>c</i> .8%), <i>Pinus</i> , and <i>Betula</i> present (2-4%). A continuous but low presence of <i>Hedera helix</i> (2%) and <i>Salix</i> (1%). NAP values very low, traces of Poaceae, Cyperaceae and <i>Filipendula</i> . Charcoal very low.	OM ranges from <i>c</i> .35to <i>c</i> .40% across the LPAZ. Average <i>c</i> .37%
			http:	://mc.manuscriptcentral.com/holocene	

Lab Code	Dated Material	¹⁴ C Age (BP)	Depth (m)	¹³ δC
UBA-22422	Fragment of willow branch	2526±32	1.78	-20.
SUERC-70812	Organic sediment (humic fraction)	1141±32	2.19	-29.
SUERC-70813	Organic sediment (humin fraction)	1198±29	2.19	-30.
Statistic	ally consistent measurements (T'=1.9; T' (5%)=3.8%; v=1). Weighte	d mean 1170±21	
Beta-423349	Organic sediment (humin fraction)	1420±30	2.29	-29
Beta-423350	Organic sediment (humic fraction)	1500±30	2.29	-29
Statistic	ally consistent measurements (T'=3.6; T' (5%)=3.8%; v=1). Weighte	d mean 1460±22	
SUERC-70814	Organic sediment (humic fraction)	1661±27	2.37	-29
SUERC-70815	Organic sediment (humin fraction)	1635±29	2.37	-30
Statistic	ally consistent measurements (T'=0.4; T' (5%)=3.8%; v=1). Weighte	d mean 1622±20	
Beta-426773	Organic sediment (humin fraction)	1910±30	2.45	-29
SUERC-70816	Organic sediment (humic fraction)	1959±29	2.49	-29
SUERC-70817	Organic sediment (humin fraction)	1996±29	2.49	-29.
Statistic	ally consistent measurements (T'=0.8; T' (5%)=3.8%; v=1). Weighte	d mean 1978±21	
SUERC-70821	Organic sediment (humic fraction)	2003±29	2.53	-29
SUERC-70822	Organic sediment (humin fraction)	2074±29	2.53	-29
Statistic	ally consistent measurements (T'=3.0; T' (5%)=3.8%; v=1). Weighte	d mean 2039±21	
SUERC-72575	Organic sediment (humic fraction)	2114±29	2.55	-29
SUERC-72576	Organic sediment (humin fraction)	2135±29	2.55	-29
Statistic	ally consistent measurements (T'=0.2; T' (5%)=3.8%; v=1). Weighte	d mean 2125±24	
Beta-426774	Organic sediment (humin fraction)	2150±30	2.60	-29
SUERC-72571	Organic sediment (humin fraction)	2181±33	2.61	-29
GU43402	Organic sediment (humic fraction)	FAILED	2.61	-
SUERC-70823	Organic sediment (humic fraction)	2250±29	2.65	-29
SUERC-70824	Organic sediment (humin fraction)	2336±29	2.65	-30
	Statistically inconsistent measurements (T'	=4.397; T' (5%) =3.8%	; v=1).	
SUERC-70825	Organic sediment (humic fraction)	2328±29	2.71	-29
SUERC-70826	Organic sediment (humin fraction)	2436±26	2.71	-29
	Statistically inconsistent measurements (T	"=7.7; T' (5%) =3.8%;	v=1)	
UBA-22423	Organic sediment (humin fraction)	2413±31	2.77	-25
SUERC-70827	Organic sediment (humic fraction)	2467±29	2.81	-29
SUERC-70831	Organic sediment (humin fraction)	2506±29	2.81	-29.
Statistica	ally consistent measurements (T'=1.0; T' (5%) =3.8%; v=1). Weighte	ed mean 2489±20	
SUERC-70832	Organic sediment (humic fraction)	2446±27	2.915	-29
SUERC-70833	Organic sediment (humin fraction)	2514±29	2.915	-29
Statistica	ally consistent measurements (T'=3.0; T' (5%) =3.8%; v=1). Weighte	ed mean 2478±20	
Beta-426775	Organic sediment (humin fraction)	2620±30	2.94	-29
Beta-426776	Organic sediment (humin fraction)	2750±30	3.04	-28
SUERC-70834	Organic sediment (humic fraction)	2794±26	3.08	-28
SUERC-70835	Organic sediment (humin fraction)	2821±29	3.08	-29
Statistica	ally consistent measurements (T'=0.5; T' (5%) =3.8%; v=1). Weighte	ed mean 2806±20	
Beta-423351	Organic sediment (humic fraction)	3050±30	3.18	-28
Beta-423352	Organic sediment (humin fraction)	3020±30	3.18	-30
Statistica	ally consistent measurements (T'=0.5; T' (5%) =3.8%; v=1). Weighte	ed mean 3035±22	
SUERC-76326	Organic sediment (humic fraction)	3026±24	3.26	-29
SUERC-76666	Organic sediment (humin fraction)	3132±32	3.26	-29
	Statistically inconsistent measurements (T			<u> </u>
	Organic sediment (humic fraction),	3162±24	3.34	-29.

http://mc.manuscriptcentral.com/holocene

	SUERC-76328	Organic sediment (humin fraction)	3168±24	3.34	-28.5			
	Statistically consistent measurements (T'=0.0; T' (5%) =3.8%; v=1). Weighted mean 3165±17							
	SUERC-76329	Organic sediment (humic fraction)	3469±24	3.56	-27.6			
	SUERC-76667	Organic sediment (humin fraction)	3473±32	3.56	-29.9			
	Statistically	consistent measurements (T'=0.0; T' (5%) =3.8%; v=1). Weighted mea	an 3470±20				
	UBA-22424	Organic sediment (humin fraction)	3800±36	3.79	-28.2			
	SUERC-70836	Organic sediment (humic fraction)	4459±29	4.31	-29.7			
	SUERC-70837	Organic sediment (humin fraction)	4520±27	4.31	-30.1			
	Statisticall	y consistent measurements (T'=2.4; T'	(5%) =3.8; v=1). Weighted mean	4459±29				
	UBA-22425	Organic sediment (humin fraction)	4710±39	4.60	*			
	SUERC-70841	Organic sediment (humic fraction)	4755±29	4.69	-29.3			
	SUERC-70842	Organic sediment (humin fraction)	4795±26	4.69	-29.4			
	Statistically consistent measurements (T'=1.1; T'(5%)=3.8; v=1). Weighted mean 4777±20							
	UBA-33721	Organic sediment (humin fraction)	5080±56	4.87	*			
	UBA-33722	Organic sediment (humin fraction)	5076±51	4.97	*			
	UBA-30917	Organic sediment (humin fraction)	5210±34	5.01	*			
	SUERC-70843	Organic sediment (humic fraction)	5675±27	5.34	-29.4			
	SUERC-70844	Organic sediment (humin fraction)	5788±26	5.34	-30.0			
	Statistically inconsistent measurements (T'=9.1; T'(5%) =3.8; v=1).							
	UBA-22426	Organic sediment (humin fraction)	6154±47	5.46	*			
	UBA-30918	Organic sediment (humin fraction)	7358±37	5.91	*			
_	UBA-22427	Organic sediment (humin fraction)	9057±48	6.21	*			

Organic sediment (humin fraction) 905 / == 40

http://mc.manuscriptcentral.com/holocene




Figure 1 Location map of Lough Cullin. Inset map indicating the location of selected sites referenced in this paper. 1-Lough Muckno, 2-Lough Arrow, 3-Lough Inchiquin, 4-Lough Dargan & 5-Lough Meenachrinna.

404x316mm (118 x 118 DPI)






Figure 2 Pollen percentage diagram from Lough Cullin.

625x281mm (118 x 118 DPI)







60








Figure 4 Loss-on-ignition (LoI), geochemistry and magnetic susceptibility (MS) diagram from Lough Cullin.

569x402mm (87 x 87 DPI)



HOLOCENE



	R_Date SUERC-70824 [A:106] R_Date SUERC-70826 [A:109] R_Date UBA-22423 [A:93] R_Combine LC281 [A:108] R_Combine LC291.5 [A:96] R_Date BETA-426775 [A:98]				
		1000 Posterior density e	800 600 estimate (cal BC)	400	200
Figure 6	5 Unmodelled (light grey) and m dates are derived fro	odelled (dark gr m the Bayesian	ey) dates across tl age-model shown	he `Hallstatt' pla in Figure 5.	teau. Modelled
	4	00x127mm (87	x 87 DPI)		
	http://m	c manuscriptcon	tral.com/holocene		
	n.cp.//m	emanascripteen			