

This is a repository copy of *A cofactor consumption screen identifies promising NfsB family nitroreductases for dinitrotoluene remediation*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/183112/>

Version: Accepted Version

---

**Article:**

Williams, Elsie, Sharrock, Abigail, Rylott, Liz [orcid.org/0000-0002-1609-414X](https://orcid.org/0000-0002-1609-414X) et al. (3 more authors) (2019) A cofactor consumption screen identifies promising NfsB family nitroreductases for dinitrotoluene remediation. BIOTECHNOLOGY LETTERS. pp. 1155-1162. ISSN: 0141-5492

<https://doi.org/10.1007/s10529-019-02716-z>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Biotechnology Letters

## A cofactor consumption screen identifies promising NfsB family nitroreductases for dinitrotoluene remediation --Manuscript Draft--

<b>Manuscript Number:</b>		
<b>Full Title:</b>	A cofactor consumption screen identifies promising NfsB family nitroreductases for dinitrotoluene remediation	
<b>Article Type:</b>	Original Article	
<b>Section/Category:</b>	Microbial and Enzyme Technology	
<b>Keywords:</b>	Bioremediation; Dinitrotoluene; NADPH depletion assay; NfsA; NfsB; Nitroreductase; YfkO	
<b>Corresponding Author:</b>	David Ackerley Victoria University of Wellington Wellington, NEW ZEALAND	
<b>Corresponding Author Secondary Information:</b>		
<b>Corresponding Author's Institution:</b>	Victoria University of Wellington	
<b>Corresponding Author's Secondary Institution:</b>		
<b>First Author:</b>	Elsie M Williams, PhD	
<b>First Author Secondary Information:</b>		
<b>Order of Authors:</b>	Elsie M Williams, PhD Abigail V Sharrock, PhD Elizabeth L Rylott, PhD Neil C Bruce, PhD Joanna K MacKichan, PhD David Ackerley	
<b>Order of Authors Secondary Information:</b>		
<b>Funding Information:</b>	Marsden Fund (VUW0704) Marsden Fund (VUW1502) Biotechnology and Biological Sciences Research Council (BB/P005713/1) Strategic Environmental Research and Development Program (ER-2723)	Professor David Ackerley Professor David Ackerley Professor Neil C Bruce Professor Neil C Bruce
<b>Abstract:</b>	<p><b>Objectives</b></p> <p>To survey a library of over-expressed nitroreductases to identify those most active with 2,4- and 2,6-dinitrotoluene substrates, as promising candidates for phytoremediation of soils and groundwater contaminated with poly-nitro toluene pollutants.</p> <p><b>Results</b></p> <p>To indirectly monitor dinitrotoluene reduction we implemented a nitroblue tetrazolium dye screen to compare relative rates of NADPH consumption for 58 nitroreductase candidates, over-expressed in a nitroreductase-deleted strain of Escherichia coli. Although the screen only provides activity data at a single substrate concentration, by</p>	

	<p>altering the substrate concentration and duration of incubation we showed we could first distinguish between more-active and less-active enzymes and then discriminate between the relative rates of reduction exhibited by the most active nitroreductases in the collection. We observed that members of the NfsA and NfsB nitroreductase families were the most active with 2,4-dinitrotoluene, but that only members of the NfsB family were proficient with 2,6-dinitrotoluene. Two NfsB family members, YfkO from <i>Bacillus subtilis</i> and NfsB from <i>Vibrio vulnificus</i>, appeared especially effective with these substrates. Purification of both enzymes as His6-tagged recombinant proteins enabled in vitro determination of Michaelis-Menten kinetic parameters with each dinitrotoluene substrate.</p> <p>Conclusions</p> <p><i>V. vulnificus</i> NfsB is a particularly promising candidate for bioremediation applications, being ca. 5-fold more catalytically efficient with 2,4-dinitrotoluene and over 26-fold more active with 2,6-dinitrotoluene than the benchmark <i>E. coli</i> nitroreductases NfsA and NfsB.</p>
<b>Suggested Reviewers:</b>	<p>Chris D Gwenin, PhD Bangor University c.d.gwenin@bangor.ac.uk Expertise in nitroreductase chemistry</p> <p>Maria D Roldan, PhD Universidad Católica de Córdoba bb2rorum@uco.es Expertise in nitroreductases for bioremediation</p>

## **A cofactor consumption screen identifies promising NfsB family nitroreductases for dinitrotoluene remediation**

Elsie M Williams<sup>†,1,2</sup>, Abigail V Sharrock<sup>†,1,3</sup>, Elizabeth L Rylott<sup>4</sup>, Neil C Bruce<sup>4</sup>, Joanna K  
MacKichan<sup>1,3</sup>, David F Ackerley<sup>\*,1,3</sup>

<sup>1</sup>School of Biological Sciences, Victoria University of Wellington, Wellington, New Zealand

<sup>2</sup>Current address: Department of Chemistry, Emory University, Atlanta, Georgia, USA

<sup>3</sup>Centre for Biodiscovery, Victoria University of Wellington, Wellington, New Zealand

<sup>4</sup>Centre for Novel Agricultural Products, Department of Biology, University of York,  
York YO10 5DD, UK

<sup>†</sup>These authors contributed equally to this work

\*Correspondence to: david.ackerley@vuw.ac.nz

Journal Section: Microbial and Enzyme Technology

## Abstract

**Objectives:** To survey a library of over-expressed nitroreductases to identify those most active with 2,4- and 2,6-dinitrotoluene substrates, as promising candidates for phytoremediation of soils and groundwater contaminated with poly-nitro toluene pollutants.

**Results:** To indirectly monitor dinitrotoluene reduction we implemented a nitroblue tetrazolium dye screen to compare relative rates of NADPH consumption for 58 nitroreductase candidates, over-expressed in a nitroreductase-deleted strain of *Escherichia coli*. Although the screen only provides activity data at a single substrate concentration, by altering the substrate concentration and duration of incubation we showed we could first distinguish between more-active and less-active enzymes and then discriminate between the relative rates of reduction exhibited by the most active nitroreductases in the collection. We observed that members of the NfsA and NfsB nitroreductase families were the most active with 2,4-dinitrotoluene, but that only members of the NfsB family were proficient with 2,6-dinitrotoluene. Two NfsB family members, YfkO from *Bacillus subtilis* and NfsB from *Vibrio vulnificus*, appeared especially effective with these substrates. Purification of both enzymes as His<sub>6</sub>-tagged recombinant proteins enabled *in vitro* determination of Michaelis-Menten kinetic parameters with each dinitrotoluene substrate.

**Conclusions:** *V. vulnificus* NfsB is a particularly promising candidate for bioremediation applications, being *ca.* 5-fold more catalytically efficient with 2,4-dinitrotoluene and over 26-fold more active with 2,6-dinitrotoluene than the benchmark *E. coli* nitroreductases NfsA and NfsB.

**Keywords:** Bioremediation; Dinitrotoluene; NADPH depletion assay; NfsA; NfsB; Nitroreductase; YfkO.

## Introduction

2,4-dinitrotoluene and 2,6-dinitrotoluene, widely used as propellants or as precursors in the manufacture of 2,4,6-trinitrotoluene (TNT), are toxic and persistent contaminants of soil and groundwater (Dontsova et al, 2009). The strongly electron-withdrawing nitro groups on poly-nitro toluenes delocalise the electronic charge distribution in the aromatic rings, rendering them resistant to oxidative degradation by other enzymes (Roldan et al, 2008). Bacterial type I nitroreductases, which reduce nitro substituents via concerted two-electron transfer steps, yield derivatives that are far more amenable substrates for ring degrading enzymes (Williams et al, 2015; Roldan et al, 2008).

Plants engineered for high-level expression of bacterial nitroreductases have shown enhanced transformation and tolerance of poly-nitro toluenes, offering prospects for phytoremediation of contaminated environments (Zhang et al, 2019; Van Dillewijn et al, 2008; Hannink et al, 2007). However, despite there being a great diversity of bacterial nitroreductases available in nature (Akiva et al, 2017), very few candidates have been surveyed to assess their relative potential for these applications. A dearth of characterised nitroreductase candidates is particularly evident in the context of dinitrotoluene remediation.

Effective surveying of novel enzyme candidates requires an efficient activity screen. We previously described how *Escherichia coli* cells over-expressing nitroreductase enzymes that are active with dinitrotoluene substrates became more sensitive to these toxins, and that IC<sub>50</sub> data were somewhat predictive of relative levels of catalytic efficiency ( $k_{cat}/K_M$ ) for the purified enzymes (Rich et al, 2018). However, as IC<sub>50</sub> calculations require measurements to be made across a broad range of substrate concentrations, this is not an approach

amenable to large-scale screening of candidate gene libraries. Here we describe our implementation of an alternative assay, based on cofactor consumption, to rapidly identify nitroreductase enzymes proficient in reduction of both 2,4- and 2,6-dinitrotoluene.

## Materials and Methods

### *Chemicals, plasmids, bacterial strains and nitroreductases*

All reagents were purchased from Sigma-Aldrich (Saint Louis, MO). For protein purification, the genes encoding YfkO from *Bacillus subtilis* and NfsB from *Vibrio vulnificus* were expressed from the His<sub>6</sub>-tag expression vector pET28a+ in *E. coli* strain BL21(DE3) (Novagen, Merck Millipore; Billerica, MA) as previously described (Prosser et al, 2013). For library screening and IC<sub>50</sub> assays, nitroreductase candidates were over-expressed from plasmid pUCX in *E. coli* 7NT, a gene-deleted derivative of strain W3110 that lacks seven endogenous oxidoreductase genes (*nfsA*, *nfsB*, *azoR*, *nema*, *yieF*, *ycaK* and *mdaB*) to minimise background nitroreductase activity and an efflux pump gene (*tolC*) to maximise substrate retention (Copp et al, 2014). The 58-membered nitroreductase gene library was constructed as previously described (Mowday et al, 2016) and the full list of candidate genes (with UniProt default identifiers) is summarised in Table 1.

<Insert Table 1 here>

### *Nitroblue tetrazolium assay*

Cells of *E. coli* 7NT pUCX:nitroreductase strains were used to inoculate wells of a sterile 96-well microplate containing 200  $\mu$ L lysogeny broth (LB) supplemented with ampicillin (100  $\mu$ g.mL<sup>-1</sup>) and glucose (0.2% w/v). Cultures were incubated at 30 °C, 200 rpm for 16 hours,

after which 10  $\mu\text{L}$  of each overnight culture was used to inoculate wells of a sterile 96-well microplate containing 190  $\mu\text{L}$  induction medium (LB medium supplemented with ampicillin (100  $\mu\text{g}.\text{mL}^{-1}$ ), glucose (0.2% w/v) and IPTG (0.05 mM). Induced cultures were incubated at 30 °C, 200 rpm for 6 hours. Crude cell lysates were prepared by incubating 100  $\mu\text{L}$  of each culture with BugBuster® Protein Extraction Reagent at a 1:1 (v/v) ratio at room temperature for 30 min. To monitor cell lysate NADPH consumption, duplicate 200  $\mu\text{L}$  reactions were prepared consisting of 100 mM  $\text{KPO}_4$  buffer pH 8.0, crude cell lysate (40  $\mu\text{L}$  for full library screening assays or 5  $\mu\text{L}$  for NfsB-family screening assays) and 150  $\mu\text{M}$  substrate (2,4-dinitrotoluene or 2,6-dinitrotoluene), and reactions were initiated by addition of 50  $\mu\text{L}$  of 800  $\mu\text{M}$  NADPH. Reactions were incubated at room temperature for 30 min for full library screening assays, or 10 min for NfsB-family screening assays. Reactions were halted by the addition of 50  $\mu\text{L}$  nitroblue tetrazolium solution (nitroblue tetrazolium (2  $\text{mg}.\text{mL}^{-1}$ ), phenazine methosulfate (0.3  $\text{mg}.\text{mL}^{-1}$ , prepared fresh prior to use) and 100 mM  $\text{KPO}_4$  buffer (pH 8.0)). Levels of formazan production, corresponding to the amount of NADPH remaining in each well, were quantified 3-5 min post addition of the nitroblue tetrazolium solution by measuring absorbance at 590 nm using an EnSpire™ 2300 Multilabel Reader (PerkinElmer, Waltham, MA). In pilot tests, four *E. coli* 7NT pUCX:nitroreductase strains (expressing the genes encoding NfrA\_Bli, NfsA\_Vv, Ydgl\_Bs and YcnD\_Bs) exhibited a substantial decrease in NADPH levels in the absence of substrate. Reasoning that this high background NADPH oxidase activity would confound measurement of substrate consumption, these strains were excluded from the full library analysis.

#### *IC<sub>50</sub> assays*



Cells of *E. coli* 7NT pUCX:*nitroreductase* strains were used to inoculate wells of sterile 96-well microplates containing 200  $\mu$ L LB medium supplemented with ampicillin (100  $\mu$ g.mL<sup>-1</sup>) and glucose (0.2% w/v). Cultures were incubated at 30 °C, 200 rpm for 16 hours, after which they were diluted 20-fold into 15 mL centrifuge tubes containing 2 mL induction medium (LB medium supplemented with ampicillin (100  $\mu$ g.mL<sup>-1</sup>), glucose (0.2% w/v) and IPTG (0.05 mM). Induced cultures were incubated at 30 °C, 200 rpm for 2.5 hours. Aliquots (30  $\mu$ L) of each culture were added to wells of a sterile 384-well microplate containing 30  $\mu$ L induction medium  $\pm$  two-fold the desired final dinitrotoluene concentration. In total, each strain was challenged in duplicate across 7 to 15 conditions containing 1.5-fold increasing titrations of 2,4-dinitrotoluene or 2,6-dinitrotoluene together with a medium-only control. The absorbance of each well at 600 nm was measured using an EnSpire™ 2300 Multilabel Reader (PerkinElmer, Waltham, MA). Cultures were incubated at 30 °C, 200 rpm for a further 4 hours, after which absorbance readings at 600 nm were recorded once more. The increase in absorbances at 600 nm of corresponding challenged and unchallenged wells for each strain were compared and used to calculate percentage growth following challenge. IC<sub>50</sub> values (the concentration of 2,4-dinitrotoluene or 2,6-dinitrotoluene estimated to reduce growth by 50% relative to the unchallenged control) were calculated using a dose-response inhibition four-parameter variable slope equation in GraphPad Prism 7.0 (GraphPad Software Inc., La Jolla, CA).

#### *His<sub>6</sub>-tagged protein purification and Michaelis-Menten kinetics*

Recombinant His<sub>6</sub>-tagged nitroreductases were purified by nickel-affinity chromatography using Ni/NTA resin (Novagen, Merck, Darmstadt, Germany), followed by incubation with a molar excess of FMN cofactor and buffer exchange as previously described (Prosser et al,

2010). Apparent steady state enzyme kinetics were determined via monitoring decrease in absorbance at 340 nm (NADPH extinction co-efficient,  $6,220 \text{ M}^{-1} \text{ cm}^{-1}$ ) at a fixed concentration of NADPH and varying concentrations of 2,4-dinitrotoluene or 2,6-dinitrotoluene. Reactions were carried out in 60  $\mu\text{L}$  volumes in UVettes™ (Eppendorf, Hamburg, Germany), using a 1 cm path length, and the wavelength was measured using a Helios  $\gamma$  UV-Vis spectrophotometer (Thermo Fisher Scientific, Waltham, MA). Reactions consisted of 10 mM Tris-Cl pH 7.0 buffer, 250  $\mu\text{M}$  NADPH, 2% DMSO and substrate concentrations ranging from 0 - 400  $\mu\text{M}$  2,4-dinitrotoluene and 0 -1600  $\mu\text{M}$  2,6-dinitrotoluene. Reactions were initiated by addition of an appropriate dilution of enzyme and the linear decrease in absorbance monitored for the first 20-40 s post enzyme addition. Non-linear regression analyses and Michaelis-Menten curve fitting was performed using GraphPad Prism v 7.0 (GraphPad Software, San Diego, California, USA).

## Results and Discussion

To identify enzymes capable of activating diverse nitroaromatic prodrugs we previously generated a 58-membered library of nitroreductase candidates in *Escherichia coli* reporter strains that express either a  $\beta$ -galactosidase or GFP gene under control of a DNA damage responsive (SOS) promoter (Mowday et al, 2016; Copp et al, 2017). We tested whether dinitrotoluene reduction could likewise be monitored via SOS assays, reasoning that reduction of TNT is known to yield genotoxic metabolites (Bolt et al, 2006). However, we found that both 2,4-dinitrotoluene and 2,6 dinitrotoluene evoked high levels of background signal in nitroreductase-expressing reporter strains that precluded accurate screening (not shown). We therefore sought to implement an alternative screening procedure based on

consumption of the NAD(P)H cofactor that acts as an electron source for bacterial Type I nitroreductases. For this, we adapted a colorimetric nitroblue tetrazolium assay, previously described by Mayer and Arnold (2002) for directed evolution of 6-phosphogluconate dehydrogenase enzymes.

In the presence of NADPH yellow nitroblue tetrazolium is reduced to a purple formazan dye that can be quantified at 590 nm. This allowed us to measure the relative levels of dinitrotoluene reduction in lysates derived from *E. coli* strains over-expressing nitroreductase candidates. To minimise background nitroreductase activity, screening of the full 58-membered nitroreductase candidate library was performed in a strain of *E. coli* lacking endogenous *nfsA*, *nfsB*, *azoR*, *nemA*, *yieF*, *ycaK* and *mdaB* genes (strain 7NT; Copp et al, 2014). Lysates were incubated with NADPH and either 2,4-dinitrotoluene or 2,6-dinitrotoluene, then reactions were stopped with nitroblue tetrazolium chloride and the amount of formazan in challenged lysates compared to unchallenged duplicates.

The full library screen confirmed that, of the enzymes tested, NfsA and NfsB family nitroreductases consistently consumed the most NADPH in the presence of dinitrotoluene substrates (Figure 1A). Notably, although members of both the NfsA and NfsB families were represented among the top 25 most active enzymes with 2,4-dinitrotoluene as substrate, the NfsB family enzymes markedly outperformed those from the NfsA family with 2,6-dinitrotoluene as substrate (Figure 1A). To distinguish the relative activities of the NfsB family enzymes it was necessary to reduce the amount of culture lysate from 40  $\mu$ L for the full library screening assay to 5  $\mu$ L and the incubation time from 30 min to 10 min (Figure 1B).

<Insert Figure 1 here>

We next sought to test whether results from the nitroblue tetrazolium assay were generally congruent with IC<sub>50</sub> data by performing detailed IC<sub>50</sub> assays for all strains expressing *nfsB* family genes (Table 2) and correlating the measured IC<sub>50</sub> values (μM) with NADPH consumption (percent ΔA<sub>590</sub>) (Figure 2). The inverse correlation between IC<sub>50</sub> and NADPH consumption for 2,4-dinitrotoluene was moderate rather than strong ( $r^2 = 0.35$ ); this is likely a consequence of the nitroblue tetrazolium assay providing a ‘snapshot’ of activity at a single substrate concentration, whereas IC<sub>50</sub> data derive from a more comprehensive midpoint of activity across a titration range. However, an additional advantage of the rapid nitroblue tetrazolium assay over performing more extensive IC<sub>50</sub> measurements was evident in its ability to generate data for 2,6-dinitrotoluene, whereas IC<sub>50</sub> values could not be calculated for the majority of *nfsB*-expressing strains owing to insufficient toxicity at the achievable solubility limit (Table 2).

<Insert Table 2 here> <Insert Figure 2 here>

Based on NADPH consumption data, the NfsB family enzyme *Bacillus subtilis* YfkO (YfkO\_Bs) appeared most active with 2,4-dinitrotoluene, whereas *Vibrio vulnificus* NfsB (NfsB\_Vv) appeared most active with 2,6-dinitrotoluene (Figure 1B). We purified each enzyme as a His<sub>6</sub>-tagged recombinant protein by nickel affinity chromatography and measured their Michaelis-Menten kinetic parameters with either 2,4-dinitrotoluene or 2,6-dinitrotoluene as the reductive substrate (Table 3). In terms of catalytic efficiency, the  $k_{cat}/K_M$  values for NfsB\_Vv and YfkO\_Bs with 2,4-dinitrotoluene were 4- to 7-fold higher than those previously reported for either of the canonical Type I nitroreductases NfsA\_Ec and NfsB\_Ec (Rich et al, 2018). Their superiority with 2,6-dinitrotoluene was even more profound, with the measured  $k_{cat}/K_M$  values for NfsB\_Vv and YfkO\_Bs being 20- to 37-fold higher than those previously reported for NfsA\_Ec and NfsB\_Ec (Rich et al, 2018). NfsB\_Vv was particularly

effective with each substrate, and appears a promising candidate for expression in transgenic dinitrotoluene remediation systems.

<Insert Table 3 here>

## Conclusions

We have previously used activation of the *E. coli* SOS response to report on genotoxicity as a measure of nitroreductase-mediated activation of prodrug substrates. Here, to monitor activity with dinitrotoluene substrates that did not yield a clear SOS signal upon reduction, we implemented a more generally applicable screen for nitroreductase activity based on consumption of exogenously added NADPH that provided a dominant electron source for the reaction. In addition to providing much higher throughput than IC<sub>50</sub> assays, we showed that our screen could provide relative activity data with substrates such as 2,6-dinitrotoluene that are insufficiently toxic at achievable concentrations for IC<sub>50</sub> values to be calculated. The linear range of the assay was insufficient to discriminate between the more active NfsB family nitroreductases while still detecting the less active NfsA and AzoR family enzymes. However, by reducing both the substrate concentration and duration of incubation we were able to discriminate between the different NfsB family nitroreductases and identified *V. vulnificus* NfsB and *B. subtilis* YfkO as two of the most active dinitrotoluene reducing enzymes in our collection. Purification of these nitroreductases as His<sub>6</sub>-tagged proteins enabled steady state kinetic assays to be performed, demonstrating *V. vulnificus* NfsB to be over 5-fold more active with 2,4-dinitrotoluene and over 26-fold more active with 2,6-dinitrotoluene than either of the canonical nitroreductases *E. coli* NfsA and NfsB.

## Acknowledgements

212 This work was supported by grants from the Royal Society of New Zealand Marsden Fund  
213 (VUW0704 and VUW1502), the UK Biotechnology and Biological Sciences Research Council  
214 (BB/P005713/1), and the Strategic Environmental Research and Development Program (ER-  
215 2723).

216

## 217 **References**

218 Akiva E, Copp JN, Tokuriki N, Babbitt PC (2017) Evolutionary and molecular foundations of  
219 multiple contemporary functions of the nitroreductase superfamily. *Proc Natl Acad Sci U S A*  
220 114:E9549-E9558.

221 Bolt HM, Degen GH, Dorn SB, Plottner S, Harth V (2006) Genotoxicity and potential  
222 carcinogenicity of 2,4,6-trinitrotoluene: structural and toxicological considerations. *Rev*  
223 *Environ Health* 21:217–228.

224 Copp JN, Williams EM, Rich MH, Patterson AV, Smaill JB, Ackerley DF (2014) Toward a high-  
225 throughput screening platform for directed evolution of enzymes that activate genotoxic  
226 prodrugs. *Protein Eng Des Sel* 27:399–403.

227 Copp JN et al. (2017) Engineering a multifunctional nitroreductase for improved activation  
228 of prodrugs and PET probes for cancer gene therapy. *Cell Chem Biol* 24:391-403

229 Dontsova KM, Pennington JC, Hayes C, Simunek J, Williford CW (2009) Dissolution and  
230 transport of 2,4-DNT and 2,6-DNT from M1 propellant in soil. *Chemosphere* 77:597–603.

231 Hannink NK, Subramanian M, Rosser SJ, Basran A, Murray JA, Shanks JV, Bruce NC (2007)  
232 Enhanced transformation of TNT by tobacco plants expressing a bacterial nitroreductase. *Int*  
233 *J Phytoremediation* 9:385-401.

234 Mayer KM, Arnold FH (2002) A colorimetric assay to quantify dehydrogenase activity in  
235 crude cell lysates. *J Biomol Screen* 7:135-140.

236 Mowday AM et al. (2016) Rational design of an AKR1C3-resistant analog of PR-104 for  
 237 enzyme-prodrug therapy. *Biochem Pharmacol* 116:176-187.  
 238 Prosser GA et al. (2010) Discovery and evaluation of *Escherichia coli* nitroreductases that  
 239 activate the anti-cancer prodrug CB1954. *Biochem Pharmacol* 79:678-687.  
 240 Prosser GA et al. (2013) Creation and screening of a multi-family bacterial oxidoreductase  
 241 library to discover novel nitroreductases that efficiently activate the bio-reductive prodrugs  
 242 CB1954 and PR-104A. *Biochem Pharmacol* 85:1091-1103.  
 243 Roldan MD, Perez-Reinado E, Castillo F, Moreno-Vivian C (2008) Reduction of  
 244 polynitroaromatic compounds: the bacterial nitroreductases. *FEMS Microbiol Rev* 32:474-  
 245 500.  
 246 Rich MH, Sharrock AV, Hall KR, Ackerley DF, MacKichan JK (2018) Evaluation of NfsA-like  
 247 nitroreductases from *Neisseria meningitidis* and *Bartonella henselae* for enzyme-prodrug  
 248 therapy, targeted cellular ablation, and dinitrotoluene bioremediation. *Biotechnol Lett*  
 249 40:359-367.  
 250 Van Dillewijn P, Couselo JL, Corredoira E, Delgado A, Wittich RM, Ballester A, Ramos JL  
 251 (2008) Bioremediation of 2,4,6-trinitrotoluene by bacterial nitroreductase expressing  
 252 transgenic aspen. *Environ Sci Technol* 42:7405–7410.  
 253 Williams EM et al. (2015) Nitroreductase gene-directed enzyme prodrug therapy: insights  
 254 and advances toward clinical utility. *Biochem J* 471:131-153.  
 255 Zhang L, Rylott EL, Bruce NC, Strand SE (2019) Genetic modification of western wheatgrass  
 256 (*Pascopyrum smithii*) for the phytoremediation of RDX and TNT. *Planta* 249:1007-1015.

**Table 1. Nitroreductase candidate library**

<b><i>Bacterial strain nitroreductase sourced from</i></b>	<b>Nitroreductase gene</b>	<b>UniProt identifier</b>
<i>Bacillus amyloliquefaciens</i> CAMR 0454	<i>ywrO</i>	Q8VSR5
<i>Bacillus coagulans</i> ATCC 7050	<i>nfsA</i>	G2TLJ0
<i>Bacillus licheniformis</i> ATCC14580	<i>nfrA</i>	Q65DM9
	<i>ycnD</i>	Q65ND8
	<i>yfkO</i>	Q65MG6
<i>Bacillus subtilis</i> NZ isolate	<i>nfrA</i>	P39605
	<i>ycnD</i>	P94424
	<i>ydgl</i>	P96707
	<i>yfkO</i>	O34475
	<i>ywrO</i>	P80871
<i>Bacillus thuringiensis</i> serovar konkukian str. 97-27	<i>nfsA</i>	Q6HKT9
<i>Citrobacter koseri</i> ATCC 27156	<i>nfsA</i>	A8AIR6
	<i>nfsB</i>	A8AJP2
<i>Cronobacter</i> (previously <i>Enterobacter</i> ) <i>sakazakii</i> ATCC 29544	<i>nfsA</i>	A7MF39
	<i>nfsB</i>	A7MK61
<i>Erwinia carotovora</i> subsp. <i>atroseptica</i> SCRI1043	<i>nfsA</i>	Q6D3R4
<i>Escherichia coli</i> W3110	<i>azoR</i>	P41407
	<i>mdaB</i>	P0AEY5
	<i>nemaA</i>	P77258
	<i>nfsA</i>	P17117
	<i>nfsB</i>	P38489
	<i>wrbA</i>	A1A9Q9
	<i>ycaK</i>	P43340
	<i>ycdI</i>	P75894
	<i>ydjA</i>	P0ACY1
	<i>yieF</i>	P0AGE6
<i>Klebsiella pneumoniae</i> ATCC 13883	<i>nemaA</i>	A6T9Z9
	<i>nfsA</i>	A6TT6U7
	<i>nfsB</i>	A6T5Y2
	<i>ycdI</i>	A6T798
<i>Lactobacillus sakei</i> subsp. <i>sakei</i> 23K	<i>nfsA</i>	Q38UW5
<i>Listeria innocua</i> Clip11262	<i>nfsA</i>	Q92D83
	<i>ywrO</i>	Q92AM4
<i>Listeria welshimeri</i> ATCC 35897	<i>nfsA</i>	A0AH51
<i>Mycobacterium smegmatis</i> mc <sup>2</sup> 155	<i>nfsA</i>	A0QWF2
<i>Neisseria meningitidis</i> strain H44/76	<i>nfsB</i>	E6N0E6
<i>Nostoc punctiforme</i> PCC73102	<i>nfsA</i>	B2J8C5
<i>Pseudomonas aeruginosa</i> PAO1	<i>ycaK</i>	Q9I4B3
	<i>ydgl</i>	Q9HTZ9
	<i>yieF</i>	O31038
<i>Pseudomonas putida</i> KT2440	<i>azoR</i>	Q88EC8
	<i>nfsA</i>	Q88K03



	<i>nfsB</i>	Q88K59
<i>Pseudomonas savastanoi</i> pv. <i>phaseolicola</i>	<i>mdaB</i>	Q48NL7
	<i>wrbA</i>	E7PSG4
<i>Salmonella typhi</i> ATCC 19430	<i>azoR</i>	P63462
	<i>nema</i>	Q8Z6P3
	<i>nfsA</i>	Q8Z848
	<i>nfsB</i>	Q8Z8M5
<i>Vibrio fischeri</i> ATCC 7744	<i>frasel</i>	P46072
	<i>nfsA</i>	B5EVB7
<i>Vibrio harveyi</i> ATCC 14126	<i>frp</i>	Q56691
	<i>nfsB</i>	A6AP86
<i>Vibrio harveyi</i> KCTC 2720	<i>frp</i> , herein referred to as <i>frp2</i>	AAM73721 (Genbank)
<i>Vibrio vulnificus</i> ATCC 27562	<i>azoR</i>	Q8DA68
	<i>nfsA</i>	Q8D4B5
	<i>nfsB</i>	Q7MCD2
	<i>nema</i>	Q8D549

**Table 2. Dinitrotoluene IC<sub>50</sub> values for *E. coli* 7NT pUCX:*nitroreductase* strains**

Nitroreductase	IC <sub>50</sub> (μM)	
	2,6-dinitrotoluene	2,4-dinitrotoluene
YfkO_Bs	670 ± 60	330 ± 60
NfsB_Nme	680 ± 70	310 ± 60
NfsB_Vv	800 ± 70	510 ± 110
NfsB_Cs	940 ± 130	530 ± 60
YfkO_Bli	>1000	470 ± 40
NfsB_Pp	>1000	700 ± 140
NfsB_Kp	>1000	740 ± 20
FraseI_Vf	>1000	750 ± 50
NfsB_Vh	>1000	770 ± 60
NfsB_St	>1000	790 ± 20
NfsB_Ck	>1000	800 ± 60
NfsB_Ec	>1000	840 ± 60

**Table 3. Michaelis-Menten kinetic parameters for reduction of 2,4-dinitrotoluene and 2,6-dinitrotoluene by purified His<sub>6</sub>-tagged nitroreductases**

Nitroreductase	Substrate	$K_M^1$ ( $\mu\text{M}$ )	$k_{cat}^1$ ( $\text{s}^{-1}$ )	$k_{cat}/K_M$ ( $\text{M}^{-1} \text{s}^{-1}$ )
<i>B. subtilis</i> YfkO	2,4-dinitrotoluene	62 ± 7	140 ± 42	440,000 ± 140,000
	2,6-dinitrotoluene	510 ± 80	22 ± 2	43,000 ± 7,000
<i>V. vulnificus</i> NfsB	2,4-dinitrotoluene	320 ± 50	200 ± 20	640,000 ± 110,000
	2,6-dinitrotoluene	1600 ± 110	90 ± 4	57,000 ± 5,000

<sup>1</sup>Apparent kinetic parameters as measured at 250  $\mu\text{M}$  NADPH

## Figure Legends

### **Figure 1. NADPH consumption by *E. coli* 7NT pUCX:*nitroreductase* cell lysates in the**


**presence of 2,4- or 2,6-dinitrotoluene compounds. A.** Crude cell lysates (40  $\mu$ l) of nitroreductase-expressing *E. coli* 7NT strains were incubated with 200  $\mu$ M NADPH and 150  $\mu$ M of either 2,4-dinitrotoluene (2,4-DNT) or 2,6-dinitrotoluene (2,6-DNT) for 30 min. Addition of nitroblue tetrazolium post-incubation yielded formazan dye in proportion to the remaining NADPH, which was quantified by measuring absorbance at 490 nm. Plotted values indicate the percentage of NADPH consumed by dinitrotoluene-challenged lysates during the reaction compared to unchallenged duplicate controls. Individual *E. coli* 7NT pUCX:*nitroreductase* strains are labelled according to the name of the nitroreductase (NTR) followed by a two- or three-letter abbreviation that defines the genus and species of the bacteria that the enzyme was derived from (a full list of all NTRs tested in this study and their accession codes is provided in the Materials and Methods). Data are the average of three independent assays  $\pm$  SD. **B.** To delineate the activities of NfsB-family NTRs, the assay was repeated as in panel A, only using a smaller volume of crude cell lysate (5  $\mu$ l) and shorter incubation time (10 min).

### **Figure 2. Scatter diagram and correlation analysis of NfsB-family nitroreductase activities**


**with 2,4-dinitrotoluene in nitroblue tetrazolium and IC<sub>50</sub> assays.** Correlation analysis showed that 2,4-dinitrotoluene (2,4-DNT) IC<sub>50</sub> values for *E. coli* 7NT pUCX:*nitroreductase* strains were negatively correlated ( $p < 0.05$ , Pearson's correlation test) with the percentage of NADPH consumed by the corresponding strain lysates in nitroblue tetrazolium assays ( $r^2=0.35$ ).

**Figure 1**


**A.**



**B.**



**Figure 2**



## **AUTHOR'S QUICK CHECKLIST FOR PREPARING MANUSCRIPTS FOR SUBMISSION**

**A copy of this, appropriately checked and signed, should accompany each submitted paper.**

Your paper must follow the advice given in our **Instructions for Authors** and **Guidance Notes** that are available online (<http://www.springer.com/life+sciences/microbiology/journal/10529>). If you are uncertain about the relevancy of your paper for publication in this Journal, please read our **Aims and Scope** that is also available online.

Authors are strongly advised to consult some recent issues of this journal to familiarize themselves with the style used.

Papers that have not been correctly prepared or written to a high standard of English will be returned to the authors for revision without scientific evaluation.

Authors whose first language is not English and may be inexperienced in writing for an international journal are urged to have their manuscript thoroughly checked before submission. The use of a professional paper writing agency is strongly recommended for those authors who are doubtful of their command of the English language.

We normally allow incorrectly presented papers to be resubmitted on one occasion only.

**Please do not tick the following boxes until you have complied with the instructions indicated.**

- ☒ All authors have seen a copy of the paper AND have approved its submission.
- N/A ☒ Any animal experiments, including those involving humans, must have been approved by the correct ethics agency. A statement to this effect should be given in the text at the appropriate place.
- N/A ☒ Experiments involving plants or microorganisms taken from outside the authors' country should have been with the correct authorization.
- ☒ Authors must indicate in which section of the Journal they would like their papers to appear.
- ☒ Please indicate the corresponding author of the paper. NOTE: To avoid possible confusion, only one person should be nominated.
- ☒ All text, including Abstract, tables, references, footnotes and figure legends, is in double-spacing (**NOT 1½ spacing**). The typing must be 15 cm width. The font size should be such that there are 22-25 lines per page.
- ☒ All pages (including references, tables, figure legends, figures and any Supplementary Data) are numbered sequentially. Lines on each page are also numbered either sequentially throughout the paper or on each individual page.
- ☒ The title should be informative and clear. (A running title is not used.) Do not use unspecified, non-standard abbreviations in the title. Please do not begin the title of your paper with expressions such as "The effect of..." or "A study of...". Please ensure that the biological origin of the system under investigation is given in the title.
- ☒ Key words or phrases (but not abbreviations) are given in **alphabetical order** after the Abstract. Up to seven can be used.
- ☒ A structured Abstract for research papers is to be given. (Abstracts for reviews should be in a single paragraph and no more than 150 words.) The structured Abstract should comprise: **Objectives**: up to 35 words in a single sentence; **Results**: up to 120 words; **Conclusions**: up to 35 words in a single sentence. The writing must be clear and concise and give quantitative values of key results.

(Comparative values, that is % values, are of little use and should not be used.) Please do not begin with phrases such as "This study shows..." or "In this study...". (If you use such a phrase, it indicates you have not read these notes with sufficient care!)

- ☒ Sub-headings in all sections are clearly indicated but are **NOT** numbered.
- ☒ All non-standard abbreviations should be kept to a minimum but need to be defined at first mention **in each section** including tables and figure legends.
- ☒ Organisms must be named according to the latest available lists of plants and microorganisms. If appropriate, a synonym may be used.
- ☒ Ensure you follow the Journal style and give correct abbreviations of standard units (for example: ml, l, g, kg, h, min, s, kDa etc. Concentrations are given as 10 g glucose l<sup>-1</sup> **NOT** 10 g l<sup>-1</sup> glucose. Centrifugal force (g) is used and not rpm. Mixtures of compounds are given as, for example, Tris/HCl or chloroform/methanol/water.
- ☒ Tables and figures are to be self-explanatory so that they can be understood without recourse to the text. Key details of procedures should be given in the legend or footnotes. 100% relative values **MUST** be given as the actual (absolute) values in the legend or footnote. It is recommended that procedures used only once should be detailed in the legends to figures or footnotes to tables rather than being placed in the **Methods** section.
- ☒ Tables, each on a separate sheet, are suitable either for single (7.5 cm) or double column (16 cm) printing. Titles of tables are included with the table and appear above the table itself.
- ☒ Values in tables and text **MUST NOT** exaggerate the accuracy of the methods used. Rarely can values given to four or more significant figures be justified. Please also do not use non-significant zeros: e.g. 15.00 is given as 15.
- ☒ Figures, each on a separate sheet, are suitable for printing at 7.5 cm width; exceptionally 16 cm. All symbols are clear, all lines are of adequate thickness and equal blackness, labelling is accurate and clear. Large areas of blank spaces should be avoided. Each figure is numbered on its page.
- ☒ Legends to figures are grouped together on a separate sheet(s) which precede the figures themselves.
- ☒ Position for each table and figure should be indicated in the text by a marginal note or a clear note between paragraphs in the text.
- N/A ☒ Please ensure that photographs will print satisfactorily at the indicated column width with clear resolution of detail.
- N/A ☒ Micrographs must have a clear scale bar on the photograph with its size given in the legend.
- ☒ References agree between text and list. (Please cross-check carefully.) References in the text and list conform to the Journal style. References are given in alphabetical order. For citation of references in the list, please do not give issue numbers of journal volumes or doi numbers unless these are to papers in press that have been accepted for publication or are to papers in e-journals. For a letter-type paper, such as you are writing for this journal, it is recommended that the total number of references should not exceed 15 but this is not an absolute maximum. All authors up to a maximum of 10 should be given for each reference; above 10, please give the first four authors followed by et al.
- N/A ☒ A list of mathematical notations, if used, appears before the Introduction.



N/A

- ☒ **Supporting information.** Results supplied in the form of supplementary tables or figures, or as additional methods, should be listed as a separate section after the Acknowledgements and before the list of references. Citation of this information should be given in the text at the appropriate place using “Supplementary Table 1”, “Supplementary Figure 1” etc. There is no limit to the amount of information that can be given in this way though its usefulness for readers must be evident. The supplementary tables and figures etc. are placed at the very end of the paper. Please only give a simple but clear title of the table or figure in the list of Supporting Information.

For submission of your manuscript the following order should be observed:

- Title page giving the section in which the paper is to be considered, the title plus list of all authors plus affiliations with corresponding author (one only) indicated.
- Abstract
- Key words
- Mathematical notations (if used)
- Introduction
- Materials and methods
- Results
- Discussion (the latter two may be combined)
- Conclusions (optional)
- Acknowledgements
- Supporting information – titles only of tables, figures etc.
- List of references
- Tables
- Figure legends
- Figures
- Supplementary material as headlined in Supporting Information

David Akerley 25-May-2019