UNIVERSITY of York

This is a repository copy of Alcohol use among young Australian adults in May-June 2020 during the COVID-19 pandemic:a prospective cohort study.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/185345/</u>

Version: Accepted Version

Article:

Clare, Philip J, Aiken, Alexandra, Yuen, Wing See et al. (10 more authors) (2021) Alcohol use among young Australian adults in May-June 2020 during the COVID-19 pandemic:a prospective cohort study. Addiction. pp. 3398-3407. ISSN 1360-0443

https://doi.org/10.1111/add.15599

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

Alcohol use among young Australian adults in May–June 2020 during the COVID-19 pandemic: a prospective cohort study

Philip J. Clare^{1,2}, Alexandra Aiken², Wing See Yuen², Emily Upton², Kypros Kypri³, Louisa Degenhardt², Raimondo Bruno⁴, Jim McCambridge⁵, Nyanda McBride⁶, Delyse Hutchinson^{2,7,8,9}, Tim Slade¹⁰, Richard Mattick² & Amy Peacock²

Prevention Research Collaboration, University of Sydney, Camperdown, NSW, Australia,¹ National Drug and Alcohol Research Centre, University of New South Wales, Sydney, NSW, Australia,² School of Medicine and Public Health, University of Newcastle, Callaghan, NSW, Australia,³ School of Psychological Sciences, University of Tasmania, Hobart, Tasmania, Australia,⁴ Department of Health Sciences, University of York, York, VK,⁵ National Drug Research Institute, Curtin University, Perth, WA, Australia,⁶ Centre for Social and Early Emotional Development, Deakin University, Melbourne, VIC, Australia,⁷ Department of Paediatrics, University of Melbourne, Melbourne, VIC, Australia,⁸ Murdoch Children's Research Institute, Royal Children's Hospital, University of Melbourne, Melbourne, VIC, Australia,⁹ and The Matilda Centre for Research in Mental Health and Substance Use, University of Sydney, Sydney, NSW, Australia¹⁰

ABSTRACT

Aims To estimate change in young people's alcohol consumption during COVID-19 restrictions in Australia in early-mid 2020, and test whether those changes were consistent by gender and level of consumption prior to the pandemic. **Design** Prospective longitudinal cohort. **Setting** Secondary schools in New South Wales, Tasmania and Western Australia. **Participants** Subsample of a cohort (n = 443) recruited in the first year of secondary school in 2010–11. Analysis data included three waves collected in September 2017–July 2018, September 2018–May 2019 and August 2019–January 2020), and in May–June 2020. **Measurements** The primary predictors were time, gender and level of consumption prior to the pandemic. Outcome variables, analysed by mixed-effects models, included frequency and typical quantity of alcohol consumption, binge drinking, peak consumption, alcohol-related harm and drinking contexts. **Findings** Overall consumption (frequency × quantity) during the restrictions declined by 17% [incidence rate ratio (IRR) = 0.83; 95% confidence interval (CI) = 0.73, 0.95] compared to February 2020, and there was a 35% decline in the rate of alcohol-related harms in the same period (IRR = 0.66; 95% CI = 0.54, 0.79). Changes in alcohol consumption were largely consistent by gender. **Conclusions** From a survey of secondary school students in Australia, there is evidence for a reduction in overall consumption and related harms during the COVID-19 restrictions.

Keywords Alcohol, COVID-19, epidemiology, prospective cohort, public health, young adults.

Correspondence to: Dr Philip Clare, Prevention Research Collaboration, Level 6 The Hub, The Charles Perkins Centre (D17), The University of Sydney, Camperdown, NSW 2006, Australia. E-mail: philip.clare@sydney.edu.au

Submitted 4 October 2020; initial review completed 20 January 2021; final version accepted 26 May 2021

INTRODUCTION

The COVID-19 pandemic has had consequences beyond disease morbidity and mortality, with social and travel restrictions imposed world-wide. There are questions regarding the impact of COVID-19 on alcohol. Restrictions on gatherings could result in declining consumption within social contexts [1–3]. Conversely, the pandemic and related restrictions have generated financial and psychological distress for many Australians [4]: known risk factors for elevated alcohol consumption [5]. Further, some governments have temporarily relaxed liquor licensing

restrictions, facilitating on-line purchasing and home delivery [1].

Financial analyses showed increased alcohol purchasing in Australia in March 2020 when restrictions were implemented [6], although industry reported declines in total alcohol volume sold [1,7]. Similarly, wastewater analysis showed a decline in alcohol consumption during initial COVID-19 restrictions in Adelaide, Australia [8]. By contrast, survey research from Australia and other countries suggest that the majority of adults perceived their alcohol use as stable since COVID-19 restrictions [9–11]. Other Australian research indicated that self-reported alcohol consumption may be elevated relative to benchmark data collected before COVID-19 restrictions [12]. However, research to date largely does not account for pre-COVID-19 levels of alcohol use, highlighting the necessity for prospective, longitudinal data comparing alcohol consumption patterns before and after COVID-19 restrictions.

Alcohol use is very common among young adults in Australia: 18–24-year-olds typically report the highest rates of drinking, binge drinking and alcohol-related harms of all Australians [13]. There is evidence of an ongoing decline in consumption overall among young Australians [13]; however, it is not clear whether this trend has been impacted by the COVID-19 pandemic. Young adults are typically less financially stable [14] than older adults, and over-represented in service industries which have been particularly impacted by COVID-19 restrictions [9]. Thus, restrictions may have disproportionately impacted younger people. Conversely, young people tend to consume more alcohol outside the home [15] than older adults, so consumption may have declined with restrictions preventing gatherings.

Emerging research also suggests gender differences in the response to the COVID-19 pandemic and associated restrictions. Men in the United States were more likely to drink following restrictions than women, but their drinking was largely unaffected by stress caused by the pandemic [16]. Australian research [17] also found gender differences in alcohol use, mediated by other factors such as employment, exercise, diet and sleep. However, that study occurred early in the pandemic, when stress from isolation and economic factors may not have fully emerged.

This study used data from the Australian Parental Supply of Alcohol Longitudinal Study (APSALS) to investigate changes in alcohol consumption among young adults in May to June 2020 (i.e. after the implementation of restrictions and the first wave of the pandemic in Australia had peaked in late March), relative to consumption before March 2020. Specifically, we aimed to: (1) evaluate changes in alcohol consumption and harm, compared with pre-COVID-19; (2) test whether changes in alcohol consumption and harm were consistent in men and women; and (3) test whether changes in alcohol consumption and harm were consistent across levels of alcohol consumption prior to the COVID-19 pandemic. We also explored drivers for change in alcohol consumption.

METHODS

Sample

We used the APSALS cohort (registered at ClinicalTrials. gov: NCT02280551) of 1927 families [18]. Participants and a parent/guardian were recruited via an opt-in process in 2010 and 2011 from grade 7 classes (mean age 12.9 years) in Australian schools across three jurisdictions (New South Wales, Tasmania and Western Australia), with signed consent obtained from parents. The target sample was $n \sim 1800$ [18]. Participants completed annual surveys, with parents/guardians surveyed until wave 5. Participants were reimbursed for survey completion (\$AUD20 for wave 8; \$AUD50 for waves 9/10) and could enter a draw to win one of eight \$AUD500 vouchers. Response rates for each wave are presented in Supporting information, Appendix B.

Wave 10 data collection was under way when the first COVID-19 fatality was recorded in Australia on 15 March 2020 and social distancing restrictions came into effect in late March. We invited participants who had completed wave 10 up to April 30 and had provided an e-mail address to complete a 10-minute online survey via Qualtrics assessing alcohol use, other substance use, mental and physical health during COVID-19 restrictions (hereafter 'COVID-19 survey'). Participants could win a \$AUD500 gift voucher. Of 813 participants invited, 443 completed COVID-19 surveys (54% response rate) between 18 May and 25 June 2020 (see Supporting information, Appendix A for time-line of data collection and the COVID-19 pandemic in Australia and Supporting information, Appendix B for study flow-chart).

The study was approved by the UNSW Sydney Research Ethics Committee and ratified by the universities of Tasmania, Newcastle and Queensland and Curtin University. Findings are reported according to the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement (Supporting information, Appendix B).

Measures

Outcome variables

Five prospective outcome measures were used in addition to retrospective outcomes assessed in the COVID-19 survey. Further details of the measures are shown in Supporting information, Appendix C.

Frequency of use. Participants were asked how often they consumed alcohol. In the COVID-19 survey, participants were asked first about the month of February, and then about the past month (not at all, 1 day/month, 2–3 days/ month, 1–2 days/week, 3–4 days/week, 5–6 days/week or daily). In the APSALS annual survey they were asked about the past year (daily, 5–6 days/week, 3–4 days/week, 1–2 days/week, 2–3 days/month, 1 day/month, less often or never). This was pro-rated to past month use (see Supporting information, Appendix C for details).

Typical quantity consumed. Participants who reported any alcohol consumption were asked how many standard drinks (equivalent to 10 g of alcohol) they typically

consumed on a drinking occasion, with categorical responses in waves 8-10 (none; a sip or a taste; one to two drinks; three to four drinks; five to siz drinks; seven to 10 drinks; 11-12 drinks; or 13 or more drinks) and a continuous response in the COVID-19 survey, which was coded to the same categories as the waves 8-10 data. To limit impossibly high responses without making assumptions about maximum levels of consumption, the continuous response was capped at 100 drinks.

Frequency of binge drinking. Participants who reported any alcohol consumption were asked how often they consumed ≥ 5 standard drinks, with the same response options as for frequency of use. Responses in the annual APSALS surveys were pro-rated to the past month (Supporting information, Table C2).

Overall consumption. We multiplied frequency of use and typical quantity consumed at each time-point to create a proxy for overall consumption of alcohol in the month [19].

Maximum consumption. Those who reported any alcohol consumption in the COVID-19 survey were asked the maximum number of standard drinks they consumed on a single occasion in the past month and in February 2020.

Alcohol-related harms. The COVID-19 survey included a subset of the harms from the School Health and Alcohol Harm Reduction Project (SHAHRP) scale [20] (range = 0-8; see Supporting information, Appendix C for list of items).

Drinking contexts. The COVID-19 survey included questions about time spent drinking alone, physically with

 Table 1
 Total number of harms, report of any alcohol-related harms and report of individual alcohol-related harms, prior to and during the COVID-19 restrictions.

Alcohol-related harms	February 2020	May–June 2020
Total number of harms (mean/SD)	0.70	0.47
	(0.05)	(0.04)
Reported any harms	36.8%	26.2%
Individual harms		
Were sick after drinking	16.0%	10.8%
Had a hangover	33.9%	24.6%
Were unable to remember what	13.1%	7.4%
happened		
Got into a physical fight	0.9%	0.5%
Damaged something	1.6%	2.3%
Got into trouble with friends	1.4%	0.9%
Had sex that they later regretted	2.9%	0.5%
Had an accident, injury or fall	1.8%	1.6%

Total number of harms is calculated as the total number of individual harms experienced. Information on harms in February 2020 was obtained in the COVID-19 survey conducted in May–June 2020. Results are based on multiple imputation with estimates combined using Rubin's rules. SD = standard deviation.

others or virtually with others, coded as the proportion of time drinking from 0 to 100.

Use of delivery services. Participants were asked in the COVID-19 survey how often they had alcohol delivered to their home, coded as a binary (no/yes) variable of any use of alcohol delivery services.

Self-reported change in alcohol use. Participants were asked to self-report whether their alcohol use had changed since the COVID-19 restrictions commenced in March. Those reporting increased or decreased consumption were asked to select from a list of possible reasons why the change occurred (with multiple selections possible; Table 1).

Predictors

The primary predictor was time to assess changes before and during the COVID-19 restrictions. The primary time variable was a categorical variable with five time-periods: wave 8 (September 2017–July 2018), wave 9 (September 2018–May 2019), wave 10 (August 2019–January 2020), COVID-19 survey pre-restrictions (February 2020; reported retrospectively) and COVID-19 survey post-restrictions (May–June 2020). For sensitivity analyses we coded the quarter and year in which the survey was completed (or to which data referred to). We also coded a binary variable of alcohol consumption in wave 8 (above or below/equal to the median for consumption).

Other variables

Participants were asked about their experiences during the COVID-19 pandemic, including whether they had been tested for/diagnosed with the disease, and a range of other potential impacts, such as being required to quarantine or losing employment. Further details are available in Supporting information, Appendix C.

Statistical analysis

We included a range of socio-demographic characteristics of the sample, with means and standard deviations (for continuous measures and ns and percentages for categorical measures. We compared the characteristics of participants who completed the COVID-19 survey to all participants who completed ASPALS wave 8, with *t*-tests and χ^2 tests to assess differences between responders and non-responders (including those not invited to the COVID-19 survey). We reported descriptive statistics on alcohol consumption of participants over time. We compared the subjective report of change in alcohol use to a measure of change derived from change in overall consumption between February 2020 and May–June 2020.

We conducted analyses of change in alcohol use using mixed effects models to control for repeated observations for each participant. Primary analyses were conducted using negative binomial regression, with results reported as incidence rate ratios (IRRs). Because the primary difference of interest was with alcohol use immediately prior to the pandemic, we used February 2020 (or quarter 1, 2020 for analyses using calendar time) as the referent in all analyses. Although participants who completed wave 10 up to April 30 were invited to participate in the COVID-19 survey, to maintain separation between the wave 10 data and the COVID-19 restrictions only wave 10 responses to the end of January 2020 were included. We analysed alcohol outcomes that were only asked in the COVID-19 survey with mixed-effects models across two time-periods (February 2020 and May–June 2020) using linear regression (maximum consumption and drinking contexts), negative binomial regression (alcohol-related harms) and logistic regression (use of delivery services). Results were reported as changes in mean, IRRs and odds ratios (ORs), respectively.

To examine whether changes over time differed by gender, we reconducted all models with a gender \times time interaction effect. Two participants identified as a gender other than male or female and were

Table 2Comparison of sociodemographic characteristics of the COVID-19 survey subsample to full sample of all those who completedWave 8

		Full sample, Wave 8	COVID Survey Subsample	Statistic; p-value
Mean monthly alcohol consumption (SD)		18.7 (19.8)	17.5 (19.7)	t(913)=1.13;
Mean age (SD)		19.8 (0.5)	19.7 (0.5)	t(842)=1.18; p=0.239
Gender	Male	49.3%	38.6%	$\chi^2(1)=27.75;$
	Female	50.7%	61.4%	p=0.000
Income	\$1-\$12,999	42.8%	46.6%	χ2(3)=3.82; p=0.281
	\$13,000-\$31,199	39.8%	38.1%	
	\$31,200-\$67,599	15.8%	14.0%	
	\$67.600+	1.5%	1.3%	
Single parent household	No	62.8%	64.4%	$\chi^{2(1)=0.57}$; p=0.449
	Yes	37.2%	35.6%	
Older siblings	No	48.1%	48.5%	$\chi^{2}(1)=0.02; p=0.888$
5	Yes	51.9%	51.5%	X (C) IIIII
Parent born in Australia	No	26.5%	27.9%	$\chi 2(1)=0.50; p=0.481$
	Yes	73.5%	72.1%	
Parent education	High school or less	31.7%	26.3%	$\gamma 2(2) = 15.77;$
	Diploma, Trade, non-trade	32.1%	29.7%	p=0.000
	University degree	36.3%	44.0%	Ĩ
Parent religiosity	Not/a little	70.5%	65.8%	$\gamma 2(1) = 6.04$; p=0.014
5	Pretty/verv	29.5%	34.2%	X (C) IIIII
Parent employment	Employed (full-time/part- time)	81.1%	83.4%	χ2(2)=2.30; p=0.317
	Unemployed - in workforce	12.5%	11.3%	
	Unemployed - not in workforce	6.4%	5.2%	
Mean parent demandingness (SD)		23.7 (3.7)	23.7 (3.5)	t(775)=-0.01; p=0.993
Mean parent responsiveness (SD)	29.7 (4.3)	29.8 (4.3)	t(739)=-0.48; p=0.631
SEIFA	Low	17.3%	13.8%	$\gamma 2(2) = 5.57; p = 0.062$
	Medium	21.9%	21.3%	
	High	60.8%	64.9%	
Family history of alcohol	No	51.1%	54.6%	$\chi^{2}(1)=2.57$; p=0.109
problems	Yes	48.9%	45.4%	
Mean peer substance use (SD)		14.7 (4.3)	14.4 (4.1)	t(884)=1.48; p=0.140
Mean peer disapproval of substance use (SD)		2.6 (1.8)	2.8 (1.8)	t(817)=-1.92; p=0.055

excluded from all analyses. To examine whether changes differed as a function of the amount of alcohol consumed, we conducted secondary analyses using an interaction between time and wave 8 consumption. As a *post-hoc* sensitivity analysis, we conducted an analysis based on Australian government guidelines [21], with participants classified based on reported binge drinking more than once in the month or heavy drinking (10+ drinks per week) at wave 8.

We conducted sensitivity analyses with time considered as continuous, grouped into years/quarters (e.g. quarter 1, 2020) rather than split categorically into waves. Another sensitivity analysis was conducted using the same continuous time measure, but excluding the retrospective data collected concerning February 2020. For both sensitivity analyses, all wave 10 data up to 30 April 2020 were included.

All regression models controlled for confounding, using waves 1 and 8 variables. The confounding variables included: child gender and age [22,23], socio-economic status of area of residence [24], family history of alcohol problems [25], having older siblings [22] and peer substance use and disapproval of alcohol and tobacco use [22,23,25]. More details on the confounding variables are included in Supporting information, Appendix C. Analysis was conducted using the 'mi' commands and mixed effects models (mixed, meologit, menbreg) in Stata version 16.1 [26]. We used a critical P-value of P < 0.05, and all results of inferential statistics are presented with 95% confidence intervals (CIs). The analysis code is available at https://www.philipclare. com/code/apsals/. The specific analyses in this study were not pre-registered and should be considered exploratory.

Missing data

To reduce the chance of bias due to missing data, we used multiple imputation, implemented using the R package 'mice' [27] and combined using the package 'Amelia' [28] in R version 4.0.3 [29]. Further details of missing data and the imputation are included in Supporting information, Appendix D.

RESULTS

Sample characteristics

Participants who completed the COVID-19 survey (n = 443) largely recruited from Tasmania (37.5%), Western Australia (26.9%) and New South Wales (25.7%) (Supporting information, Table E1). The COVID-19 sample was broadly similar to the APSALS cohort, although there was a higher proportion of women (60.8 compared with 49.3%, respectively; Table 2). The COVID-19 sample also came from slightly more educated (44.0% of parents with university education versus 36.3%) and religious (34.2% of parents' religious versus 29.5%) backgrounds (Table 2).

Six per cent of COVID-19 survey respondents reported having been tested for SARS-CoV-2, and one participant reported having received a positive test result. Three-quarters (74%) reported voluntary home isolation (i.e. only leaving their primary residence for governmentspecified essential reasons, including grocery shopping and medical appointments), and one in 10 (9%) reported that they had been in quarantine (i.e. at risk of contracting COVID-19 and required to quarantine for 14 days). Further details of restrictions during the COVID-19 pandemic in Australia are included in Supporting information, Appendix A [30]. There was relatively little concern about

		September 2017–July 2018	September 2018–May 2019	August 2019–January 2020	February 2020	May–June 2020
Frequency of alcohol	None	23.4%	21.9%	21.1%	14.6%	21.8%
consumption	Yes: less than weekly	46.5%	46.9%	41.4%	48.6%	42.3%
	Yes: at least weekly	30.1%	31.2%	37.5%	36.8%	35.9%
Typical consumption	None	23.4%	21.9%	21.1%	14.7%	21.8%
	Up to four standard drinks	31.0%	35.0%	46.3%	53.3%	60.3%
	5 or more standard drinks	45.5%	43.1%	32.6%	32.0%	17.9%
Binge drinking	None	40.3%	42.0%	45.7%	35.3%	50.0%
	Yes: less than weekly	43.4%	43.3%	36.9%	50.6%	40.8%
	Yes: at least weekly	16.2%	14.7%	17.5%	14.1%	9.2%

Table 3Raw trends in frequency of alcohol consumption, typical consumption and binge drinking reported by the sample over the fivewaves of the study.

Results are based on multiple imputation with estimates combined using Rubin's rules.

(a) Frequency of consumption

(b) Typical quantity consumed





(d) Total consumption (Frequency x quantity)



Figure 1 Analysis of change in frequency, typical quantity, binge drinking and total consumption of alcohol consumption over time: results of negative binomial mixed effects models. Outcomes were analysed using negative binomial mixed-effects models, with February 2020 as the reference category, and results presented as exponented coefficients corresponding to incidence rate ratios (IRRs), with bounds of 95% confidence intervals shown in dashed lines. Models are adjusted for covariates; full results are included in Supporting information, Table E2. Results are based on multiple imputation with estimates combined using Rubin's rules

COVID-19 itself, with 17% reporting that they were at least moderately worried about contracting the disease.

Alcohol use

Between 76.6 and 85.4% of respondents reported consuming alcohol in each period from September 2017 to June 2020 (Table 3), while 50.0–64.7% reported binge drinking.

There was evidence of increased frequency of consumption which did not abate during the pandemic, with frequency of alcohol consumption in May-June 2020 31% higher than in wave 8 (Fig. 1a, Supporting information, Table E3), but not different to February 2020 (IRR = 1.08; 95% CI = 0.98, 1.19). In contrast, the typical quantity consumed per occasion in May-June 2020 declined by 27% from February 2020 (Fig. 1b), continuing a downward trend since wave 8 (41% lower than wave 8). Binge drinking was stable throughout the period to February 2020, but then declined by 28% in May-June 2020 (IRR = 0.72; 95% CI = 0.63, 0.83; Fig. 1c). Overall consumption (frequency \times quantity) declined 17% from February 2020 (IRR = 0.83; 95% CI = 0.73, 0.95; Fig. 1d).

Respondents reported a lower maximum number of drinks consumed in May-June 2020 compared with February (coeff. = -1.58; 95% CI = -1.99, -1.16; Table 4. and Supporting information, Table E3). More than a third (36.8%) reported experiencing an alcohol-related harm in February 2020, which declined to 26.2% in May-June 2020 (Table 5). Similarly, there was an approximately 30% decline in the number of harms reported (IRR = 0.65; 95% CI = 0.54, 0.79; Table 4.).

In May-June 2020, compared to February 2020, there were increases in the proportion of drinking alone reported by participants (coeff. = 2.15; 95% CI = 0.50, 3.81) and in drinking with others virtually (coeff. = 7.80; 95% CI = 5.57, 10.02) and decreases in drinking with others in person (coeff. = -13.80; 95% CI = -17.56, -10.04) and in the use of alcohol delivery services (OR = 0.68; 95% CI = 0.43, 1.08).

Gender differences

Changes in alcohol consumption were relatively consistent by gender, although men showed higher incidence rates of

(c) Frequency of binge drinking

	Maximum alcohol	Number of	Drinking	Drinking with	Drinking with	Use of alcohol
	consumed in month ^a	alcohol-related harms ^b	alone ^c	others in person ^c	others virtually ^c	delivery services ^d
	Coeff.	IRR	Coeff.	Coeff.	Coeff.	OR
	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)
February 2020 May/June 2020	Ref -1.58 (-1.99, -1.16)	Ref 0.65 (0.54, 0.79)	Ref 2.15 (0.50, 3.81)	Ref -13.80 (-17.56, -10.04)	Ref 7.80 (5.57, 10.02)	Ref 0.68 (0.43, 1.08)

Table 4Change in maximum alcohol consumed, number of alcohol-related harms, drinking context and use of delivery services duringCOVID-19 restrictions compared to February 2020: results of mixed-effects models.

^aAnalysed using linear mixed-effects models, with results presented as a coefficient equating to the difference in the mean maximum alcohol consumed. ^bAnalysed using negative binomial mixed-effects models, with results presented as incidence rate ratios (IRRs). ^cAnalysed using linear mixed-effects models with results presented as coefficients equating to the difference in the mean proportion of consumption in each setting. ^dAnalysed using logistic mixed-effects models, and presented as odds ratios (ORs). Models are adjusted for covariates; full results are included in Supporting information, Table E4. Results are based on multiple imputation with estimates combined using Rubin's rules.

 Table 5
 Self-reported change in alcohol consumption due to

 COVID-19
 restrictions, with reasons for the change.

Self-reported change in drinking since before the COVID-19 restrictions	%
Increased	19.4
Staved the same	57.0
Decreased	23.5
Reasons for increased consumption $(n = 75)$	
More bored/less things to occupy time	78.0
More time to use alcohol	55.7
To cope with anxiety/stress of day-to-day activities (e.g. looking after children, work)	40.2
Greater anxiety/depression with COVID-19	37.0
Larger amounts available on hand because I stocked up	29.0
To cope with loneliness	26.3
More money to buy alcohol	17.9
Buying alcohol more on-line/via delivery services	14.5
Reasons for decreased consumption $(n = 116)$	
Fewer opportunities to be with people/go out to drink	91.2
Didn't feel like using alcohol	52.5
Because of work/study commitments	34.5
Less money to spend on alcohol or saving money	27.8
Avoiding leaving home to obtain alcohol	26.1
Worried about effects on my mental health	26.0
Worried about effects on my physical health	22.3
Living arrangements made it difficult to use alcohol	17.5
To avoid the hangover	17.4
Worried that my use was out of control	13.3
Worried effects of alcohol may make symptoms worse if get COVID-19	12.7
Other people were worried about my use	12.6

Results are based on multiple imputation with estimates combined using Rubin's rules.

drinking across the period than women (Supporting information, Fig. E1). However, while there were significant declines in typical quantity consumed and binge drinking for both men and women (Supporting information, Figs E1b and E1c), there was a decline in overall consumption only among women (IRR = 0.72; 95% CI = 0.55, 0.94; Supporting information, Fig. E1d and Table E4).

Both men and women showed declines in the maximum amount of alcohol they consumed on an occasion, with mean declines of 1.74 drinks among men (95% CI = -2.41, -1.07) and 1.47 among women (95% CI = -1.99, -0.95; Supporting information, Fig. E2a and Table E5). Patterns of alcohol-related harms were consistent across men and women (Supporting information, Fig. E2b). Men drank alone more in May–June than February (Supporting information, Fig. E2c and Table E6), but women did not. Otherwise, the results for drinking context were similar to the overall results. There was no change in the use of alcohol delivery services by gender (Supporting information, Fig. E2f).

Alcohol use by level of consumption prior to the COVID-19 pandemic

Those in the upper half of drinkers drank more frequently, and in greater quantity, than the lower half of drinkers. Both groups showed similar trends during the course of the study, with no change in frequency, but declines in typical quantity consumed and frequency of binge drinking. However, only those in the upper half of drinkers showed a decline in overall consumption (Supporting information, Fig. E3 and Table E7). Similarly, both the upper and lower half of drinkers showed declines in the maximum number of drinks consumed (Supporting information, Fig. E4a and Table E8) and the number of alcohol-related harms experienced (Supporting information, Fig. E4b). Neither group reported changes in use of alcohol delivery services (Supporting information, Fig. E4f and Table E9). Sensitivity analysis split by 'high-risk' drinking based on binge drinking or consuming > 10 drinks per week were very

similar to the primary analysis (Supporting information, Figs E5–E6 and Tables E10-E12).

Drivers of change in alcohol use

Approximately one in five participants (19.4%) perceived that their alcohol use had increased since the COVID-19 restrictions, while approximately a quarter (23.5%) said that their use had decreased. Compared to a measure of change derived from the amount of alcohol reported consumed in in February 2020 and May-June 2020 there was a relatively high degree of discrepancy, with 61% in the same category (alcohol use decreased on both, alcohol use stayed the same on both or alcohol use increased on both measures; Supporting information, Table E13). Of those who reported increased consumption, the most common reasons endorsed related to boredom and having fewer things to occupy them (78.0%) and having more time to drink (55.7%; Table 5). The most common reason for decreased consumption was having fewer opportunities to go out or be with people (91.2%).

Sensitivity analyses using continuous time

Results of the sensitivity analyses using continuous rather than categorical time were broadly consistent with the primary analyses, with increases in the frequency of consumption, but decreases in typical quantity consumed (Supporting information, Fig. E7). These results were similar when split by gender (Supporting information, Fig. E8) and overall drinking in wave 8 (Supporting information, Fig. E9). These results were largely unaffected by the omission of the retrospective data for February 2020 (Supporting information, Figs E10–E12).

DISCUSSION

We examined the impact of the COVID-19 pandemic and associated restrictions on alcohol consumption among a cohort of young adults, finding declines in typical quantity consumed, overall consumption and frequency of binge drinking.

The use of alcohol is the leading risk for disabilityadjusted life-years in 10–24-year-olds [31], as well as being linked to a range of other health burdens [32,33]. As such, altered patterns of alcohol consumption during the pandemic, with young people less likely to drink in large quantities, is of considerable interest. The number of acute alcohol-related harms (which are more strongly associated with quantity consumed per drinking occasion than frequency of drinking) [34] saw a similar decline. This decline may be driven by the fact that drinking was more likely to occur alone or 'virtually' with others due to the need to Our findings broadly align with research showing that alcohol consumption in Australia declined due to restrictions early in the pandemic [35], although other research has shown an increase in quantity of alcohol consumed during the pandemic in Australia and internationally [36]. In contrast, past research has shown greater increases in alcohol consumption among women than men during pandemics [37] while this study found relatively similar changes in both men and women, with pre-existing gender differences remaining.

The changes observed in drinking context also align with other research showing that young people consume more alcohol outside the home [15]. With restrictions preventing people from going out, 'drinking with others' was expected to decline. However, this was matched largely by increases in drinking 'virtually' with others. That is, it appears that alcohol consumption in this demographic remains a social activity, with only the medium changing.

There were differences based on level of alcohol consumption pre-COVID-19, with declines in overall consumption among those drinking at heavier levels before COVID-19 but not those drinking at lower levels. This suggests that studies examining population-level changes may miss important nuances in individual-level changes in consumption. While some of this may be attributable to regression to the mean, the changes in the higher consumption group mirrored the overall changes in the sample while the lower consumption group was more stable, suggesting that declines in overall consumption were driven by reduced consumption by heavier drinkers.

Importantly, the sample reported relatively little concern about the COVID-19 disease itself, and only one respondent had been diagnosed with the disease. Rather, changes in consumption appear to be driven by the COVID-19 restrictions although, given the range of resulting social and economic impacts, the precise cause remains unclear. In addition, while restrictions have softened since these data were collected, the pandemic is ongoing and some restrictions will probably remain for some time. As such, it is likely that other impacts may emerge that were not present at the time of this survey and may have implications for alcohol consumption.

Strengths and limitations

A major strength of our study is that we used prospective longitudinal data, and thus do not rely upon delayed recall of alcohol consumption prior to the pandemic, an issue underlined by the discrepancy between self-reported change and changed derived from prospective data. However, not all items were included in our prospective survey, and thus some analyses rely upon recall concerning February 2020, a period 3–4 months prior to the survey completion. Nevertheless, sensitivity analyses were similar regardless of whether or not they included the data for February 2020, providing some reassurance to the reliability of the data collected. In addition, the period covered by the COVID-19 survey was shorter than the annual APSALS surveys—thus, seasonal differences may affect the COVID-19 survey that are less present in the main data.

We also note that the APSALS cohort were recruited using an opt-in process, and thus results may not generalize to the general population of young adults. However, APSALS has similar levels of alcohol use and a comparable demographic profile to the Australian population, although families of lower socio-economic status are under-represented [18]. While the APSALS sample was broadly representative, the COVID-19 survey subsample was more likely to be female. However, as the aim of the study was to investigate trends rather than prevalence, this should not impact the reliability of the findings.

CONCLUSIONS

We present evidence indicating decreased overall alcohol consumption, binge drinking and alcohol-related harms among young Australians during the COVID-19 restrictions, although this differed by levels of alcohol consumption prior to the pandemic.

Acknowledgements

This research includes computations conducted using the computational cluster Katana supported by Research Technology Services at UNSW Sydney.

Author contributions

Philip Clare: Conceptualization; data curation; formal analvsis; funding acquisition; investigation; methodology; project administration; visualization. Alexandra Aiken: Data curation; funding acquisition; investigation; project administration. Wing See Yuen: Data curation; investigation; project administration. Emily Upton: Investigation; project administration. Kypros Kypri: Conceptualization; funding acquisition; investigation; project administration. Louisa Degenhardt: Conceptualization; investigation; project administration. Raimondo Bruno: Conceptualization: funding acquisition; investigation; project administration. Jim McCambridge: Funding acquisition; investigation; project administration. Nyanda McBride: Conceptualization; funding acquisition; investigation. Delyse Hutchinson: Conceptualization; funding acquisition; investigation; project administration. Tim Slade: Conceptualization; funding acquisition; investigation. Richard Mattick: Conceptualization; funding acquisition; investigation; project administration; supervision. Amy Peacock:

Conceptualization; investigation; project administration; supervision.

References

- Colbert S., Wilkinson C., Thornton L., Richmond R. COVID-19 and alcohol in Australia: industry changes and public health impacts. *Drug Alcohol Rev* 2020; **39**: 435–40.
- Dietze P. M., Peacock A. Illicit drug use and harms in Australia in the context of COVID-19 and associated restrictions: anticipated consequences and initial responses. *Drug Alcohol Rev* 2020; 39: 297–300.
- Marsden J., Darke S., Hall W., Hickman M., Holmes J., Humphreys K., *et al.* Mitigating and learning from the impact of COVID-19 infection on addictive disorders. *Addiction* 2020; 115: 1007–10.
- 4. Melbourne Institute *Taking the Pulse of the Nation*. Melbourne, Australia: Melbourne Institute; 2020.
- Keyes K. M., Hatzenbuehler M. L., Hasin D. S. Stressful life experiences, alcohol consumption, and alcohol use disorders: the epidemiologic evidence for four main types of stressors. *Psychopharmacology* 2011; 218: 1–17.
- Clifton K. Global Economic and Markets Research report: CBA Card Spend—week ending 3 April 2020. Sydney, Australia: Commonwealth Bank of Australia; 2020.
- Alcohol Beverages Australia. Impact of COVID-19 on the drinks industry. Sydney, Australia: Alcohol Beverages Australia; 2020. Available at: https://www.alcoholbeverage saustralia.org.au/wp-content/uploads/ABA-Industry-reporton-coronavirus.pdf (accessed 13 July 2020).
- Bade R., Simpson B. S., Ghetia M., Nguyen L., White J. M., Gerber C. Changes in alcohol consumption associated with social distancing and self-isolation policies triggered by COVID-19 in South Australia: a wastewater analysis study. *Addiction* 2021; 116: 1600–5.
- Australian Bureau of Statistics. Household Impacts of COVID-19 Survey, cat. no. 4940.0. 2020 [13/07/2020]. Available at: https://www.abs.gov.au/ausstats/abs@.nsf/mf/ 4940.0 (accessed 13 July 2020).
- Stanton R., To Q. G., Khalesi S., Williams S. L., Alley S. J., Thwaite T. L., *et al.* Depression, anxiety and stress during COVID-19: associations with changes in physical activity, sleep, tobacco and alcohol use in Australian adults. *Int J Environ Res Public Health* 2020; 17: 4065.
- Chodkiewicz J., Talarowska M., Miniszewska J., Nawrocka N., Bilinski P. Alcohol consumption reported during the COVID-19 pandemic: the initial stage. *Int J Environ Res Public Health* 2020; 17: 4677.
- Biddle N, Edwards B, Gray M, Sollis K. Alcohol Consumption During the COVID-19 Period: May 2020. COVID-19 Briefing Paper. Canberra, Australia: Australian National University; 2020.
- Australian Institute of Health and Welfare (AIHW). National Drug Strategy Household Survey 2019. Contract No.: PHE 270. Canberra, Australia: AIHW; 2020.
- 14. Sinha G., Tan K., Zhan M. Patterns of financial attributes and behaviors of emerging adults in the United States. *Child Youth Serv Rev* 2018; **93**: 178–85.
- Callinan S., Livingston M., Room R., Dietze P. Drinking contexts and alcohol consumption: how much alcohol is consumed in different Australian locations? *J Stud Alcohol Drugs* 2016; 77: 612–9.

- Rodriguez L. M., Litt D. M., Stewart S. H. Drinking to cope with the pandemic: the unique associations of COVID-19-related perceived threat and psychological distress to drinking behaviors in American men and women. *Addict Behav* 2020; 110: 106532.
- Neill E., Meyer D., Toh W. L., van Rheenen T. E., Phillipou A., Tan E. J., et al. Alcohol use in Australia during the early days of the COVID-19 pandemic: initial results from the COLLATE project. Psychiatry Clin Neurosci 2020; https://doi.org/ 10.1111/pcn.13099
- Aiken A., Wadolowski M., Bruno R., Najman J., Kypri K., Slade T., *et al.* Cohort profile: the Australian parental supply of alcohol longitudinal study (APSALS). *Int J Epidemiol* 2017; 46: e6–e.
- Gmel G., Rehm J. Measuring alcohol consumption. *Contemp* Drug Probl 2004; 31: 467–540.
- McBride N., Farringdon F., Meuleners L., Midford R. The School Health and Alcohol Harm Reduction Project: Details of Intervention Development and Research Procedures. Perth, Australia: National Drug Research Institute; 2006.
- 21. National Health and Medical Research Council Australian Guidelines to Reduce Health Risks From Drinking Alcohol. Canberra, Australia: Department of Health; 2020.
- 22. Fisher L. B., Miles I., Austin S., Camargo C. A., Colditz G. A. Jr. Predictors of initiation of alcohol use among US adolescents: findings from a prospective cohort study. *Arch Pediatr Adolesc Med* 2007; 161: 959–66.
- 23. Swendsen J., Burstein M., Case B., Conway K. P., Dierker L., He J., *et al.* Use and abuse of alcohol and illicit drugs in US adolescents: results of the national comorbidity survey–adolescent supplement. *Arch Gen Psychiatry* 2012; 69: 390–8.
- 24. Pink B. Information Paper: An Introduction to Socio-Economic Indexes for Areas (SEIFA), 2006. Canberra, Australia: Australian Bureau of Statistics (ABS); 2008.
- Kuperman S., Chan G., Kramer J. R., Wetherill L., Bucholz K. K., Dick D., *et al.* A model to determine the likely age of an adolescent's first drink of alcohol. *Pediatrics* 2013; 131: 242–8.
- StataCorp. Stata Statistical Software: Release 16. College Station, TX: StataCorp LLC; 2019.
- van Buuren S., Groothuis-Oudshoorn K. Mice: multivariate imputation by chained equations in R. J Stat Softw 2011; 45; https://doi.org/10.18637/jss.v045.i03
- Honaker J., King G., Blackwell M. Amelia II: a program for missing data. J Stat Softw 2011; 45: 47.
- 29. R Core Team R: A Language and Environment for Statistical Computing, 4.0.3 edn. Vienna, Austria: R Foundation for Statistical Computing; 2020.
- Smith P. Hard lockdown and a 'health dictatorship': Australia's lucky escape from covid-19. BMJ 2020; 371: m4910.
- Gore F. M., Bloem P. J. N., Patton G. C., Ferguson J., Joseph V., Coffey C., et al. Global burden of disease in young people aged 10–24 years: a systematic analysis. *Lancet* 2011; 377: 2093–102.
- Patton G. C., Coffey C., Cappa C., Currie D., Riley L., Gore F., et al. Health of the world's adolescents: a synthesis of internationally comparable data. *Lancet* 2012; 379: 1665–75.
- Hall W. D., Patton G., Stockings E., Weier M., Lynskey M., Morley K. I., *et al.* Why young people's substance use matters for global health. *Lancet Psychiatry* 2016; 3: 265–79.
- Midanik L. T., Tam T. W., Greenfield T. K., Caetano R. Risk functions for alcohol-related problems in a 1988 US national sample. *Addiction* 1996; 91: 1427–37 discussion 39–56.
- Newby J. M., O'Moore K., Tang S., Christensen H., Faasse K. Acute mental health responses during the

COVID-19 pandemic in Australia. *PLOS ONE* 2020; **15**: e0236562.

- Winstock A. R., Zhuparris A., Gilchrist G., Davies E. L., Puljevic C., Potts L. *et al.* GDS Special Edition on COVID-19 Key Findings Report. London, UK: Global Drug Survey; 2020.
- Rehm J., Kilian C., Ferreira-Borges C., Jernigan D., Monteiro M., Parry C. D. H., *et al.* Alcohol use in times of the COVID 19: implications for monitoring and policy. *Drug Alcohol Rev* 2020; 39: 301–4.

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Data S1 Appendix A – Timeline of survey completion

Figure A1 Dates of survey completion of each wave included in study.

Figure A2 Dates of survey completion shown against COVID-19 cases in Australia.

Figure A3 Dates of survey completion shown against stringency of Australian response to COVID-19.

Appendix B – STROBE

 Table B1 STROBE checklist.

Figure B1 Study flowchart of recruitment and assessment of APSALS cohort.

Appendix C - Measures

 Table C1 Summary of times at which each outcome measure was assessed.

 Table C2 Equivalence of frequency response options for pro-rating of annual to monthly frequency.

Table C3 Items included in reduced SHAHRP harms scale.Appendix D – Missing data

Appendix D – Missing data

 Table D1 Patterns of missing data

Appendix E – Additional results

Table E1Location of participants who completedCOVID-19 survey at time of survey completion.

 Table E2 Predictors of frequency, typical quantity, binge

 drinking and total consumption of alcohol consumption

 over time.

Table E3 Predictors of maximum alcohol consumed, num-ber of alcohol-related harms, drinking context, and use ofdelivery services – change during COVID-19 restrictions.

Figure E1 Frequency, typical quantity, binge drinking and total consumption of alcohol consumption – time by gender interaction.

Table E4 Predictors of frequency, typical quantity, bingedrinking and total consumption of alcohol consumption –time by gender interaction.

Figure E2 Change in maximum alcohol consumed, number of alcohol-related harms, drinking context, and use of delivery services – time by gender interaction.