



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/185537/>

Version: Published Version

Article:

Rousham, E.K., Goudet, S., Markey, O. et al. (2022) Unhealthy food and beverage consumption in children and risk of overweight and obesity: a systematic review and meta-analysis. *Advances in Nutrition*, 13 (5). ISSN: 2161-8313

<https://doi.org/10.1093/advances/nmac032>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Unhealthy Food and Beverage Consumption in Children and Risk of Overweight and Obesity: A Systematic Review and Meta-Analysis

EK Rousham,¹ S Goudet,¹ O Markey,¹ P Griffiths,¹ B Boxer,¹ C Carroll,² ES Petherick,^{1,3} and R Pradeilles¹

¹Centre for Global Health and Human Development, School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, United Kingdom; ²School of Health and Related Research, The University of Sheffield, Sheffield, United Kingdom; and ³National Institute for Health Research (NIHR) Leicester Biomedical Research Centre, University Hospitals of Leicester NHS Trust and University of Leicester, Leicester, United Kingdom

ABSTRACT

This WHO-commissioned review contributed to the update of complementary feeding recommendations, synthesizing evidence on effects of unhealthy food and beverage consumption in children on overweight and obesity. We searched PubMed (Medline), Cochrane CENTRAL, and Embase for articles, irrespective of language or geography. Inclusion criteria were: 1) randomized controlled trials (RCTs), non-RCTs, cohort studies, and pre/post studies with control; 2) participants aged ≤ 10.9 y at exposure; 3) studies reporting greater consumption of unhealthy foods/beverages compared with no or low consumption; 4) studies assessing anthropometric and/or body composition; and 5) publication date ≥ 1971 . Unhealthy foods and beverages were defined using nutrient- and food-based approaches. Risk of bias was assessed using the ROBINS-I (risk of bias in nonrandomized studies of interventions version I) and RoB2 [Cochrane RoB (version 2)] tools for nonrandomized and randomized studies, respectively. Narrative synthesis was complemented by meta-analyses where appropriate. Certainty of evidence was assessed using Grading of Recommendations Assessment, Development, and Evaluation. Of 26,542 identified citations, 60 studies from 71 articles were included. Most studies were observational (59/60), and no included studies were from low-income countries. The evidence base was low quality, as assessed by ROBINS-I and RoB2 tools. Evidence synthesis was limited by the different interventions and comparators across studies. Evidence indicated that consumption of sugar-sweetened beverages (SSBs) and unhealthy foods in childhood may increase BMI/BMI z-score, percentage body fat, or odds of overweight/obesity (low certainty of evidence). Artificially sweetened beverages and 100% fruit juice consumption make little/no difference to BMI, percentage body fat, or overweight/obesity outcomes (low certainty of evidence). Meta-analyses of a subset of studies indicated a positive association between SSB intake and percentage body fat, but no association with change in BMI and BMI z-score. High-quality epidemiological studies that are designed to assess the effects of unhealthy food consumption during childhood on risk of overweight/obesity are needed to contribute to a more robust evidence base upon which to design policy recommendations. This protocol was registered at <https://www.crd.york.ac.uk/PROSPERO> as CRD42020218109. *Adv Nutr* 2022;00:1–28.

Statement of Significance: This systematic review was the first to synthesize evidence on the effects of consuming unhealthy foods and beverages in children aged ≤ 10.9 y across low-, middle-, and high-income country settings. This review identified a major evidence gap from low- and middle- income countries. Consumption of sugar-sweetened beverages and unhealthy foods in childhood may increase BMI, percentage body fat, or odds of overweight/obesity, whereas consumption of artificially sweetened beverages and 100% fruit juice has less demonstrable effects on these outcomes.

Keywords: complementary food, infant and young child nutrition, diet, ultraprocessed foods, sugar-sweetened beverages, infant, child, low- and middle-income countries, cohort, GRADE approach

Introduction

Infants and children are consuming increasing amounts of foods with added sugars, high in salt, and high in saturated or *trans* fats (1, 2). Commercially prepared foods

are more likely to be high in energy, low in nutrients (energy-dense, nutrient-poor), and ultraprocessed (3, 4). Globally, the consumption of sugary and savory snacks and refined foods has been increasing across all socioeconomic groups

(5). These foods can have direct consequences on health, as well as indirect consequences through displacement of healthy foods in the diet (6). Consumption of foods that are energy-dense and nutrient-poor is a particular risk for malnutrition in socioeconomically disadvantaged groups and urban communities in low- and middle-income countries (LMIC) leading to increasing disparities in health globally (7). Exposures to sugar-sweetened beverages (SSBs) and sweet foods in childhood can contribute to sweet food preferences in later life (7). Diet quality in early life is also important for child development (6), and suboptimal diet is a preventable risk factor for noncommunicable diseases (NCDs) (8).

Existing complementary feeding guidelines were developed when the prevention of undernutrition was a primary concern (9, 10). In response to increasing rates of childhood overweight/obesity and the rising prevalence of NCDs (6), however, complementary feeding guidelines need to consider all forms of malnutrition including undernutrition, micronutrient deficiencies, and overweight or obesity (5, 11, 12).

A previous systematic review examined the impact of consuming unhealthy foods and beverages in the complementary feeding period (age 6–23 mo) in countries ranked high or very high on the Human Development Index (13). Limited evidence suggested that SSB consumption was associated with greater obesity risk in children aged <2 y, but not other growth or body composition indicators (13). A systematic review of 32 studies in high-income countries (HIC) concluded that SSB consumption promotes weight gain in children, adolescents, and adults (14). A systematic review of 100% fruit juice consumption in longitudinal studies reported non-clinically significant BMI *z*-score increases and a lack of evidence for children under 7 y (15). Positive associations were reported between ultraprocessed food (UPF) consumption and percentage body fat in children and adolescents in a systematic review including both cross-sectional and longitudinal study designs (16). Findings from cross-sectional studies provide weak evidence of associations because of the potential for reverse causality (17).

Existing studies and reviews highlight the paucity of evidence on effects of unhealthy food consumption in the complementary feeding period (1, 13). There has also been very little consideration of these effects in LMIC settings. A

review of cross-sectional and longitudinal studies in LMIC found limited and inconclusive evidence on the relation between snack food, SSB consumption, and child growth and dietary adequacy in children aged 6–23 mo (7), meaning a review of all country settings is needed.

The WHO, as part of the process of updating the guiding principles for complementary feeding (9, 10), commissioned a series of systematic review reports, one of which examined the impact of unhealthy food and beverage consumption on prespecified critical (growth, overweight/obesity and body composition; diet-related NCD indicators (cardiometabolic disease risk outcomes); displacement of healthy foods or breastmilk intake; dietary quality and diversity) and important (food or taste preferences later in life; oral health/dental caries; micronutrient deficiencies; and child development) outcomes. There is limited evidence on the effects of unhealthy food and beverage consumption in infants and young children. Furthermore, the entire childhood period is a critical window for reducing malnutrition and obesity-related NCD risk in later life (18). Hence, the aim of the current systematic review was to examine, in children aged ≤ 10.9 y, the risks of greater consumption of unhealthy foods and beverages compared with no or low consumption on overweight and obesity.

Methods

Review typology

A systematic review and meta-analysis were chosen to systematically search for, appraise, and synthesize quantitative research evidence (19). We followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses 2020 reporting guidelines (20). The study protocol for the review was registered on the International Prospective Register of Systematic Reviews, PROSPERO (<https://www.crd.york.ac.uk/prospéro/>; registration number: CRD42020218109).

Eligibility criteria

Study eligibility and inclusion criteria are shown in **Table 1** for population/participants, interventions or exposures, comparators, and outcomes (PI/ECO), and study design. We included quantitative human studies of children where age at intervention or exposure was between birth and ≤ 10.9 y, published from January 1971 with no restriction on publication language. Non-English language records were screened by native speakers with subject-specific knowledge.

Unhealthy foods and beverages were defined using both nutrient- and food-based approaches. Four main measures were used to classify foods and beverages as unhealthy, including: 1) UPFs based on the NOVA classification (22); 2) unhealthy foods and beverages according to WHO infant and young child feeding indicators (21); 3) foods high in free sugars, artificial sweeteners, and salt; and 4) foods high in saturated or *trans* fats. In addition, we included studies in which authors defined unhealthy foods using terminologies such as: “junk food,” “fast food,” “snack food,” “extra food,” “non-core food,” and “convenience food,” which also met

Funding support was received from the Food and Nutrition Action in Health Systems unit, Department of Nutrition and Food Safety, WHO (EKR, RP, PG). The funders had no role in the collection, analysis, interpretation of data, or writing of the report. ESP was supported by the NIHR Leicester Biomedical Research Centre.

Author disclosures: The authors report no conflicts of interest.

Supplemental Methods 1–4, Supplemental Tables 1–11, and Supplemental Figures 1 and 2 are available from the “Supplementary data” link in the online posting of the article and from the same link in the online table of contents at <https://academic.oup.com/advances/>.

Address correspondence to EKR (e-mail: e.k.rousham@lboro.ac.uk).

Abbreviations used: aOR, adjusted odds ratio; aRR, adjusted relative risk; ASB, artificially sweetened beverage; GRADE, Grading of Recommendations Assessment, Development, and Evaluation; HIC, high-income countries; LMIC, low- and middle-income countries; MIC, middle-income countries; NCD, noncommunicable disease; NRSI, nonrandomized study of interventions; PI/ECO, population, intervention or exposure, comparator, outcome; RCT, randomized controlled trial; RoB, risk of bias; ROBINS, risk of bias in nonrandomized studies of interventions; SSB, sugar-sweetened beverage; UPF, ultraprocessed food.

TABLE 1 Inclusion and exclusion criteria for the review of the effect of unhealthy food and beverage consumption in children aged ≤ 10.9 y and risk of overweight and obesity¹

Parameter	Inclusion criteria	Exclusion criteria
Participants/population	Human studies including both boys and girls Age at intervention or exposure: infants from birth to ≤ 10.9 y	Nonhuman studies Age at intervention or exposure > 10.9 y Studies that exclusively enroll participants with a disease or with the health outcomes of interest (listed below) Studies using hospitalized patients; severely malnourished participants, or clinical populations Studies of exclusively preterm babies (< 37 wk gestation) or exclusively babies that are low birth weight (< 2500 g) or small-for-gestational age
Independent variable (intervention or exposure)	Studies reporting (greater) consumption of unhealthy foods and beverages compared with no or low consumption Unhealthy foods defined using: 1) nutrient-based approaches [foods high in added sugars, free sugars, artificial sweeteners, fats (e.g., saturated/ <i>trans</i>), salt]; and food-based approaches including: 2) ultraprocessed foods (based on NOVA classification, excluding formula and follow-on milks); 3) unhealthy foods and beverages listed in the WHO/UNICEF infant and young child feeding guide (21); 4) food items defined by authors using terms such as "fast-food," "convenience foods," "non-core foods" Consumption defined as: 1) quantities consumed (grams per day, week, or month); 2) portion sizes; 3) frequency of consumption (per week, month, year), or consumed/not consumed	Studies not reporting consumption of unhealthy foods and beverages as per the protocol definition of consumption Studies reporting only dietary patterns (i.e., data reduction techniques such as principal component analysis) or eating practices (e.g., meals per day; snacking patterns; meal times and duration of eating episodes)
Comparator	Consumption of less or no unhealthy foods and beverages: no or low added sugar, free sugars, artificial sweeteners; less fat (or less of certain types of fat), less consumption of foods high in salt or ultraprocessed/energy-dense, nutrient-poor foods	
Study design	Randomized controlled trials Nonrandomized controlled trials (including historically controlled studies) Prospective cohort studies (including interrupted time series analyses) Retrospective cohort studies	Cross-sectional studies Trials without a control group Narrative reviews, systematic reviews and meta-analyses Case-control studies: i.e., cases with disease (e.g., diabetes) vs. controls without disease Pre/post studies without a control
Dependent variable (outcome)	Pre/post studies with a control Growth and body composition: stunting; length-for-age or height-for-age; underweight or weight-for-age; wasting or weight-for-length/weight-for-height; BMI; BMI z-score; waist circumference; prevalence of overweight or obesity; percentage body fat	
Country	All contexts (high-, middle-, and low-income countries)	NA
Date range	Articles published from 1971 onwards	Articles published before 1971
Publication status	Reports published in peer-reviewed journals	Conference abstracts, conference proceedings, unpublished data, reports, letters, editorials
Language	All languages	NA

¹NA, not applicable.

inclusion criteria based on consumption of foods high in sugar, salt, fat, or UPFs. Studies reporting dietary patterns were not included because these only provide evidence on overall combinations of foods such as "unhealthy" diets compared with "healthier" diets. Eligible studies were subsequently grouped into exposures of unhealthy beverages [SSBs; artificially sweetened beverages (ASBs), and 100% fruit juice separately] and unhealthy foods. For detailed

definitions of unhealthy foods and beverages see **Supplemental Method 1** and **Supplemental Table 1**.

Search strategy

A literature search strategy was developed and checked by an academic librarian. Scoping searches were conducted to refine the search strategy ensuring that known relevant studies had been identified. Systematic searches were run

in PubMed (MEDLINE), Cochrane CENTRAL, and Embase during December 17–23, 2020. Gray literature was not included in the systematic search due to time and budgetary constraints.

The search results were imported into Covidence software (Veritas Health Innovation), which was employed for screening titles, abstracts, and full texts. Citation alerts were set up in PubMed to flag new potentially relevant items. Additional supplementary searches included reference checking of included publications and relevant systematic reviews and consultation with subject experts for relevant published studies. Supplementary searches continued until April 30, 2021. The search strategies for each database are included in **Supplemental Table 2**.

Study selection

Duplicate records were identified automatically by the review software prior to screening. Half of the duplicate records were checked and no incorrect duplicates were identified.

All reviewers underwent training by screening the same test sample of 25 records selected at random. The results of the test screening were combined and discussed across the review team; this informed further guidance on inclusion and exclusion criteria. We amended the protocol with a change in the age inclusion criteria from <10 y to ≤ 10.9 y to ensure consistency in screening and greater inclusion of evidence.

All records included at title/abstract and full-text stage were screened by 2 reviewers independently (OM, RP, SG, PG, EKR, Natalie Pearson, Kathrin Burdenski, or Megan Stanley). Conflicting votes were considered by a third reviewer (RP or NP) and a fourth reviewer in cases of uncertainty (EKR). Two reviewers each checked 2 distinct random 10% samples of excluded records at title/abstract and full-text stage (OM, RP, SG, or EKR). Reasons for exclusions at full-text screening were recorded. Studies that met all inclusion criteria but reported data for a wider age range (e.g., 8–13 y) were included. The review team e-mailed study investigators (with a follow-up e-mail to nonresponders) to request disaggregated data for participants aged ≤ 10.9 y. Of the 16 study authors contacted, 8 responded to say that the datafile was no longer available. One study author provided disaggregated data (23).

Data collection process, data items, and effect measures

An Excel data extraction form was developed and piloted by all data extractors using a selection of 6 included articles covering different review outcomes. This resulted in modifications, and a second pilot data extraction was undertaken with all reviewers extracting data from a single article and comparing notes. Further modifications were made to finalize the data extraction form. One reviewer undertook data extraction independently (OM, RP, SG, BB, or EKR). Any data extraction queries were discussed among the team. A second reviewer (EKR) checked 50% of all records extracted for completeness and accuracy.

Full details of the information extracted from eligible studies are presented in **Supplemental Table 3**. Data extracted from studies included: 1) general information [study ID, title, authors, start and end date, study location (country, urban compared with rural), study design, study aim, aim of intervention, study funding sources, conflicts of interest, ethical approval reported]; 2) study eligibility (participant selection and randomization process), sample size, participant characteristics (age, number, and sex), duration of intervention, exposure measures (type of food consumption data, unit of measurement, and dietary assessment methods) and critical and important review outcomes, and the method of assessment of outcomes; and 3) study findings.

We extracted data on all ages of follow-up to assess longer-term outcomes. We recorded the variables adjusted for in analyses, such as education, socioeconomic status, sex, maternal age, race and/or ethnicity, other feeding practices (breast milk, infant formula, or both), and birth weight/length. Study protocols and supplementary materials were searched for data extraction if the required data were not presented in the included articles. For studies not in English, data extraction and risk of bias (RoB) were conducted with 1 review team member working alongside a researcher proficient in the native language with relevant subject expertise.

We extracted the measures of intervention/exposure effect (mean differences, ORs, β coefficients, RRs with 95% CIs, and/or P value) for the outcome of interest from all studies. We extracted data from fully adjusted models where available. If unadjusted effect measures only were reported, these were extracted. Where multiple articles from the same study were included, we extracted data that were unique to each article. In some instances, this resulted in a greater number of articles than studies (i.e., 2 articles from the same study were included if different outcomes or exposures were reported). If the same data were reported in >1 included article, we extracted data from the article that most closely addressed the review question.

RoB assessment

RoB was assessed by 2 reviewers (OM, RP, SG, BB, or EKR) independently using Covidence to ensure blinding. Reviewers noted justifications for any domains that were assessed as serious, critical, or no information. Information was checked from study protocols, clinical trial registers, and supplementary files if not presented in the included reports. Reviewers discussed discrepancies and reached consensus on each domain of the RoB tool. If agreement could not be reached, a third reviewer (RP or EKR) assessed RoB and a consensus was reached. RoB was conducted at the outcome level.

RoB for nonrandomized studies of the effects of interventions (prospective cohort studies).

The ROBINS-I tool (risk of bias in nonrandomized studies of interventions version I) was applied in accordance with Cochrane and Grading of Recommendations Assessment,

Development, and Evaluation (GRADE) considerations of observational studies as nonrandomized studies of interventions (NRSIs) (24, 25). Each of the 7 domains of the ROBINS-I tool were rated as being at low, moderate, serious, or critical RoB, or no information (26). After completing consensus on the 7 domains, the overall RoB for each study was assessed using the criteria in **Supplemental Table 4**.

RoB for RCTs.

Cochrane RoB (Version 2: RoB 2.0) was used for RCTs (25, 27). RoB 2.0 addresses 5 domains. Each domain was rated as low, some concerns, high RoB, or no information. Supporting information and justifications for judgments in each domain were recorded. After reaching consensus on the 5 domains, the overall RoB was assessed using Cochrane guidance, as presented in **Supplemental Table 5** (25).

Some studies undertook a secondary analysis of data from a previous RCT to address a research question unrelated to the original trial (28–33). The trials had either reported no significant effects of the intervention and therefore pooled the intervention and control group, or assessed the control group only. We assessed these studies as NRSIs and applied the ROBINS-I tool. Individual and summary RoB tables were produced using the “robvis” tool (34).

Data synthesis

Initial data synthesis processes.

We synthesized findings using the PI/ECO framework, first grouping studies by outcome and then by intervention/exposure. For synthesis relating to participant characteristics, we stratified by age (0 to <2 y; 2 to <5 y, and 5 to ≤10.9 y) where there were sufficient studies. For completeness, we included all estimates in summary tables of results, including studies with critical RoB. In narrative synthesis, meta-analysis, and when grading the evidence, however, we did not include results from studies assessed as critical RoB, in line with guidance (24, 35).

For the synthesis of growth, body composition, and overweight/obesity outcomes, we prioritized studies that reported BMI, BMI z-score, BMI change, BMI z-score change (or for children <2 y, weight-for-length), or prevalence of overweight/obesity because these are the most widely used indicators of growth and overweight or obesity. We then collated studies with effect estimates for percentage body fat because this was a relatively homogeneous outcome across studies. For completeness, data for other indicators such as waist circumference, central adiposity, waist-to-height ratio, and sums of skinfold thicknesses were extracted and included in summary tables, but not described in detail.

Exposures were synthesized using 2 overarching groups: 1) unhealthy beverages—this was disaggregated into SSBs alone, ASBs only, and 100% fruit juice only, where studies specifically reported these items separately (any fruit drink that was not 100% fruit juice was included within the SSB category); and 2) unhealthy foods.

Data synthesis methods.

Heterogeneity across studies arose primarily from measurement of exposure (including the dietary assessment methods, recall period, definition of food items/food groups, or units of measurement). Data reporting varied from dichotomous, multiple categories or continuous measures of consumption. Where exposures could be harmonized for the same study outcomes, we included in meta-analyses. A priori, we set a minimum requirement of 2 studies reporting the same outcomes and the same study design to produce a forest plot.

For SSB and 100% juice consumption, there were studies that could be harmonized based on the reported quantities of consumption, or number of servings, in relation to the reported outcomes, hence these were pooled for meta-analyses and corresponding forest plots. Further information on the harmonization process can be found in **Supplemental Method 2**. I^2 values were generated as indicators of heterogeneity, although these should be interpreted with caution when there are few studies in a meta-analysis. We adopted interpretative guidance for heterogeneity of 0% to 40% as not important; 30% to 60% moderate; 50% to 90% substantial, and 75% to 100% considerable heterogeneity (25). These ranges overlap because these are not absolute cut-offs. Reported β coefficients and SEs were either multiplied or divided to achieve the common serving size estimate. Random effects models were performed as recommended where heterogeneity is likely. Analyses were undertaken using the meta command in Stata SE 16 (StataCorp).

For unhealthy food consumption, we examined all studies to identify those that could be harmonized. Data on exposures and comparators for unhealthy food consumption could not be harmonized from ≥2 studies (see **Supplemental Method 3** for further details). We therefore performed narrative synthesis. We followed synthesis without meta-analysis guidelines for data synthesis without meta-analysis (35).

Reporting bias assessment

Funnel plots were not undertaken to examine potential publication bias in the meta-analysis given we could not meet the recommended number of ≥10 studies included in meta-analyses (25). Bias due to missing participants was considered within the RoB assessment using ROBINS-I for NRSIs.

Certainty of evidence

We used GRADE criteria to assess the certainty of evidence for the effect of exposures on the critical outcomes. Two reviewers (SG, EKR) independently graded the evidence as high, moderate, low, and very low, then agreed ratings through discussion and consensus. Statements defining the certainty for each grade are provided in **Supplemental Table 6**. Detailed information on the grading of the evidence can be found in **Supplemental Method 4**. Evidence profile tables were produced for each outcome using GRADEpro GDT software (GRADEpro Guideline Development Tool)

following recommended guidance (36) and disaggregated by age (<2 y; 2 to <5 y; 5 to ≤10.9 y) where there were sufficient numbers of studies. Studies of critical RoB were excluded from GRADE evidence profiles in line with guidance (25). We used standard statements to report the results of the review interventions (exposures) based on guidelines (37).

Results

Study selection

The wider search for the WHO review retrieved 35,433 studies with 8841 duplicate records detected. **Figure 1** presents the study search and selection process (20). We screened 26,542 studies of which 581 were eligible for full-text review. Of these, 579 studies were assessed for eligibility because 2 studies could not be retrieved (38, 39). After full-text screening, 161 articles from 115 studies were included. Of the included studies, 89 articles from 76 studies focused on growth and body composition outcomes. Data could not be extracted from 13 studies because they included participants younger and older than 10.9 y and results were not age-stratified (40–52). There were 3 further articles from 2 studies where data could not be extracted because of age range but data were extracted from other articles from the respective studies (53–55). Characteristics of the studies and articles where data could not be extracted are presented in **Supplemental Table 7**.

Study and participant characteristics

We summarized characteristics of 71 articles from 60 included studies. Some studies had >1 included article because different outcomes or exposures were reported in each article. The extracted data are summarized in **Supplemental Table 8**, with the country, setting, study design, baseline age, exposure details, and outcomes assessed. Studies were published from 1993 to 2020. Around 88.5% (53/60) of studies were conducted in HIC and 11.5% (7/60) in middle-income countries (MIC); no studies were from low-income countries, based on the current gross national income per capita (56). Studies in MIC were conducted in Belarus, Brazil, China, Mexico, Peru, and South Africa. One study was a pre/post design with a control (57), 1 was a retrospective cohort design (58), and 1 was an RCT (59). The remaining studies ($n = 57$) used prospective cohort designs. About 43.5% of studies included participants from urban settings; 13.3% from both rural and urban areas; and only 8.3% from rural areas. Twenty-one studies (35.0%) did not report the residence/location of participants. Sample size at baseline ranged from 72 (60) to 16,058 (61). Baseline participant age ranged from 1 mo to 10.8 y. Two studies recruited only girls as participants (62, 63). The oldest ages at follow-up were 20–21 y (63) and 21 y (64).

RoB assessment

Sixty-seven articles from 60 studies reported on growth and body composition. One study was an RCT (59), and 59

were observational studies (NRSIs). Sixty-six articles from 59 observational studies (NRSIs) were assessed for RoB. Four articles were not assessed for RoB because they did not provide any additional effect estimates to the selected article from the same study (65–68). No articles had low RoB; 32 articles (48.5%) had moderate RoB (29, 57, 58, 64, 69–96), 25 articles (37.9%) had serious RoB (60, 62, 97–119), and 8 articles (12.1%) had critical RoB (61, 120–126) (**Supplemental Figure 1**). One article (1.6%) (127) was classed as having “no information.” The ROBINS-I domains that most contributed to moderate, serious, or critical RoB were confounding bias (D1), participant selection bias (D2), bias due to intervention protocol deviations (D4), and missing data bias (D5) (**Figure 2**). The 1 included RCT study (59) was assessed as “some concerns” (**Supplemental Figure 2**).


Synthesis

SSBs and growth, body composition, and overweight/obesity outcomes.

Forty-five studies reported on SSB consumption and growth, body composition, and overweight/obesity outcomes. Some studies analyzed sodas, juice drinks, or other sweetened beverages separately (57, 63, 83, 122) whereas other studies examined multiple types of SSB as a single category. Studies were predominantly from HIC. Studies from MIC were conducted in China, Brazil, Peru, Mexico, and Belarus (70, 77, 91, 110, 115).

SSB consumption and BMI and overweight/obesity outcomes (narrative synthesis). Thirty-five of the 45 studies reported BMI outcomes (raw values, percentiles or z-scores, or change in raw/z-score values) and/or overweight/obesity prevalence (**Supplemental Table 9**).

Children aged <2 y were examined in 10 studies, 2 of which had critical RoB and are not reported on further (61, 123). Of the remaining 8 studies, 3 reported a positive association. Consumption of SSBs in early life was associated with significantly higher odds of obesity at ages 8–14 y [adjusted odds ratio (aOR) = 2.99; 95% CI: 1.27, 7.00] (serious RoB) (115). Similarly, SSB consumption >1/wk compared with ≤1/wk in infancy was associated with greater odds of overweight and obesity at age 17 mo (aOR = 1.6; 95% CI: 1.04, 1.93; $P < 0.01$) (serious RoB) (110). Pan et al. (104) reported that SSB intake at 10–12 mo was associated with significantly greater odds of obesity in the highest intake group (≥3 times/wk) compared with no consumption, but not in the intermediate intake groups (<1 or 1 to <3 times/wk) compared with no consumption (serious RoB). The same study compared “any” with “no” consumption of SSB from 1 to 12 mo and observed a higher prevalence of obesity at 6 y in the group that consumed SSBs (aOR = 1.71; 95% CI: 1.09, 2.68) (104). Three studies reported different effects based on either the time point of assessment, differences between boys or girls, or the outcome assessed. Flores and Lin (97) reported no association between SSB consumption at age 2 y and severe obesity at 5 y; only SSB



¹This term is used throughout this flowchart as Covidence software calculated numbers at the study level.

²No automation tools were used, all screened by review team.

FIGURE 1 Flow chart of study search and selection for the review of effects of unhealthy food or beverage consumption in children aged <10.9 y on risk of overweight/obesity.




FIGURE 2 Summary risk of bias assessment of nonrandomized studies reporting unhealthy food or beverage consumption and growth, body composition, and overweight/obesity outcomes assessed using ROBINS-I (risk of bias in nonrandomized studies of interventions version I) tool.

consumption at 5 y was associated with severe obesity at 5 y (aOR = 2.3; 95% CI: 1.4, 3.7) (serious RoB). Quah et al. (105) reported that SSB intake at 18 mo was not associated with BMI z-score or overweight/obesity at 6 y, but intake at 5 y was significantly associated with both outcomes ($\beta = 0.34$; 95% CI: 0.11, 0.58; $P = 0.004$; RR = 1.54; 95% CI: 1.03, 2.30; $P = 0.033$, respectively) (serious RoB). Leermakers et al. (128) found a significant association between SSB intake and BMI z-score in girls, but not in boys (girls: $\beta = 0.11$; 95% CI: 0.00, 0.23; $P = 0.04$; boys: $\beta = 0.05$; 95% CI: -0.08, 0.18; $P = 0.42$) at 6 y (moderate RoB). Two studies reported no association between consumption of SSBs and BMI or overweight outcomes (both serious RoB) (111, 112).

In children aged 2 to <5 y, 11 studies reported on SSB intake and BMI or overweight/obesity (Supplemental Table 9). In 1 study, results were not extractable (117). Four of the remaining 10 studies reported an association. In US children, SSB intakes in children aged 2–4.7 y at baseline and followed up at age 12.3–15 y were significantly positively associated with BMI z-score ($\beta = 0.05$; 95% CI: 0.022, 0.079; $P = 0.001$) (moderate RoB) (81). In Australian children, SSB intake per day was significantly positively associated with BMI z-score ($\beta = 0.017$; 95% CI: 0.007, 0.027; $P < 0.01$) (moderate RoB) (73). Consuming SSBs above compared with below the median intake (>65 mL/d compared with <65 mL/d) at 18 and 30 mo was associated with increased odds of overweight/obesity at 18-mo follow-up (aOR = 1.92; 95% CI: 1.19, 3.11; $P \leq 0.01$) and at 30-mo follow-up (aOR = 1.82; 95% CI: 1.11, 3.00; $P \leq 0.05$) (serious RoB) (118). In 1 study, total daily consumption of SSBs was not associated with obesity prevalence, but regular consumers of SSBs between meals compared with those who did not consume between meals at 2.5 y had greater odds of obesity at 4.5 y (aOR = 2.36; 95% CI: 1.03, 5.39; $P \leq 0.05$) (moderate RoB) (96). Five studies (6 articles) reported no association (all moderate RoB) (58, 70, 72, 74, 83, 93). One study reported no association between SSB consumption and odds of overweight/obesity, but significantly greater odds for

obesity alone (obese: OR = 1.65; 95% CI: 1.12, 2.44; $P = 0.01$) (moderate RoB) (80).

In children aged 5 to ≤ 10.9 y, 16 studies reported on SSB and BMI or overweight/obesity, 1 study did not report effect estimates (107), and 2 had critical RoB (120, 121) (Supplemental Table 9). Of the 13 studies with included effect estimates, all were observational designs except 1 RCT (59). In a cluster RCT in Germany, SSB intake in children was associated with significantly greater odds of obesity (aOR = 1.22; 95% CI: 1.04, 1.44; $P = 0.014$) but not overweight. There was also an association with SSB intake (per 200-mL glass/d) and BMI change ($\beta = 0.02$; 95% CI: 0.00, 0.03). In this RCT, SSB intake was a secondary outcome of the intervention (RoB: some concerns) (59). Among observational studies, 1 study reported that SSB intake per 100 g/d at age 8 y was significantly associated with BMI z-score change at 11.5 y ($\beta = 0.10$; SE = 0.03; $P = 0.003$) (serious RoB) (98). In Peru, daily compared with no intake of SSBs in the past 30 d was associated with greater BMI change ($\beta = 0.74$; 95% CI: 0.15, 1.33) and greater risk of overweight/obesity from age 8 y to 12 y [adjusted relative risk (aRR) = 2.12; 95% CI: 1.05, 4.28] (moderate RoB) (91). In US children, SSB intake at 3–5 y was associated with significantly greater odds of overweight/obesity at follow-up (aOR = 1.04; 95% CI: 1.01, 1.07; $P < 0.05$) (moderate RoB) (75). One study examined fruit drinks, non-100% fruit juice, and sodas, with only soda intake (grams per day) significantly associated with BMI ($\beta = 0.011$; SE = 0.005; $P < 0.05$) (serious RoB) (63). Eight studies reported no association between SSB intake and outcomes, 5 moderate RoB (57, 64, 76, 88, 129) and 3 serious RoB (62, 99, 116).

SSB consumption and BMI and overweight/obesity outcomes (meta-analysis). A meta-analysis of 3 studies reporting effects of SSB consumption on change in BMI from baseline to follow-up showed no effect (pooled effect estimate: $\beta = 0.01$; 95% CI: -0.00, 0.02) (57, 78, 83 (Figure 3A). Heterogeneity across studies was high




FIGURE 3 Forest plot of the effect of sugar-sweetened beverage consumption in children aged <10.9 y on BMI. (A) Effect of sugar-sweetened beverage consumption in children aged <10.9 y on BMI change (baseline to follow-up). (B) Effect of sugar-sweetened beverage consumption in children aged <10.9 y on BMI z-score values. REML, residual maximum likelihood.

($I^2 = 73.66\%$). A meta-analysis of SSB consumption and BMI z-score values showed no association (pooled effect estimate: $\beta = 0.10$; 95% CI: $-0.11, 0.31$) (3 included studies) (81, 105, 116) (Figure 3B). There was no heterogeneity across individual studies ($I^2 = 0.0\%$).

SSB consumption and percentage body fat outcomes (narrative synthesis). Seven studies examined SSB consumption and percentage body fat across all age groups (Supplemental Table 10). Three of 7 studies reported a significant positive association. SSB intake ≥ 2 servings/d compared with <1 serving/d at age 5 y was positively associated with higher percentage body fat (ANOVA group: $P < 0.01$; age: $P < 0.01$; group \times age interaction: $P < 0.01$; no F statistic reported) (serious RoB) (62). High compared with low SSB intake at 6.7 y was associated with higher percentage body fat at 2-y follow-up ($\beta = 1.40$; 95% CI: 0.09, 2.72; $P = 0.036$) (serious RoB) (116). Zheng et al. (98) also reported a positive association between SSB intake (per 100 g/d) at 9 y and percentage body fat at 11.5 y ($\beta = 1.04$; SE = 0.32; $P = 0.001$) (serious RoB). Four studies reported no association between

SSB consumption and percentage body fat, 3 with moderate RoB (72, 95, 128) and 1 with serious RoB (99).

SSB consumption and percentage body fat outcomes (meta-analysis). Figure 4 shows the effect estimates for the consumption of SSBs (per 250-mL serving) on percentage body fat. There was a significant positive association between consumption of SSBs and percentage body fat at follow-up (pooled effect estimate: $\beta = 1.86$; 95% CI: 0.38, 3.34) in 3 studies (98, 99, 116). Heterogeneity was low ($I^2 = 22.8\%$).

Certainty of evidence: SSB consumption. GRADE evidence profiles for the effect of SSB consumption and BMI, overweight/obesity, and percentage body fat are presented in Table 2. The certainty of evidence was low for all outcomes except for children aged <2 y where the certainty was very low for overweight/obesity (Table 2). Among observational studies, RoB across studies was assessed as very serious for most age groups due to nonrandomization leading to a likelihood of confounding and selection bias. Inconsistency was judged as not serious, but it was noted that interventions




FIGURE 4 Forest plot of the effect of sugar-sweetened beverage consumption in children aged <10.9 y on percentage body fat. REML, residual maximum likelihood.

and comparators were different across studies. Indirectness and imprecision were judged as not serious. Evidence was downrated by 1 further level for overweight/obesity in children <2 y because the included studies were all at serious RoB. For the single RCT, the certainty of evidence was also low. In sum, in children ≤ 10.9 y, the body of evidence indicates that SSB consumption may increase BMI, percentage body fat, or the risk of overweight/obesity (low certainty).

ASB consumption and BMI, overweight/obesity, and body composition outcomes.

Seven studies reported ASB consumption in relation to all child growth, body composition, and overweight/obesity outcomes. Four studies defined the exposure as diet sodas (63, 83, 98, 120), 2 used the term “ASB” (72, 80), and 1 referred to reduced sugar, or sugar-free fruit squashes, cordials, and diet sodas (95).

ASB consumption and BMI and overweight/obesity outcomes (narrative synthesis). Six studies examined ASB consumption and BMI or overweight/obesity (Supplemental Table 9). One was assessed as critical RoB (120). No included studies examined ASB consumption in children aged <2 y. Of the 5 studies with included results, 1 observed an inverse association between ASB intake (grams per day) and BMI z-score change ($\beta = -0.20$; $SE = 0.07$; $P = 0.01$) (serious RoB) (98). Three of 5 studies reported no association between ASB intake and BMI, 2 with moderate RoB (72, 83), 1 with serious RoB (63). One study reported no difference in odds of overweight/obesity combined but greater odds of obesity with high ASB consumption (once per day) compared with low (less than once per week or never) (aOR = 1.57; 95% CI: 1.05, 2.36; $P = 0.03$) (moderate RoB) (80).

ASB consumption and percentage body fat outcomes (narrative synthesis). Three studies examined ASB intake in relation to body fat (Supplemental Table 10). One reported a negative association (per 100 g/d) (serious RoB) ($\beta = -1.41$;

$SE = 0.70$; $P = 0.046$) (98) and 2 reported no association, both moderate RoB (72, 95).

Certainty of evidence: ASB consumption. There was no evidence on the effects of ASB consumption on children <2 y. The certainty of evidence for effects of ASB consumption in children aged 2 to <5 y was low for BMI and overweight/obesity (Table 3). In children aged 5 to ≤ 10.9 y, the certainty of evidence was very low for BMI and there was no evidence for overweight/obesity. Certainty of evidence for ASB consumption and percentage body fat was low. Therefore, the body of evidence for all age groups ≤ 10.9 y indicates that ASB consumption makes little or no difference to increased BMI, percentage body fat, or the risk of overweight/obesity (low certainty).

One hundred percent fruit juice consumption and BMI, overweight/obesity, and body composition outcomes.

Seventeen studies reported effects of fruit juice consumption. In 16 studies, the exposure was specified as 100% juice. One study examined unsweetened fruit juice and small intakes of sweetened fruit and vegetable juice (72). This study was placed with 100% fruit juice for the synthesis because this was the closest match. Two studies were judged as critical RoB (120, 124).

One hundred percent fruit juice consumption and BMI and overweight/obesity outcomes (narrative synthesis). Ten studies assessed fruit juice consumption and BMI or overweight/obesity (Supplemental Table 9). Nine of the 10 studies reported no association (5 moderate, 4 serious RoB) (58, 72, 81, 83, 87, 98, 114, 116, 130). One study reported mixed results, with fruit juice intake from 2 to 4 y associated with greater BMI z-score change from baseline to 4 y [mean change = 0.282, $SE = 0.028$, compared with 0.030, $SE = 0.037$; $P = 0.0003$ for groups ≥ 1 serving (236.5 mL)/d compared with <1 serving/d], but not from 4 to 5 y (mean change = 0.034, $SE = 0.031$, compared with 0.020, $SE = 0.021$; $P = 0.6778$) (moderate RoB) (86). In the same study,

TABLE 2 GRADE evidence profile for the effects of sugar-sweetened beverage consumption in children aged ≤ 10.9 y and BMI, body composition, and overweight/obesity outcomes¹
Question: High consumption of SSBs compared with low or no consumption for increased risk of overweight/obesity in children aged ≤ 10.9 y.
Setting: All countries, community settings.

Total studies (references)	Study design	Certainty assessment					Other considerations	Impact	Certainty	Importance
		Risk of bias	Inconsistency	Indirectness	Imprecision					
Mean BMI/BMI z-score or change in BMI/BMI z-score in children <2 y at exposure 3 (105, 112, 128)	Observational studies	Very serious ²	Not serious ³	Not serious ⁴	Not serious ⁵	None	Increased BMI (0 studies) Different effects (2 studies, $n = 3138$); different effect in boys vs. girls (105); different effects by age of follow-up: from age 18 mo to 6 y, β : 0.06; 95% CI: -0.20, 0.31; $P = 0.67$; and from age 5 y to 6 y, β : 0.34; 95% CI: 0.11, 0.58; $P = 0.004$ (58) No association (1 study, $n = 743$): mean BMIz difference -0.10; 95% CI: -0.36, 0.16 (112) ^{6,7}	⊕⊕○○ Low	Critical	
Mean BMI/BMI z-score or change in BMI/BMI z-score in children 2 to <5 y at exposure 6 (70, 72, 73, 81, 83, 93)	Observational studies	Very serious ⁸	Not serious ³	Not serious ⁴	Not serious ⁵	None	Increased BMI (2 studies, $n = 4792$): β : 0.05; 95% CI: 0.022, 0.079; $P = 0.001$ (81); β : 0.017; 95% CI: 0.007, 0.027; $P < 0.01$ (73); No association (4 studies, $n = 2163$): β : -0.01; 95% CI: -0.05, 0.04; $P = 0.852$ (70); ANCOVA $P = 0.0626$ (72); β : -0.01; SE: 0.02; $P = 0.50$ (83); $P > 0.05$ (93) ^{6,7}	⊕⊕○○ Low	Critical	
Mean BMI/BMI z-score or change in BMI/BMI z-score in children 5 to ≤ 10.9 y at exposure 10 (57, 62-64, 76, 91, 98, 99, 116, 129)	Observational studies	Very serious ⁹	Not serious ³	Not serious ⁴	Not serious ⁵	None	Increased BMI (2 studies, $n = 158$) β : 0.74; 95% CI: 0.15, 1.33 (91); β : 0.10; SE: 0.03; $P = 0.003$ (98); Different effects (1 study, $n = 2371$); positive association for sodas (β : 0.011; SE: 0.005; $P < 0.05$) but not other SSBs (β : 0.009; SE: 0.007; $P > 0.05$) (63)	⊕⊕○○ Low	Critical	

(Continued)

TABLE 2 (Continued)

Total studies (references)	Certainty assessment					Other considerations	Impact	Certainty	Importance
	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision				
Prevalence of overweight and obesity or prevalence of obesity only in children aged 2 to <5 y (assessed with: %) 5 (58, 74, 80, 96, 118)	Observational studies	Very serious ¹⁴	Not serious ³	Not serious ⁴	Not serious ⁵	None	Increased risk (1 study, $n = 473$); aOR: 1.92; 95% CI: 1.19, 3.11; $P \leq 0.01$ (118); Different effects (1 study, $n = 2986$); overweight/obesity no association, obesity only aOR: 1.65; 95% CI: 1.12, 2.44; $P = 0.01$ (58); No association (3 studies, $n = 17083$); no association (no estimate) (96); aOR: 1.3; 95% CI: 0.8, 2.1 (58); boys aOR: 1.01; 95% CI: 0.8, 1.29; girls aOR: 1.08; 95% CI: 0.87, 1.35 (74)	⊕⊕○○ Low	Critical
Prevalence of overweight and obesity or prevalence of obesity only in children aged 5 to ≤ 10.9 y (assessed with: %) 3 (75, 88, 91)	Observational studies	Very serious ⁸	Not serious ³	Not serious ⁴	Not serious ⁵	None	Increased risk (2 studies, $n = 1668$): aRR: 2.12; 95% CI: 1.05, 4.28 (91); aOR: 1.04; 95% CI: 1.01, 1.07; $P < 0.05$ (75); No association (1 study, $n = 1250$); overweight only, aOR: 1.29; 95% CI: 0.84, 1.96; $P = 0.246$; obese only, aOR: 1.57; 95% CI: 0.82, 3.03; $P = 0.177$ (88)	⊕⊕○○ Low	Critical
Prevalence of overweight and obesity or prevalence of obesity only in children aged 5 to ≤ 10.9 y (assessed with: %) 1 (59)	Randomized trial	Serious ¹⁰	Not serious ¹¹	Serious ¹²	Not serious ⁵	None	Increased risk (1 study, $n = 1987$); each 200-mL glass/d of sugar-sweetened beverage consumption increased the odds of obesity (aOR: 1.22; 95% CI: 1.04, 1.44; $P = 0.014$) but not overweight ($P = 0.83$)	⊕⊕○○ Low	Critical

(Continued)

TABLE 3 GRADE evidence profile for the effects of artificially sweetened beverage consumption in children aged ≤ 10.9 y and BMI, body composition, and overweight/obesity outcomes¹
Question: High consumption of artificially sweetened beverages compared with low or no consumption of artificially sweetened beverages for increased risk of overweight/obesity in children ≤ 10.9 y.
Setting: All countries, community settings.

Total studies (references)	Certainty assessment					Other considerations	Impact	Certainty	Importance
	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision				
Mean BMI/BMI z-score or change in BMI/BMI z-score in children <2 y at exposure 0							No included studies		
Mean BMI/BMI z-score or change in BMI/BMI z-score in children 2 to <5 y at exposure 2 (72, 83)	Observational studies	Very serious ²	Not serious ³	Not serious ⁴	Not serious ⁵	None	Increased BMI (0 studies); No association (2 studies, $n = 1443$): ANCOVA $P = 0.444$ (72); β : 0.01; SE: 0.02; $P = 0.83$ (83)	⊕⊕○○ Low	Critical
Mean BMI/BMI z-score or change in BMI/BMI z-score in children 5 to ≤ 10.9 y at exposure 2 (63, 98)	Observational studies	Extremely serious ⁶	Not serious ³	Not serious ⁴	Not serious ⁵	None	Increased BMI (0 studies); No association (1 study, $n = 2371$); β : 0.01; SE: 0.013; $P > 0.05$ (63); Decreased BMI (1 study, $n = 158$); β : -0.20; SE: 0.07; $P = 0.01$ (98)	⊕○○○ Very low	Critical
Prevalence of overweight and obesity or prevalence of obesity only in children aged <2 y (assessed with: %) 0							No included studies	—	
Prevalence of overweight and obesity or prevalence of obesity only in children aged 2 to <5 y (assessed with: %) 1 (80)	Observational studies	Very serious ²	Not serious ⁷	Not serious ⁴	Not serious ⁵	None	Increased risk (0 studies); Different effects (1 study, $n = 2986$) no association with overweight/obesity (aOR: 0.85; 95% CI: 0.63, 1.15; $P = 0.85$) but increased risk of obesity (aOR: 1.57; 95% CI: 1.05, 2.36; $P = 0.03$) (58)	⊕⊕○○ Low	Critical
Prevalence of overweight and obesity or prevalence of obesity only in children aged 5 to ≤ 10.9 y (assessed with: %) 0							No included studies	—	
Mean percentage body fat in children aged ≤ 10 y (assessed with: %) 3 (72, 95, 98)	Observational studies	Very serious ⁸	Not serious ³	Not serious ⁴	Not serious ⁵	None	Increased percentage body fat (1 study, $n = 362$) β : 0.26; 95% CI: -0.004, 0.52; $P = 0.05$ (95); No association (1 study, $n = 98$): ANCOVA $P = 0.584$ (72)	⊕⊕○○ Low	Critical

(Continued)

TABLE 3 (Continued)

Total studies (references)	Certainty assessment					Impact	Certainty	Importance
	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision			
						Negative association (1 study, $n = 158$): $\beta: -1.41$; SE: 0.70; $P = 0.046$ (98)		

¹aOR, adjusted odds ratio; GRADE, Grading of Recommendations Assessment, Development, and Evaluation.

²Risk of bias was moderate for all studies. Downgraded by 2 levels for inherent bias due to nonrandomization in observational studies leading to confounding and selection bias.

³Not downgraded for inconsistency but note that there were differences between interventions and comparators across studies.

⁴Not downgraded because study populations, exposures, and comparators were relevant to review question, although no studies were from low-income country populations.

⁵Not downgraded because no evidence of imprecision (i.e., not wide CIs, small sample size, or low number of events).

⁶Risk of bias was serious for all studies. Downgraded by 2 levels for risk of bias due to nonrandomization (confounding and selection bias) and 1 further level for serious risk of bias across the body of evidence.

⁷Not downgraded because only 1 study.

⁸Risk of bias was moderate in 2 studies (72, 95) and serious in 1 study (98). Downgraded by 2 levels due to risk of bias due to nonrandomization in observational studies leading to confounding and selection bias.

odds of overweight were not associated with juice intake in normal weight or risk-of-overweight children at baseline (moderate RoB) (86).

One hundred percent fruit juice consumption and BMI and overweight/obesity outcomes (meta-analysis). Figure 5 shows the effect estimates for the consumption of 100% fruit juice (per 250-mL serving) on BMI z-score for 3 studies (81, 116, 119). The pooled effect estimate was positive but small ($\beta=0.01$; 95% CI: 0.00, 0.01). There was no heterogeneity across individual studies ($I^2 = 0.0\%$).

One hundred percent juice consumption and percentage body fat outcomes. Four studies reported effects of 100% juice consumption on percentage body fat (Supplemental Table 10); all 4 studies reported no association (2 moderate, 2 serious RoB) (72, 95, 98, 116).

Certainty of evidence: 100% juice consumption. GRADE evidence profiles for effects of 100% fruit juice consumption are shown in Table 4. The certainty of evidence for BMI, overweight/obesity, and percentage body fat was low, with the exception of children aged 5 to ≤ 10 y, where the certainty was very low for BMI and there was no evidence for overweight/obesity (Table 4). The body of evidence for all age groups ≤ 10.9 y indicates that 100% fruit juice consumption makes little or no difference to increased BMI, percentage body fat, or the risk of overweight/obesity (low certainty).

Unhealthy food item consumption and BMI, overweight/obesity, and body composition outcomes.

Twenty-six studies reported effects of unhealthy food consumption on growth, body composition, or overweight/obesity outcomes with a range of exposures. Consumption of high-fat foods was assessed in 4 studies (5 articles) (73, 74, 103, 108, 116). Six studies (7 articles) examined the intake of free sugars or added sugar or sweetened foods (85, 99, 101, 102, 112, 121, 127). Fast food consumption was examined in 5 studies (69, 71, 76, 111, 113). Three studies reported on UPF consumption (70, 90, 131). Other exposures included salty snacks (91), sweets (125), or combinations of both (82, 118). Studies were predominantly conducted in HIC. Studies from MIC were conducted in Brazil and Peru (70, 89–91, 131). Four of the 26 studies were assessed as being at critical RoB and are not reported further (61, 121, 125, 126).

Unhealthy foods and BMI and overweight/obesity outcomes (narrative synthesis). Of the 22 included studies examining unhealthy food consumption, 16 studies reported BMI outcomes or overweight and obesity prevalence (Supplemental Table 9).

In children aged < 2 y, 4 studies examined unhealthy foods. Of these, 1 observed a positive association between sweet foods consumption from 3 to 12 mo and weight-for-length z-scores at 3 y (ANOVA, $F = 3.23$, $P = 0.03$), but




FIGURE 5 Forest plot of the effect of 100% juice consumption in children aged <10.9 y on BMI z-score values. REML, residual maximum likelihood.

no association with snack foods (moderate RoB) (82). The remaining 3 studies (1 at moderate and 2 at serious RoB) found no associations between unhealthy foods (“extra food,” fast food and snacks, sweetened first foods) and BMI or overweight/obesity (29, 111, 112). In children aged 2 to <5 y, there were 7 studies (10 articles). Two studies reported a positive association with unhealthy food consumption and outcomes. Consumption of added sugar to milk and fruits was associated with significantly higher BMI in boys and girls aged 2 to <6 y at baseline, but in older children (6 to <10 y) the association was only significant in boys (no effect estimate available) (moderate RoB) (85). Frequency of fast food intake (high or low) was associated with increased risk of change in BMI status (normal to overweight, or overweight to obese) in children aged 3–5 y followed up 1 y later (RR: 1.38; 95% CI: 1.13, 1.67; $P < 0.01$) (moderate RoB) (69).

In children aged 2 to <5 y, 3 of the 7 studies presented results that differed by quantity consumed, outcome, or time point. Consumption of high-fat food was associated with significantly higher BMI z-scores (73), but not with odds of overweight and obesity (moderate RoB) (74). In a study in Brazil, frequency of energy-dense food consumption was not associated with BMI z-scores (89), but the percentage energy intake from UPFs at age 4 y was significantly associated with BMI z-score at 7 y, whereas intake at 7 y was not (moderate RoB) (90). One study reported no effects of added sugar at age 2 y on change in BMI z-score at 5 and 6 y of age. A separate analysis from the same study found that consumption at age 1 y was not associated with change in BMI z-score at 7 y, but change in intake from 1 to 7 y was significantly associated with change in BMI z-scores (serious RoB) (101, 102). The remaining 2 of 7 studies reported no association between unhealthy food consumption and BMI or overweight and obesity (1 moderate, 1 serious RoB) (70, 118).

Five studies examined effects of unhealthy food consumption in children aged 5 to ≤ 10.9 y. One reported an association between salty, high-fat snack frequency with change in BMI from 8 y to 12 y ($\beta = 0.71$; 95% CI: 0.14,

1.28; $P < 0.05$) (moderate RoB) (91). Bel-Serrat et al. (113) found significantly lower odds of overweight/obesity with savory snack intake some days per week (aOR = 0.48; 95% CI: 0.23, 0.99; $P < 0.05$) or never (OR = 0.27; 95% CI: 0.10, 0.72; $P < 0.01$) compared with every day, but no association between fast food intake and overweight/obesity (serious RoB). Three of the 5 studies of ages 5 to ≤ 10.9 y reported no association between unhealthy food intake and BMI or overweight/obesity outcomes (1 moderate, 2 serious RoB) (76, 99, 116).

Unhealthy food consumption and percentage body fat outcomes. Four studies (5 articles) examined unhealthy food consumption in relation to body fat. Three studies measured percentage body fat and reported no association with unhealthy food consumption (all serious RoB) (99, 101, 102, 116) (Supplemental Table 10). One study examined fat mass index and reported an association between annual consumption of UPFs (in grams via 12-mo recall) in children aged 6 y at baseline and higher fat mass index at 5-y follow-up ($\beta = 0.05$; 95% CI: 0.04, 0.06; $P < 0.001$) (moderate RoB) (131).

Certainty of evidence: unhealthy foods. GRADE evidence profiles for the effects of consumption of unhealthy foods are presented in Table 5. The certainty of evidence for BMI/BMI z-score and overweight/obesity was low, with the exception of children aged <2 y, where the certainty was very low for overweight/obesity. The certainty of evidence for percentage body fat across all ages ≤ 10.9 y was very low (Table 5).

Synthesis of results of unhealthy food and beverage consumption and other growth and body composition outcomes can be found in Supplemental Table 11.

Discussion

Summary of evidence

In this review of the effects of unhealthy food and beverage consumption on risk of overweight and obesity, we found no

TABLE 4 GRADE evidence profile for the effects of 100% fruit juice consumption in children aged ≤ 10.9 y and BMI, body composition, and overweight/obesity outcomes¹**Question:** High consumption of 100% fruit juice compared with low or no consumption of 100% fruit juice for increased risk of overweight/obesity in children ≤ 10.9 y.**Setting:** All countries, community settings.

Total studies (references)	Study design	Certainty assessment					Other considerations	Impact	Certainty	Importance
		Risk of bias	Inconsistency	Indirectness	Imprecision					
Mean BMI/BMI z-score or change in BMI/BMI z-score in children <2 y at exposure 1 (87)	Observational study	Very serious ²	Not serious ³	Not serious ⁴	Not serious ⁵	None	Increased BMI (0 studies); No association (1 study, $n = 1038$) β : 0.30; 95% CI: -0.01, 0.61 at 2.1 y follow-up; β : 0.27; 95% CI: -0.05, 0.59 at 6.7 y follow-up (87) ⁶	⊕⊕○○ Low	Critical	
Mean BMI/BMI z-score or change in BMI/BMI z-score in children 2 to <5 y at exposure 5 (67, 72, 81, 83, 86)	Observational studies	Very serious ⁷	Not serious ⁸	Not serious ⁴	Not serious ⁵	None	Increased BMI (0 studies); Different effects (1 study, $n = 6250$): mean BMI z-score change = 0.282 (SE: 0.028) vs. 0.030 (SE: 0.037), $P = 0.0003$ at 2–4 y, 0.034 (SE: 0.031) 0.020 (SE: 0.021) $P = 0.6778$ at 4–5 y (86); No association (4 studies, $n = 2138$): ANCOVA $P = 0.062$ (72); β : 0.01; SE: 0.00; $P = 0.20$ (83); β : -0.001; 95% CI: -0.059, 0.057; $P = 0.97$ (81); β : -0.057; $P = 0.099$ (SE not stated) (67) ⁶	⊕⊕○○ Low	Critical	
Mean BMI/BMI z-score or change in BMI/BMI z-score in children 5 to ≤ 10.9 y at exposure 2 (98, 116)	Observational studies	Extremely serious ⁹	Not serious ⁸	Not serious ⁴	Not serious ⁵	None	Increased BMI (0 studies); No association (2 studies, $n = 412$): β : -0.04, 95% CI: -0.21, 0.13; $P = 0.631$ (116); β : 0.07; SE: 0.05; $P = 0.12$ (98) ⁶	⊕○○○ Very low	Critical	

(Continued)

TABLE 4 (Continued)

Total studies (references)	Certainty assessment					Other considerations	Impact	Certainty	Importance
	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision				
1 (87)	Observational study	Extremely serious ⁹	Not serious ³	Not serious ⁴	Not serious ⁵	None	Increased risk (0 studies); No association (1 study, $n = 1076$); odds of overweight including obesity, aOR: 1.0; 95% CI: 0.5, 2.0; $P = 0.916$ (114)	⊕○○○ Very low	Critical
2 (58, 86)	Observational studies	Very serious ²	Not serious ⁸	Not serious ⁴	Not serious ⁵	None	Increased risk (0 studies); Different effects (1 study, $n = 6250$); overweight/obesity aOR: 1.30; 95% CI: 1.06, 1.59; $P = 0.0129$ at 2–4 y follow-up; aOR: 0.80; 95% CI: 0.43, 1.49; $P = 0.473$ at 4–5 y follow-up) (86); No association (1 study, $n = 10,904$): high vs. low intake in normal weight at baseline, aOR: 1.2; 95% CI: 0.8, 1.7); high vs. low intake in at risk of overweight at baseline, aOR: 0.8; 95% CI: 0.5, 1.1 (58)	⊕⊕○○ Low	Critical
0							No included studies	—	
4 (72, 95, 98, 116)	Observational studies	Very serious ¹⁰	Not serious ⁸	Not serious ⁴	Not serious ⁵	None	Increased percentage body fat (0 studies); No association (4 studies, $n = 872$); β : -1.06 ; 95% CI: $-2.70, 0.57$; $P = 0.202$ (116); ANCOVA $P = 0.119$ (72); β : -0.11 ; 95% CI:	⊕⊕○○ Low	

(Continued)

TABLE 4 (Continued)

Total studies (references)	Certainty assessment					Impact	Certainty	Importance
	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision			

−0.61, 0.38; $P = 0.66$
(95); β : −0.05; SE: 0.44;
 $P = 0.91$ (98)

¹aOR, adjusted odds ratio; GRADE, Grading of Recommendations Assessment, Development, and Evaluation.

²Risk of bias was moderate in all studies. Downgraded by 2 levels due to nonrandomization in observational studies leading to confounding and selection bias.

³Not downgraded because only 1 study.

⁴Not downgraded because study populations, exposures, and comparators were relevant to review question, although no studies were from low-income country populations.

⁵Not downgraded because no evidence of imprecision (i.e., not wide CIs, small sample size, or low number of events).

⁶Meta-analysis of 3 studies across age groups on BMI z-score effect size: 0.01; 95% CI: 0.00, 0.01.

⁷Risk of bias was moderate in 4 studies (72, 81, 83, 86) and serious in 1 study (67). Downgraded by 2 levels due to nonrandomization in observational studies leading to confounding and selection bias.

⁸Not downgraded for inconsistency but note that interventions and comparators were not the same across studies.

⁹Risk of bias was serious in all studies. Downgraded by 2 levels for inherent risk of bias due to nonrandomization and 1 further level due to body of evidence based on studies at serious risk of bias.

¹⁰Risk of bias was moderate in 2 studies (72, 95) and serious in 2 studies (98, 116). Downgraded by 2 levels due to nonrandomization leading to bias due to confounding and selection bias.

studies in low-income countries, and a paucity of evidence in children aged 0–2 y. Previous survey data from 18 countries indicate that consumption of SSBs and sugary snacks is high in LMIC, with $\leq 75\%$ of children in Asia and 46% of children in Africa consuming such foods at ages 12–23 mo (1). Despite this, the effect of unhealthy food and beverage consumption in infancy and childhood remains poorly understood, particularly in LMIC settings where diets are rapidly changing, and where multiple forms of malnutrition coexist (5). Prospective studies are needed in LMIC on the amounts and types of foods consumed in relation to nutritional outcomes. This review also highlights a lack of robust studies purposefully designed to assess the effects of unhealthy food and beverage consumption in childhood on growth outcomes. High-quality and standardized data are needed in order to make nutritional recommendations to prevent all forms of malnutrition.

The largest body of evidence in this review was on the effects of SSB consumption. For children ≤ 10.9 y, the body of evidence indicates that SSB consumption may increase BMI, percentage body fat, and risk of overweight/obesity (low certainty). This accords with review findings for infants aged 6–23 mo (13). Meta-analyses in the present review indicated a positive association between SSB intake and percentage body fat, but no association with change in BMI and BMI z-score; however, the number of pooled studies was small. A previous systematic review reported that BMI increased by 0.07 (95% CI: 0.01, 0.12) for each additional daily serving (~ 354 mL) of SSBs in children and adolescents, but heterogeneity was high ($I^2 = 91.6\%$; $P < 0.001$) (14). Some RCTs have examined the effects of SSB consumption by comparing with a group receiving ASBs (132–134). One 18-mo RCT reported lower BMI increases in children receiving ASBs compared with SSBs, but on an intention-to-treat basis there was no significant difference in BMI z-score increase between the 2 groups (132). Such studies did not meet eligibility criteria for this review because they compared 2 items on the review list of exposures (SSBs and ASBs) with no control group.

We found that consumption of ASBs and consumption of 100% fruit juice in children ≤ 10.9 y makes little or no difference to increased BMI, percentage body fat, or risk of overweight/obesity (low certainty). For ASB consumption, no evidence was available for children < 2 y. For 100% juice consumption, the pooled estimate from meta-analysis in this review ($\beta = 0.01$; 95% CI: 0.00, 0.01) accords with a systematic review of longitudinal studies of 100% fruit juice consumption in children aged 1 to 6 y where a 1 serving increment was associated with a 0.087 (95% CI: 0.008, 0.167) unit increase in BMI z-score, considered not clinically significant (15). Importantly, our review ensured that all included evidence was for 100% juice consumption only, whereas some reviews have included evidence from juice drinks where the proportion of fruit juice varied or was unstated (13).

TABLE 5 GRADE evidence profile for the effects of consumption of unhealthy food items in children aged ≤ 10.9 y and growth, body composition, and overweight/obesity outcomes¹

Question: High consumption of unhealthy food items compared with low or no consumption of unhealthy food items for increased risk of overweight/obesity in children ≤ 10 y.

Setting: All countries, community settings.

Total studies (references)	Certainty assessment						Other considerations	Impact	Certainty	Importance
	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision					
Mean BMI/BMI z-scores or change in BMI/BMI z-scores in children aged <2 y 3 (29, 82, 112)	Observational studies	Very serious ²	Not serious ³	Not serious ⁴	Not serious ⁵	None	Increased BMI (1 study, $n = 666$): candies, ANOVA $F = 3.23$, $P = 0.03$ (82); No association (2 studies, $n = 1105$); "extra foods" β : -0.10 ; 95% CI: $-0.30, 0.11$; $P = 0.36$ (29); sweetened foods BMIz mean difference = 0.03 ; 95% CI: $-0.12, 0.19$ (112)	⊕⊕○○ Low	Critical	
Mean BMI/BMI z-scores or change in BMI/BMI z-scores in children aged 2 to <5 y at exposure 6 (69, 70, 73, 85, 90, 101)	Observational studies	Very serious ⁶	Not serious ³	Not serious ⁴	Not serious ⁵	None	Increased BMI (3 studies, $n = 11639$); fast foods, aRR: 1.38; 95% CI: 1.13, 1.67; $P < 0.01$ (69); high-fat foods, β : 0.021; 95% CI: 0.014, 0.029; $P < 0.001$ (73); sugar-added to foods 2 < 6 y: boys, $P = 0.005$; girls, $P = 0.03$; 6 < 10 y: boys, $P = 0.001$; girls, $P > 0.05$ (85); Different effects (1 study, $n = 1175$); ultraprocessed food intake at 4 y, β : 0.028; 95% CI: 0.006, 0.051; intake at 7 y β : 0.014; 95% CI: $-0.007, 0.036$ (90); No association (2 studies, $n = 695$); added sugar, β : -0.001 ; SE: 0.010; $P = 0.9$ (101); ultraprocessed foods, β : 0.05; 95% CI: $-0.04, 0.15$; $P = 0.282$ (70)	⊕⊕○○ Low	Critical	

TABLE 5 (Continued)

Total studies (references)	Study design	Certainty assessment					Other considerations	Impact	Certainty	Importance
		Risk of bias	Inconsistency	Indirectness	Imprecision					
Percentage body fat ≤ 10 y 4 (99, 101, 116, 131)	Observational studies	Extremely serious ¹²	Not serious ³	Not serious ⁴	Not serious ⁵	None	everyday, aOR: 0.27; 95% CI: 0.10, 0.72; $P < 0.01$; fast food never vs. everyday, aOR: 0.91; 95% CI: 0.19, 4.31; $P > 0.05$ (113); No association (1 study, $n = 1414$); savory snacks, aRR: 1.43; 95% CI: 0.78, 2.69 (91)	⊕○○○ Very low	Critical	
							Increased percentage body fat (1 study, $n = 3514$); ultraprocessed foods, β : 0.05; 95% CI: 0.04, 0.06; $P < 0.001$ (NOTE: fat mass index, not % body fat) (131); No association (3 studies, $n = 1239$); added sugar, β : 0.048; SE: 0.046; $P = 0.3$ (101); high-fat foods, β : -0.38 , 95% CI: -0.81 , 0.05; $P = 0.081$ (116); other sugars, β : 0.83; SE: 0.72; $P > 0.05$ (99)			

¹ aOR, adjusted odds ratio; aRR, adjusted relative risk; BMIz, body mass index z-score; GRADE, Grading of Recommendations Assessment, Development, and Evaluation.
² Risk of bias was moderate in 2 studies (29, 82), serious in 1 study (112). Downgraded by 2 levels for risk of bias due to nonrandomization in observational studies leading to confounding and selection bias.
³ Not downgraded for inconsistency but note that interventions and comparators were different across studies.
⁴ Not downgraded because study populations, exposures, and comparators were relevant to review question, although no studies were from low-income country populations.
⁵ Not downgraded because no evidence of imprecision (i.e., not wide CIs, small sample size, or low number of events).
⁶ Risk of bias was moderate in 5 studies (69, 70, 73, 85, 90) and serious in 1 study (101). Downgraded by 2 levels for risk of bias due to nonrandomization in observational studies leading to confounding and selection bias.
⁷ Risk of bias was moderate in 2 studies (76, 91) and serious in 2 studies (99, 116). Downgraded by 2 levels for risk of bias due to nonrandomization in observational studies leading to confounding and selection bias.
⁸ Risk of bias was serious in all studies (111). Downgraded by 2 levels for nonrandomization in observational studies leading to confounding and selection bias, and 1 level further due to body of evidence all from studies with serious risk of bias.
⁹ Risk of bias was moderate in 1 study and serious in 1 study. Downgraded by 2 levels for inherent risk of bias due to nonrandomization in observational studies.
¹⁰ Risk of bias was moderate in 1 study (91) and serious in 1 study (113). Downgraded by 2 levels for risk of bias due to nonrandomization leading to confounding and selection bias.
¹¹ Not downgraded because only 1 study.
¹² Risk of bias was moderate for 1 study (131) and serious for 3 studies (99, 101, 116). Downgraded by 2 levels for risk of bias due to nonrandomization leading to confounding and selection bias and 1 level further due to majority of the body of evidence had serious risk of bias.

Unhealthy food consumption

Studies reporting unhealthy food consumption assessed salty, high-fat food consumption (73, 74, 91, 113), UPFs (90, 131), fast food or “extra foods” (29, 69, 111), and added sugars or foods high in sugars (85, 101, 112). We found consumption of unhealthy foods in children ≤ 10.9 y may increase BMI, percentage body fat, or risk of overweight/obesity (low to very low certainty). However, no meta-analysis was possible due to the high heterogeneity of reporting exposures and comparators. A systematic review of complementary feeding (6–23 mo) found insufficient evidence of effects of consumption of unhealthy foods (13), concurring with the findings of the present study for all children ≤ 10.9 y. A previous systematic review of UPF consumption and body fat in children and adolescents included both longitudinal and cross-sectional study designs and therefore reverse causality was a possible underlying factor in observed associations (16).

Limitations of the evidence

A major limitation of the evidence was that all included studies, except 1, were observational cohorts. Moreover, very few studies were designed purposively to examine the effect of unhealthy food and beverage consumption on indicators of overweight/obesity. Although RCTs could provide greater certainty of evidence, purposeful consumption of unhealthy foods and beverages is precluded for ethical reasons. The results from meta-analyses were limited by the small number of studies that could be harmonized. Further, the pooled studies included different baseline ages and varying duration of follow-up across studies, hence pooled estimates should be interpreted with caution. The high heterogeneity of reporting of dietary intakes [i.e., differences in dietary assessment methods, recall periods, units of measurements, and definition of the exposure (typology of food item/food group)] prevented further data harmonization and limited the extent of meta-analyses.

Strengths and limitations of the review

Strengths of the review are the inclusion of studies dating from 1993, with no restrictions on language or country. Other systematic reviews of unhealthy food consumption in infants and young children have been confined to countries classified as high on the Human Development Index and English language only (13, 135). The inclusion of infants and young children in this review also added valuable insights because existing reviews have predominantly examined later childhood and adolescence (e.g., reference 14). We followed Cochrane recommended methods for RoB and grading of evidence (25). We used a comprehensive food-based and nutrient-based approach in addition to the NOVA classification, to consider all types of unhealthy foods and beverages. We searched 3 databases and did not search gray literature, which could be a potential limitation of the review.

Recommendations for future research

Studying the effect of unhealthy food consumption on the risk of overweight/obesity is challenging due to the high heterogeneity in measuring and reporting dietary intakes. More robust nutritional epidemiological intervention or prospective studies are needed to enhance our understanding of the relation between unhealthy food and beverage consumption and overweight/obesity. Evidence could be strengthened by collecting the highest quality dietary data possible and by standardizing data collection and reporting measures of diet in studies investigating the relation between unhealthy food consumption and health. A clear definition and conceptualization of the dietary risk factors for overweight/obesity and nutrition-related NCDs is key to ensuring standardization and hence comparison of exposure measures across studies. Dietary assessment approaches are now recognizing the importance of capturing information on unhealthy food and beverage intake, as reflected by the updated infant and young child feeding indicators (58), which include sentinel unhealthy foods and SSBs. Furthermore, the recently published diet quality questionnaire (internationally standardized survey instrument) (136) provides a list of food groups to limit or avoid (i.e., baked sweets; other sweets; sodas, energy drinks, sports drinks; fruit juice and fruit-flavored drinks; sweet tea, coffee, cocoa; packaged ultraprocessed salty snacks; instant noodles; deep-fried foods; fast foods). These food group classifications could be applied in nutritional epidemiological studies. Diet quality questionnaires aligned with the WHO and UNICEF indicators for infants and young children are soon to be released. Wider adoption of the STROBE-nut reporting guidelines in future studies would help enhance evidence syntheses (137). In addition, future work should focus on children ≤ 2 y in LMIC settings where diets are rapidly changing, and multiple forms of malnutrition coexist.

Conclusion

In children ≤ 10.9 y, consumption of SSBs and unhealthy foods may increase BMI, percentage body fat, or odds of overweight/obesity (low to very low certainty). Consumption of ASBs and 100% fruit juice makes little or no difference to BMI, percentage body fat, or overweight/obesity outcomes (low certainty). High-quality nutritional epidemiological studies that are designed to assess the effects of unhealthy food consumption during childhood on risk of overweight/obesity are needed to contribute to a more robust evidence base upon which to develop policy recommendations. This is key to address the growing burden of overweight and obesity that children are experiencing worldwide. Evidence from low-income countries is also needed.

Acknowledgments

We gratefully acknowledge academic librarian, N Rush (Loughborough University), for his support with database searches and article retrieval, N Pearson (Loughborough University) for her assistance with searching and screening, K Burdenski, M Stanley, and Y Todorova (Loughborough

University) for their assistance with screening and article retrieval, and K Baye (Addis Ababa University) for his role as advisor on the review. We gratefully acknowledge feedback from the WHO Technical Advisors and the WHO Guideline Development Group on the review protocol and the draft report of the review.

The authors' responsibilities were as follows—RP, SG, PG, CC, and EKR: protocol development and searches; OM, RP, SG, PG, and EKR: screening; OM, RP, SG, BB, and EKR: data extraction, risk of bias assessment, and synthesis; ESP: data harmonization and meta-analysis; SG and EKR: grading the evidence; EKR and RP: wrote the first draft of the manuscript; SG, OM, BB, PG, CC, and ESP: provided critical review of important intellectual content; EKR: had primary responsibility for the final content; and all authors: reviewed and commented on versions of the manuscript and read and approved the final manuscript.

References

- Huffman SL, Piwoz EG, Vosti SA, Dewey KG. Babies, soft drinks and snacks: a concern in low- and middle-income countries? *Matern Child Nutr* 2014;10(4):562–74.
- Pries AM, Rehman AM, Filteau S, Sharma N, Upadhyay A, Ferguson EL. Unhealthy snack food and beverage consumption is associated with lower dietary adequacy and length-for-age z-scores among 12–23-month-olds in Kathmandu Valley. *J Nutr* 2019;149(10):1843–51.
- Monteiro CA, Cannon G, Levy R, Moubarac J-C, Jaime P, Martins AP, et al. NOVA. The star shines bright. *World Nutr* 2016;7(1–3): 28–38.
- Monteiro CA, Moubarac JC, Cannon G, Ng SW, Popkin B. Ultra-processed products are becoming dominant in the global food system. *Obes Rev* 2013;14(S2):21–8.
- Development Initiatives. 2018 Global nutrition report: shining a light to spur action on nutrition. Bristol (UK): Development Initiatives; 2018.
- Lutter CK, Grummer-Strawn L, Rogers L. Complementary feeding of infants and young children 6 to 23 months of age. *Nutr Rev* 2021;79(8):825–46.
- Pries AM, Filteau S, Ferguson EL. Snack food and beverage consumption and young child nutrition in low- and middle-income countries: a systematic review. *Matern Child Nutr* 2019;15(S4):e12729.
- Afshin A, Sur PJ, Fay KA, Cornaby L, Ferrara G, Salama JS, et al. Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2019;393(10184):1958–72.
- Pan American Health Organization/World Health Organization. Guiding principles for complementary feeding of the breastfed child. Washington (DC): Pan American Health Organization; 2003.
- World Health Organization. Guiding principles for feeding non-breastfed children 6–24 months of age. Geneva (Switzerland): World Health Organization; 2005.
- Development Initiatives. 2020 global nutrition report: action on equity to end malnutrition. Bristol (UK): Development Initiatives; 2020.
- FAO. The state of food security and nutrition in the world 2020. FAO, IFAD, UNICEF, WFP, WHO; 2020.
- English LK, Obbagy JE, Wong YP, Butte NF, Dewey KG, Fox MK, et al. Types and amounts of complementary foods and beverages consumed and growth, size, and body composition: a systematic review. *Am J Clin Nutr* 2019;109(Suppl 1):956S–77S.
- Malik VS, Pan A, Willett WC, Hu FB. Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis. *Am J Clin Nutr* 2013;98(4):1084–102.
- Auerbach BJ, Wolf FM, Hikida A, Vallila-Buchman P, Littman A, Thompson D, et al. Fruit juice and change in BMI: a meta-analysis. *Pediatrics* 2017;139(4):e20162454.
- Costa CS, Del-Ponte B, Assunção MCF, Santos IS. Consumption of ultra-processed foods and body fat during childhood and adolescence: a systematic review. *Public Health Nutr* 2018;21(1):148–59.
- Bucher Della Torre S, Keller A, Laure Depeyre J, Kruseman M. Sugar-sweetened beverages and obesity risk in children and adolescents: a systematic analysis on how methodological quality may influence conclusions. *J Acad Nutr Diet* 2016;116(4):638–59.
- Mozaffarian D. Dietary and policy priorities for cardiovascular disease, diabetes, and obesity. *Circulation* 2016;133(2):187–225.
- Grant MJ, Booth A. A typology of reviews: an analysis of 14 review types and associated methodologies. *Health Info Libr J* 2009;26(2):91–108.
- Page MJ, Moher D, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ* 2021;372:n160.
- WHO, UNICEF. Indicators for assessing infant and young child feeding practices: definitions and measurement methods. Geneva (Switzerland): WHO, UNICEF; 2021.
- Monteiro CA, Levy RB, Claro RM, de Castro IRR, Cannon G. A new classification of foods based on the extent and purpose of their processing. *Cad Saude Publica* 2010;26(11):2039–49.
- Rompay MIV, McKeown NM, Goodman E, Eliasziw M, Chomitz VR, Gordon CM, et al. Sugar-sweetened beverage intake is positively associated with baseline triglyceride concentrations, and changes in intake are inversely associated with changes in HDL cholesterol over 12 months in a multi-ethnic sample of children. *J Nutr* 2015;145(10):2389–95.
- Sterne J, Hernán M, Reeves B, Savović J, Berkman N, Viswanathan M, et al. Risk of bias in non-randomized studies of interventions (ROBINS-I): detailed guidance. [Internet]. *BMJ* 2016;355:i4919. [Accessed 2021 Dec 20]. Available from: <http://www.riskofbias.info>.
- Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. *Cochrane handbook for systematic reviews of interventions*. 2nd ed. Hoboken (NJ): Wiley-Blackwell; 2019.
- Sterne JA, Hernán MA, Reeves BC, Savović J, Berkman ND, Viswanathan M, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016;355:i4919.
- Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;366:i4898.
- Okronipa H, Arimond M, Arnold CD, Young RR, Adu-Afarwah S, Tamakloe SM, et al. Exposure to a slightly sweet lipid-based nutrient supplement during early life does not increase the level of sweet taste most preferred among 4-to 6-year-old Ghanaian children: follow-up of a randomized controlled trial. *Am J Clin Nutr* 2019;109(4):1224–32.
- Garden FL, Marks GB, Almqvist C, Simpson JM, Webb KL. Infant and early childhood dietary predictors of overweight at age 8 years in the CAPS population. *Eur J Clin Nutr* 2011;65(4):454–62.
- Feldens CA, Giugliani E, Vigo Á, Vítolo MR. Early feeding practices and severe early childhood caries in four-year-old children from southern Brazil: a birth cohort study. *Caries Res* 2010;44(5):445–52.
- Chaffee BW, Feldens CA, Rodrigues PH, Vítolo MR. Feeding practices in infancy associated with caries incidence in early childhood. *Community Dent Oral Epidemiol* 2015;43(4):338–48.
- Ruottinen S, Karjalainen S, Pienihäkkinen K, Lagström H, Niinikoski H, Salminen M, et al. Sucrose intake since infancy and dental health in 10-year-old children. *Caries Res* 2004;38(2):142–8.
- Marshall TA, Levy SM, Broffitt B, Warren JJ, Eichenberger-Gilmore JM, Burns TL, et al. Dental caries and beverage consumption in young children. *Pediatrics* 2003;112(3 Pt 1):e184–91.
- McGuinness LA, Higgins JPT. Risk-of-bias VISualization (robvis): an R package and Shiny web app for visualizing risk-of-bias assessments. *Res Synth Methods* 2021;12(1):55–61.

35. Campbell M, McKenzie JE, Sowden A, Katikireddi SV, Brennan SE, Ellis S, et al. Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline. *BMJ* 2020;368:l6890.
36. Schünemann H, Brozek J, Guyatt G, Oxman A. Handbook for grading the quality of evidence and the strength of recommendations using the GRADE approach. [Internet]. Updated October 2013. [Cited 2021 Jul 20]. Available from: <https://gdt.gradepro.org/app/handbook/handbook.html>.
37. Cochrane Effective Practice and Organisation of CARE (EPOC). Reporting the effects of an intervention in EPOC reviews [Internet]. 2018. [Cited 2022 Apr 6]. Available from: https://epoc.cochrane.org/sites/epoc.cochrane.org/files/public/uploads/Resources-for-authors2017/how_to_report_the_effects_of_an_intervention.pdf.
38. Ishihara T, Takeda Y, Mizutani T, Okamoto M, Koga M, Tamura U, et al. Relationships between infant lifestyle and adolescent obesity. The Enzan maternal-and-child health longitudinal study. *Nihon Koshu Eisei Zasshi* 2003;50(2):106–17.
39. Zhang Y, Cheng R, Cheng M, Li Y. The prevalence of dental caries in primary dentition and the risk factors of 5-year-old children in Northeast of China. *Shanghai Kou Qiang Yi Xue* 2007;16(6):570–3.
40. Berkey CS, Rockett HRH, Field AE, Gillman MW, Colditz GA. Sugar-added beverages and adolescent weight change. *Obes Res* 2004;12(5):778–88.
41. Bisset S, Gauvin L, Potvin L, Paradis G. Association of body mass index and dietary restraint with changes in eating behaviour throughout late childhood and early adolescence: a 5-year study. *Public Health Nutr* 2007;10(8):780–9.
42. Seferidi P, Millett C, Lavery AA. Sweetened beverage intake in association to energy and sugar consumption and cardiometabolic markers in children. *Pediatr Obes* 2018;13(4):195–203.
43. Shroff MR, Perng W, Baylin A, Mora-Plazas M, Marin C, Villamor E. Adherence to a snacking dietary pattern and soda intake are related to the development of adiposity: a prospective study in school-age children. *Public Health Nutr* 2014;17(7):1507–13.
44. Xue H, Wu Y, Wang X, Wang Y. Time trends in fast food consumption and its association with obesity among children in China. *PLoS One* 2016;11(3):1–14.
45. Field AE, Austin SB, Gillman MW, Rosner B, Rockett HR, Colditz GA. Snack food intake does not predict weight change among children and adolescents. *Int J Obes* 2004;28(10):1210–6.
46. Jensen BW, Nichols M, Allender S, De Silva-Sanigorski A, Millar L, Kremer P, et al. Inconsistent associations between sweet drink intake and 2-year change in BMI among Victorian children and adolescents. *Pediatr Obes* 2013;8(4):271–83.
47. Johnson BA, Kremer PJ, Swinburn BA, De Silva-Sanigorski AM. Multilevel analysis of the Be Active Eat Well intervention: environmental and behavioural influences on reductions in child obesity risk. *Int J Obes* 2012;36(7):901–7.
48. Lee EY, Kang B, Yang Y, Yang HK, Kim HS, Lim SY, et al. Study time after school and habitual eating are associated with risk for obesity among overweight Korean children: a prospective study. *Obes Facts* 2018;11(1):46–55.
49. Mrdjenovic G, Levitsky DA. Nutritional and energetic consequences of sweetened drink consumption in 6- to 13-year-old children. *J Pediatr* 2003;142(6):604–10.
50. Mundt CA, Baxter-Jones A, Whiting SJ, Bailey DA, Faulkner RA, Mirwald RL. Relationships of activity and sugar drink intake on fat mass development in youths. *Med Sci Sports Exerc* 2006;38(7):1245–54.
51. Nissinen K, Mikkilä V, Männistö S, Lahti-Koski M, Räsänen L, Viikari J, et al. Sweets and sugar-sweetened soft drink intake in childhood in relation to adult BMI and overweight. The Cardiovascular Risk in Young Finns Study. *Public Health Nutr* 2009;12(11):2018–26.
52. Phillips SM, Bandini LG, Naumova EN, Cyr H, Colclough S, Dietz WH, et al. Energy-dense snack food intake in adolescence: longitudinal relationship to weight and fatness. *Obes Res* 2004;12(3):461–72.
53. Alexy U, Libuda L, Mersmann S, Kersting M. Convenience foods in children's diet and association with dietary quality and body weight status. *Eur J Clin Nutr* 2011;65(2):160–6.
54. Dong D, Bilger M, van Dam RM, Finkelstein EA. Consumption of specific foods and beverages and excess weight gain among children and adolescents. *Health Aff* 2015;34(11):1940–8.
55. Libuda L, Alexy U, Sichert-Hellert W, Stehle P, Karaolis-Danckert N, Buyken AE, et al. Pattern of beverage consumption and long-term association with body-weight status in German adolescents—results from the DONALD study. *Br J Nutr* 2008;99(6):1370–9.
56. World Bank. World Bank country and lending groups. [Internet]. 2021. [Cited 2021 May 29]. Available from: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>.
57. Jensen BW, Nielsen BM, Husby I, Bugge A, El-Naaman B, Andersen LB, et al. Association between sweet drink intake and adiposity in Danish children participating in a long-term intervention study. *Pediatr Obes* 2013;8(4):259–70.
58. Welsh JA, Cogswell ME, Rogers S, Rockett H, Mei Z, Grummer-Strawn LM. Overweight among low-income preschool children associated with the consumption of sweet drinks: Missouri, 1999–2002. *Pediatrics* 2005;115(2):e223–9.
59. Muckelbauer R, Gortmaker SL, Libuda L, Kersting M, Clausen K, Adelberger B, et al. Changes in water and sugar-containing beverage consumption and body weight outcomes in children. *Br J Nutr* 2016;115(11):2057–66.
60. Skinner JD, Carruth BR. A longitudinal study of children's juice intake and growth: the juice controversy revisited. *J Am Diet Assoc* 2001;101(4):432–7.
61. Huus K, Brekke HK, Ludvigsson JFF, Ludvigsson JFF. Relationship of food frequencies as reported by parents to overweight and obesity at 5 years. *Acta Paediatr* 2009;98(1):139–43.
62. Fiorito LM, Marini M, Francis LA, Smiciklas-Wright H, Birch LL. Beverage intake of girls at age 5 y predicts adiposity and weight status in childhood and adolescence. *Am J Clin Nutr* 2009;90(4):935–42.
63. Striegel-Moore RH, Thompson D, Affenito SG, Franko DL, Obarzanek E, Barton BA, et al. Correlates of beverage intake in adolescent girls: the National Heart, Lung, and Blood Institute Growth and Health Study. *J Pediatr* 2006;148(2):183–7.
64. Zheng M, Rangan A, Olsen NJ, Bo Andersen L, Wedderkopp N, Kristensen P, et al. Sugar-sweetened beverages consumption in relation to changes in body fatness over 6 and 12 years among 9-year-old children: the European Youth Heart Study. *Eur J Clin Nutr* 2014;68(1):77–83.
65. Garden FL, Marks GB, Simpson JM, Webb KL. Body mass index (BMI) trajectories from birth to 11.5 years: relation to early life food intake. *Nutrients* 2012;4(10):1382–98.
66. Marshall TA, Curtis AM, Cavanaugh JE, Warren JJ, Levy SM. Higher longitudinal milk intakes are associated with increased height in a birth cohort followed for 17 years. *J Nutr* 2018;148(7):1144–9.
67. Skinner JD, Carruth BR, Moran J, Houck K, Coletta F. Fruit juice intake is not related to children's growth. *Pediatrics* 1999;103(1):58–64.
68. Wheaton N, Millar L, Allender S, Nichols M. The stability of weight status through the early to middle childhood years in Australia: a longitudinal study. *BMJ Open* 2015;5(4):e006963.
69. Emond JA, Longacre MR, Titus LJ, Hendricks K, Drake KM, Carroll JE, et al. Fast food intake and excess weight gain over a 1-year period among preschool-age children. *Pediatr Obes* 2020;15(4):1–9.
70. Costa CS, Rauber F, Leffa PS, Sangalli CN, Campagnolo P, Vitolo MR. Ultra-processed food consumption and its effects on anthropometric and glucose profile: a longitudinal study during childhood. *Nutr Metab Cardiovasc Dis* 2019;29(2):177–84.
71. Guerrero AD, Mao C, Fuller B, Bridges M, Franke T, Kuo AA. Racial and ethnic disparities in early childhood obesity: growth trajectories in body mass index. *J Racial Ethn Health Disparities* 2016;3(1):129–37.

72. Hasnain SR, Singer MR, Bradlee ML, Moore LL. Beverage intake in early childhood and change in body fat from preschool to adolescence. *Child Obes* 2014;10(1):42–9.
73. Millar L, Rowland B, Nichols M, Swinburn B, Bennett C, Skouteris H, et al. Relationship between raised BMI and sugar sweetened beverage and high fat food consumption among children. *Obesity* 2014;22(5):E96–E103.
74. Zulfiqar T, Strazdins L, Dinh H, Banwell C, D'Este C, D'Este C, et al. Drivers of overweight/obesity in 4–11 year old children of Australians and immigrants; evidence from growing up in Australia. *J Immigr Minor Health* 2019;21(4):737–50.
75. Lim S, Zoellner JM, Lee JM, Burt BA, Sandretto AM, Sohn W, et al. Obesity and sugar-sweetened beverages in African-American preschool children: a longitudinal study. *Obesity* 2009;17(6):1262–8.
76. Jackson SL, Cunningham SA. The stability of children's weight status over time, and the role of television, physical activity, and diet. *Prev Med* 2017;100:229–34.
77. Kramer MS, Guo T, Platt RW, Vanilovich I, Sevkovskaya Z, Dzikovich I, et al. Feeding effects on growth during infancy. *J Pediatr* 2004;145(5):600–5.
78. Laurson K, Eisenmann JC, Moore S. Lack of association between television viewing, soft drinks, physical activity and body mass index in children. *Acta Paediatr* 2008;97(6):795–800.
79. Leermakers ETM, Felix JF, Jaddoe VWV, Raat H, Franco OH, Kiefte-de Jong JC. Sugar-containing beverage intake at the age of 1 year and cardiometabolic health at the age of 6 years: the Generation Study. *Int J Behav Nutr Phys Act* 2015;12(1):114.
80. Macintyre AK, Marrayat L, Chambers S. Exposure to liquid sweetness in early childhood: artificially-sweetened and sugar-sweetened beverage consumption at 4–5 years and risk of overweight and obesity at 7–8 years. *Pediatr Obes* 2018;13(12):755–65.
81. Marshall TA, Curtis AM, Cavanaugh JE, Warren JJ, Levy SM. Sweetened beverage intakes are longitudinally associated with higher body mass index z scores in a birth cohort followed 17 years. *J Acad Nutr Diet* 2019;119(3):425–34.
82. Moore AM, Vadiveloo M, Tovar A, McCurdy K, Østbye T, Benjamin-Neelon SE. Associations of less healthy snack food consumption with infant weight-for-length z-score trajectories: findings from the nurture cohort study. *Nutrients* 2019;11(11):2752.
83. Newby PK, Peterson KE, Berkey CS, Leppert J, Willett WC, Colditz GA. Beverage consumption is not associated with changes in weight and body mass index among low-income preschool children in North Dakota. *J Am Diet Assoc* 2004;104(7):1086–94.
84. Olafsdottir S, Berg C, Eiben G, Lanfer A, Reisch L, Ahrens W, et al. Young children's screen activities, sweet drink consumption and anthropometry: results from a prospective European study. *Eur J Clin Nutr* 2014;68(2):223–8.
85. Russo M D, Ahrens W, De Henauw S, Eiben G, Hebestreit A, Kourides Y, et al. The impact of adding sugars to milk and fruit on adiposity and diet quality in children: a cross-sectional and longitudinal analysis of the Identification and Prevention of Dietary-and Lifestyle-Induced Health Effects in Children and Infants (IDEFICS) study. *Nutrients* 2018;10(10):1350.
86. Shefferly A, Scharf RJ, Deboer MD. Longitudinal evaluation of 100% fruit juice consumption on BMI status in 2–5-year-old children. *Pediatr Obes* 2016;11(3):221–7.
87. Sonneville KR, Long MW, Rifas-Shiman SL, Kleinman K, Gillman MW, Taveras EM. Juice and water intake in infancy and later beverage intake and adiposity: could juice be a gateway drink? *Obesity* 2015;23(1):170–6.
88. Traub M, Lauer R, Kesztytis T, Wartha O, Steinacker JM, Kesztytis D, et al. Skipping breakfast, overconsumption of soft drinks and screen media: longitudinal analysis of the combined influence on weight development in primary schoolchildren. *BMC Public Health* 2018;18(1):1–10.
89. Durão C, Severo M, Oliveira A, Moreira P, Guerra A, Barros H, et al. Evaluating the effect of energy-dense foods consumption on preschool children's body mass index: a prospective analysis from 2 to 4 years of age. *Eur J Nutr* 2015;54(5):835–43.
90. Vedovato GM, Vilela S, Severo M, Rodrigues S, Lopes C, Oliveira A. Ultra-processed food consumption, appetitive traits and BMI in children: a prospective study. *Br J Nutr* 2021;125:1427–36.
91. Alviso-Orellana C, Estrada-Tejada D, Carrillo-Larco RM, Bernabé-Ortiz A. Sweetened beverages, snacks and overweight: findings from the Young Lives cohort study in Peru. *Public Health Nutr* 2018;21(9):1627–33.
92. Arcan C, Hannan PJ, Fulkerson JA, Himes JH, Rock BH, Smyth M, et al. Associations of home food availability, dietary intake, screen time and physical activity with BMI in young American-Indian children. *Public Health Nutr* 2013;16(1):146–55.
93. Byrne R, Zhou Y, Perry R, Mauch C, Magarey A. Beverage intake of Australian children and relationship with intake of fruit, vegetables, milk and body weight at 2, 3.7 and 5 years of age. *Nutr Diet* 2018;75(2):159–66.
94. Costa D, Warkentin S, Oliveira A. The effect of sugar-sweetened beverages at 4 years of age on appetitive behaviours of 7-year-olds from the Generation XXI birth cohort. *Br J Nutr* 2021;126:790–800.
95. Johnson L, Mander AP, Jones LR, Emmett PM, Jebb SA. Is sugar-sweetened beverage consumption associated with increased fatness in children? *Nutrition* 2007;23(7–8):557–63.
96. Dubois L, Farmer A, Girard M, Peterson K. Regular sugar-sweetened beverage consumption between meals increases risk of overweight among preschool-aged children. *J Am Diet Assoc* 2007;107(6):924–34.
97. Flores G, Lin H. Factors predicting severe childhood obesity in kindergarteners. *Int J Obes* 2013;37(1):31–9.
98. Zheng M, Allman-Farinelli M, Heitmann BL, Toelle B, Marks G, Cowell C, et al. Liquid versus solid energy intake in relation to body composition among Australian children. *J Hum Nutr Diet* 2015;28(s2):70–9.
99. Hur YI, Park H, Kang JH, Lee H, Song HJ, Lee H, et al. Associations between sugar intake from different food sources and adiposity or cardio-metabolic risk in childhood and adolescence: the Korean child-adolescent cohort study. *Nutrients* 2015;8(1):20.
100. Hwang IT, Ju Y-S, Lee HJ, Shim YS, Jeong HR, Kang MJ. Body mass index trajectories and adiposity rebound during the first 6 years in Korean children: based on the National Health Information Database, 2008–2015. *PLoS One* 2020;15(10):e0232810.
101. Buyken AE, Cheng G, Günther A, Liese AD, Remer T, Karaolis-Danckert N. Relation of dietary glycemic index, glycemic load, added sugar intake, or fiber intake to the development of body composition between ages 2 and 7 y. *Am J Clin Nutr* 2008;88(3):755–62.
102. Herbst A, Diethelm K, Cheng G, Icks UA, Buyken AE. Direction of associations between added sugar intake in early childhood and body mass index at age 7 years may depend on intake levels. *J Nutr* 2011;141(7):1348–54.
103. Newby PK, Peterson KE, Berkey CS, Leppert J, Willett WC, Colditz GA. Dietary composition and weight change among low-income preschool children. *Arch Pediatr Adolesc Med* 2003;157(8):759–64.
104. Pan L, Li R, Park S, Galuska DA, Sherry B, Freedman DS. A longitudinal analysis of sugar-sweetened beverage intake in infancy and obesity at 6 years. *Pediatrics* 2014;134(Suppl 1):S29–35.
105. Quah PL, Kleijweg J, Chang YY, Toh JY, Lim HX, Sugianto R, et al. Association of sugar-sweetened beverage intake at 18 months and 5 years of age with adiposity outcomes at 6 years of age: the Singapore GUSTO mother-offspring cohort. *Br J Nutr* 2019;122(11):1303–12.
106. Ritchie LD, Spector P, Stevens MJ, Schmidt MM, Schreiber GB, Striegel-Moore RH, et al. Dietary patterns in adolescence are related to adiposity in young adulthood in black and white females. *J Nutr* 2007;137(2):399–406.
107. Bayer O, Nehring I, Bolte G, Von Kries R. Fruit and vegetable consumption and BMI change in primary school-age children: a cohort study. *Eur J Clin Nutr* 2014;68(2):265–70.
108. Thurber KA, Dobbins T, Neeman T, Banwell C, Banks E. Body mass index trajectories of Indigenous Australian children and relation

- to screen time, diet, and demographic factors. *Obesity* 2017;25(4): 747–56.
109. Wan L, Jakkilinki PD, Singer MR, Bradlee ML, Moore LL. A longitudinal study of fruit juice consumption during preschool years and subsequent diet quality and BMI. *BMC Nutr* 2020;6:25.
 110. Wang N, Huang J, Li K, Zhao Y, Wen J, Ye Y, et al. Prevalence and risk factors of overweight and obesity among infants in Chongqing urban area. *Zhongguo Dang Dai Er Ke Za Zhi [Chin J Contemp Pediatr]* 2013;15(3):207–11. In Chinese.
 111. Wijga AH, Scholtens S, Bemelmans W, Kerkhof M, Koppelman GH, Brunekreef B, et al. Diet, screen time, physical activity, and childhood overweight in the general population and in high risk subgroups: prospective analyses in the PIAMA birth cohort. *J Obes* 2010;2010:423496.
 112. Santorelli G, Fairley L, Petherick ES, Cabieses B, Sahota P. Ethnic differences in infant feeding practices and their relationship with BMI at 3 years of age—results from the Born in Bradford birth cohort study. *Br J Nutr* 2014;111(10):1891–7.
 113. Bel-Serrat S, Heinen MM, Mehegan J, O'Brien S, Eldin N, Murrin CM, et al. Predictors of weight status in school-aged children: a prospective cohort study. *Eur J Clin Nutr* 2019;73(9):1299–306.
 114. Budree S, Goddard E, Brittain K, Cader S, Myer L, Zar HJ. Infant feeding practices in a South African birth cohort—a longitudinal study. *Matern Child Nutr* 2017;13(3):1–9.
 115. Cantoral A, Téllez-Rojo MM, Ettinger AS, Hu H, Hernández-Ávila M, Peterson K. Early introduction and cumulative consumption of sugar-sweetened beverages during the pre-school period and risk of obesity at 8–14 years of age. *Pediatr Obes* 2016;11(1):68–74.
 116. Carlson JA, Crespo NC, Sallis JF, Patterson RE, Elder JP. Dietary-related and physical activity-related predictors of obesity in children: a 2-year prospective study. *Child Obes* 2012;8(2):110–5.
 117. De Boer MD, Scharf RJ, Demmer RT. Sugar-sweetened beverages and weight gain in 2- to 5-year-old children. *Pediatrics* 2013;132(3):413–20.
 118. De Coen V, De Bourdeaudhuij I, Verbestel V, Maes L, Vereecken C. Risk factors for childhood overweight: a 30-month longitudinal study of 3- to 6-year-old children. *Public Health Nutr* 2014;17(9):1993–2000.
 119. Faith MS, Dennison BA, Edmunds LS, Stratton HH. Fruit juice intake predicts increased adiposity gain in children from low-income families: weight status-by-environment interaction. *Pediatrics* 2006;118(5):2066–75.
 120. Blum JW, Jacobsen DJ, Donnelly JE. Beverage consumption patterns in elementary school aged children across a two-year period. *J Am Coll Nutr* 2005;24(2):93–8.
 121. Olsen NJ, Andersen LB, Wedderkopp N, Kristensen PL, Heitmann BL. Intake of liquid and solid sucrose in relation to changes in body fatness over 6 years among 8- to 10-year-old children: the European Youth Heart Study. *Obes Facts* 2012;5(4):506–12.
 122. Tam CS, Garnett SP, Cowell CT, Campbell K, Cabrera G, Baur LA. Soft drink consumption and excess weight gain in Australian school students: results from the Nepean study. *Int J Obes* 2006;30(7): 1091–3.
 123. Weijs P, Kool LM, Van Baar NM, Van Der Zee SC. High beverage sugar as well as high animal protein intake at infancy may increase overweight risk at 8 years: a prospective longitudinal pilot study. *Nutr J* 2011;10(1):1–8.
 124. Alexy U, Sichert-Hellert W, Kersting M, Manz F, Schöch G. Fruit juice consumption and the prevalence of obesity and short stature in German preschool children: results of the DONALD study. *J Pediatr Gastroenterol Nutr* 1999;29(3):343.
 125. Lissau I, Breum L, Sorensen TIA. Maternal attitude to sweet eating habits and risk of overweight in offspring: a ten-year prospective population study. *Int J Obes* 1993;17(3):125–9.
 126. Sugimori H, Yoshida K, Izuno T, Miyakawa M, Suka M, Sekine M, et al. Analysis of factors that influence body mass index from ages 3 to 6 years: a study based on the Toyama cohort study. *Pediatr Int* 2004;46(3):302–10.
 127. Jardí C, Aranda N, Bedmar C, Ribot B, Elias I, Aparicio E, et al. Consumption of free sugars and excess weight in infants. A longitudinal study. *An Pediatr(Engl Ed)* 2019;90(3):165–72.
 128. Leermakers ETM, Felix JF, Erler NS, Čerimagić A, Wijtzes AI, Hofman A, et al. Sugar-containing beverage intake in toddlers and body composition up to age 6 years: the Generation R Study. *Eur J Clin Nutr* 2015;69(3):314–21.
 129. Laurson K, Eisenmann JC, Moore S. Lack of association between television viewing, soft drinks, physical activity and body mass index in children. *Acta Paediatr* 2008;97(6):795–800.
 130. Skinner JD, Carruth BR. A longitudinal study of children's juice intake and growth: the juice controversy revisited. *J Am Diet Assoc* 2001;101(4):432–7.
 131. Costa CD S, Assunção MCF, Loret de Mola C, Cardoso JS, Matijasevich A, Barros AJD, et al. Role of ultra-processed food in fat mass index between 6 and 11 years of age: a cohort study. *Int J Epidemiol* 2021;50:256–65.
 132. de Ruyter JC, Olthof MR, Seidell JC, Katan MB. A trial of sugar-free or sugar-sweetened beverages and body weight in children. *N Engl J Med* 2012;367(15):1397–406.
 133. Ebbeling CB, Feldman HA, Chomitz VR, Antonelli TA, Gortmaker SL, Osganian SK, et al. A randomized trial of sugar-sweetened beverages and adolescent body weight. *N Engl J Med* 2012;367(15):1407–16.
 134. Ebbeling CB, Feldman HA, Osganian SK, Chomitz VR, Ellenbogen SJ, Ludwig DS. Effects of decreasing sugar-sweetened beverage consumption on body weight in adolescents: a randomized, controlled pilot study. *Pediatrics* 2006;117(3):673–80.
 135. Obbagy JE, English LK, Psota TL, Wong YP, Butte NF, Dewey KG, et al. Complementary feeding and micronutrient status: a systematic review. *Am J Clin Nutr* 2019;109(Suppl 1):852S–71S.
 136. Global Diet Quality Project. Homepage. [Internet]. [Cited 2021 Dec 21]. Available from: <https://www.globaldietquality.org/>.
 137. Lachat C, Hawwash D, Ocké MC, Berg C, Forsum E, Hörnell A, et al. Strengthening the Reporting of Observational Studies in Epidemiology—Nutritional Epidemiology (STROBE-nut): an extension of the STROBE statement. *PLoS Med* 2016;13(6):e1002036.