



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/186408/>

Version: Published Version

Article:

Rimmer, M.P., Howie, R.A., Subramanian, V. et al. (2022) Outcome reporting across randomized controlled trials evaluating potential treatments for male infertility: a systematic review. *Human Reproduction Open*, 2022 (2). hoac010. ISSN: 2399-3529

<https://doi.org/10.1093/hropen/hoac010>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>


Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Outcome reporting across randomized controlled trials evaluating potential treatments for male infertility: a systematic review

Michael P. Rimmer ^{1,†}, Ruth A. Howie ^{2,†}
Venkatesh Subramanian ³, Richard A. Anderson ^{1,2}
Ricardo Pimenta Bertolla ⁴, Yusuf Beebejaun ³
Pietro Bortoletto ⁵, Sesh K. Sunkara ⁶, Rod T. Mitchell ¹
Allan Pacey ⁷, Madelon van Wely ⁸, Cindy M. Farguhar ^{9,10}
James M.N. Duffy ^{3,*} and Craig Niederberger ^{11,12}

¹MRC Centre for Reproductive Health, Queens Medical research Institute, University of Edinburgh, Edinburgh, UK ²Edinburgh Fertility Centre, Simpsons Centre for Reproductive Health, Royal Infirmary of Edinburgh, Edinburgh, UK ³King's Fertility, The Fetal Medicine Research Unit, King's College London, London, UK ⁴Division of Urology, Department of Surgery, Universidade Federal de Sao Paulo, São Paulo, Brazil ⁵The Ronald O. Perelman and Claudia Cohen Center for Reproductive Medicine, Weill Cornell Medicine, New York, NY, USA ⁶Division of Women's Health, Faculty of Life Sciences and Medicine, King's College London, London, UK ⁷Department of Oncology and Metabolism, University of Sheffield, Sheffield, UK ⁸Amsterdam University Medical Centers, Amsterdam, The Netherlands ⁹Cochrane Gynaecology and Fertility Group, Auckland, New Zealand ¹⁰Department of Obstetrics and Gynaecology, University of Auckland, Auckland, New Zealand ¹¹Department of Urology, University of Illinois at Chicago, Chicago, IL, USA ¹²Department of Bioengineering, University of Illinois at Chicago College of Engineering, Chicago, IL, USA

*Correspondence address. King's Fertility, The Fetal Medicine Research Unit, King's College London, London, UK. Tel: +44-7949-066806; E-mail: james.duffy@kingsfertility.co.uk  <https://orcid.org/0000-0001-6127-9859>

Submitted on November 05, 2021; resubmitted on February 18, 2022; editorial decision on February 28, 2022

STUDY QUESTION: What are the primary outcomes and outcome measures used in randomized controlled trials (RCTs) evaluating potential treatments for male infertility in the last 10 years?

SUMMARY ANSWER: Outcome reporting across male infertility trials is heterogeneous with numerous definitions and measures used to define similar outcomes.

WHAT IS KNOWN ALREADY: No core outcome set for male infertility trials has been developed. Male infertility trials are unique in that they have potentially three participants, a man, a female partner and their offspring and this will likely lead to significant variation in outcome reporting in randomized trials.

STUDY DESIGN, SIZE, DURATION: A systematic review of RCTs mapping outcomes and outcome measures evaluating potential treatments for men with infertility registered in the Cochrane Register of Controlled Trials (CENTRAL) between January 2010 and July 2021.

PARTICIPANTS/MATERIALS, SETTING, METHODS: Abstract screening and study selection was undertaken in duplicate using a review protocol that was developed prior to commencing the review. No risk of bias assessment was undertaken as this review aims to report on outcome reporting only.

MAIN RESULTS AND THE ROLE OF CHANCE: One hundred and seventy-five RCTs were identified, and given the large number of studies we limited our review to the 100 largest trials. Seventy-nine different treatments were reported across the 100 largest RCTs including vitamin and dietary supplements (18 trials), surgical treatments (18 trials) and sperm selection techniques (22 trials). When considering the largest 100 trials (range: 80–2772 participants), 36 primary and 89 secondary outcomes were reported. Forty-seven trials reported a primary outcome and 36 trials clearly defined their primary outcome. Pregnancy outcomes were inconsistently reported and included pregnancy rate (51 trials), pregnancy loss including miscarriage, ectopic pregnancy, stillbirth (9 trials) and live birth (13 trials). Trials

[†]The authors consider that the first two authors should be regarded as joint First Authors.

© The Author(s) 2022. Published by Oxford University Press on behalf of European Society of Human Reproduction and Embryology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

consistently reporting the same outcome frequently used different definitions. For example, semen quality was reported by 75 trials and was defined in 7 different ways, including: the World Health Organization (WHO) 2010 criteria (32 trials), WHO 1999 criteria (18 trials), WHO 1992 criteria (3 trials), WHO 1999 and 1992 criteria (1 trial) and the Kruger strict morphology criteria (1 trial).

LIMITATIONS, REASONS FOR CAUTION: We only evaluated the 100 largest trials published in the last 10 years and did not report outcomes on the remaining 75. An outcome was included as a primary outcome only if clearly stated in the manuscript and we did not contact authors to clarify this. As our review mapped outcomes and outcome measures, we did not undertake an integrity assessment of the trials included in our review.

WIDER IMPLICATIONS OF THE FINDINGS: Most randomized trials evaluating treatments for male infertility report different outcomes. Only half of the RCTs reported pregnancy rate and even fewer reported live birth; furthermore, the definitions of these outcomes varies across trials. Developing, disseminating and implementing a minimum data set, known as a core outcome set, for male infertility research could help to improve outcome selection, collection and reporting.

STUDY FUNDING/COMPETING INTEREST(S): A.P.—chairman of external scientific advisory committee of Cryos International Denmark ApS, member of the scientific advisory board for Cytoswim LDT and ExSeed Health. Guest lecture at the 'Insights for Fertility Conference', funded by MERK SERONO Limited. M.v.W.—holds a ZON-MW research grant. No external funding was obtained for this study.

Key words: clinical practice guidelines / core outcome set / male infertility / outcome reporting / randomized controlled trials / systematic review

WHAT DOES THIS MEAN FOR PATIENTS?

This study looks at what information randomized controlled trials (RCTs) collect and report, to help evaluate possible treatments for male infertility.

Male infertility affects millions of men worldwide, and many different treatments have been proposed for this. Treatments with the potential to reduce this health burden require robust evaluation. When assessing new treatments, RCTs are considered the 'gold-standard' method. How effective these treatments are can only be truly understood if clinical trials report the same outcomes, which are measured and defined in the same way.

We identified many RCTs that reported different outcomes, for example semen parameters, pregnancy rate or live birth, making it challenging to combine the results of these trials. Even when trials did report the same outcome, for example pregnancy rate, the outcome was either undefined or defined in numerous different ways. This means that when new RCTs are published to evaluate a treatment for male infertility, researchers and clinicians may not be able to truly understand its potential benefit for patients, in the context of previously published research.

Introduction

Infertility affects 50 million couples globally (Martinez et al., 2012; Vander Borgh and Wyns, 2018). Male factor infertility affects up to 18 million men worldwide (Winters and Walsh, 2014; Agarwal et al., 2015) and is recognized as a contributing factor in up to one-third of cases (Thonneau et al., 1991; Agarwal et al., 2015; Tamrakar and Bastakoti, 2019). Treatments with the potential to reduce this health burden require robust evaluation. When assessing new treatments, randomized controlled trials (RCTs) are considered the gold-standard method to determine the efficacy and safety of potential treatments (Liberati et al., 2009). However, despite their potentially robust design, methodology and conduct, RCTs are only as meaningful as the outcomes they collect and report (Ioannidis et al., 2014; Duffy et al., 2019).

Complex issues, including a failure to consider the perspectives of people with fertility problems when selecting outcomes, variations in outcome definitions and measurement instruments as well as outcome reporting bias can make the selection, collection and reporting of outcomes challenging. The unique nature of male infertility research can add further complexity as outcomes will often need to consider three

research participants, namely the male, his female partner or gestational carrier, and their subsequent offspring.

Little is known about outcome reporting in male infertility clinical trials. To understand the heterogeneity in outcome reporting of RCTs in male infertility, and provide a basis for more consistent reporting to the highest possible standards, we undertook a systematic review of the outcomes and outcome measures reported by the 100 largest RCTs published over the last 10 years. Reporting on the outcomes, outcome measures and consistency of these outcomes across trials will enable us to identify how outcome reporting could be standardized in future trials. This will allow researchers to better understand the true efficacy of interventions assessed in RCTs to address male infertility.

Materials and methods

A protocol was developed prior to commencing the review and included clearly defined objectives, including search criteria, study selection criteria and extraction of data (Supplementary Data). We

followed the reporting guidelines for systematic reviews of RCTs, as outlined by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement (Minebois *et al.*, 2017).

The objective of our review was to characterize outcome reporting across RCTs evaluating interventions for male infertility. Our main outcome of interest was primary outcome reporting in these trials and the definition of this outcome. RCTs were identified by searching the Cochrane Register of Controlled Trials (CENTRAL) for RCTs published between 1 January 2010 and 24 July 2021. CENTRAL is populated by the Cochrane Collaboration by regularly searching the Cumulative Index to Nursing and Allied Health Literature (CINAHL), EMBASE, MEDLINE and PsycINFO (Supplementary Data). Two authors (M.P.R. and R.A.H.) independently performed the screening of each potentially relevant record, based on the title and abstract and reviewed the full text of each selected study to assess eligibility. Where data were reported twice, such as a conference abstract and peer reviewed paper published at a later date, extracted data from the peer reviewed paper was used. Discrepancies between the authors were resolved through discussion and a consensus being agreed.

We included all RCTs which evaluated potential treatments for male factor infertility. We excluded systematic reviews and non-randomized trials. We limited our search to publications written in English. The largest 100 RCTs based on the number of participants were included in our analysis.

Using a standardized data extraction form, two authors (M.P.R. and R.A.H.) independently extracted study characteristics, nature of the intervention and both the primary and secondary outcomes reported. We reported the definitions used for commonly reported outcomes, including semen quality, pregnancy rate and live birth, to illustrate how these definitions varied. An outcome was considered to be a primary outcome only if this was clearly stated in the method section. Discrepancies between authors were resolved through discussion and a consensus being achieved. A comprehensive inventory of outcomes was developed. We used descriptive statistics to characterize outcome reporting across included RCTs. No risk of bias was undertaken as the scope of this review was to report outcome reporting across RCTs and not to assess the quality of the trials.

Results

We identified 1620 records. After excluding 12 duplicate records, 1608 titles and abstracts were screened to identify RCTs evaluating interventions for male infertility (Fig. 1). We excluded 1411 records as they were either non-randomized studies, systematic reviews, did not report an intervention for male infertility or did not report on a male infertility cohort. Two independent reviewers evaluated the remaining 197 potentially relevant trials of which 175 were deemed to be relevant. From these, the largest 100 RCTs reporting data from 24 542 men (range: 80–2772 men) were used to identify and report outcomes (Abdel-Maguid and Othman, 2010; Fang *et al.*, 2010; Fayez *et al.*, 2010; Kovacic *et al.*, 2010; Abdel-Meguid *et al.*, 2011; Azadi *et al.*, 2011; Balaban *et al.*, 2011; Figueira Rde *et al.*, 2011; Hafeez *et al.*, 2011; Safarinejad, 2011; Safarinejad *et al.*, 2011; Selice *et al.*, 2011; Turhan *et al.*, 2011; Wilding *et al.*, 2011; Amirzargar *et al.*, 2012; Colacurci *et al.*, 2012; El-Khayat *et al.*, 2012; Lee *et al.*, 2012; Mansour

Ghanaie *et al.*, 2012; Parmegiani *et al.*, 2012; Rago *et al.*, 2012; Safarinejad *et al.*, 2012; Velaers *et al.*, 2012; Azizollahi *et al.*, 2013; De Vos *et al.*, 2013; Gopinath *et al.*, 2013; Kang *et al.*, 2013; Leandri *et al.*, 2013; Majumdar and Majumdar, 2013; Pan *et al.*, 2013; WorriLOW *et al.*, 2013; Akin *et al.*, 2014; Karamahmutoglu *et al.*, 2014; Kolahdooz *et al.*, 2014; Moslemi Mehni *et al.*, 2014; Nematollahi-Mahani *et al.*, 2014; Pourmand *et al.*, 2014; Raigani *et al.*, 2014; Romany *et al.*, 2014; Wang *et al.*, 2014; Calogero *et al.*, 2015; Cyrus *et al.*, 2015; Ding *et al.*, 2015; ElSheikh *et al.*, 2015; Farrag *et al.*, 2015; Guo *et al.*, 2015; Haje and Naom, 2015; Hou *et al.*, 2015; Peivandi *et al.*, 2015; Sikka *et al.*, 2015; Youssef and Abdalla, 2015; Hosseini *et al.*, 2016; Jin *et al.*, 2016; Nasr Esfahani *et al.*, 2016; Pan *et al.*, 2016; Park *et al.*, 2016; Bryniarski *et al.*, 2017; Guo *et al.*, 2017; Milardi *et al.*, 2017; Qu *et al.*, 2017; Rosety *et al.*, 2017; Taiyeb *et al.*, 2017; Hajizadeh Maleki and Tartibian, 2017a,b; Babak *et al.*, 2018; Blomberg Jensen *et al.*, 2018; Bodin *et al.*, 2018; Busetto *et al.*, 2018; Habous *et al.*, 2018; Hajizadeh Maleki and Tartibian, 2018; Ketabchi and Salajegheh, 2018; Ketabchi *et al.*, 2018; Nasimi Doost Azgomi *et al.*, 2018; Sun *et al.*, 2018; Tsounapi *et al.*, 2018; Almekaty *et al.*, 2019; Chen *et al.*, 2019; De Geyter *et al.*, 2019; Hajizadeh Maleki *et al.*, 2019; Kizilay and Altay, 2019; Lin *et al.*, 2019; Mangoli *et al.*, 2019; Miller *et al.*, 2019; Tehrani *et al.*, 2019; Yetkinel *et al.*, 2019; Yu *et al.*, 2019; Zhao *et al.*, 2019; Chen *et al.*, 2020; Degirmenci *et al.*, 2020; Eslamian *et al.*, 2020; Hajizadeh Maleki and Tartibian, 2020; Hasanen *et al.*, 2020; Huang *et al.*, 2020; Joseph *et al.*, 2020; Karimi *et al.*, 2020; Kopets *et al.*, 2020; Liu *et al.*, 2020; Schisterman *et al.*, 2020; Bozhedomov *et al.*, 2021; Salas-Huetos *et al.*, 2021).

Seventy-nine different treatments were reported across the 100 RCTs (Table I). These included trials reporting vitamin or dietary supplements or nutraceuticals (n = 18), surgical procedures (n = 18) and sperm selection or modification techniques (n = 22).

Primary and secondary outcomes

One hundred and four outcomes were reported across the included trials (Tables II and III).


Thirty-six different primary outcomes were reported by 47 trials with 13 of these 47 trials (28%) reporting a definition of these outcomes. Commonly reported primary outcomes included semen quality (16 trials; 34%), pregnancy rate (13 trials; 28%) and live birth (4 trials; 9%).

Ninety-six trials reported 89 different secondary outcomes. Reported secondary outcomes were heterogeneous and included semen quality (52 trials; 54%), pregnancy rate (39 trials; 41%), pregnancy loss (9 trials, 10%) and live birth (9 trials; 10%) (Table III). Primary and secondary outcomes reported by the 25 largest RCTs are outlined in Table IV.

Definitions

Nine trials defined live birth as a primary or secondary outcome in two different ways: birth >37 weeks' gestation; and birth <37 weeks' gestation. The remaining seven trials did not define this term.

Pregnancy rate was reported by 51 trials as either a primary or secondary outcome, with 12 different definitions used by 21 trials. The remaining 26 trials did not define pregnancy rate, and 4 definitions were unclear. Definitions varied greatly, from a threshold of serum hCG >25IU/l and the presence of a gestational sac on ultrasound scan (USS) to a viable foetus on transvaginal USS (Table V).



Semen quality was reported by 75 trials as either a primary or secondary outcome and there was a comparatively higher level of consensus between trials. A total of 57/75 trials defined these by the World Health Organization (WHO) criteria, having used either [WHO 2010](#) (n = 32), [WHO 1999](#) (n = 18), [WHO 1992](#) (n = 3), [WHO 1999](#) and [1992](#) (n = 1), in three the WHO semen analysis edition was not specified. Of the remaining studies, one used the Kruger strict morphology ([Ketabchi et al., 2018](#)) and the remaining 17 trials did not define this outcome ([Table V](#)).

Not all of the studies included in our review used the most up to date edition of the WHO criteria available when conducting their trial. Studies defining semen analysis parameters using [WHO 1992](#) criteria were commenced in 2012 and 2006 and could have utilized the [WHO 1999](#) criteria when conducting the trial ([Selice et al., 2011](#); [Sikka et al., 2015](#)). A similar issue was identified with some studies defining semen analysis criteria using [WHO 1999](#), where the trial was commenced in 2013 or 2016 after the introduction of [WHO 2010](#) ([Haje and Naoom, 2015](#); [Hosseini et al., 2016](#); [Tehrani et al., 2019](#)). Although these trials may initially appear to use an outdated version of the WHO semen analysis manual, their design may have occurred

prior to the publication of an updated WHO criteria. Deviation from the initial analysis plan, potentially using two different WHO criteria or favouring one criterion in particular, may have been considered a violation of the trial protocol. To achieve consistent outcome reporting, should these trials have used new WHO criteria, they may no longer be comparable to older trials evaluating similar interventions for male infertility.

Discussion

This systematic evaluation of the literature of RCTs in male factor infertility identified a range of primary and secondary outcomes relevant to male, maternal and neonatal participants. Many trials omitted important information about the primary outcome of the trial and how this was defined. Of the 100 randomized trials included in our review, only 47 clearly stated a primary outcome in their methodology. This lack of clear outcome reporting is not uncommon and has been identified as a problem in other areas, including in IVF, neonatal and

Table 1 Characteristics of the 100 largest trials included in this review evaluating interventions for male infertility.

Study	Intervention group one	Intervention group two	Participants (n)
Miller <i>et al.</i> (2019)	Physiological intracytoplasmic sperm injection	Intracytoplasmic sperm injection	2772
Schisterman <i>et al.</i> (2020)	Folic acid and zinc sulphate	Placebo	2370
WorriLOW <i>et al.</i> (2013)	Hyaluronic binding prior to intracytoplasmic sperm injection	Intracytoplasmic sperm injection	802
Huang <i>et al.</i> (2020)	Folic acid	Placebo	769
Kovacic <i>et al.</i> (2010)	Embryo culture in 5% oxygen	Embryo culture in 20% oxygen	647
Hajizadeh Maleki and Tartibian (2017a)	Exercise	No intervention	556
Hajizadeh Maleki and Tartibian (2020)	Exercise	No intervention	441
Hajizadeh Maleki and Tartibian (2018)	Exercise	No intervention	430
Hajizadeh Maleki and Tartibian(2017b)	Exercise	No intervention	419
Hasanen <i>et al.</i> (2020)	Physiological intracytoplasmic sperm injection	Magnetic activated cell sorting	413
Sun <i>et al.</i> (2018)	Bilateral varicocelectomy	Unilateral varicocelectomy	358
Ding <i>et al.</i> (2015)*	Recombinant FSH	Sodium chloride injection	354
Turhan <i>et al.</i> (2011)	Double sperm wash	Single sperm wash	341
De Vos <i>et al.</i> (2013)	Intracytoplasmic morphologically selected sperm injection	Intracytoplasmic sperm injection	340
Chen <i>et al.</i> (2019)*	Chymotrypsin treatment	Vitamin C, E, zinc gluconate and a spermatogenic tablet	337
Almekaty <i>et al.</i> (2019)	Artery preserving varicocelectomy	Artery ligating varicocelectomy	330
Blomberg Jensen <i>et al.</i> (2018)	Vitamin D and calcium	Placebo	330
Zhao <i>et al.</i> (2019)	hCG and hMG	Placebo	316
Velaers <i>et al.</i> (2012)	Single touch sperm immobilization	Triple touch sperm immobilization	290
Hajizadeh Maleki <i>et al.</i> (2019)*	Exercise	No intervention	283
Habous <i>et al.</i> (2018)*	Clomiphene citrate	hCG injections	282
Romany <i>et al.</i> (2014)	Sperm swim up and removal of annexin V positive sperm	Sperm swim up	263
Safarinejad <i>et al.</i> (2011)	Saffron	Placebo	260
Leandri <i>et al.</i> (2013)	Intracytoplasmic morphologically selected sperm injection	Intracytoplasmic sperm injection	255
Safarinejad (2011)	Pentoxifylline	Placebo	254
Wilding <i>et al.</i> (2011)	Intracytoplasmic morphologically selected sperm injection	Intracytoplasmic sperm injection	250
Taiyeb <i>et al.</i> (2017)	Prednisolone	Placebo	241
Moslemi Mehni <i>et al.</i> (2014)*	Pentoxifylline and L-carnitine	Placebo	235
Bodin <i>et al.</i> (2018)	Counselling	No intervention	229
Safarinejad <i>et al.</i> (2012)	Oral ubiquinol	Placebo	228
Karamahmutoglu <i>et al.</i> (2014)	Density gradient centrifugation	Swim up sperm preparation	223
Sikka <i>et al.</i> (2015)	Pregabalin	Placebo	222
Karimi <i>et al.</i> (2020)	Density gradient centrifugation and zeta selection	Density gradient centrifugation	220
Lin <i>et al.</i> (2019)	GnRH	hCG and hMG	220
Fang <i>et al.</i> (2010)*	Spermatic vein ligation, vitamin E, pentoxifylline and clomiphene	Vitamin E, pentoxifylline and clomiphene	219
Tsounapi <i>et al.</i> (2018)*	Phosphodiesterase type-5 inhibitor	No intervention	217
Rago <i>et al.</i> (2012)	Vardenafil	No intervention	205
Nasr Esfahani <i>et al.</i> (2016)	Density gradient centrifugation and zeta selection	Density gradient centrifugation	203
Joseph <i>et al.</i> (2020)	Vitamin C, Vitamin E, Zing	No antioxidants	200

(continued)

Table I Continued

Study	Intervention group one	Intervention group two	Participants (n)
Calogero et al. (2015)	Myoinositol and folic acid	Folic acid	194
Babak et al. (2018)	Varicocelectomy and hCG	Varicocelectomy	193
Guo et al. (2015)	Doppler ultrasound assisted subinguinal microscopic varicocelectomy	Microscopic varicocelectomy	180
Eslamian et al. (2020)	DHA vitamin, Vitamin E	Placebo	180
Balaban et al. (2011)	Intracytoplasmic morphologically selected sperm injection	Intracytoplasmic sperm injection	168
Abdel-Maguid and Othman (2010)	Microsurgical subinguinal varicocelectomy	Subinguinal varicocelectomy	162
Bozhedomov et al. (2021)	Hydrophilic nutrients	Lipophilic nutrients	160
Nematollahi-Mahani et al. (2014)*	Zinc sulphate and folic acid	Placebo	160
Azizollahi et al. (2013)*	Varicocelectomy and zinc sulphate	Varicocelectomy and placebo	160
Majumdar and Majumdar (2013)	Physiological intracytoplasmic sperm injection	Intracytoplasmic sperm injection	156
Fayez et al. (2010)*	Varicocelectomy Ivanissevich technique	Varicocelectomy subinguinal sclerotherapy	155
Abdel-Meguid et al. (2011)	Microsurgical subinguinal varicocelectomy	Subinguinal varicocelectomy	150
Mangoli et al. (2019)	Intracytoplasmic morphologically selected sperm injection	Intracytoplasmic sperm injection	150
Guo et al. (2017)	Doppler ultrasound at laparoscopic varicocelectomy	Laparoscopic varicocelectomy	147
Pan et al. (2016)*	Dietary supplement, Chinese herbal medicine and zinc selenium	Chinese herbal medicine	147
Ketabchi and Salajegheh (2018)*	Microscopic varicocelectomy and acupuncture	Sham acupuncture	140
Gopinath et al. (2013)*	Antioxidants	Placebo	138
Ketabchi et al. (2018)	Microsurgical subinguinal varicocelectomy	No intervention	138
Ghanaie et al. (2012)	Varicocele repair	No intervention	136
Colacurci et al. (2012)	FSH	Vitamin supplement	129
Haje and Naoom (2015)*	Tamoxifen and L-carnitine	Placebo	128
Yetkinel et al. (2019)	Microfluidic sperm selection	Conventional swim up technique	122
Yu et al. (2019)*	Transcutaneous electrical acupuncture point stimulation 2 hertz	Lifestyle advice	121
El-Khayat et al. (2012)	Fallopian tube sperm perfusion	IUI	120
Figueira Rde et al. (2011)	Intracytoplasmic morphologically selected sperm injection	Intracytoplasmic sperm injection	120
Pan et al. (2013)	Inguinal varicocelectomy	Subinguinal varicocelectomy	120
Liu et al. (2020)	Green model lifestyle intervention	Conventional nursing	120
Salas-Huetos et al. (2021)	60 g mixed nuts	Nuts	119
Chen et al. (2020)	Yishen tongluo recipe	Minimally invasive surgery	116
Cyrus et al. (2015)	Varicocele and vitamin C	Varicocele and placebo	115
Amirzargar et al. (2012)*	Varicocelectomy and hCG	Varicocelectomy	113
De Geyter et al. (2019)	Sperm preparation and deselecting sperm with fragmented DNA	Conventional sperm preparation	111
Hosseini et al. (2016)	Ginger	Placebo	106
Selice et al. (2011)	FSH	No intervention	105
Busetto et al. (2018)	Nutritional supplement	Placebo	104
Azadi et al. (2011)	Varicocelectomy and zaditen	Varicocelectomy and placebo	103

(continued)

Table I Continued

Study	Intervention group one	Intervention group two	Participants (n)
Akin <i>et al.</i> (2014)*	Varicocelectomy and ligation with titanium clips	Varicocelectomy and ligation with surgical silk	100
Nasimi Doost Azgomi <i>et al.</i> (2018)	Withania somnifera	Pentoxifylline	100
Hafeez <i>et al.</i> (2011)	Herbal medicine	Allopathic medicine	100
Hou <i>et al.</i> (2015)	Microsurgical subinguinal varicocelectomy with testicular delivery	Microsurgical subinguinal varicocelectomy without testicular delivery	100
Parmegiani <i>et al.</i> (2012)	Physiological intracytoplasmic sperm injection	Intracytoplasmic sperm injection with sperm slow selection device	100
Pourmand <i>et al.</i> (2014)	Varicocelectomy	Varicocelectomy and L-Carnitine	100
Kizilay and Altay (2019)	Varicocelectomy and antioxidant	Varicocelectomy	93
Youssef and Abdalla (2015)	Single laparoscopic varicocelectomy	Transperitoneal varicocelectomy	93
ElSheikh <i>et al.</i> (2015)*	Vitamin E	Clomiphene citrate	90
Milardi <i>et al.</i> (2017)*	Prednisolone 5 mg	Prednisolone 12.5 mg	90
Peivandi <i>et al.</i> (2015)	Intrauterine insemination	Intrauterine insemination with fallopian tube sperm transfer	90
Rosety <i>et al.</i> (2017)	Exercise	No intervention	90
Wang <i>et al.</i> (2014)	Laparoscopic varicocelectomy	Transperitoneal varicocelectomy	90
Degirmenci <i>et al.</i> (2020)*	0–2 days sexual abstinence	2–3 days sexual abstinence; >4 days sexual abstinence	90
Qu <i>et al.</i> (2017)	Varicocelectomy and xuanju	Varicocelectomy	88
Bryniarski <i>et al.</i> (2017)	Laparoscopic varicocelectomy	Microsurgical varicocelectomy	84
Jin <i>et al.</i> (2016)	Intracytoplasmic sperm injection, selecting sperm bound to zona pellucida	Intracytoplasmic sperm injection	84
Raigani <i>et al.</i> (2014)*	Folic acid and zinc sulphate	Placebo	83
Farrag <i>et al.</i> (2015)	Recombinant FSH and intracytoplasmic sperm injection	Intracytoplasmic sperm injection	82
Lee <i>et al.</i> (2012)	Transperitoneal laparoscopic varicocele ligation	Laparoscopic single-site varicocele ligation	82
Kopets <i>et al.</i> (2020)	L-carnitine/acetyl-L-carnitine, L-arginine, glutathione, co-enzyme Q10, zinc, vitamin B9, vitamin B12, selenium	Placebo	83
Kang <i>et al.</i> (2013)	Varicocele ligation with vessel and lymphatic preservation	Varicocele ligation without vessel and lymphatic preservation	80
Kolahdooz <i>et al.</i> (2014)	<i>Nigella sativa</i> oil	Liquid paraffin	80
Park <i>et al.</i> (2016)*	Varicocelectomy and Chinese herbal medicine	Placebo	80
Tehrani <i>et al.</i> (2019)*	Hypo-osmotic swelling test and intracytoplasmic sperm injection	Intracytoplasmic sperm injection	80

*Multiarm trial.

endometriosis trials (Hirsch *et al.*, 2016; Wilkinson *et al.*, 2016; Webbe *et al.*, 2020).

Interpretation

RCTs can be challenging to undertake and expensive to conduct; as such there is an ethical imperative to conduct them to the highest possible standards (Macleod *et al.*, 2014). Less than half of the 100 largest trials included in our review reported a clearly defined primary outcome, which represents a lost opportunity to obtain further robust data to inform clinical decision-making. Where trials reported the

same primary outcome, often different measurement tools and endpoints were used to define these, which precludes pooling data from these trials. Even trials with seemingly consistent primary outcome reporting and definitions are not without their limitations. We identified 57 trials using WHO semen analysis methods to report, with primary or secondary outcomes. Although WHO semen analysis is a robustly developed standard, there have now been six different editions, of which three were used in our identified trials, although the most up to date edition was used in the majority of trials at the time these they were conducted (World Health Organization, 1992, 1999, 2010). Furthermore, semen is highly variable, even within individuals,

Table II Primary outcomes reported in the 100 largest randomized trials evaluating interventions for male infertility.

Hormonal

Serum oestradiol
Serum FSH
Serum LH
Serum sex hormone-binding globulin
Serum testosterone

Metabolic

Assessment of endothelial function
Bioelectrical impedance analysis
Blood pressure
BMI
Waist circumference
Serum markers of metabolic function

Semen

Semen pH
Semen volume
Sperm concentration
Sperm count
Sperm density
Sperm morphology
Sperm motility
Total motile sperm count
Sperm DNA fragmentation index

Embryological

Fertilization rate
Embryo development
Embryo quality

Pregnancy outcomes

Spontaneous pregnancy
Pregnancy following ART
Intrauterine pregnancy confirmed by ultrasound
Ongoing pregnancy confirmed by ultrasound (from 12 weeks onwards)
Ongoing pregnancy (>20 weeks)
Cumulative pregnancy rate
Live birth
Live birth at term

Other

Fertility awareness knowledge
Awareness of lifestyle factors affecting fertility
Satisfaction with sexual life
Sexual intercourse frequency
Patient-reported symptoms of androgen deficiency
Testicular pain

with improved pregnancy outcomes. This finding challenges the assumption that improved pregnancy outcomes are always associated with improved semen quality and not achieved through other factors including the population under study and the intervention used. Once such example is [Huang et al. \(2020\)](#), who demonstrated that folic acid supplementation was only effective at improving semen quality and pregnancy outcomes in a subgroup of patients with the homozygous polymorphism of the MTHFR 677 gene, while all other MTHFR polymorphisms studied showed no effect. Variable response in semen quality and pregnancy outcome was demonstrated by [Hajizadeh Maleki et al. \(2020\)](#) who investigated high-intensity interval training, reporting both semen parameters and live births. Patients categorized as asthenozoospermic, asthenoteratozoospermic, oligospermic or oligoasthenozoospermic demonstrated significantly improved semen quality following their exercise regime. Analysis of pregnancy outcomes in these cohorts, however, did not reveal a significant increase in live births. Another trial included in this review ([Haje and Naoom, 2015](#)) reported the impact of tamoxifen and L-carnitine on semen parameters and pregnancy outcomes. Although semen parameters, including sperm count, sperm motility and sperm morphology, were found to be improved in men receiving tamoxifen or tamoxifen with L-carnitine compared to placebo or L-carnitine only, these improvements did not translate into a significant increase in pregnancy rate.

In addition to outcome selection, inconsistent outcome reporting may result from a lack of validated instruments or poorly defined endpoints. One example is the assessment of sperm DNA fragmentation, for which at least eight different methods are available, with variable results obtained based on the test used and the laboratory undertaking the assessment ([Agarwal et al., 2016a,b](#); [Pacey, 2018](#)). Despite the large number of trials published on male factor infertility and the range of primary and secondary outcomes reported on, this inconsistency fundamentally limits their clinical utility and value to inform decision-making and patient care. In addition to difficulties in pooling results of trials, a lack of agreed core outcomes presents challenges for researchers designing future trials when selecting the outcomes to report, further compounded when considering factors such as sample size, cost and time.

Our systematic review is the first to report on the primary and secondary outcomes reported in male factor infertility trials and the definitions used for the primary outcome. It builds on work undertaken in other areas of reproductive health to identify causes of subfertility and harmonize the way these data are reported ([Duffy et al., 2017a,b](#); [Lee et al., 2020](#); [Turner et al., 2020](#); [Rimmer et al., 2021](#)). At present, there is no consensus on definitions to be used for outcomes relevant to male factor infertility. To address inconsistencies in outcome reporting across male and female infertility trials, an international working group of healthcare professionals and researchers have developed the Core Outcome Measures for Infertility Trials (COMMIT) initiative ([Core Outcomes in Women's and Newborn Health Initiative, 2014](#)). This initiative will develop stakeholder-driven development of core outcome sets relevant to clinicians, researchers, and patients and has developed a consensus strategy for reporting core outcomes and standardizing their definitions ([Duffy et al., 2020, 2021](#)).

As no core outcome set for male fertility trials has been developed, it is therefore not surprising that we identified little consistency between outcome reporting and definitions used. This is further compounded by the nature of male infertility trials and the interventions

which furthers the argument that semen quality may, in itself, not be an informative primary endpoint ([Oshio et al., 2004](#); [Castilla et al., 2006](#)). This is demonstrated in some of the trials included in our review, which showed that improved semen quality did not correlate

Table III Secondary outcomes reported in the 100 largest randomized trials evaluating interventions for male infertility.**Clinical examination**

Testicular volume
 Varicocele grade
 Spermatic vein diameter
 Physical fitness assessed by continuous maximal incremental test
 Bioelectrical impedance analysis
 Body mass index
 Waist circumference

Hormonal

Serum oestradiol
 Serum FSH
 Serum inhibin B
 Serum LH
 Serum testosterone
 Serum inhibin B to FSH ratio
 Prostate-specific antigen
 Haematocrit
 Serum alanine aminotransferase
 Serum aspartate aminotransferase

Semen

Semen liquefaction time
 Semen pH
 Semen volume
 Sperm concentration
 Sperm count
 Sperm density
 Sperm morphology
 Sperm motility
 Sperm DNA fragmentation index
 Time to initiation of spermatogenesis
 Acrosome integrity
 Sperm penetration assay
 Levels of reactive oxygen species
 Malondialdehyde levels in seminal plasma

Embryological

Fertilization rate
 Number of embryos
 Embryo quality
 Number of embryos available for transfer
 Number of embryos cryopreserved
 Number of euploid embryos
 Number of blastocysts
 Blastocyst quality

Pregnancy and childbirth

Gestational diabetes
 Pre-eclampsia
 Stillbirth
 Gestational age at delivery
 Live birth
 Pregnancies to term
 Preterm birth
 Caesarean delivery

Maternal complications

Anaemia requiring blood transfusion
 Haemolysis, elevated liver enzymes, low platelet count syndrome
 Postpartum haemorrhage
 Seizure
 Sepsis

Neonatal outcomes

Birthweight
 Small for gestational age
 Neonatal mortality
 Bronchopulmonary dysplasia
 Chromosomal anomalies
 Necrotizing enterocolitis
 Periventricular leucomalacia
 Retinopathy of prematurity
 Severe intraventricular haemorrhage
 Structural malformations

Intraoperative outcomes

Operating time
 Number of internal spermatic veins ligated
 Number of internal spermatic arteries preserved
 Haematoma formation
 Hydrocele
 Infection
 Pain
 Pyrexia
 Testicular atrophy

Postoperative outcomes

Patient satisfaction
 Time to return to normal activities
 Recurrence of varicocele

Resource utilization

Length of hospital stay
 Cost

(continued)

Table III Continued**Early pregnancy**

Spontaneous pregnancy
 Pregnancy following ART
 β hCG detected pregnancy
 Intrauterine pregnancy confirmed by ultrasound
 Singleton pregnancy
 Multiple pregnancy
 Early pregnancy loss
 Ectopic pregnancy
 Late pregnancy loss
 Time to conception
 Cumulative pregnancy rate

Other

Testosterone deficiency symptoms
 Prostatic symptoms
 Sexual dysfunction

studied. Trials reporting on exercise or dietary supplements to improve semen quality may not report the same outcomes as techniques to select sperm to be used in ART to achieve a pregnancy. Researchers planning future male infertility trials should consider either using a core outcome set for these trials or, in the absence of this, consider reporting outcomes and outcome measures previously used in the literature to improve pooling of data across trials.

This review highlights inconsistencies in outcome reporting across male infertility trials but can also be used to identify commonly reported outcomes, when designing future trials. Using previously reported outcomes in new trials when evaluating interventions for male infertility may allow data from these trials to be pooled and meta-analysed. The outcomes identified in this review can be used to develop a core outcome set following discussion by a group of multinational, multiprofessional stakeholders as has been done for general infertility trials (Duffy et al., 2021). Development of this core outcome set would guide researchers in which core outcomes to report, allow data from several trials to be pooled better inform patient care and reduce research waste (Duffy et al., 2017a,b). We plan to develop a core outcome set for future male infertility research using outcomes reported in this review, using a modified Delphi method and modified Nominal Group Technique to identify relevant outcomes, their measurement and definitions.

Strength and limitations

Our review has several strengths. The comprehensive search strategy and methodological design gives us confidence in the results we have identified. Collecting outcomes reported by 100 trials means that the outcomes and definitions identified reflect a significant body of work and are representative of the field of male infertility trials. To avoid bias, abstract screening and data extraction were undertaken by two independent reviewers, utilizing a third to resolve any queries and reach a consensus. However, our review is not without its limitations. For example, owing to the large number of RCTs published in male factor infertility, we only included the 100 largest trials. This means smaller trials were excluded, and inclusion of the data in these trials may have altered the results obtained and the conclusions drawn. Many trials reported on outcomes but did not clearly state it was the primary outcome or base a sample size calculation on this; as such,

they were not included as a primary outcome in our review, despite being reported on by the authors. We did not contact the authors to clarify the primary outcomes where it was not clearly stated or if the researchers extracting data were unsure. We were also unable to validate the quality of the outcomes reported, as there is no validated tool to do this. As our review assessed outcome reporting and the definitions of these outcomes, we did not undertake an integrity check of the trials included in our review. However, we did identify that the conduct of some of the included trials could have been improved upon. These include one trial which was not registered (Karimi et al., 2020) and four that were retrospectively registered (Moslemi Mehni et al., 2014; Youssef and Abdalla, 2015; Taiyeb et al., 2017; Ketabchi et al., 2018). One trial obtained ethics committee approval after trial registration and the proposed recruitment period (according to the clinical trial registration), although specific recruitment dates were not included in the manuscript (Ketabchi et al., 2018). One study had an enrolment period that ended 8 months prior to submission of the manuscript but also reported live birth as an outcome (Majumdar and Majumdar, 2013).

The conduct and integrity of RCTs are central to their ability to produce robust high-quality evidence (Li et al., 2020). This would be improved by the implementation of a core outcome set for future male infertility research and assessment of trial integrity when undertaking systematic reviews and meta-analysis.

Conclusion

Randomized trials reporting on interventions for male factor infertility frequently omit a primary outcome and often report these outcomes differently. This hinders the utility of these trials in how their results can be combined to inform health care professionals' clinical decision-making and improve patient outcomes. Developing a core outcome set for male infertility trials will help inform how primary outcome measures are selected and reported on and translate into meaningful improvements in patient care.

Supplementary data

Supplementary data are available at *Human Reproduction Open* online.

Table IV Detailed primary and secondary outcomes reported in the largest 25 randomized trials reevaluating interventions for male infertility.

Study	Primary outcomes														Secondary outcomes																					
	No. of participants	Serum testosterone levels	Semen analysis*	Sperm DNA fragmentation index	Fertilization rate	Embryo development	Spontaneous pregnancy rate	Pregnancy following ART	Ongoing pregnancy confirmed by USS	Livebirth	Livebirth at term	Reproductive hormones	Semen analysis*	Malondialdehyde levels	Sperm DNA fragmentation index	Fertilization rate	No. of embryos	Embryo quality	No. of embryos cryopreserved	Spontaneous pregnancy	Pregnancy following ART	Implantation rate	βhCG detected pregnancy	Intrauterine pregnancy confirmed by USS	Multiple pregnancy	Early pregnancy loss	Ectopic pregnancy	Pregnancy rate	Cumulative pregnancy rate	Time to conception	Pregnancy outcomes*	Gestational age at delivery	Livebirth	Preterm birth	Testosterone deficiency symptoms	
Miller et al. (2019)	2772									●														●		●									●	
Schisterman et al. (2020)	2370	●	●						●														●	●	●	●	●			●				●		
Worilow et al. (2013)	802							●														●				●										
Huang et al. (2020)	769												●	●	●									●		●						●	●			
Kovacic et al. (2010)	647							●								●	●					●	●	●		●		●								
Hajizadeh Maleki and Tartibian (2017a)	556	●	●																															●		
Hajizadeh Maleki and Tartibian (2020)	441												●		●																			●		
Hajizadeh Maleki and Tartibian (2018)	430												●		●																			●		
Hajizadeh Maleki and Tartibian (2017b)	419												●		●																			●		
Hasanen et al. (2020)	413							●									●					●	●													
Sun et al., (2018)	358						●						●																							
Ding et al. (2015)	354	●					●	●																												
Turhan et al. (2011)	341						●						●																							
De Vos et al. (2013)	340				●	●															●															
Chen et al. (2019)	337																											●		●						
Almekaty et al. (2019)	330	●																										●								
Jensen et al. (2018)	330	●																															●			

(continued)

Table IV Continued

Study	Primary outcomes													Secondary outcomes																							
	No. of participants	Serum testosterone levels	Semen analysis*	Sperm DNA fragmentation index	Fertilization rate	Embryo development	Spontaneous pregnancy rate	Pregnancy following ART	Ongoing pregnancy confirmed by USS	Livebirth	Livebirth at term	Reproductive hormones	Semen analysis*	Malondialdehyde levels	Sperm DNA fragmentation index	Fertilization rate	No. of embryos	Embryo quality	No. of embryos cryopreserved	Spontaneous pregnancy	Pregnancy following ART	Implantation rate	βhCG detected pregnancy	Intrauterine pregnancy confirmed by USS	Multiple pregnancy	Early pregnancy loss	Ectopic pregnancy	Pregnancy rate	Cumulative pregnancy rate	Time to conception	Pregnancy outcomes*	Gestational age at delivery	Livebirth	Preterm birth	Testosterone deficiency symptoms		
Zhao <i>et al.</i> (2019)	316										●	●		●					●																		
Velaers <i>et al.</i> (2012)	290				●											●	●					●	●														
Hajizadeh Maleki <i>et al.</i> (2019)	283			●																																	
Habous <i>et al.</i> (2018)	282	●																																		●	
Romany <i>et al.</i> (2014)	263								●						●								●			●											
Safarinejad (2011)	260											●																									
Leandri <i>et al.</i> (2013)	255											●		●	●	●		●			●		●										●				
Safarinejad (2011)	254										●	●																									

● denotes if outcome included in trial.

*Semen analysis includes: semen volume; sperm concentration; sperm motility; sperm morphology; sperm count; and total motile sperm count.

*Pregnancy outcomes includes: caesarean section; pre-eclampsia; gestational diabetes; gestational age at delivery; birth weight; small for gestational age; severe postpartum maternal morbidity (including postpartum haemorrhage, anaemia requiring transfusion, sepsis, seizure, HELLP [haemolysis, elevated level of liver enzymes, low platelet count] syndrome, and pre-eclampsia with pulmonary oedema), major neonatal complications (including structural malformations, chromosomal anomalies, bronchopulmonary dysplasia, necrotizing enterocolitis, severe intraventricular haemorrhage, periventricular leukomalacia, and retinopathy of prematurity), still-birth and neonatal death.

USS, ultrasound scan.

Table V Variation in outcome reporting definitions across the 100 largest trials evaluating interventions for male infertility.**Semen parameters (n = 75)**

- WHO 2010 criteria (n = 32)
- WHO 1999 criteria (n = 18)
- WHO 1992 criteria (n = 3)
- WHO 1999 and 1992 criteria (n = 1)
- WHO edition not specified (n = 3)
- Kruger strict morphology test (n = 1)
- Undefined (n = 17)

Pregnancy (n = 51)

- Serum hCG
- Positive hCG test (n = 1)
- >25 IU/l (n = 1)
- >50 IU/l (n = 1)
- >60 IU/l (n = 1)
- Serum hCG >25 IU/l and USS confirmation (n = 1)

Ultrasound examination

- Presence of one or more gestational sacs (n = 5)
- Presence of a gestational sac with or without foetal heartbeat (n = 1)
- Presence of a gestational sac with foetal heartbeat (n = 2)
- Presence of a gestational sac or foetal heartbeat (n = 1)
- >1 embryo with a foetal heartbeat (n = 4)
- Foetal heart beat (n = 2)
- Presence of a gestational sac or histological assessment confirming PoC (n = 1)
- Unclear (n = 4)
- Undefined (n = 26)

Live birth (n = 9)

- Birth >37 weeks gestation (n = 1)
- Birth <37 weeks gestation (n = 1)
- Undefined (n = 7)

Three outcomes were selected to demonstrate variation across studies in how common outcomes were defined differently. The outcomes selected for this were semen analysis, pregnancy and live birth.

PoC, product of conception; USS, ultrasound scan; WHO, World Health Organization.

Data availability

The data underlying this article will be shared on reasonable request to the corresponding author.

Authors' roles

M.P.R., R.A.H., V.S. and J.M.N.D. undertook the searches, data extraction and drafted the manuscript. M.P.R., R.A.H., V.S., R.A.A., Y.B., R.P.B., S.K.S., A.P., B.P., R.T.M., A.P., M.v.W., C.M.F., C.N. and J.M.N.D. participated in data analysis and interpretation, preparation of the manuscript and critically revising the paper. C.M.F., C.N. and J.M.N.D. conceived the idea of the manuscript. All authors approved the final version of the manuscript.

Funding

This work received no funding.

Conflict of interest

A.P.—chairman of external scientific advisory committee of Cryos International Denmark ApS, member of the scientific advisory board for Cytoswim LDT and ExSeed Health. Guest lecture at the 'Insights for Fertility Conference'. M.v.W.—holds a ZON-MW research grant.

References

- Abdel-Maguid AF, Othman I. Microsurgical and nonmagnified subinguinal varicocelectomy for infertile men: a comparative study. *Fertil Steril* 2010;**94**:2600–2603.
- Abdel-Meguid TA, Al-Sayyad A, Tayib A, Farsi HM. Does varicocele repair improve male infertility? An evidence-based perspective from a randomized, controlled trial. *Eur Urol* 2011;**59**:455–461.
- Agarwal A, Cho CL, Esteves SC. Should we evaluate and treat sperm DNA fragmentation? *Curr Opin Obstet Gynecol* 2016a;**28**:164–171.
- Agarwal A, Majzoub A, Esteves SC, Ko E, Ramasamy R, Zini A. Clinical utility of sperm DNA fragmentation testing: practice

- recommendations based on clinical scenarios. *Transl Androl Urol* 2016b;**5**:935–950.
- Agarwal A, Mulgund A, Hamada A, Chyatte MR. A unique view on male infertility around the globe. *Reprod Biol Endocrinol* 2015;**13**: 37.
- Akin Y, Ateş M, Yücel S, Başara I, Çelik O, Bozkurt A, Nuhoglu B. Comparison of different ligation techniques in laparoscopic varicocelectomy. *Turk J Med Sci* 2014;**44**:273–278.
- Almekaty K, Zahran MH, Zoeir A, Minhas S, Salem K. The role of artery-preserving varicocelectomy in subfertile men with severe oligozoospermia: a randomized controlled study. *Andrology* 2019;**7**: 193–198.
- Amirzargar MA, Yavangi M, Basiri A, Hosseini Moghaddam SM, Babbolhavaeji H, Amirzargar N, Amirzargar H, Moadabshoar L. Comparison of recombinant human follicle stimulating hormone (rhFSH), human chorionic gonadotropin (HCG) and human menopausal gonadotropin (HMG) on semen parameters after varicocelectomy: a randomized clinical trial. *Iran J Reprod Med* 2012;**10**: 441–452.
- Azadi L, Abbasi H, Deemeh MR, Tavalaei M, Arbabian M, Pilevarian AA, Nasr-Esfahani MH. Zaditen (Ketotifen), as mast cell blocker, improves sperm quality, chromatin integrity and pregnancy rate after varicocelectomy. *Int J Androl* 2011;**34**:446–452.
- Azizollahi G, Azizollahi S, Babaei H, Kianinejad M, Baneshi MR, Nematollahi-Mahani SN. Effects of supplement therapy on sperm parameters, protamine content and acrosomal integrity of varicocelectomized subjects. *J Assist Reprod Genet* 2013;**30**:593–599.
- Babak J, Behruz F, Mohammadreza Y, Morteza FK. The effect of human chorionic gonadotropin therapy on semen parameters and pregnancy rate after varicocelectomy. *Curr Urol* 2018;**11**:92–96.
- Balaban B, Yakin K, Alatas C, Oktem O, Isiklar A, Urman B. Clinical outcome of intracytoplasmic injection of spermatozoa morphologically selected under high magnification: a prospective randomized study. *Reprod Biomed Online* 2011;**22**:472–476.
- Blomberg Jensen M, Lawaetz JG, Petersen JH, Juul A, Jørgensen N. Effects of vitamin D supplementation on semen quality, reproductive hormones, and live birth rate: a randomized clinical trial. *J Clin Endocrinol Metab* 2018;**103**:870–881.
- Bodin M, Tydén T, Käll L, Larsson M. Can reproductive life plan-based counselling increase men's fertility awareness? *Ups J Med Sci* 2018;**123**:255–263.
- Bozhedomov VA, Epanchintseva EA, Bozhe-Domova GE, Kamarina RA, Rokhlikov IM, Kamalov AA. [Hydrophilic and lipophilic nutrients for the treatment of male idiopathic infertility: a randomized, comparative, open-label, multicenter, prospective, controlled study]. *Urol Mosc* 2021;**70**–78.
- Bryniarski P, Taborowski P, Rajwa P, Kaletka Z, Życzkowski M, Paradysz A. The comparison of laparoscopic and microsurgical varicocelectomy in infertile men with varicocele on paternity rate 12 months after surgery: a prospective randomized controlled trial. *Andrology* 2017;**5**:445–450.
- Busetto GM, Agarwal A, Virmani A, Antonini G, Ragonesi G, Del Giudice F, Micic S, Gentile V, De Berardinis E. Effect of metabolic and antioxidant supplementation on sperm parameters in oligoastheno-teratozoospermia, with and without varicocele: a double-blind placebo-controlled study. *Andrology* 2018;**50**:e12927.
- Calogero AE, Gullo G, La Vignera S, Condorelli RA, Vaiarelli A. Myo-inositol improves sperm parameters and serum reproductive hormones in patients with idiopathic infertility: a prospective double-blind randomized placebo-controlled study. *Andrology* 2015;**3**:491–495.
- Castilla JA, Álvarez C, Aguilar J, González-Varea C, Gonzalvo MC, Martínez L. Influence of analytical and biological variation on the clinical interpretation of seminal parameters. *Hum Reprod* 2006;**21**: 847–851.
- Chen X, Sun ZX, Zhao SP, Zhang XH, Chen JS, Wang R, Men B. [Yishen Tongluo Recipe combined with minimally invasive surgery for the treatment of varicocele-associated asthenospermia]. *Zhonghua Nan Ke Xue* 2020;**26**:341–345.
- Chen Z, Song T, Yan Y, Chen C, Liu T, Wen X, Yao Y, Zou C, Li X, Xu Q et al. Prodom: a new assisted reproductive device to treat male infertility caused by impaired semen liquefaction. *Int J Clin Exp Med* 2019;**12**:4088–4099.
- Colacurci N, Monti MG, Fornaro F, Izzo G, Izzo P, Trotta C, Mele D, De Franciscis P. Recombinant human FSH reduces sperm DNA fragmentation in men with idiopathic oligoastheno-teratozoospermia. *J Androl* 2012;**33**:588–593.
- Core Outcomes in Women's and Newborn Health Initiative. The CROWN Initiative: journal editors invite researchers to develop core outcomes in women's health. *Hum Reprod* 2014;**29**:1349–1350.
- Cyrus A, Kabir A, Goodarzi D, Moghimi M. The effect of adjuvant vitamin C after varicocele surgery on sperm quality and quantity in infertile men: a double blind placebo controlled clinical trial. *Int Braz J Urol* 2015;**41**:230–238.
- De Geyter C, Gobrecht-Keller U, Ahler A, Fischer M. Removal of DNA-fragmented spermatozoa using flow cytometry and sorting does not improve the outcome of intracytoplasmic sperm injection. *J Assist Reprod Genet* 2019;**36**:2079–2086.
- De Vos A, Van de Velde H, Bocken G, Eylenbosch G, Franceus N, Meersdom G, Tistaert S, Vankelecom A, Tournaye H, Verheyen G. Does intracytoplasmic morphologically selected sperm injection improve embryo development? A randomized sibling-oocyte study. *Hum Reprod* 2013;**28**:617–626.
- Degirmenci Y, Demirdag E, Guler I, Yildiz S, Erdem M, Erdem A. Impact of the sexual abstinence period on the production of seminal reactive oxygen species in patients undergoing intrauterine insemination: a randomized trial. *J Obstet Gynaecol Res* 2020;**46**: 1133–1139.
- Ding YM, Zhang XJ, Li JP, Chen SS, Zhang RT, Tan WL, Shi XJ. Treatment of idiopathic oligozoospermia with recombinant human follicle-stimulating hormone: a prospective, randomized, double-blind, placebo-controlled clinical study in Chinese population. *Clin Endocrinol (Oxf)* 2015;**83**:866–871.
- Duffy J, Bhattacharya S, Herman M, Mol B, Vail A, Wilkinson J, Farquhar C. Reducing research waste in benign gynaecology and fertility research. *BJOG* 2017a;**124**:366–369.
- Duffy JMN, AlAhwany H, Bhattacharya S, Collura B, Curtis C, Evers JLH, Farquharson RG, Franik S, Giudice LC, Khalaf Y et al. Developing a core outcome set for future infertility research: an international consensus development study. *Hum Reprod* 2020;**35**: 2725–2734.

- Duffy JMN, Bhattacharya S, Bhattacharya S, Bofill M, Collura B, Curtis C, Evers JLH, Giudice LC, Farquharson RG, Franik S et al. Standardizing definitions and reporting guidelines for the infertility core outcome set: an international consensus development study. *Fertil Steril* 2021; **115**:201–212.
- Duffy JMN, Hirsch M, Gale C, Pealing L, Kawsar A, Showell M, Williamson PR, Khan KS, Ziebland S, McManus RJ; International Collaboration to Harmonize Outcomes for Pre-eclampsia (iHOPE). A systematic review of primary outcomes and outcome measure reporting in randomized trials evaluating treatments for pre-eclampsia. *Int J Gynaecol Obstet* 2017b; **139**:262–267.
- Duffy JMN, Ziebland S, von Dadelszen P, McManus RJ. Tackling poorly selected, collected, and reported outcomes in obstetrics and gynecology research. *Am J Obstet Gynecol* 2019; **220**: 71.e1–71.e4.
- El-Khayat W, El-Mazny A, Abou-Salem N, Moafy A. The value of fallopian tube sperm perfusion in the management of mild-moderate male factor infertility. *Int J Gynaecol Obstet* 2012; **117**:178–181.
- ElSheikh MG, Hosny MB, Elshenoufy A, Elghamrawi H, Fayad A, Abdelrahman S. Combination of vitamin E and clomiphene citrate in treating patients with idiopathic oligoasthenozoospermia: a prospective, randomized trial. *Andrology* 2015; **3**:864–867.
- Eslamian G, Amirjannati N, Noori N, Sadeghi MR, Hekmatdoost A. Effects of coadministration of DHA and vitamin E on spermogram, seminal oxidative stress, and sperm phospholipids in asthenozoospermic men: a randomized controlled trial. *Am J Clin Nutr* 2020; **112**:707–719.
- Fang Y, Zhao L, Yan F, Xia X, Xu D, Cui X. Escin improves sperm quality in male patients with varicocele-associated infertility. *Phytomedicine* 2010; **17**:192–196.
- Farrag A, Sagnella F, Pappalardo S, Costantini A, Lisi F, Carfagna P, Manna C. The use of r-hFSH in treatment of idiopathic male factor infertility before ICSI. *Eur Rev Med Pharmacol Sci* 2015; **19**: 2162–2167.
- Fayez A, El Shantaly KM, Abbas M, Hauser S, Müller SC, Fathy A. Comparison of inguinal approach, scrotal sclerotherapy and subinguinal antegrade sclerotherapy in varicocele treatment: a randomized prospective study. *Urol Int* 2010; **85**:200–203.
- Figueira Rde C, Braga DP, Setti AS, Iaconelli A Jr, Borges E. Morphological nuclear integrity of sperm cells is associated with preimplantation genetic aneuploidy screening cycle outcomes. *Fertil Steril* 2011; **95**:990–993.
- Gopinath P, Kalra B, Saxena A, Malik S, Kochhar K, Kalra S, Zaveri H. Fixed dose combination therapy of antioxidants in treatment of idiopathic oligoasthenozoospermia: results of a randomized, double-blind placebo-controlled clinical trial. *Int J Infertil Fetal Med* 2013; **4**:6–13.
- Guo L, Sun W, Shao G, Song H, Ge N, Zhao S, Liu Y, Zhang X, Xiao Z, Yuan M. Outcomes of microscopic subinguinal varicocelectomy with and without the assistance of doppler ultrasound: a randomized clinical trial. *Urology* 2015; **86**:922–928.
- Guo LQ, Zhang XL, Liu YQ, Sun WD, Zhao ST, Yuan MZ. The application of laparoscopic Doppler ultrasound during laparoscopic varicocelectomy in infertile men. *Asian J Androl* 2017; **19**:214–218.
- Habous M, Giona S, Tealab A, Aziz M, Williamson B, Nassar M, Abdelrahman Z, Remeah A, Abdelkader M, Binsaleh S et al. Clomiphene citrate and human chorionic gonadotropin are both effective in restoring testosterone in hypogonadism: a short-course randomized study. *BJU Int* 2018; **122**:889–897.
- Hafeez M, Ahmed A, Usmanhani K, Mohiuddin E, Asif M, Akram MRU, Rehman M. Clinical evaluation of herbal medicine for oligospermia. *Pak J Nutr* 2011; **10**:238–240.
- Haje M, Naoom K. Combined tamoxifen and L-carnitine therapies for the treatment of idiopathic male infertility attending intracytoplasmic sperm injection: a randomized controlled trial. *Int J Infertil Fetal Med* 2015; **6**:20–24.
- Hajizadeh Maleki B, Tartibian B, Chehrizi M. Effects of aerobic, resistance, and combined exercise on markers of male reproduction in healthy human subjects: a randomized controlled trial. *J Strength Cond Res* 2019; **33**:1130–1145.
- Hajizadeh Maleki B, Tartibian B. Combined aerobic and resistance exercise training for improving reproductive function in infertile men: a randomized controlled trial. *Appl Physiol Nutr Metab* 2017a; **42**:1293–1306.
- Hajizadeh Maleki B, Tartibian B. High-intensity interval training modulates male factor infertility through anti-inflammatory and antioxidative mechanisms in infertile men: a randomized controlled trial. *Cytokine* 2020; **125**:154861.
- Hajizadeh Maleki B, Tartibian B. Moderate aerobic exercise training for improving reproductive function in infertile patients: a randomized controlled trial. *Cytokine* 2017b; **92**:55–67.
- Hajizadeh Maleki B, Tartibian B. Resistance exercise modulates male factor infertility through anti-inflammatory and antioxidative mechanisms in infertile men: a RCT. *Life Sci* 2018; **203**:150–160.
- Hasanen E, Elqusi K, ElTanbouly S, Hussin AE, AlKhadr H, Zaki H, Henkel R, Agarwal A. PICSI vs. MACS for abnormal sperm DNA fragmentation ICSI cases: a prospective randomized trial. *J Assist Reprod Genet* 2020; **37**:2605–2613.
- Hirsch M, Duffy JM, Kuzsnir JO, Davis CJ, Plana MN, Khan KS, Farquhar C, Johnson N, Khan K; International Collaboration to Harmonize Outcomes and Measures for Endometriosis. Variation in outcome reporting in endometriosis trials: a systematic review. *Am J Obstet Gynecol* 2016; **214**:452–464.
- Hosseini J, Mardi Mamaghani A, Hosseiniifar H, Sadighi Gilani MA, Dadkhah F, Sepidarkish M. The influence of ginger (*Zingiber officinale*) on human sperm quality and DNA fragmentation: a double-blind randomized clinical trial. *Int J Reprod Biomed* 2016; **14**: 533–540.
- Hou Y, Zhang Y, Zhang Y, Huo W, Li H. Comparison between microsurgical subinguinal varicocelectomy with and without testicular delivery for infertile men: is testicular delivery an unnecessary procedure. *Urol J* 2015; **12**:2261–2266.
- Huang WJ, Lu XL, Li JT, Zhang JM. Effects of folic acid on oligozoospermia with MTHFR polymorphisms in term of seminal parameters, DNA fragmentation, and live birth rate: a double-blind, randomized, placebo-controlled trial. *Andrology* 2020; **8**:110–116.
- Ioannidis JP, Greenland S, Hlatky MA, Khoury MJ, Macleod MR, Moher D, Schulz KF, Tibshirani R. Increasing value and reducing waste in research design, conduct, and analysis. *Lancet* 2014; **383**: 166–175.
- Jin R, Bao J, Tang D, Liu F, Wang G, Zhao Y, Bai G, Liu Y, Wang Y, Liu L et al. Outcomes of intracytoplasmic sperm injection using the zona pellucida-bound sperm or manually selected sperm. *J Assist Reprod Genet* 2016; **33**:597–601.

- Joseph T, Mascarenhas M, Karuppusami R, Karthikeyan M, Kunjummen AT, Kamath MS. Antioxidant pretreatment for male partner before ART for male factor subfertility: a randomized controlled trial. *Hum Reprod Open* 2020;**2020**:hoaa050.
- Kang DH, Lee JY, Chung JH, Jo JK, Lee SH, Ham WS, Cho KS, Lee KS, Kim TH, Lee SW. Laparoendoscopic single site varicocele ligation: comparison of testicular artery and lymphatic preservation versus complete testicular vessel ligation. *J Urol* 2013;**189**:243–249.
- Karamahmutoglu H, Erdem A, Erdem M, Mutlu MF, Bozkurt N, Oktem M, Ercan DD, Gumuslu S. The gradient technique improves success rates in intrauterine insemination cycles of unexplained subfertile couples when compared to swim up technique; a prospective randomized study. *J Assist Reprod Genet* 2014;**31**:1139–1145.
- Karimi N, Mohseni Kouchesfahani H, Nasr-Esfahani MH, Tavalae M, Shahverdi A, Choobineh H. DGC/Zeta as a new strategy to improve clinical outcome in male factor infertility patients following intracytoplasmic sperm injection: a randomized, single-blind, clinical trial. *Cell J* 2020;**22**:55–59.
- Ketabchi AA, M, Salehi, S, Salajghah, S. The effect of varicolectomy on assisted reproductive technique indications and outcomes based on kruger strict morphology test: a randomized clinical trial. *J Kerman Univ Med Sci* 2018;**25**:519–527.
- Ketabchi AA, Salajegheh S. The effects of acupuncture treatment in infertile patients with clinical varicocele. *Nephro-Urol Mon* 2018;**10**:e65451.
- Kizilay F, Altay B. Evaluation of the effects of antioxidant treatment on sperm parameters and pregnancy rates in infertile patients after varicolectomy: a randomized controlled trial. *Int J Impot Res* 2019;**31**:424–431.
- Kolahdooz M, Nasri S, Modarres SZ, Kianbakht S, Huseini HF. Effects of *Nigella sativa* L. seed oil on abnormal semen quality in infertile men: a randomized, double-blind, placebo-controlled clinical trial. *Phytomedicine* 2014;**21**:901–905.
- Kopets R, Kuibida I, Chernyavska I, Cherepanyn V, Mazo R, Fedevych V, Gerasymov S. Dietary supplementation with a novel l-carnitine multi-micronutrient in idiopathic male subfertility involving oligo-, astheno-, teratozoospermia: a randomized clinical study. *Andrologia* 2020;**8**:1184–1193.
- Kovacic B, Sajko MC, Vlaisavljević V. A prospective, randomized trial on the effect of atmospheric versus reduced oxygen concentration on the outcome of intracytoplasmic sperm injection cycles. *Fertil Steril* 2010;**94**:511–519.
- Leandri RD, Gachet A, Pfeffer J, Celebi C, Rives N, Carre-Pigeon F, Kulski O, Mitchell V, Parinaud J. Is intracytoplasmic morphologically selected sperm injection (IMSI) beneficial in the first ART cycle? a multicentric randomized controlled trial. *Andrology* 2013;**1**:692–697.
- Lee D, Kim SK, Lee JR, Jee BC. Management of endometriosis-related infertility: considerations and treatment options. *Clin Exp Reprod Med* 2020;**47**:1–11.
- Lee SW, Lee JY, Kim KH, Ha US. Laparoendoscopic single-site surgery versus conventional laparoscopic varicocele ligation in men with palpable varicocele: a randomized, clinical study. *Surg Endosc* 2012;**26**:1056–1062.
- Li W, van Wely M, Gurrin L, Mol BW. Integrity of randomized controlled trials: challenges and solutions. *Fertil Steril* 2020;**113**:1113–1119.
- Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol* 2009;**62**:e1–e34.
- Lin J, Mao J, Wang X, Ma W, Hao M, Wu X. Optimal treatment for spermatogenesis in male patients with hypogonadotropic hypogonadism. *Medicine (Baltimore)* 2019;**98**:e16616.
- Liu J, Wang JG, Xu HX, Sun Y, Peng L, Sun J, Xie S. [Application of the Green Model to lifestyle intervention in male sterility patients]. *Zhonghua Nan Ke Xue* 2020;**26**:441–445.
- Macleod MR, Michie S, Roberts I, Dirnagl U, Chalmers I, Ioannidis JP, Al-Shahi Salman R, Chan AW, Glasziou P. Biomedical research: increasing value, reducing waste. *Lancet* 2014;**383**:101–104.
- Majumdar G, Majumdar A. A prospective randomized study to evaluate the effect of hyaluronic acid sperm selection on the intracytoplasmic sperm injection outcome of patients with unexplained infertility having normal semen parameters. *J Assist Reprod Genet* 2013;**30**:1471–1475.
- Mangoli E, Khalili MA, Talebi AR, Agha-Rahimi A, Soleimani M, Faramarzi A, Pouretezari M. IMSI procedure improves clinical outcomes and embryo morphokinetics in patients with different aetiologies of male infertility. *Andrologia* 2019;**51**:e13340.
- Mansour Ghanaie M, Asgari SA, Dadrass N, Allahkhah A, Iran-Pour E, Safarinejad MR. Effects of varicocele repair on spontaneous first trimester miscarriage: a randomized clinical trial. *Urol J* 2012;**9**:505–513.
- Martinez G, Daniels K, Chandra A. Fertility of men and women aged 15–44 years in the United States: National Survey of Family Growth, 2006–2010. *Natl Health Stat Report* 2012;**51**:1–28.
- Milardi D, Luca G, Grande G, Ghezzi M, Caretta N, Brusco G, De Filpo G, Marana R, Pontecorvi A, Calafiore R et al. Prednisone treatment in infertile patients with oligozoospermia and accessory gland inflammatory alterations. *Andrology* 2017;**5**:268–273.
- Miller D, Pavitt S, Sharma V, Forbes G, Hooper R, Bhattacharya S, Kirkman-Brown J, Coomarasamy A, Lewis S, Cutting R et al. Physiological, hyaluronan-selected intracytoplasmic sperm injection for infertility treatment (HABSelect): a parallel, two-group, randomised trial. *Lancet* 2019;**393**:416–422.
- Minebois H, De Souza A, Mezan de Malartic C, Agopianz M, Guillet May F, Morel O, Callec R. [Endometriosis and miscarriage: systematic review]. *Gynecol Obstet Fertil Senol* 2017;**45**:393–399.
- Moslemi Mehni N, Ketabchi AA, Hosseini E. Combination effect of Pentoxifylline and L-carnitine on idiopathic oligoasthenoteratozoospermia. *Iran J Reprod Med* 2014;**12**:817–824.
- Nasimi Doost Azgomi R, Nazemiyeh H, Sadeghi Bazargani H, Fazljou SMB, Nejatbakhsh F, Moini Jazani A, Ahmadi AsrBadr Y, Zomorodi A. Comparative evaluation of the effects of Withania somnifera with pentoxifylline on the sperm parameters in idiopathic male infertility: a triple-blind randomised clinical trial. *Andrologia* 2018;**50**:e13041.
- Nasr Esfahani MH, Deemeh MR, Tavalae M, Sekhavati MH, Gourabi H. Zeta sperm selection improves pregnancy rate and

- alters sex ratio in male factor infertility patients: a double-blind, randomized clinical trial. *Int J Fertil Steril* 2016;**10**:253–260.
- Nematollahi-Mahani SN, Azizollahi GH, Baneshi MR, Safari Z, Azizollahi S. Effect of folic acid and zinc sulphate on endocrine parameters and seminal antioxidant level after varicocelelectomy. *Andrologia* 2014;**46**:240–245.
- Oshio S, Ashizawa Y, Yotsukura M, Tohyama Y, Iwabuchi M, Adachi Y, Matsuda H, Tomomasa H, Yoshida S, Takeda K *et al*. Individual variation in semen parameters of healthy young volunteers. *Arch Androl* 2004;**50**:417–425.
- Pacey A. Is sperm DNA fragmentation a useful test that identifies a treatable cause of male infertility? *Best Pract Res Clin Obstet Gynaecol* 2018;**53**:11–19.
- Pan F, Pan L, Zhang A, Liu Y, Zhang F, Dai Y. Comparison of two approaches in microsurgical varicocelelectomy in chinese infertile males. *Urol Int* 2013;**90**:443–448.
- Pan Z, Deng N, Zou Z. Clinical observation of effect of diosmin combined with jinshuibao capsule treatment on improvement of semen quality in patients with varicocele. *Med Sci Tech* 2016;**57**:42–46.
- Park HJ, Choe S, Park NC. Effects of Korean red ginseng on semen parameters in male infertility patients: a randomized, placebo-controlled, double-blind clinical study. *Chin J Integr Med* 2016;**22**:490–495.
- Parmegiani L, Cognigni GE, Bernardi S, Troilo E, Taraborrelli S, Arnone A, Maccarini AM, Filicori M. Comparison of two ready-to-use systems designed for sperm-hyaluronic acid binding selection before intracytoplasmic sperm injection: PICSI vs. Sperm Slow: a prospective, randomized trial. *Fertil Steril* 2012;**98**:632–637.
- Peivandi S, Ebadi A, Modanlu S. The comparison between intrauterine insemination and fallopian tube sperm perfusion using FAST[®] system in patients with unexplained infertility. *Int J Fertil Steril* 2015;**8**:379–384.
- Pourmand G, Movahedin M, Dehghani S, Mehraei A, Ahmadi A, Pourhosein M, Hoseini M, Ziloochi M, Heidari F, Beladi L *et al*. Does L-carnitine therapy add any extra benefit to standard inguinal varicocelelectomy in terms of deoxyribonucleic acid damage or sperm quality factor indices: a randomized study. *Urology* 2014;**84**:821–825.
- Qu X, Shan Z, Zhang N, Guo L. Curative effect of surgery in combination with compound xuanju capsule in treating subclinical varicocele induced infertility. *Biomed Res* 2017;**28**:1247–1250.
- Rago R, Salacone P, Caponecchia L, Marcucci I, Fiori C, Sebastianelli A. Effect of vardenafil on semen parameters in infertile men: a pilot study evaluating short-term treatment. *J Endocrinol Invest* 2012;**35**:897–900.
- Raigani M, Yaghmaei B, Amirjannti N, Lakpour N, Akhondi MM, Zeraati H, Hajhosseinal M, Sadeghi MR. The micronutrient supplements, zinc sulphate and folic acid, did not ameliorate sperm functional parameters in oligoasthenoteratozoospermic men. *Andrologia* 2014;**46**:956–962.
- Rimmer MP, Fishwick K, Henderson I, Chinn D, Al Wattar BH, Quenby S. Quantifying CD138+ cells in the endometrium to assess chronic endometritis in women at risk of recurrent pregnancy loss: a prospective cohort study and rapid review. *J Obstet Gynaecol Res* 2021;**47**:689–697.
- Romany L, Garrido N, Motato Y, Aparicio B, Remohi J, Meseguer M. Removal of annexin V-positive sperm cells for intracytoplasmic sperm injection in ovum donation cycles does not improve reproductive outcome: a controlled and randomized trial in unselected males. *Fertil Steril* 2014;**102**:1567–1575.e1.
- Rosety M, Díaz AJ, Rosety JM, Pery MT, Brenes-Martín F, Bernardi M, García N, Rosety-Rodríguez M, Ordoñez FJ, Rosety I. Exercise improved semen quality and reproductive hormone levels in sedentary obese adults. *Nutr Hosp* 2017;**34**:603–607.
- Safarinejad MR, Safarinejad S, Shafiei N, Safarinejad S. Effects of the reduced form of coenzyme Q10 (ubiquinol) on semen parameters in men with idiopathic infertility: a double-blind, placebo controlled, randomized study. *J Urol* 2012;**188**:526–531.
- Safarinejad MR. Effect of pentoxifylline on semen parameters, reproductive hormones, and seminal plasma antioxidant capacity in men with idiopathic infertility: a randomized double-blind placebo-controlled study. *Int Urol Nephrol* 2011;**43**:315–328.
- Safarinejad MR, Shafiei N, Safarinejad S. A prospective double-blind randomized placebo-controlled study of the effect of saffron (*Crocus sativus* Linn.) on semen parameters and seminal plasma antioxidant capacity in infertile men with idiopathic oligoasthenoteratozoospermia. *Phytother Res* 2011;**25**:508–516.
- Salas-Huetos A, James ER, Salas-Salvadó J, Bulló M, Aston KI, Carrell DT, Jenkins TG. Sperm DNA methylation changes after short-term nut supplementation in healthy men consuming a Western-style diet. *Andrology* 2021;**9**:260–268.
- Schisterman EF, Sjaarda LA, Clemons T, Carrell DT, Perkins NJ, Johnstone E, Lamb D, Chaney K, Van Voorhis BJ, Ryan G *et al*. Effect of folic acid and zinc supplementation in men on semen quality and live birth among couples undergoing infertility treatment: a randomized clinical trial. *JAMA* 2020;**323**:35–48.
- Selice R, Garolla A, Pengo M, Caretta N, Ferlin A, Foresta C. The response to FSH treatment in oligozoospermic men depends on FSH receptor gene polymorphisms. *Int J Androl* 2011;**34**:306–312.
- Sikka SC, Chen C, Almas M, Dula E, Knapp LE, Hellstrom WJ. Pregabalin does not affect sperm production in healthy volunteers: a randomized, double-blind, placebo-controlled, noninferiority study. *Pain Pract* 2015;**15**:150–158.
- Sun XL, Wang JL, Peng YP, Gao QQ, Song T, Yu W, Xu ZP, Chen Y, Dai YT. Bilateral is superior to unilateral varicocelelectomy in infertile males with left clinical and right subclinical varicocele: a prospective randomized controlled study. *Int Urol Nephrol* 2018;**50**:205–210.
- Taiyeb AM, Ridha-Albarzanchi MT, Taiyeb SM, Kanan ZA, Alatrakchi SK, Kjelland ME, Muhsen-Alanssari SA. Improvement in pregnancy outcomes in couples with immunologically male infertility undergoing prednisolone treatment and conventional in vitro fertilization preceded by sperm penetration assay: a randomized controlled trial. *Endocrine* 2017;**58**:448–457.
- Tamrakar SR, Bastakoti R. Determinants of infertility in couples. *J Nepal Health Res Counc* 2019;**17**:85–89.
- Tehrani M, Amirian M, Jalali M, Attaranzadeh A, Fazel A, Ebrahimzadeh-Bideskan A. Comparison of intracytoplasmic sperm injection outcome with sperm selection techniques in oligoasthenozoospermic males: a randomized controlled trial. *Iran Red Crescent Med J* 2019;**21**. doi: 10.5812/ircmj.70656.

- Thonneau P, Marchand S, Tallec A, Ferial ML, Ducot B, Lansac J, Lopes P, Tabaste JM, Spira A. Incidence and main causes of infertility in a resident population (1,850,000) of three French regions (1988-1989). *Hum Reprod* 1991;**6**:811–816.
- Tsouanpi P, Honda M, Dimitriadis F, Koukos S, Hikita K, Zachariou A, Sofikitis N, Takenaka A. Effects of a micronutrient supplementation combined with a phosphodiesterase type 5 inhibitor on sperm quantitative and qualitative parameters, percentage of mature spermatozoa and sperm capacity to undergo hyperactivation: a randomised controlled trial. *Andrologia* 2018;**50**:e13071.
- Turhan N, Pekel A, Onaran Y, Candan Z, Duvan C, Bayrak Ö. Single or double sperm wash processing by density gradient centrifugation: effect on clomiphene citrate induced intrauterine insemination cycle outcomes. *Turk J Med Sci* 2011;**41**:39–44.
- Turner KA, Rambhatla A, Schon S, Agarwal A, Krawetz SA, Dupree JM, Avidor-Reiss T. Male infertility is a women's health issue—research and clinical evaluation of male infertility is needed. *Cells* 2020;**9**:990.
- Vander Borght M, Wyns C. Fertility and infertility: definition and epidemiology. *Clin Biochem* 2018;**62**:2–10.
- Velaers A, Paternot G, Debrock S, D'Hooghe T, Spiessens C. Triple touch sperm immobilization vs. single touch sperm immobilization in ICSI—a randomised trial. *Reprod Biol Endocrinol* 2012;**10**:65.
- Wang J, Xue B, Shan YX, Cui Y, Tao W, Zhu J, Liu X, Yao Q. Laparoendoscopic single-site surgery with a single channel versus conventional laparoscopic varicocele ligation: a prospective randomized study. *J Endourol* 2014;**28**:159–164.
- Webbe JWH, Ali S, Sakonidou S, Webbe T, Duffy JMN, Brunton G, Modi N, Gale C; COIN Project Steering Committee. Inconsistent outcome reporting in large neonatal trials: a systematic review. *Arch Dis Child Fetal Neonatal Ed* 2020;**105**:69–75.
- Wilding M, Coppola G, di Matteo L, Palagiano A, Fusco E, Dale B. Intracytoplasmic injection of morphologically selected spermatozoa (IMSI) improves outcome after assisted reproduction by deselection physiologically poor quality spermatozoa. *J Assist Reprod Genet* 2011;**28**:253–262.
- Wilkinson J, Roberts SA, Showell M, Brison DR, Vail A. No common denominator: a review of outcome measures in IVF RCTs. *Hum Reprod* 2016;**31**:2714–2722.
- Winters BR, Walsh TJ. The epidemiology of male infertility. *Urol Clin North Am* 2014;**41**:195–204.
- World Health Organization. *WHO Laboratory Manual for the Examination of Human Semen and Sperm-Cervical Mucus Interaction*. Cambridge: Cambridge University Press, 1992.
- World Health Organization. *WHO Laboratory Manual for the Examination of Human Semen and Sperm-Cervical Mucus Interaction*. Cambridge: Cambridge University Press, 1999.
- World Health Organization. *WHO Laboratory Manual for the Examination and Processing of Human Semen*. Geneva: WHO, 2010.
- Worriolow KC, Eid S, Woodhouse D, Perloe M, Smith S, Witmyer J, Ivani K, Khoury C, Ball GD, Elliot T et al. Use of hyaluronan in the selection of sperm for intracytoplasmic sperm injection (ICSI): significant improvement in clinical outcomes—multicenter, double-blinded and randomized controlled trial. *Hum Reprod* 2013;**28**:306–314.
- Yetkinel S, Kılıçdag EB, Aytac PC, Haydardedeoglu B, Simsek E, Cok T. Effects of the microfluidic chip technique in sperm selection for intracytoplasmic sperm injection for unexplained infertility: a prospective, randomized controlled trial. *J Assist Reprod Genet* 2019;**36**:403–409.
- Youssef T, Abdalla E. Single incision transumbilical laparoscopic varicocelectomy versus the conventional laparoscopic technique: a randomized clinical study. *Int J Surg* 2015;**18**:178–183.
- Yu Y, Sha SB, Zhang B, Guan Q, Liang M, Zhao LG, Zhang QY, Wen J, Sun W. Effects and mechanism of action of transcutaneous electrical acupuncture point stimulation in patients with abnormal semen parameters. *Acupunct Med* 2019;**37**:25–32.
- Zhao N, Lu XL, Li JT, Zhang JM. Treatment of idiopathic oligozoospermia with combined human chorionic gonadotropin/human menopausal gonadotrophin: a randomised, double-blinded, placebo-controlled clinical study. *Andrologia* 2019;**51**:e13271.