



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/186542/>

Version: Published Version

Article:

Athavan, Gayathri, Tanner, Theo F.N., Whitwood, Adrian C. et al. (2022) Direct Evidence for Competitive C-H Activation by a Well-Defined Silver XPhos Complex in Palladium-Catalyzed C-H Functionalization. *Organometallics*. ISSN: 0276-7333

<https://doi.org/10.1021/acs.organomet.2c00063>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Direct Evidence for Competitive C–H Activation by a Well-Defined Silver XPhos Complex in Palladium-Catalyzed C–H Functionalization

Gayathri Athavan, Theo F. N. Tanner, Adrian C. Whitwood, Ian J. S. Fairlamb,* and Robin N. Perutz*



Cite This: <https://doi.org/10.1021/acs.organomet.2c00063>



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: Increasing evidence indicates that silver salts can play a role in the C–H activation step of palladium-catalyzed C–H functionalization. Here we isolate a silver(I) complex by C–H bond activation and demonstrate its catalytic competence for C–H functionalization. We demonstrate how silver carbonate, a common but highly insoluble additive, reacts with pentafluorobenzene in the presence of a bulky phosphine, XPhos, to form the C–H bond activation product $\text{Ag}(\text{C}_6\text{F}_5)(\text{XPhos})$. By isolating and fully characterizing this complex and the related carbonate and iodide complexes, $[\text{Ag}(\text{XPhos})]_2(\mu\text{-}\kappa^2, \kappa^2\text{-CO}_3)$ and $[\text{AgI}(\text{XPhos})]_2$, we show how well-defined Ag(I) complexes can operate in conjunction with palladium complexes to achieve C–H functionalization even at ambient temperature. Reactions are tested against the standard cross-coupling of $\text{C}_6\text{F}_5\text{H}$ with 4-iodotoluene, catalyzed by palladium acetate at 60 °C in the presence of silver carbonate and Xphos. Key observations are that (a) $\text{PdI}(\text{C}_6\text{H}_5)(\text{XPhos})$ reacts stoichiometrically with $\text{Ag}(\text{C}_6\text{F}_5)(\text{XPhos})$ to form $\text{Ph-C}_6\text{F}_5$ instantly at room temperature; (b) catalytic cross coupling can be achieved using 5% $\text{Ag}(\text{C}_6\text{F}_5)(\text{XPhos})$ as the sole silver source; and (c) palladium acetate (typical precatalyst) can be replaced for catalytic cross coupling by the expected oxidative addition compound $\text{PdI}(\text{C}_6\text{H}_5)(\text{XPhos})$. These investigations lead to a catalytic cycle in which Ag(I) plays the C–H bond activation role and palladium plays the coupling role. Moreover, we show how the phosphine can be exchanged between silver complexes, ensuring that it is recycled even though silver carbonate is consumed during catalytic cross-coupling.



INTRODUCTION

The functionalization of C–H bonds of low reactivity has emerged as a powerful and selective strategy for the installation of a plethora of organic motifs, increasing structural complexity and diversity in applied chemical synthesis.^{1–3} A variety of transition metals can be used to activate a C–H bond, from precious metals through to earth-abundant metals, each with their individual merits in terms of practical synthetic procedures.^{4,5} There has been significant interest in palladium salts,^{6,7} which catalyze the functionalization of a myriad of (hetero)arenes, with varying electronic and steric differences. Of interest are the electron-deficient fluorinated arenes,^{8–11} which undergo reaction with iodoarenes in the presence of phosphines and bases such as carbonates and acetates. Extensive mechanistic studies have led to a consensus that C–H bond activation occurs by the concerted metalation deprotonation (CMD) mechanism,^{9,12–15} alternatively described as ambiphilic metal–ligand assistance (AMLA-6).^{16,17} Typically, these reactions are thought to involve mononuclear Pd species, operating homogeneously, with the caveat that there is some experimental evidence implicating hybrid homogeneous–heterogeneous behavior and higher order Pd species.^{18–21} In early work, Fagnou et al. demonstrated the coupling of pentafluorobenzene to iodotoluene and studied its

mechanism; we also select this as an exemplar reaction. Fagnou et al. employed potassium carbonate as the base with $\text{P}^t\text{Bu}_2\text{Me}$ as the ligand,⁹ but later a variety of other bases were used, often including silver carbonate. Bulky phosphines continue to be employed, but even PPh_3 is active.²² In the past few years, Ag(I) has been demonstrated to play an active role in the critical C–H bond activation step in several reactions, starting with independent discoveries in 2016 by Lee and Hartwig²³ and by Larrosa et al.²⁴ Silver(I) can form several dinuclear complexes with triphenylphosphine with bridging carbonate or bicarbonate,²⁵ and silver(phosphine)halide complexes with bulky phosphines have been known since 1978. Notably, these silver complexes behave as neutral molecular species in dichloromethane solution.²⁶ In NMR spectra, they are characterized by large $^{107}\text{Ag}\text{--}^{31}\text{P}$ and $^{109}\text{Ag}\text{--}^{31}\text{P}$ coupling

Special Issue: Organometallic Chemistry Inspired by Maurice Brookhart

Received: February 2, 2022

constants that decrease as the coordination number of Ag increases.^{27–30}

A series of papers implicated silver salts in the C–H bond activation of fluoroarenes, several of which are summarized in reviews.^{7,31–33} We first consider the evidence in the absence of palladium. Simple silver salts have been shown to effect H/D exchange in pentafluorobenzene (or heptafluorotoluene) with D₂O + AgOPiv (Piv = pivalate).^{34–36} H/D exchange with D₂O in the position ortho to the fluorine of a fluoroarene (or a metal complex of fluoroarene) was also reported, employing Ag₂CO₃ (or Ag₂O) and a bulky phosphine (Scheme 1, eq 1).^{37–39} Similar studies including DFT calculations have been reported with AgOC(O)Ad + PPh₃ (Ad = adamantyl).²⁴


Such H/D exchange implicates the occurrence of a C–H bond activation process between a silver salt, a phosphine, and a substrate. Identification of such intermediates is important in understanding the catalytic exchange. A recent paper (published while our work was in progress) demonstrated that selective catalytic H/D exchange occurs between CH₃OD and numerous heteroarenes in the presence of silver carbonate and certain bulky phosphines. Moreover, silver carbonate reacts with ^tBuXPhos to form a dinuclear silver phosphine carbonate complex which catalyzes H/D exchange of thiophene with CH₃OD (Scheme 1, eq 2).⁴⁰ Thus, this complex is a likely intermediate in catalysis.

We now consider reactions in the presence of both silver and palladium. In the initial 2016 work, it was shown that a silver phosphine fluoroaryl complex reacts stoichiometrically with a palladium allyl complex to form the corresponding allyl arene. Notably, this silver complex was formed, not by reaction with the neutral arene but via lithium salt (Scheme 1, eq 3).²³ Nevertheless, this complex is again a plausible intermediate in catalysis. A further study focused on the reaction of palladium complexes with pentafluorobenzene in the presence of various silver salts, revealing the formation of Ag(C₆F₅) identified by its ¹⁹F ortho-fluorine resonance. The same species could be formed by reaction of Ag(OPiv) (Piv = pivalate) with C₆F₅H in the absence of palladium complex. Silver pentafluorophenyl was shown to be capable of transmetalation at palladium and catalytic H/D exchange (Scheme 1, eq 4).³⁴ Ag(OPiv) has also been employed in the catalytic carboxylation of thiophene.⁴¹

We now turn to mechanisms of cross-coupling involving silver salts and palladium precatalysts. Such reactions are exemplified by the coupling of C₆F₅H and iodotoluene in the presence of Ag₂CO₃, PPh₃, and Pd(OAc)₂ (Scheme 1, eq 5).²² In one approach, supported by computational studies, there is direct formation of heterobimetallic Ag–Pd complexes;⁴² although Ag–Pd clusters are known,⁴³ their role in C–H functionalization has not been established experimentally. In a second approach of more relevance here, silver(I) effects C–H bond activation as implicated by the ortho-selective H/D exchange mentioned above⁴⁴ and is followed by transmetalation. Several key observations support participation of silver in the second type of pathway. In addition to the evidence already cited (Scheme 1, eq 1–4), there is strong evidence from reaction kinetics, including kinetic isotope effects and rates dependent on concentrations of silver complex (Scheme 1, eq 6 and eq 7).^{33,39,24,23,45,46} These studies are also supported by H/D exchange in the presence of both silver salts and palladium salts.

Notably, the catalytic cycles proposed in these papers address the use of alkyl carboxylic acids or acidic alcohols to displace the aryl group from silver, but the mechanism of

Scheme 1. Involvement of Ag Salts in C–H Functionalization Chemistry



action of the silver carbonate was not clarified until 2021.⁴⁰ The cycles also implicate a requirement for sufficient

phosphine for coordination to both palladium and silver. In at least one proposed cycle, there is no mention that AgI may act as a sink for phosphine with potential implications for productive catalysis.²⁴ Furthermore, dynamic behavior involving phosphine ligand at both silver and palladium has not been explored, although it may have an important role in the reaction mechanism.

In this paper, we report the isolation of silver aryl and silver carbonate complexes derived directly from reaction of silver carbonate with a sterically hindered and commonly used phosphine, XPhos, and pentafluorobenzene (Scheme 1, eq 8). The aryl derivative can be formed directly or via isolation of the carbonate complex and is competent to form a cross-coupled product by reaction with a matching palladium phosphine complex. We also exemplify cross-coupling in the presence of catalytic quantities of the silver aryl complex without another silver source (Scheme 1, eq 9). These silver complexes undergo dynamic exchange with free phosphine. These studies lead to a proposal for a catalytic cycle in which the key steps have been verified stoichiometrically.

RESULTS AND DISCUSSION

An initial survey of reactions of silver carbonate with pentafluorobenzene in acetonitrile or DMF in the presence of different phosphines at 60 °C was conducted to test for C–H bond activation. With PPh₃ and P(4-OMeC₆H₄)₃, the ¹⁹F NMR spectrum showed evidence of a new peak around δ –100 consistent with the ortho fluorine of an Ag(C₆F₅) complex,⁴⁷ but the ³¹P{¹H} NMR spectrum showed a singlet with no evidence for coupling to ¹⁰⁷Ag or ¹⁰⁹Ag at room temperature. Upon cooling to –100 °C, the sample with PPh₃ showed the presence of numerous species, each observed as a pair of concentric doublets with Ag–P coupling (Figure 1a). A more electron-withdrawing phosphine, P(furyl)₃, did not react at all. The results show that use of PPh₃ as a ligand in C–H functionalization chemistry involving Ag may be complicated by the presence of multiple Ag–PPh₃ species. In contrast, the corresponding reaction with the strongly electron-donating and sterically hindered phosphine Xphos (Figure 1b) revealed a pair of doublets of triplets in the ³¹P{¹H} NMR spectrum at room temperature characteristic of a product **1** containing an Ag(C₆F₅)(Xphos) unit. We therefore selected this reaction for more detailed study. Without Xphos, only traces of an Ag(C₆F₅) product were formed.

Reaction of Ag₂CO₃ with Xphos (2 equiv) and C₆F₅H (10 equiv) at 80 °C yielded **1** with 66% yield which crystallized on cooling (Scheme 2). The ³¹P{¹H} NMR spectrum recorded in CD₂Cl₂ exhibited a pair of doublets of triplets at δ 13.0 (¹J_{Ag¹⁰⁷P} = 444, ¹J_{Ag¹⁰⁹P} = 512, and ³J_{PF} = 3.0 Hz; Figure 1b). The ¹⁹F NMR spectrum showed three resonances at δ –107.0 (m, 2F), –61.5 (t, J_{FF} = 19.8 Hz, 1F), –163.3 (m, 2F). The ³¹P–¹⁰⁹Ag HMQC spectrum revealed the ¹⁰⁹Ag chemical shift as δ 713. The LIFDI mass spectrum showed a parent radical cation at *m/z* = 752.251 (¹⁰⁷Ag isotopologue) with an isotope pattern consistent with monomeric Ag(C₆F₅)(Xphos).

The molecular structure of **1** determined by single-crystal X-ray crystallography confirmed that Ag(C₆F₅)(Xphos) is monomeric with Ag–C and Ag–P distances of 2.124(2) and 2.380(6) Å, respectively (Figure 2a). The coordination at Ag(I) is close to linear [C–Ag–P 168.86(6)°]. The distal ring of the biphenyl shows an approach between the ipso carbon C(7) and Ag [3.062(2) Å]. Such interactions are longer and hence weaker in our Ag complexes containing the Xphos





Figure 1. (a) ³¹P{¹H} NMR spectrum (202.5 MHz) at –100 °C in toluene-*d*₈: CD₂Cl₂ (20:80), following reaction of Ag₂CO₃ with pentafluorobenzene in the presence of PPh₃; (b) ³¹P{¹H} NMR spectrum at room temperature of **1** in CD₂Cl₂ obtained by reaction of Ag₂CO₃ with pentafluorobenzene in the presence of Xphos. (Cooling to –70 °C caused slight broadening but no new species were observed.)

Scheme 2. Synthesis of Ag(C₆F₅)(Xphos) **1**



ligand than in related Pd(II) complexes [C(ipso)-Pd interactions ca. 2.45 Å].⁴⁸ The C₆F₅ ring exhibits π -stacking with the adjacent molecule, with a centroid-to-centroid distance of 3.336(2) Å. Other Ag(C₆F₅)L (L = EtCN) complexes are known but have polymeric structures.⁴⁹ Hartwig et al. have isolated a monomeric silver fluoroaryl phosphine, Ag{C₆H₃(2-OMe)(4-F)}{P(o-anisyl)₃}, with a similar structure to **1**.²³ Notably, their complex was not synthesized by a C–H bond activation process but by reaction of AgBr with the corresponding lithiated arene and phosphine.

The synthesis of **1** demonstrated that silver carbonate is capable of C–H bond activation with pentafluorobenzene. We now wanted to test whether this product was competent for cross-coupling under stoichiometric conditions. Employing a literature procedure,⁵⁰ we prepared the Pd(II) complex





Figure 2. Molecular structures of (a) $\text{Ag}(\text{C}_6\text{F}_5)(\text{Xphos})$, **1**, (b) $[\text{Ag}(\text{Xphos})_2](\mu\text{-}\kappa^2\text{-CO}_3)$, **3**, (c) $[\text{Ag}(\text{Xphos})_2](\mu\text{-I})$, **4**. Hydrogen atoms are omitted; thermal ellipsoids are shown at the 50% level. Principal bond lengths and angles ($\text{\AA}/\text{deg}$): **1** $\text{Ag}(1)\text{-P}(1)$ 2.3801(5), $\text{Ag}(1)\text{-C}(34)$ 2.124(2), $\text{Ag}(1)\text{-C}(7)$ 3.062(2), $\text{C}(34)\text{-Ag}(1)\text{-P}(1)$ 168.86(6); **3** $\text{Ag}(1)\text{-P}(1)$ 2.3328(9), $\text{Ag}(1)\text{-O}(1)$ 2.249(2), $\text{Ag}(1)\text{-O}(2)$ 2.3572(2), $\text{Ag}(1)\text{-C}(34)$ 2.678(2), $\text{C}(34)\text{-O}(1)$ 1.281(3), $\text{C}(34)\text{-O}(2)$ 1.303(5), $\text{Ag}(1)\text{-C}(7)$ 3.071(3), $\text{P}(1)\text{-Ag}(1)\text{-C}(34)$ 167.29(3), $\text{P}(1)\text{-Ag}(1)\text{-O}(1)$ 150.85(7), $\text{C}(34)\text{-Ag}(1)\text{-O}(1)$ 28.48(10), $\text{O}(1)\text{-Ag}(1)\text{-O}(2)$ 57.56(8), $\text{Ag}(1)\text{-O}(2)\text{-Ag}(1')$ 178.444(13); **4** $\text{Ag}(1)\text{-I}(1)$ 2.8357(3), $\text{Ag}(1)\text{-P}(1)$ 2.4255(7), $\text{Ag}(1)\text{-C}(7)$ 3.277(3), $\text{I}(1)\text{-Ag}(1)\text{-I}(1')$ 100.057(8), $\text{P}(1)\text{-Ag}(1)\text{-I}(1)$ 128.803(18), $\text{P}(1)\text{-Ag}\text{-I}(1)\text{-I}(1')$ 151.26(3).

$\text{PdI}(\text{C}_6\text{H}_5)(\text{Xphos})$ **2** analogous to $\text{PdI}(\text{C}_6\text{H}_4\text{-F})(\text{Xphos})$ described previously [synthesized from $(\text{COD})\text{Pd}(\text{CH}_2\text{SiMe}_3)_2$]⁵¹ and reacted it stoichiometrically with $\text{Ag}(\text{C}_6\text{F}_5)(\text{Xphos})$ in CD_2Cl_2 at room temperature. Complete conversion to $\text{Ph-C}_6\text{F}_5$, as determined by NMR, occurred immediately (Scheme 3) with concomitant formation of $[\text{AgI}(\text{Xphos})]_2$ (see below).

Scheme 3. Stoichiometric Reaction of **1** and **2**



Furthermore, a mixture of **1** and **2** was prepared at $-70\text{ }^\circ\text{C}$ in CD_2Cl_2 and analyzed by NMR spectroscopy at -70 , -50 , and $-30\text{ }^\circ\text{C}$. Formation of $\text{Ph-C}_6\text{F}_5$ was observed from $-30\text{ }^\circ\text{C}$, but no transmetalation product was detected. For comparison, Hartwig et al. reacted $\text{Ag}\{\text{C}_6\text{H}_3(2\text{-OMe})(4\text{-F})\}\{\text{P}(\text{o-anisyl})_3\}$ with $\text{Pd}(\eta^3\text{-allyl-Ph})(\text{Opiv})\{\text{P}(\text{o-anisyl})_3\}$ at room temperature to obtain the cross-coupling product.²³ As a control, we attempted cross-coupling of **1** with iodotoluene but found no reaction up to $130\text{ }^\circ\text{C}$ showing the requirement for Pd.⁵² A further control reaction of $\text{C}_6\text{F}_5\text{H}$ (10 equiv) with **2** without silver complexes or base showed no reaction at room temperature.

Since silver carbonate is highly insoluble in acetonitrile, we anticipated that the first step in the reaction with pentafluorobenzene and Xphos may be formation of a silver Xphos carbonate complex. Reaction of Ag_2CO_3 with 2 equiv Xphos at $80\text{ }^\circ\text{C}$ in MeCN yielded an amber solution from which brown crystals formed (yield 24%, Scheme 4). The product was identified as $[\text{Ag}(\text{Xphos})_2](\mu\text{-}\kappa^2, \kappa^2\text{-CO}_3)$ **3** by

Scheme 4. Synthesis and Reactivity of $[\text{Ag}(\text{Xphos})_2](\mu\text{-}\kappa^2\text{-CO}_3)$ **3**





Table 1. Catalytic Reactions



variation of reaction conditions from standard above

entry	base or Ag source	ligand	Pd catalyst	temp (°C)	time (h)	NMR yield (isolated yield ^a)/%
1	standard	standard	standard	60	3	96 (84)
2	standard	standard	standard	20	23	75
3	standard	standard	no Pd(OAc) ₂	60	3	0
4	Ag ₂ CO ₃ (10 mol %)	standard	standard	60	3	21
5	Ag ₂ CO ₃ (10 mol %)	standard	standard	60	23	23
6	Ag ₂ CO ₃ (10 mol %) + K ₂ CO ₃ (0.75 equiv)	standard	standard	60	23	65
7	1 (5 mol %) + K ₂ CO ₃ (0.75 equiv)	XPhos (5 mol %)	standard	60	23	57
8	Cs ₂ CO ₃	standard	standard	60	3	<10
9	AgOAc	standard	standard	60	3	<10
10	standard	standard	intermediate Pd complex 2 (5 mol %)	20	23	63
11	standard	standard	intermediate Pd complex 2 (5 mol %)	60	3	86

^aFor methods of determining NMR yields, see the [Supporting Information](#); yield of isolated product obtained following chromatography on silica-gel.

NMR, CHN analysis, and X-ray crystallography.⁵³ The ³¹P{¹H} NMR spectrum recorded in C₆D₆ showed a pair of doublets at δ 15.9 (¹J_{Ag³¹P} 634, ¹J_{Ag¹⁰⁹P³¹} 731 Hz). The molecular structure of **3** (Figure 2b) shows a C₂-symmetric complex with an Ag(μ-CO₃)Ag unit that is almost planar with one oxygen bridging the two silver atoms [Ag(1)–O(2)–Ag(1)′ = 178.444(13)°] and the O(1) oxygens bonded to one silver each. The Ag–O(2) distances [2.3572(2) Å] are longer than the Ag–O(1) distances [2.249(2) Å]. The C(34)–O(1) and C(34)–O(2) distances are 1.281(3) and 1.303(5) Å, respectively. The sum of the angles around C(34) is 360.0(5), indicating trigonal planar coordination. The coordination at Ag is Y-shaped with O(1)–Ag–O(2) = 57.56(8)°. The ipso carbon on the distal benzene ring C(7) lies in close contact [3.071(3) Å] with the silver atom bearing the corresponding phosphorus. The CHN analysis shows one acetonitrile of crystallization per Ag, in agreement with the crystal structure. During the course of our work, Tlahuext-Aca and Hartwig reported two direct analogues of **3**, one with a P^tBuXPhos ligand which was characterized crystallographically and one with a Johnphos ligand. Their ³¹P–Ag coupling constants and metrics are similar to those of **3**.⁴⁰ The M₂(μ-κ²,κ²-CO₃) motif is quite common among first row transition metal carbonates from Mn to Zn and has also been observed for both Pd as well as Ag.^{54–61}

Complex **3** reacts with pentafluorobenzene (5 equiv wrt Ag) in C₆D₆ at room temperature to form **1** as the sole product containing the Ag(Xphos) moiety (Scheme 4). Benzene was used as the solvent since mass spectra show that **3** reacts with chlorine sources such as chloroform or dichloromethane to form [Ag(Xphos)]₂(μ-Cl)₂.

At this stage, we needed to establish Xphos as a suitable phosphine for catalytic C–H bond arylation. Pentafluorobenzene (1.5 equiv) was reacted with iodotoluene (1 equiv) with palladium(II) acetate (5 mol % wrt Pd, nitrite free Pd₃(OAc)₆),⁶² Xphos (10 mol %), and Ag₂CO₃ (0.75 equiv) in DMF at 60 °C for 3 h resulting in 96% conversion

(measured by both ¹H and ¹⁹F NMR) to 4-MeC₆H₄–C₆F₅ (Table 1, entry 1). Repetition of the reaction at room temperature resulted in 72% (¹H) and 77% (¹⁹F) conversion after 23 h (entry 2). A fully worked-up reaction gave 84% isolated yield after 3 h reaction at 60 °C and separation by column chromatography on silica gel (entry 1), whereas a control reaction without palladium(II) acetate at 60 °C gave no product (entry 3). When the quantity of Ag₂CO₃ was reduced to 0.1 equiv, the yield after 3 h was still 21%, but it barely increased after 23 h indicating that carbonate became limiting (entries 4 and 5). Use of a reduced quantity of Ag₂CO₃ (10 mol %) with K₂CO₃ (0.75 equiv) at 60 °C (23 h) restored the conversion to 65% (¹H NMR, entry 6). When a similar experiment was performed with **1** (5 mol %) as the sole source of Ag(I) and K₂CO₃ (0.75 equiv) at 60 °C (23 h), the conversion was 57% (entry 7). Replacement of Ag₂CO₃ by either Cs₂CO₃ or AgOAc gave conversions under 10% (entries 8 and 9).

These studies demonstrated that XPhos is active in catalytic cross-coupling as is **1**. Moreover, catalytic quantities of silver carbonate or **1** are sufficient in the presence of potassium carbonate. For comparison, Fagnou performed his C–H arylation reactions of pentafluorobenzene with PtBu₂Me and K₂CO₃ in DMA at 120 °C,⁹ while Zhang et al. used PPh₃ and Ag₂CO₃ in water at 70 °C for 16 h.²² As mentioned above, Hartwig et al. demonstrated that their silver complex Ag{C₆H₃(2-OMe)(4-F)}{P(o-anisyl)₃} is competent for cross-coupling in a stoichiometric reaction, but we did not find mention of its use in catalysis.²³

The reaction of **1** with **2** yielded a silver iodide complex as a byproduct identified in the ³¹P{¹H} NMR (CD₂Cl₂) spectrum by a pair of concentric doublets. The same product [Ag(XPhos)(μ-I)]₂ **4** was formed by reaction of AgI with XPhos in CD₂Cl₂ at room temperature for 24 h (³¹P{¹H} NMR δ 16.89, pair of d, ¹J_{Ag³¹P} = 542 Hz, ¹J_{Ag¹⁰⁹P³¹} = 627 Hz, ¹⁰⁹Ag NMR δ 590) (Scheme 3). The LIFDI mass spectrum of **4** dissolved in dichloromethane shows *m/z* = 1295.459 (100%)




Figure 3. Proposed catalytic mechanisms. (a) Catalytic cycle with C–H bond activation by silver. (b) Silver carbonate acting as a base in the CMD mechanism. Note that **1** may be generated by reaction of C_6F_5H with **3** (route A) or by reaction with **4** (route B).

with isotope pattern consistent with two silver atoms (quoted for $^{107}Ag^{109}Ag$) and a much smaller peak at $m/z = 710.183$ (12%) for the monomer (quoted for ^{107}Ag). This spectrum suggests that this complex exists as a dimer in solution or as an equilibrium between monomer and dimer. The molecular structure determined crystallographically reveals a C_2 -symmetric dimer with bridging iodides. The central $Ag_2(\mu-I)_2$ unit is planar, but the phosphorus atoms lie out of that plane with a $P(1)-Ag-I(1)-I(1')$ torsional angle of $151.26(3)^\circ$. Similar interactions between the distal benzene ring C(7) and the Ag atom [$3.277(3)$ Å] are found to those in complexes **1** and **3** (Figure 2c).

Reaction of **4** with excess Ag_2CO_3 (10 equiv) and a large excess of pentafluorobenzene (100 equiv) generates **1** and precipitates silver iodide (eq 1). As controls, we attempted to react C_6F_5H with **4** but found no reaction in either DMF at $60^\circ C$ or CH_2Cl_2 at room temperature. We also showed that **4** does not react with Ag_2CO_3 in DMF at $60^\circ C$.


In addition, we investigated the acetate complex, $Ag(OAc)(XPhos)$ **5**, formed by reaction of XPhos with $AgOAc$ in CH_3CN at $80^\circ C$. Complex **5** exhibits a characteristic $^{31}P\{^1H\}$ NMR resonance δ 16.55 (pair of d, $^1J_{^{107}Ag}^{31P} = 643.9$ Hz, $^1J_{^{109}Ag}^{31P} = 743.2$ Hz). The crystal structure of this complex revealed a monomer with κ^1 -coordination of the acetate ligand (see the Supporting Information). In contrast to the behavior of **3**, the reaction of $AgOAc$ in the presence of XPhos (1 equiv) and C_6F_5H (10 equiv) did not produce any **1** (none detected by $^{31}P\{^1H\}$ or $^{19}F\{^1H\}$ NMR spectroscopy) but generated **5** instead.


Further information was obtained from kinetic isotope effects measured both for the stoichiometric and for the catalytic reaction. For the stoichiometric reaction of **3** with C_6F_5H (C_6F_5D), we employed a low temperature in order to

ensure a reasonable number of data points for the reaction with C_6F_5H which was monitored in situ by NMR. A temperature of $-40^\circ C$ in toluene proved optimal, but, under these conditions, the reaction did not proceed to completion. A fast process lasting ca. 20 min was recorded, followed by a much slower process with KIEs were determined as 3.7 ± 0.3 and 0.91 ± 0.01 , respectively, for the fast and slow processes. The KIE for the fast process is consistent with rate-determining C–H bond breaking in this reaction of this silver–carbonate complex. The product of reaction was $4-MeC_6H_4-C_6F_5$ as usual, but nature of the slow process has not been determined.

In separate experiments, catalytic reactions were conducted under the standard conditions (see Table 1) at $60^\circ C$, and the products were analyzed by NMR at different reaction times. The KIE for the catalytic reaction was determined from the initial rates of reaction as 4.5 ± 0.8 .




The recycling of XPhos was clarified by synthesizing $Ag(C_6F_5)$ independently.⁴⁹ With the authentic sample of $Ag(C_6F_5)$, we confirmed that this species is formed in low yield (ca. 10%) upon reaction of Ag_2CO_3 with pentafluorobenzene in the absence of XPhos. Additionally, as expected, $Ag(C_6F_5)$ reacts rapidly with XPhos at room temperature to form **1**. Of particular interest is the observation that $Ag(C_6F_5)$ reacts with **4** to form **1** and a precipitate of AgI (eq 2). The reactions of eq 1 and eq 2 provide opportunities to recycle phosphine in the catalytic reactions and explain how AgI may be formed as a final product. The reactions of $Ag(C_6F_5)$ in CD_3CN and **1** in CD_2Cl_2 with D_2O at room temperature both give C_6F_5D ; however, the reaction of **1** takes several days.





The observations described above lead us to postulate the catalytic cycle in Figure 3a in which Ag performs C–H bond activation and Pd effects cross-coupling.⁶³ According to this cycle, PdI(C₆H₅)(XPhos) (**2**, 5 mol %) should act as a catalyst for the coupling of 4-iodotoluene with pentafluorobenzene in the presence of 5 mol % XPhos and Ag₂CO₃ (0.75 equiv). On testing this hypothesis at room temperature, we obtained yields of tol–C₆F₅ after 3 and 23 h of 6% and 63%, respectively; at 60 °C, they increase to 86% and 100%, respectively (Table 1, entries 10, 11). As expected, Ph–C₆F₅ was also formed with yield 5% wrt tol–C₆F₅. This cycle is also supported by the catalytic reaction with 5 mol % **1** as the source of Ag(I) (entry 7). All the steps in the catalytic cycle have been verified stoichiometrically other than the formation and reaction of a Pd(0) species which is not expected to be detectable under our conditions (oxidative addition being fast). The cycle shows two alternative routes from Ag₂CO₃ to form **1** reflecting the experiments described above. The experiments with reduced Ag₂CO₃ with and without K₂CO₃ confirm that carbonate is required for the C–H bond activation acting as the Brønsted base. The postulated alternative mechanism invoking Ag(I) as a base and Pd as involved in a CMD process is expected to involve Ag–carbonato–Pd **6** as an intermediate (Figure 3b).

A further group of experiments illuminates the ability of XPhos to undergo exchange, thereby ensuring that its supply is not exhausted prematurely. To test the lability of the silver complexes, **1**, **3**, and **4**, we reacted each of them with *t*BuXPhos, in which the isopropyl groups are replaced by *t*butyl groups. Addition of *t*BuXPhos (1 equiv) to **1** resulted in immediate formation of a new silver phosphine species at room temperature (³¹P NMR δ 38.5, pair of dt, ¹J_{Ag³¹P} = 448 Hz, ¹J_{Ag³¹P} = 516 Hz, and J_{PF} = 2.7 Hz, compared to free ligand at δ 21.4) in 50% conversion consistent with exchange of the phosphine to form the *t*BuXPhos analogue Ag(C₆F₅)-(*t*BuXPhos) **1'**. Assuming this is an equilibrium reaction, we deduce an equilibrium constant of 1.03 ± 0.05 for formation of **1'** (see eq 3). Similar behavior was observed for **3** and **4** (eqs 4 and 5).⁶⁴



4 and **4'** assumed to be monomers in solution for calcn of K_{eq}

The experiments with *t*BuXPhos encouraged us to test for exchange between Ag(XPhos) complexes and free XPhos. A ³¹P–³¹P EXSY spectrum of a sample of **1** in the presence of XPhos (5 equiv, mixing time 2 s) revealed cross peaks consistent with intermolecular exchange between **1** and XPhos on the time scale of seconds. Since there is exchange between two sites only, 1D-EXSY⁶⁵ is suitable and is more economical in instrument time. 1D-EXSY experiments on **1** in the presence of a range of concentrations of free XPhos were conducted in CD₂Cl₂ yielding evidence that the rate of exchange V_{ex} depends linearly on [XPhos] (where $V_{\text{ex}} = k_{\text{obs}}[\text{XPhos}][\mathbf{1}]/$

([XPhos] + [**1**]) and k_{obs} is derived from the variation of fractional integration of [XPhos] with mixing time τ_{m} on selective excitation of **1**).^{66,67} Variable temperature 1D-EXSY experiments with excess XPhos (5 equiv) were performed in C₆D₆ in order to access a suitable temperature range to determine of activation parameters yielding $\Delta H^\ddagger = 40.0 \pm 1.2$ kJ/mol and $\Delta S^\ddagger = -162 \pm 4$ J/K mol, providing clear evidence for associative exchange.

CONCLUSION

In conclusion, silver carbonate is capable of C–H bond activation of pentafluorobenzene in the presence of XPhos yielding discrete and well-defined Ag(C₆F₅)(XPhos) **1**. This reaction can also be carried out stepwise yielding first the carbonate complex [Ag(XPhos)]₂(μ-κ²,κ²-CO₃) **3** and then **1**. A third silver complex has been isolated by reaction of AgI with XPhos, [Ag(XPhos)(μ-I)]₂ **4**. All three silver complexes have been characterized in solution and crystallographically. They are all labile with respect to associative exchange with free XPhos or with *t*BuXPhos and all participate in cross-coupling. The viability of **1** for cross-coupling has been demonstrated stoichiometrically by reaction of **1** with PdI(C₆H₅)(XPhos) **2** yielding **4** as a byproduct; this reaction even occurs at –30 °C. Catalytic cross-coupling of 4-iodotoluene to pentafluorobenzene may be achieved in the presence of palladium(II) acetate, XPhos, and silver carbonate at 60 °C or at room temperature. Catalytic cross-coupling also occurs with **2** as the catalyst in the presence of Ag₂CO₃ and a reduced quantity of XPhos. Catalytic quantities of silver in the form of Ag₂CO₃ or **1** are active in conjunction with potassium carbonate as the stoichiometric base. A catalytic cycle involving C–H bond activation at silver has been proposed consistent with the set of tested stoichiometric reactions. It is now clear how the phosphine solubilizes silver carbonate and silver iodide; however, the phosphine is labile enough to allow precipitation of silver iodide as the final byproduct of catalysis. We stress that the study does not exclude competing C–H bond activation steps at palladium or a role for higher-order catalyst species. Nevertheless, the evidence for the C–H bond activation abilities of silver carbonate in the presence of a suitable phosphine is now conclusive and is shown to involve well-defined, soluble, mononuclear, and dinuclear silver phosphine complexes that are active in cross-coupling reactions with palladium complexes.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.organomet.2c00063>.

Synthetic procedures, catalytic reactions, stoichiometric reactions, phosphine-exchange experiments, EXSY experiments, kinetic isotope effect experiments, crystallographic data, characterization data including NMR spectra (PDF)

Accession Codes

CCDC 2110192– 2110194 and 2160367 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: + 44 1223 336033.

AUTHOR INFORMATION

Corresponding Authors

Ian J. S. Fairlamb – Department of Chemistry, University of York, York YO10 5DD, U.K.; orcid.org/0000-0002-7555-2761; Email: ian.fairlamb@york.ac.uk

Robin N. Perutz – Department of Chemistry, University of York, York YO10 5DD, U.K.; orcid.org/0000-0001-6286-0282; Email: robin.perutz@york.ac.uk

Authors

Gayathri Athavan – Department of Chemistry, University of York, York YO10 5DD, U.K.

Theo F. N. Tanner – Department of Chemistry, University of York, York YO10 5DD, U.K.; orcid.org/0000-0001-7563-9325

Adrian C. Whitwood – Department of Chemistry, University of York, York YO10 5DD, U.K.; orcid.org/0000-0002-5132-5468

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.organomet.2c00063>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We are grateful to the Wild Fund of the University of York for a scholarship for G.A., to the University of York for funding a Ph.D. studentship for TFNT and laboratory equipment, and to Professor Simon Duckett for recording the ^{31}P - ^{109}Ag NMR spectra.

DEDICATION

We dedicate this paper to Professor Maurice Brookhart in admiration for his achievements in organometallic chemistry.

REFERENCES

- (1) Baudoin, O. Multiple Catalytic C-H Bond Functionalization for Natural Product Synthesis. *Angew. Chem. Int. Ed* **2020**, *59*, 17798–17809.
- (2) Yu, C. J.; Sanjose-Orduna, J.; Patureau, F. W. W.; Perez-Temprano, M. H. Emerging unconventional organic solvents for C-H bond and related functionalization reactions. *Chem. Soc. Rev.* **2020**, *49*, 1643–1652.
- (3) Rej, S.; Ano, Y.; Chatani, N. Bidentate Directing Groups: An Efficient Tool in C-H Bond Functionalization Chemistry for the Expedient Construction of C-C Bonds. *Chem. Rev.* **2020**, *120*, 1788–1887.
- (4) Mihai, M. T.; Genov, G. R.; Phipps, R. J. Access to the meta position of arenes through transition metal catalysed C-H bond functionalisation: a focus on metals other than palladium. *Chem. Soc. Rev.* **2018**, *47*, 149–171.
- (5) Ping, L.; Chung, D. S.; Bouffard, J.; Lee, S. G. Transition metal-catalyzed site- and regio-divergent C-H bond functionalization. *Chem. Soc. Rev.* **2017**, *46*, 4299–4328.
- (6) Ackermann, L. Carboxylate-Assisted Transition-Metal-Catalyzed C-H Bond Functionalizations: Mechanism and Scope. *Chem. Rev.* **2011**, *111*, 1315–1345.
- (7) Carrow, B. P.; Sampson, J.; Wang, L. Base-Assisted C-H Bond Cleavage in Cross-Coupling: Recent Insights into Mechanism, Speciation, and Cooperativity. *Isr. J. Chem.* **2020**, *60*, 230–258.
- (8) Eisenstein, O.; Milani, J.; Perutz, R. N. Selectivity of C-H Activation and Competition between C-H and C-F Bond Activation at Fluorocarbons. *Chem. Rev.* **2017**, *117*, 8710–8753.
- (9) Lafrance, M.; Rowley, C. N.; Woo, T. K.; Fagnou, K. Catalytic intermolecular direct arylation of perfluorobenzenes. *J. Am. Chem. Soc.* **2006**, *128*, 8754–8756.
- (10) Cao, M. N.; Wu, D. S.; Su, W. P.; Cao, R. Palladium nanocrystals stabilized by cucurbit[6]uril as efficient heterogeneous catalyst for direct C-H functionalization of polyfluoroarenes. *J. Catal.* **2015**, *321*, 62–69.
- (11) Ma, X. W.; Liu, Y.; Liu, P.; Xie, J. W.; Dai, B.; Liu, Z. Y. Palladium-catalyzed direct arylation of polyfluoroarene and facile synthesis of liquid crystal compounds. *Appl. Organomet. Chem.* **2014**, *28*, 180–185.
- (12) Garcia-Cuadrado, D.; Braga, A. A. C.; Maseras, F.; Echavarren, A. M. Proton abstraction mechanism for the palladium-catalyzed intramolecular arylation. *J. Am. Chem. Soc.* **2006**, *128*, 1066–1067.
- (13) Gorelsky, S. I.; Lapointe, D.; Fagnou, K. Analysis of the concerted metalation-deprotonation mechanism in palladium-catalyzed direct arylation across a broad range of aromatic substrates. *J. Am. Chem. Soc.* **2008**, *130*, 10848–10849.
- (14) Lafrance, M.; Lapointe, D.; Fagnou, K. Mild and efficient palladium-catalyzed intramolecular direct arylation reactions. *Tetrahedron* **2008**, *64*, 6015–6020.
- (15) Pascual, S.; de Mendoza, P.; Braga, A. A. C.; Maseras, F.; Echavarren, A. M. Bidentate phosphines as ligands in the palladium-catalyzed intramolecular arylation: the intermolecular base-assisted proton abstraction mechanism. *Tetrahedron* **2008**, *64*, 6021–6029.
- (16) Davies, D. L.; Donald, S. M. A.; Macgregor, S. A. Computational study of the mechanism of cyclometalation by palladium acetate. *J. Am. Chem. Soc.* **2005**, *127*, 13754–13755.
- (17) Davies, D. L.; Macgregor, S. A.; McMullin, C. L. Computational Studies of Carboxylate-Assisted C-H Activation and Functionalization at Group 8–10 Transition Metal Centers. *Chem. Rev.* **2017**, *117*, 8649–8709.
- (18) Pla, D.; Gomez, M. Metal and metal oxide nanoparticles: A lever for C-H functionalization. *ACS Catal.* **2016**, *6*, 3537–3552.
- (19) Reay, A. J.; Fairlamb, I. J. S. Catalytic C-H bond functionalisation chemistry: the case for quasi-heterogeneous catalysis. *Chem. Commun.* **2015**, *51*, 16289–16307.
- (20) Fairlamb, I. J. S.; Scott, N. W. J. Pd Nanoparticles in C-H Activation and Cross-coupling Catalysis. *Top. Organometal. Chem.* **2020**, *66*, 171–205.
- (21) Reay, A. J.; Neumann, L. K.; Fairlamb, I. J. S. Catalyst efficacy of homogeneous and heterogeneous palladium catalysts in the direct arylation of common heterocycles. *Synlett* **2016**, *27*, 1211–1216.
- (22) Chen, F.; Min, Q. Q.; Zhang, X. G. Pd-Catalyzed direct arylation of polyfluoroarenes on water under mild conditions using PPh_3 ligand. *J. Org. Chem.* **2012**, *77*, 2992–2998.
- (23) Lee, S. Y.; Hartwig, J. F. Palladium-Catalyzed, Site-Selective Direct Allylation of Aryl C-H Bonds by Silver-Mediated C-H Activation: A Synthetic and Mechanistic Investigation. *J. Am. Chem. Soc.* **2016**, *138*, 15278–15284.
- (24) Whitaker, D.; Bures, J.; Larrosa, I. Ag(I)-Catalyzed C-H Activation: The Role of the Ag(I) Salt in Pd/Ag-Mediated C-H Arylation of Electron-Deficient Arenes. *J. Am. Chem. Soc.* **2016**, *138*, 8384–8387.
- (25) Bowmaker, G. A.; Effendy, Hanna, J. V.; Healy, P. C.; King, S. P.; Pettinari, C.; Skelton, B. W.; White, A. H. Solution and mechanochemical syntheses, and spectroscopic and structural studies in the silver(I) (bi-)carbonate: triphenylphosphine system. *Dalton Trans* **2011**, *40*, 7210–7218.
- (26) Goel, R. G.; Pilon, P. Tri-Tert-Butylphosphine Complexes of Silver(I) - Preparation, Characterization, and Spectral Studies. *Inorg. Chem.* **1978**, *17*, 2876–2879.
- (27) Muettterties, E. L.; Alegranti, C. W. Solution Structure and Kinetic Study of Metal-Phosphine and Metal-Phosphite Complexes 0.1. Silver(I) System. *J. Am. Chem. Soc.* **1972**, *94*, 6386–6391.
- (28) Bachman, R. E.; Andretta, D. F. Metal-ligand bonding in coinage metal-phosphine complexes: The synthesis and structure of some low-coordinate silver(I)-phosphine complexes. *Inorg. Chem.* **1998**, *37*, 5657–5663.


- (29) Patmore, N. J.; Hague, C.; Cotgreave, J. H.; Mahon, M. F.; Frost, C. G.; Weller, A. S. Silver phosphanes partnered with carborane monoanions: Synthesis, structures and use as highly active lewis acid catalysts in a hetero-Diels-Alder reaction. *Chem. - Eur. J.* **2002**, *8*, 2088–2098.
- (30) Keller, S.; Camenzind, T. N.; Abraham, J.; Prescimone, A.; Haussinger, D.; Constable, E. C.; Housecroft, C. E. Self-assembly of heteroleptic dinuclear silver(I) complexes bridged by bis-(diphenylphosphino)ethyne. *Dalton Trans* **2018**, *47*, 946–957.
- (31) Bay, K. L.; Yang, Y. F.; Houk, K. N. Multiple roles of silver salts in palladium-catalyzed C-H activations. *J. Organomet. Chem.* **2018**, *864*, 19–25.
- (32) Mudarra, A. L.; Martinez de Salinas, S.; Perez-Temprano, M. H. Beyond the traditional roles of Ag in catalysis: the transmetalating ability of organosilver(I) species in Pd-catalysed reactions. *Org. Biomol. Chem.* **2019**, *17*, 1655–1667.
- (33) Panigrahi, A.; Whitaker, D.; Vitorica-Yrezabal, I. J.; Larrosa, I. Ag/Pd Cocatalyzed Direct Arylation of Fluoroarene Derivatives with Aryl Bromides. *ACS Catal.* **2020**, *10*, 2100–2107.
- (34) Lotz, M. D.; Camasso, N. M.; Canty, A. J.; Sanford, M. S. Role of Silver Salts in Palladium-Catalyzed Arene and Heteroarene C-H Functionalization Reactions. *Organometallics* **2017**, *36*, 165–171.
- (35) Shimoyama, Y.; Kuwabara, J.; Kanbara, T. Mechanistic Study of Pd/Ag Dual-Catalyzed Cross-Dehydrogenative Coupling of Perfluoroarenes with Thiophenes. *ACS Catal.* **2020**, *10*, 3390–3397.
- (36) Li, W. P.; Yuan, D. D.; Wang, G. Q.; Zhao, Y.; Xie, J.; Li, S. H.; Zhu, C. J. Cooperative Au/Ag Dual-Catalyzed Cross-Dehydrogenative Biaryl Coupling: Reaction Development and Mechanistic Insight. *J. Am. Chem. Soc.* **2019**, *141*, 3187–3197.
- (37) Hu, G. Q.; Li, E. C.; Zhang, H. H.; Huang, W. Ag(I)-Mediated hydrogen isotope exchange of mono-fluorinated (hetero)arenes. *Org. Biomol. Chem.* **2020**, *18*, 6627–6633.
- (38) Wilkinson, L. A.; Pike, J. A.; Walton, J. W. C-H Activation of pi-Arene Ruthenium Complexes. *Organometallics* **2017**, *36*, 4376–4381.
- (39) Tlahuext-Aca, A.; Lee, S. Y.; Sakamoto, S.; Hartwig, J. F. Direct Arylation of Simple Arenes with Aryl Bromides by Synergistic Silver and Palladium Catalysis. *ACS Catal.* **2021**, *11*, 1430–1434.
- (40) Tlahuext-Aca, A.; Hartwig, J. F. Site-Selective Silver-Catalyzed C-H Bond Deuteration of Five-Membered Aromatic Heterocycles and Pharmaceuticals. *ACS Catal.* **2021**, *11*, 1119–1127.
- (41) Lee, M.; Hwang, Y. K.; Kwak, J. Ag(I)-Catalyzed C-H Carboxylation of Thiophene Derivatives. *Organometallics* **2021**, *40*, 3136–3144.
- (42) Bhaskararao, B.; Singh, S.; Anand, M.; Verma, P.; Prakash, P.; Athira, C.; Malakar, S.; Schaefer, H. F.; Sunoj, R. B. Is silver a mere terminal oxidant in palladium catalyzed C-H bond activation reactions? *Chem. Sci.* **2020**, *11*, 208–216.
- (43) Kozitsyna, N. Y.; Nefedov, S. E.; Klyagina, A. P.; Markov, A. A.; Dobrokhotova, Z. V.; Velikodny, Y. A.; Kochubey, D. I.; Zyubina, T. S.; Gekhman, A. E.; Vargaftik, M. N.; Moiseev, I. I. Novel heterometallic palladium-silver complex. *Inorg. Chim. Acta* **2011**, *370*, 382–387.
- (44) Clot, E.; Megret, C.; Eisenstein, O.; Perutz, R. N. Exceptional Sensitivity of Metal-Aryl Bond Energies to ortho-Fluorine Substituents: Influence of the Metal, the Coordination Sphere, and the Spectator Ligands on M-C/H-C Bond Energy Correlations. *J. Am. Chem. Soc.* **2009**, *131*, 7817–7827.
- (45) Colletto, C.; Panigrahi, A.; Fernandez-Casado, J.; Larrosa, I. Ag(I)-C-H Activation Enables Near-Room-Temperature Direct alpha-Arylation of Benzo b thiophenes. *J. Am. Chem. Soc.* **2018**, *140*, 9638–9643.
- (46) Batuecas, M.; Luo, J. F.; Gergelitsova, I.; Kramer, K.; Whitaker, D.; Vitorica-Yrezabal, I. J.; Larrosa, I. Catalytic Asymmetric C-H Arylation of (eta(6)-Arene)Chromium Complexes: Facile Access to Planar-Chiral Phosphines. *ACS Catal.* **2019**, *9*, 5268–5278.
- (47) Belt, S. T.; Helliwell, M.; Jones, W. D.; Partridge, M. G.; Perutz, R. N. Eta(2)-Coordination and C-F Activation of Hexafluorobenzene by Cyclopentadienylrhodium and Cyclopentadienyliridium Complexes. *J. Am. Chem. Soc.* **1993**, *115*, 1429–1440.
- (48) Bruno, N. C.; Tudge, M. T.; Buchwald, S. L. Design and preparation of new palladium precatalysts for C-C and C-N cross-coupling reactions. *Chem. Sci.* **2013**, *4*, 916–920.
- (49) Tyrre, W.; Wickleder, M. S. Silver compounds in synthetic chemistry. 1 - A facile preparative route for pentafluorophenylsilver, AgC₆F₅ and its use as an oxidative pentafluorophenyl group transfer reagent in reactions with group 12 to 16 elements - the single crystal structure of AgC₆F₅. EtCN, the first arylsilver derivative crystallising in infinite chains. *Z. Anorg. Allg. Chem.* **2002**, *628*, 1841–1847.
- (50) Milner, P. J.; Maimone, T. J.; Su, M. J.; Chen, J. H.; Muller, P.; Buchwald, S. L. Investigating the Dearomative Rearrangement of Biaryl Phosphine-Ligated Pd(II) Complexes. *J. Am. Chem. Soc.* **2012**, *134*, 19922–19934.
- (51) Wagschal, S.; Perego, L. A.; Simon, A.; Franco-Espejo, A.; Tocqueville, C.; Albanese-Walker, J.; Jutand, A.; Grimaud, L. Formation of XPhos-Ligated Palladium(0) Complexes and Reactivity in Oxidative Additions. *Chem. - Eur. J.* **2019**, *25*, 6980–6987.
- (52) Ag(C₆F₅) reacts with iodobenzene at 110 °C. Sun, K. K.; Miller, W. T. Perfluorophenylsilver. *J. Am. Chem. Soc.* **1970**, *92*, 6985–6987.
- (53) Soltys-Brzostek, K.; Terlecki, M.; Sokolowski, K.; Lewinski, J. Chemical fixation and conversion of CO₂ into cyclic and cage-type metal carbonates. *Coord. Chem. Rev.* **2017**, *334*, 199–231.
- (54) Sattler, W.; Parkin, G. Structural characterization of zinc bicarbonate compounds relevant to the mechanism of action of carbonic anhydrase. *Chem. Sci.* **2012**, *3*, 2015–2019.
- (55) Dietrich, J.; Heinemann, F. W.; Schrodt, A.; Schindler, S. A new carbonate bridged dinuclear zinc complex with tripodal amine ligands. *Inorg. Chim. Acta* **1999**, *288*, 206–209.
- (56) Shopov, D. Y.; Sharninghausen, L. S.; Sinha, S. B.; Borowski, J. E.; Mercado, B. Q.; Brudvig, G. W.; Crabtree, R. H. Synthesis of pyridine-alkoxide ligands for formation of polynuclear complexes. *New J. Chem.* **2017**, *41*, 6709–6719.
- (57) Roy, L.; Al-Afyouni, M. H.; DeRoshia, D. E.; Mondal, B.; DiMucci, I. M.; Lancaster, K. M.; Shearer, J.; Bill, E.; Brennessel, W. W.; Neese, F.; Ye, S. F.; Holland, P. L. Reduction of CO₂ by a masked two-coordinate cobalt(I) complex and characterization of a proposed oxodicobalt(II) intermediate. *Chem. Sci.* **2019**, *10*, 918–929.
- (58) Stewart, J. A.; McKeown, P.; Driscoll, O. J.; Mahon, M. F.; Ward, B. D.; Jones, M. D. Tuning the Thiolen: Al(III) and Fe(III) Thiolen Complexes for the Ioselective ROP of rac-Lactide. *Macromolecules* **2019**, *52*, 5977–5984.
- (59) Kwak, C. H.; Kim, J. Preparation, characterization and X-ray structure of {[(3,2,3-tet)Ni(mu-CO₃)Ni(3,2,3-tet)] [(3,2,3-tet)Cu]}-(ClO₄)₄: a heteronuclear trimetallic complex containing hydrogen bonds. *Inorg. Chem. Commun.* **2000**, *3*, 550–552.
- (60) Avdeeva, V. V.; Dziova, A. E.; Polyakova, I. N.; Malinina, E. A.; Goeva, L. V.; Kuznetsov, N. T. Copper(I), copper(II), and heterovalent copper(I,II) complexes with 1,10-phenanthroline and the closo-decaborate anion. *Inorg. Chim. Acta* **2015**, *430*, 74–81.
- (61) Muller, A.; Holzapfel, C. W. mu-carbonato bis{2'-di-tert-butylphosphanyl)biphenyl-2-yl}palladium(II) dichloromethane. *Acta Crystallogr. Sect. E: Crystallogr. Commun.* **2012**, *68*, m1574–m1575.
- (62) Bajwa, S. E.; Storr, T. E.; Hatcher, L. E.; Williams, T. J.; Baumann, C. G.; Whitwood, A. C.; Allan, D. R.; Teat, S. J.; Raitby, P. R.; Fairlamb, I. J. S. On the appearance of nitrite anion in [PdX(OAc)L₂] and [Pd(X)(C[^]N)L] syntheses (X = OAc or NO₂): photocrystallographic identification of metastable Pd(eta(1)-ONO)-(C[^]N)PPh₃. *Chem. Sci.* **2012**, *3*, 1656–1661.
- (63) Interactions between arenes and silver(I) are well-known and probably precede C-H activation. Lindeman, S.; Rahore, R.; Kochi, J. K. Silver(I) Complexation of (Poly)aromatic Ligands. Structural Criteria for Depth Penetration into cis-Stilbenoid Cavities. *Inorg. Chem.* **2000**, *39*, 5707–5716.
- (64) **3** was treated as dinuclear, but **4** was considered mononuclear for calculation of equilibrium constants. Since **3** and **4** are dinuclear, we might expect separate resonances for the complex with one XPhos and one tBuXPhos, but they were not observed. At low temperature,

the resonances of **1** broadened while those of **1'** remained sharp, consistent with fluxionality.

(65) Perrin, C. L.; Engler, R. E. Accurate rate constants for chemical exchange from improved weighted linear-least-squares analysis of multiple 1D-EXSY data. *J. Magn. Reson., Ser. A* **1996**, *123*, 188–195.

(66) Wilkins, R. G. *Kinetics and mechanism of reactions of transition metal complexes*, 2nd ed.; VCH: Weinheim, 1992.


(67) McKay, H. A. C. Kinetics of some exchange reactions of the type $RI + I^{*-}$ reversible arrow $RI^* + I^-$ in alcoholic solution. *J. Am. Chem. Soc.* **1943**, *65*, 702–706.




ACS
ENGINEERING Au
AN OPEN ACCESS JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

Editor-in-Chief: **Prof. Shelley D. Minteer**, University of Utah, USA

Deputy Editor:
Prof. Vivek Ranade
University of Limerick, Ireland

Open for Submissions 

pubs.acs.org/engineeringau  ACS Publications
Most Trusted. Most Cited. Most Read.