



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/187/>

Article:

Roberts, H, Liabo, K, Lucas, P et al. (2004) Mentoring to reduce antisocial behaviour in childhood. British medical journal. pp. 512-514. ISSN: 1756-1833

<https://doi.org/10.1136/bmj.328.7438.512>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Mentoring to reduce antisocial behaviour in childhood

Helen Roberts, Kristin Liabo, Patricia Lucas, David DuBois and Trevor A Sheldon

BMJ 2004;328;512-514
doi:10.1136/bmj.328.7438.512

Updated information and services can be found at:
<http://bmj.com/cgi/content/full/328/7438/512>

These include:

Data supplement

"Further details"
<http://bmj.com/cgi/content/full/328/7438/512/DC1>

References

This article cites 15 articles, 4 of which can be accessed free at:
<http://bmj.com/cgi/content/full/328/7438/512#BIBL>

1 online articles that cite this article can be accessed at:
<http://bmj.com/cgi/content/full/328/7438/512#otherarticles>

Rapid responses

4 rapid responses have been posted to this article, which you can access for free at:
<http://bmj.com/cgi/content/full/328/7438/512#responses>

You can respond to this article at:
<http://bmj.com/cgi/eletter-submit/328/7438/512>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

- [Sociology](#) (272 articles)
 - [Health education \(including prevention and promotion\)](#) (567 articles)
 - [Children](#) (1116 articles)
 - [Other evidence based practice](#) (344 articles)
-

Notes

To order reprints of this article go to:
<http://bmj.bmjournals.com/cgi/reprintform>

To subscribe to *BMJ* go to:
<http://www.bmjournals.com/subscriptions>

Education and debate

Mentoring to reduce antisocial behaviour in childhood

Helen Roberts, Kristin Liabo, Patricia Lucas, David DuBois, Trevor A Sheldon

The effects of social interventions need to be examined in real life situations as well as studies

Child Health
Research and Policy
Unit, Institute of
Health Sciences,
City University,
London
EC1A 7QN

Helen Roberts
professor

Kristin Liabo
research fellow
Patricia Lucas
research officer

Community Health
Sciences, School of
Public Health
(MC 923),
University of Illinois
at Chicago,
1603 W Taylor St,
Chicago,
IL 60612-4324,
USA

David DuBois
associate professor

Department of
Health Sciences,
University of York,
York YO10 5DD
Trevor A Sheldon
professor

Correspondence to:
H Roberts
h.roberts@city.ac.uk

BMJ 2004;328:512-4

Politicians and policy makers are increasingly interested in evidence based decision making. They are under pressure to look to research for solutions to policy problems and justify programmes by reference to the knowledge base. It is tempting for policy makers to grasp any research on seemingly intractable social problems, however slim, in the hope of finding simple solutions. Rolling out national programmes based on inadequate evidence can, however, do more harm than good. We use the example of mentoring for young people with, or at risk of, antisocial behaviour problems to show the potential dangers of running ahead of the evidence.

Social interventions aimed at children

Public health interventions to improve outcomes for children are an example of policy and research evolving in tandem.¹⁻³ Interventions such as Highscope,

Headstart, parenting education, home visiting, and mentoring have been well designed and robustly evaluated, some of them by randomised controlled trials.⁴⁻¹⁰ Nevertheless, parent education, home visiting, and mentoring, as their proponents and evaluators would be the first to agree, largely remain black boxes with a great many unanswered questions about what specific forms of intervention are effective and under which conditions.

Meanwhile, a climate has been created in which it is widely held that these interventions are effective and national programmes are being established. Questions about who delivers the service, the kind of young people who might benefit, and the content of services likely to be effective can be lost in the drive to get the programmes running.¹¹ These programmes can gain momentum because they have strong face validity: they look like the sort of things that should work, our instincts tell us that they will work, and we want them to work. This can not only result in premature roll out on the basis of insufficient evidence but also make it difficult to stop or change direction.

Mentoring for antisocial behaviour in childhood

Antisocial behaviour in childhood and adolescence is a problem for young people and their families, for health and welfare professionals planning multidisciplinary services, and for general practitioners approached by fraught parents.¹²⁻¹³ Behaviour problems in childhood can presage more serious problems in later life.¹⁴ Antisocial behaviour in young people is also a problem for the police, communities, and politicians. This makes finding a solution a political as well as a therapeutic imperative—a potent driver for action.

One approach is through mentoring schemes, which a meta-analysis has shown to have benefits.¹⁰ In a typical non-directive mentoring programme, a mentor will be a volunteer who provides support or guidance to someone younger or less experienced. The mentor aims to offer support, understanding, experience, and advice. Mentoring is non-invasive and does not require drug treatment. It is easy to see why it might work, and why it is attractive to politicians and policy makers.



ANDREW DUNSMORE/REX

Not all mentoring to reduce antisocial behaviour is effective



Further details of the studies are on bmj.com

In February 2003, Lord Filkin, then a minister in the Home Office, announced £850 000 (\$1.6m; €1.26m) of funding for mentoring schemes in England: "Mentors can make a real difference to ... some of the most vulnerable people ... and help to make our society more inclusive. There are ... excellent examples of schemes which really work."¹⁵

There is equal enthusiasm in the United States. In his state of the union address in January 2003, President Bush announced plans for a \$450m (£238m; €352m) initiative to expand the availability of mentoring programmes for young people. This included \$300m for mentoring at risk pupils and \$150m to provide mentors to children of prisoners.¹⁶

What does the research show?

One problem with interventions that become politically attractive, and to which large sums are attached, is that research may be used for support rather than illumination. Robust research does indicate benefits from mentoring for some young people, for some programmes, in some circumstances, in relation to some outcomes.⁸⁻¹⁰ But there are also good descriptive evaluations suggesting that those young people who stay on in programmes are inclined to report favourably on the experience.^{17, 18}

We examined the existing reviews (see bmj.com) and concluded that research on mentoring programmes does not provide evidence of measurable gains in outcomes such as truancy or other antisocial behaviours.¹⁹ When improvements have been reported, such as in reports of the Big Brothers Big Sisters mentoring programme,⁸ critical examination suggests flaws that weaken the conclusions.⁸ Although this study was well conducted, it failed to report on changes observed from administrative records. A comprehensive meta-analysis on mentoring found no effect in studies using more objective measuring tools than self report and non-blinded reports of behaviour change.¹⁰

Mentoring programmes for vulnerable young people may have a negative impact, and adverse effects associated with breakdowns of relationships with mentors have been reported.²⁰ Worryingly, a three year follow up study of one well designed scheme found that a subgroup of mentored young people, some of whom had previously been arrested for minor offences, were more likely to be arrested after the project than those not mentored.²¹ On the basis of these findings, we concluded that non-directive mentoring programmes delivered by volunteers cannot be recommended as an effective intervention for young people at risk of or already involved in antisocial behaviour or criminal activities.

We are not suggesting that mentoring cannot work. There are many different kinds of mentoring, and some show better evidence of effect than others.¹⁰ Our current state of knowledge on the effectiveness of mentoring is similar to that of a new drug that shows promise but remains in need of further research and development. There is no equivalent of the National Institute for Clinical Excellence or Food and Drug Administration for mentoring. If there were, no more than a handful of programmes might have realistic hopes of qualifying. And even then, it would have to be acknowledged that we lack full understanding of the safeguards needed to

Summary points

Mentoring of children and young people involved in antisocial behaviour has become a popular intervention

Research evidence to support the most commonly used programmes is lacking

Failed mentoring relationships may have a detrimental effect on some young people

Commitment to research based practice needs to include focus on what works in implementation as well as evidence of effect

ensure that young people are not harmed by participation. For some of the most vulnerable young people, mentoring programmes as currently implemented may fail to deliver on their promises.

How should we respond to the evidence?

Showing that something works (or not) is one thing, and difficult enough. But what happens next in real life settings? We summarised the research findings and presented them to practitioners and planners who were implementing mentoring. Unsurprisingly, their response was not to abandon the projects. Instead, they asked how they could make it work.

They asked for evidence on what seemed to have a more positive effect for the group of children with whom they were working. The research suggests that cognitive behaviour therapeutic approaches are effective in attaining some of the outcomes sought, including when these are delivered through a mentoring-like component.^{22, 23} The practitioners therefore decided to include a directive element in their approach. They also drew on practices that seem to be associated with stronger benefits for young people, such as ongoing training for volunteer mentors and involvement of parents.¹⁰ Of course, without implementing such innovations in a trial setting, we will never know whether new approaches are better, worse, or much the same.

Conclusion

Social interventions are complex and are capable of doing as much or even more harm than medical ones. They need to be as thoroughly evaluated before and after implementation.

Those of us who have been pushing policy makers and practitioners to adopt evidence based practice need to be careful that we do not sell it as a simple way to solve problems. We need a lot more work on how to help policy makers deal with complex interventions and evidence. Research and development in health and social care needs huge investment if we are to develop adequate social interventions for big problems. At present, practitioners, parents, and children and young people themselves, looking for good research evidence on common problems, will find the evidence cupboard disappointingly bare.

Young people have the right to evidence based interventions. We know from the past that many well meaning attempts to do good resulted in harm, but we now have the means through systematic review, trials, sound evaluations, and good qualitative work, to do better.

Contributors and sources: HR and TAS are grant holders of the What Works for Children? project. Under their guidance, KL and PL authored the EvidenceNugget on mentoring for young people. Drawing on the EvidenceNugget, HR drafted the original idea for this paper and coordinated comments from KL, PL, DDB, and TAS. Sarah Frost of What Works for Children? facilitated work with practitioners and Mark Anslow, who is a Children's Fund manager, provided a practitioner's views on putting research into practice.

Funding: HR and TAS hold a grant from the Economic and Social Research Council under its evidence network programme (www.evidencenetwork.org). HR also had a grant from the Health Development Agency for work on interventions that may reduce inequalities in child health.

Competing interests: All the authors are involved in research on evidence based services.

- 1 Roberts H. Socio-economic determinants of health: children, inequalities and health. *BMJ* 1997;314:1122-5.
- 2 Glass N. Sure Start: the development of an early intervention programme for young children in the UK. *Child Soc* 1999;13:257-64.
- 3 Glass N. What works for children: the political issues. *Child Soc* 2001;15:14-20.
- 4 Schweinhart L, Weikart D. *A summary of significant benefits: the HighScope Perry pre-school study through age 27*. Ypsilanti, MI: High/Scope Press, 1993.
- 5 Barlow J. *Systematic review of the effectiveness of parent-training programmes in improving behaviour problems in children aged 3-10 years*. 2nd ed. Oxford: Health Services Research Unit, 1999.
- 6 Webster-Stratton C, Hollinsworth T, Kolpacoff M. The long-term effectiveness and clinical significance of three cost-effective training programs for families with conduct-problem children. *J Consult Clin Psychol* 1989;57:550-3.
- 7 Olds DL, Robinson J, O'Brien R, Luckey DW, Pettitt LM, Henderson CR Jr, et al. Home visiting by paraprofessionals and by nurses: a randomised, controlled trial. *Pediatrics* 2002;110:486-96.
- 8 Tierney JP, Grossman JB, Resch NL. *Making a difference. An impact study of Big Brothers Big Sisters*. Philadelphia: Public/Private Ventures, 2000.
- 9 Grossman JP, Tierney JP. Does mentoring work? An impact study of the Big Brothers Big Sisters program. *Eval Rev* 1998;22:403-26.
- 10 DuBois DL, Holloway BE, Valentine JC, Cooper H. Effectiveness of mentoring programs for youth: a meta-analytic review. *Am J Community Psychol* 2002;30:157-97.
- 11 Sheldon TA. It ain't what you do but the way that you do it. *J Health Serv Res Policy* 2001;6:3-5.
- 12 Scott S, Knapp M, Henderson J, Maughan B. Financial cost of social exclusion: follow up study of antisocial children into adulthood. *BMJ* 2001;323:191.
- 13 Webster-Stratton C. Long-term follow-up of families with young conduct problem children: from preschool to grade school. *J Clin Child Psychol* 1990;19:144-9.
- 14 Scott S. Fortnightly review: aggressive behaviour in childhood. *BMJ* 1998;316:202-6.
- 15 Home Office. Mentoring capital grant round 2003/4. www.homeoffice.gov.uk/docs/capital_mentoring_grants.html (accessed 13 Aug 2003).
- 16 Bush G. State of the union speech, 2003. www.cnn.com/2003/ALLPOLITICS/01/28/sou.transcript (accessed 13 Aug 2003).
- 17 St James-Roberts I, Samlal Singh C. *Can mentors help primary school children with behaviour problems? Final report of the Thomas Coram Research Unit between March 1997 and 2000*. London: Home Office, 2001.
- 18 Tarling R, Burrows J, Clarke A. *Dalston youth project part II (11-14) an evaluation*. London: Home Office, 2001.
- 19 What works for children? www.whatworksforchildren.org.uk (accessed 13 Aug 2003).
- 20 Grossman JB, Rhodes JE. The test of time: predictors and effects of duration in youth mentoring programs. *Am J Community Psychol* 2002;30:199-206.
- 21 O'Donnell CR, Lydgate T, Fo WSO. The buddy system: review and follow-up. *Child Behavior Therapy* 1979;1:161-9.
- 22 Davidson WS, Redner R, Blakely CH, Mitchell CM, Emshoff JG. Diversion of juvenile offenders: an experimental comparison. *J Consult Clin Psychol* 1987;55:68-75.
- 23 Cavell TA, Hughes JN. Secondary prevention as context for assessing change processes in aggressive children. *J School Psychol* 2000;38:199-235.

Where is the evidence that animal research benefits humans?

Pandora Pound, Shah Ebrahim, Peter Sandercock, Michael B Bracken, Ian Roberts on behalf of the Reviewing Animal Trials Systematically (RATS) Group

Much animal research into potential treatments for humans is wasted because it is poorly conducted and not evaluated through systematic reviews

Clinicians and the public often consider it axiomatic that animal research has contributed to the treatment of human disease, yet little evidence is available to support this view. Few methods exist for evaluating the clinical relevance or importance of basic animal research, and so its clinical (as distinct from scientific) contribution remains uncertain.¹ Anecdotal evidence or unsupported claims are often used as justification—for example, statements that the need for animal research is “self evident”² or that “Animal experimentation is a valuable research method which has proved itself over time.”³ Such statements are an inadequate form of evidence for such a controversial area of research. We argue that systematic reviews of existing and future research are needed.

receives much more funding than clinical research.^{1 4 5} Given this, and because the public accepts animal research only on the assumption that it benefits humans,⁶ the clinical relevance of animal experiments needs urgent clarification.

Several methods are available to evaluate animal research. These include historical analysis,⁷ critiques of animal models,⁸ investigations into the development of treatments,⁵ surveys of clinicians' views,⁹ and citation analyses.¹⁰ However, perhaps the best way of producing evidence about the value of animal research is to conduct systematic reviews of animal studies and, where possible, compare the results of these with the results of the corresponding clinical trials. So what do studies that have done this show?

Assessing animal research

Despite the lack of systematic evidence for its effectiveness, basic animal research in the United Kingdom

Department of Social Medicine, University of Bristol, Bristol BS8 2PR
Pandora Pound research fellow
Shah Ebrahim professor
Department of Clinical Neurosciences, University of Edinburgh, Western General Hospital, Edinburgh EH4 2XU
Peter Sandercock professor
continued over

BMJ 2004;328:514-7

 Details of the search strategy and references w1-w18 are on bmj.com