



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/188201/>

Version: Accepted Version

Article:

Abu-Odah, H, Said, NB, Nair, SC et al. (2022) Identifying barriers and facilitators of translating research evidence into clinical practice: A systematic review of reviews. *Health and Social Care in the Community*, 30 (6). e3265-e3276. ISSN: 0966-0410

<https://doi.org/10.1111/hsc.13898>

© 2022 John Wiley & Sons Ltd. This is the peer reviewed version of the following article: Abu-Odah, H., Said, N. B., Nair, S. C., Allsop, M. J., Currow, D. C., Salah, M. S., Hamad, B. A., Elessi, K., Alkhatib, A., ElMokhallalati, Y., Bayuo, J., & AlKhaldi, M. (2022). Identifying barriers and facilitators of translating research evidence into clinical practice: A systematic review of reviews. *Health & Social Care in the Community*, 30, e3265– e3276., which has been published in final form at <https://doi.org/10.1111/hsc.13898>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Abstract

Translating research into clinical practice is a global priority because of its potential impact on health services delivery and outcomes. Despite the ever-increasing depth and breadth of health research, most areas across the globe seem to be slow to translate relevant research evidence into clinical practice. Thus, this review sought to synthesise existing literature to elucidate the barriers and facilitators to the translation of health research into clinical practice. A systematic review of reviews approach was utilised. Review studies were identified across PubMed, Scopus, Embase, CINAHL and Web of Science databases, from their inception to 15 March 2021. Searching was updated on 30 March 2022. All retrieved articles were screened by two authors; reviews meeting the inclusion criteria were retained. Based on the review type, two validated tools were employed to ascertain their quality: A Measurement Tool to Assess Systematic Reviews-2 and International Narrative Systematic assessment. The framework synthesis method was adopted to guide the analysis and narrative synthesis of data from selected articles. Ten reviews met the inclusion criteria. The study revealed that the translation of new evidence was limited predominantly by individual-level issues and less frequently by organisational factors. Inadequate knowledge and skills of individuals to conduct, organise, utilise and appraise research literature were the primary individual-level barriers. Limited access to research evidence and lack of equipment were the key organisational challenges. To circumvent these barriers, it is critical to establish collaborations and partnerships between policymakers and health professionals at all levels and stages of the research process. The study concluded that recognising barriers and facilitators could help set key priorities that aid in translating and integrating research evidence into practice. Effective stakeholder collaboration and cooperation should improve the translation of research findings into clinical practice.

Keywords

Barriers and facilitators, evidence translation, healthcare professionals, policymakers, systematic review

What is known about this topic

- Effective knowledge translation has the potential to improve standards of care and overall health service delivery.
- Healthcare professionals' lack of motivation, lack of continuous education, uncooperative and unsupportive organisational culture and the disintegration between knowledge producers and users are the key barriers to the translation of research into clinical practice.

What this review adds

- Various barriers, predominantly individual-related issues and organisational factors, affect the translation of health research findings to clinical environments.
- Inadequate knowledge and skills of healthcare staff to conduct, organise, utilise and appraise research literature were significant barriers to the translation process.
- Establishing collaborations and partnerships between policymakers and health professionals at all levels and stages of the research process were the main facilitators of the knowledge translation process.
- A novel application of the Innovative Care for Chronic Conditions model facilitated categorisation of barriers and facilitators to evidence translation in the clinical settings.

Introduction

The utilisation of knowledge generated from health research to inform and guide clinical practice is recognised as a high global priority (Brownson et al., 2018; Conalogue et al., 2017; Poot et al., 2018). In response to rapid increases in evidence production and knowledge base development, the restructuring of healthcare, insufficient resources and increased professional accountability, many governmental and policymaking institutions and funding agencies require evidence-informed clinical practice approaches (Milner et al., 2006). Effective translation of knowledge can improve standards of care and overall health service delivery (Barratt et al., 2017; Langlois et al., 2016).

Knowledge translation (KT) is a term used to explain the activities involved in translating health-related research findings into clinical practice (Gervais et al., 2015). It attempts to assure that stakeholders or 'knowledge users' are familiar with, using and accessing research findings and engaging them as active participants in the research process (Engebretsen et al., 2017; Kreindler, 2018). KT refers to an 'iterative process that includes the synthesis, dissemination, exchange and ethically sound application of knowledge to improve health, provide more effective health services and products and strengthen the healthcare system' (The Canadian Institutes of Health Research, 2016). Understanding and applying the process of KT is therefore essential for alleviating barriers to translating health research evidence into clinical practice.

Despite the increasing amount of health research globally, the process of translating health-related research findings into clinical practice remains slow (Grimshaw et al., 2012; Straus et al., 2011) and the gap between findings and clinical practice is widening (LaRocca et al., 2012). This gap could be related to a wide range of challenges that have been reported in previous studies. Insufficient time (Wallis, 2012), lack of healthcare professionals'

motivation (Curtis et al., 2017; Ramón et al., 2022), lack of continuing education (Curtis et al., 2017; Wallis, 2012), uncooperative and unsupportive organisational culture (Ramón et al., 2022; Wallis, 2012) and the disintegration between knowledge producers and users (Norström et al., 2020) are the key barriers to translating research to clinical practice. A clear understanding and assessment of the barriers related to KT in practice can help health systems optimize their use of research evidence and improve the quality of health services provided to patients.

Numerous studies have explored the barriers to translating health research evidence into clinical practice. Oliver et al., (2014) identified challenges that policymakers faced when utilising health research findings; key obstacles were lack of access to relevant research and lack of timely research output. Lawrence et al., (2019) explored decision-makers' involvement in KT. Findings underscored that lack of motivation to update knowledge and lack of experience were the key barriers to utilising KT in practice. Despite these reviews' important findings, most focused on policymakers and gave limited attention to other stakeholders. Further, reviews tended to emphasise barriers to KT as opposed to facilitators. Limited systematic reviews comprehensively address barriers and facilitators to KT. Therefore, the current review aimed to comprehensively synthesise available evidence and provide an overview of barriers and facilitators that enable the translation of health research findings into clinical practice.

The translation process was defined as utilising the Innovative Care for Chronic Conditions (ICCC) model (World Health Organization, 2016). The ICCC model is widely used across studies that include guidelines on how to translate health research evidence effectively. Although the ICCC model targets chronic diseases, it covers three levels of determinants: micro, meso and macro. These three levels allow for the stratification and understanding of KT barriers and facilitators. The model is based on understanding the barriers and

facilitators from the policy environment, healthcare organisations and healthcare professionals. It was utilised in previous studies as an analytical framework for identifying barriers and facilitators to provide services to health systems (Abu-Odah et al., 2020). Utilising such an approach could help address barriers and promote the facilitators that alleviate the translation of health research evidence into practice, and the ICCC was adopted as an analytical framework in this systematic review.

Methods

Study design

The study adopted a review of reviews approach to provide a systematic, comprehensive synopsis of the field and identify the current barriers and facilitators to KT. This design allows for the creation and synthesis of high-level evidence generated from reviews (e.g. systematic, scoping, integrative and narrative reviews) in a single document. It also enables the researchers to combine large amounts of data from multiple sources from which they can distil key messages to guide future research. The Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidance was utilised to structure this review (Page et al., 2021).

Search sources and strategies

PubMed, Web of Science, Scopus, Embase and CINAHL were searched for reviews published in English from inception to 15 March 2021. Searching was updated on 30 March 2022. The following search keywords, developed based on population, concept and context framework, were utilised for searching: (“translational research” OR “knowledge translation” OR “evidence to practice” OR “evidence-informed” OR “evidence-based practice” OR “knowledge exchange” OR “knowledge interaction” OR “research utilisation” OR “research dissemination” OR “knowledge uptake” OR “knowledge-to-

action” OR “research diffusion”) AND (medicine OR nursing OR “public health” OR “health”) AND (“challenges” OR “obstacles” OR “limitations” OR “problems” OR “barriers ”). Expressions for “review*” were also included (see Table S1).

Eligibility criteria

Inclusion criteria

- Reviews (systematic, scoping, narrative and integrative reviews) with a defined and systematic approach to identifying literature (i.e., developing a search strategy and using it to identify data across multiple databases) and reporting barriers or/and facilitators associated with research evidence and clinical practice.
- Reviews incorporating any methodological approach, including those employing qualitative, quantitative, and mixed methods approaches.
- Reviews focusing on the translation of knowledge evidence into clinical practice.
- Reviews were written in English.

Exclusion criteria

- Informal literature reviews without defined research questions and do not have defined search process or discussion papers.
- Other types of articles such as letters, editorials, conference abstracts, short surveys.
- Publications where full-text versions could not be obtained.

Data extraction

All papers retrieved from the databases were exported into EndNote X9 software. After removing duplications, the remaining papers were screened independently for eligibility criteria by the first and second authors. The potentially eligible full-text papers were then located for screening by the two main authors, and any disagreements were resolved by the third author. Reasons for excluding papers are reported in Figure 1.

Ten papers were independently extracted by two authors using data extraction sheets developed for this study (Table 1). The data sheets were used to report the following data: (1) citation information; (2) number of studies included; (3) aim of review; (4) main findings (barriers and facilitators) summarised based on the ICCC model.

Quality assessment of included reviews

Two authors (HAO and NS) independently utilised two separate validated tools for evaluating the selected reviews. The first is A Measurement Tool to Assess systematic Reviews-2 (AMSTAR-2) (Shea et al., 2017), which is used to assess systematic and scoping reviews. It includes 16 appraisal items; each item is ranked into three levels: yes, no, or partial yes. Each review's final score was rated as 'high,' which meant the review provided a comprehensive summary of findings, 'moderate,' which meant the review had some weaknesses but not critical flaws, while 'low' scores indicated that the review did not provide comprehensive findings and had critical flaws (Shea et al., 2017). The second tool is the International Narrative Systematic assessment (INSA), which is used to assess narrative review papers (La Torre et al., 2015). The INSA tool contains seven appraisal items where each item is rated as 'yes' or 'no'. A paper with a score higher than 5 points is judged as a 'good' quality paper (La Torre et al., 2015).

Data analysis

The framework synthesis method (Barnett-Page & Thomas, 2009; Pope et al., 2000) was adopted to guide the analysis and narrative synthesis of data from selected articles. This synthesis method was used because it offers a robust approach to shaping and synthesising large amounts of textual data (Barnett-Page & Thomas, 2009; Pope et al., 2000) including where varying methodologies may have been used. The initial framework used to explore and organise data was the multilevel World Health Organization ICCC model that is widely

used across studies for data analysis considering multilevel determinants, such as the micro-, meso- and macro-levels (World Health Organization, 2016). The framework includes three levels: the micro-level focusing on individual users, the meso-level focusing on healthcare institutions and systems and, finally, the macro-level related to national policies and strategies.

Results

Study characteristics

Of 986 articles indexed in five databases, 834 articles were excluded due to not matching the inclusion criteria. The full texts of the remaining articles were screened. Five articles were eligible for inclusion in the review. References from each of the five articles were scanned to check for any additional eligible articles. Five additional articles were included from the screened references. As a result, ten articles underwent quality appraisal and analysis (Figure 1).

The study revealed that there are different types of included research articles. Of the ten included articles, five were systematic reviews (Edwards et al., 2019; Légaré et al., 2008; Oliver et al., 2014; Sadeghi-Bazargani et al., 2014; Shayan et al., 2019), four were narrative reviews (Athanasakis, 2013; Derman & Jaeger, 2018; Kalassian et al., 2002; Straus et al., 2011), and one was a scoping review (Lawrence et al., 2019). Three reviews targeted developing countries (Derman & Jaeger, 2018; Edwards et al., 2019; Shayan et al., 2019), one addressed the KT topic from the global perspective (Straus et al., 2011), and the other reviews did not specify the geographic scope (Athanasakis, 2013; Kalassian et al., 2002; Lawrence et al., 2019; Légaré et al., 2008; Oliver et al., 2014; Sadeghi-Bazargani et al., 2014; Straus et al., 2011).

Most reviews primarily assessed policymaker and healthcare system perspectives on KT (Edwards et al., 2019; Lawrence et al., 2019; Oliver et al., 2014), although two reviews investigated nurses' views (Athanasakis, 2013; Shayan et al., 2019) and two investigated healthcare professionals' perspectives on KT (Légaré et al., 2008; Sadeghi-Bazargani et al., 2014). A single review dealt with clinicians (Athanasakis, 2013). Most reviews were published in the last ten years (Athanasakis, 2013; Derman & Jaeger, 2018; Edwards et al., 2019; Oliver et al., 2014; Sadeghi-Bazargani et al., 2014; Shayan et al., 2019; Straus et al., 2011; Woolf et al., 2015). Two reviews were published before 2010 (Kalassian et al., 2002; Légaré et al., 2008).

Six reviews specified the number of studies they included (Edwards et al., 2019; Lawrence et al., 2019; Légaré et al., 2008; Oliver et al., 2014; Sadeghi-Bazargani et al., 2014; Shayan et al., 2019). Two reviews reported being comprised of over 100 studies (Oliver et al., 2014; Sadeghi-Bazargani et al., 2014), one review included 62 studies (Edwards et al., 2019) and the other reviews included fewer than 50 studies; ranging between 16 and 38 studies (Lawrence et al., 2019; Légaré et al., 2008; Shayan et al., 2019).

Quality appraisal of reviews

The methodological quality of the included systematic reviews ranged from moderate to high, meaning that they showed robust and accurate summaries of their findings (Edwards et al., 2019; Lawrence et al., 2019; Légaré et al., 2008; Oliver et al., 2014; Sadeghi-Bazargani et al., 2014; Shayan et al., 2019). Of the four included narrative reviews, three reviews scored ≥ 5 points, reflecting a good quality paper (Athanasakis, 2013; Kalassian et al., 2002; Straus et al., 2011). Although three reviews were assessed as being of low to moderate quality, quality assessment was not used as a means of determining inclusion in the review. Instead, quality appraisal was used to inform the current state of the evidence. The three reviews received low-quality grades because they did not report any potential

sources of conflict of interest, including funding they received for conducting the reviews. A detailed methodological quality assessment of the included reviews is outlined in Table S2.

Barriers and facilitators

This section presents barriers and facilitators to translating KT into practice. Key findings of reviews were structured into the three ICCC levels: micro, meso and macro. These findings are presented as follows: 1) barriers to KT; 2) facilitators or contributing factors to KT (Figure 2).

Micro-level (individuals)

Individual barriers focused on healthcare professionals' abilities to conduct, read, use or translate evidence into clinical practice. Findings revealed that limited professional engagement in the research process, lack of time, insufficient critical appraisal skills and an inability of healthcare professionals to use the research findings and recommendations in clinical practice were the most common individual-level challenges (Edwards et al., 2019; Shayan et al., 2019). Moreover, low technology literacy levels and unfamiliarity with research databases were barriers to healthcare professionals' understanding and use of research (Athanasakis, 2013; Lawrence et al., 2019). These factors created an obstacle in the translation of research evidence into clinical practice.

Other barriers, such as professionals' unfamiliarity with evidence-based practice concepts (Shayan et al., 2019), lack of interest to update knowledge on emerging best practices (Lawrence et al., 2019), and underestimation of the value of research were additional barriers identified at the individual level (Athanasakis, 2013; Lawrence et al., 2019). Not only did healthcare professionals lack the capabilities to interpret findings from studies, but also to understand statistical measures utilised in research (Sadeghi-Bazargani et al., 2014).

The facilitators for overcoming micro-level challenges healthcare professionals' motivation and interest (Légaré et al., 2008) in addressing and studying research findings, given suitable packaging and targeted communication of results (Derman & Jaeger, 2018).

Meso-level (systems or organisations)

This level includes systemic or organisational barriers, such as education, training, resources, services, policies, and organisation culture. Time constraints, insufficient organisational resources, poor knowledge dissemination and lack of access to evidence and research were the most commonly reported barriers (Athanasakis, 2013; Edwards et al., 2019; Lawrence et al., 2019; Légaré et al., 2008; Oliver et al., 2014; Shayan et al., 2019). Insufficient resources (e.g., materials and equipment) required for the implementation of research and inadequate facilities to conduct research were reported in three studies (Derman & Jaeger, 2018; Edwards et al., 2019; Kalassian et al., 2002; Lawrence et al., 2019; Légaré et al., 2008; Sadeghi-Bazargani et al., 2014).

Inadequate organisational, political, technical and financial support was the second most frequently cited barrier to translating evidence into clinical practice. Workforce shortage was reported as an additional organisational challenge (Shayan et al., 2019). Inappropriate management, organisation of staff, and workload density made it difficult for staff to allocate work time to search and read research papers or synthesise evidence and guidelines (Shayan et al., 2019). There were additional barriers specific to KT in nursing practice, including lack of training and educational opportunities for research (Shayan et al., 2019) and limited cooperation among educational and clinical environments or entities (Shayan et al., 2019).

Institutional support for the translation of evidence into practice can come through a policy or plan to implement various capacity-building activities, such as workshops.

Dissemination of primary research findings across organisations (Derman & Jaeger, 2018; Oliver et al., 2014) and budgeting for research activities were supplementary capacity-building facilitating actions to support institutional KT (Derman & Jaeger, 2018; Oliver et al., 2014). A unique but important facilitator was the development and establishment of mechanisms and channels for effective, sustainable communication to build a shared understanding and experience of the work (Lawrence et al., 2019).

Macro-level (economic and political)

This level focused on policymakers' issues that support translating evidence into clinical practice, such as guidelines, partnerships and regulations. Three reviews revealed that political challenges are a key impediment to the translation of health research evidence into clinical practice (Kalassian et al., 2002; Lawrence et al., 2019; Oliver et al., 2014). Policymakers were not sufficiently trained and skilled in research methods (Oliver et al., 2014) and did not perceive or observe alignment or integration between research and policy (Lawrence et al., 2019). In addition, they remain doubtful about the utility of research findings (Kalassian et al., 2002).

The main macro-level facilitators reported in the two reviews were identifying the stakeholders and developing robust collaboration and connections between policymakers and research staff (Lawrence et al., 2019; Oliver et al., 2014). These can be achieved by building technology, such as web-based conferencing platforms, and supporting its capacity. This technology provides policymakers with updated information and knowledge in research and engages them in all research priorities, which helps them make evidence-based decisions (Lawrence et al., 2019). Additional facilitators play a significant role in translating evidence into clinical practice, including developing trust across policymakers and researchers (Lawrence et al., 2019) and developing guidelines that promote clinical best practices (Kalassian et al., 2002). It is also important to involve stakeholders early in

the research design and initiation process, as they are most likely to be affected by research output (Derman & Jaeger, 2018).

Discussion

This review sought to provide a comprehensive and systematic mapping of the barriers related to translating research findings to clinical practice. It further articulates possible facilitators that strengthen this process. The included primary reviews highlighted that the translation of health research findings into a clinical environment is affected by various barriers, predominantly individual-related issues and those relating to organisational factors. It was identified that inadequate healthcare professionals' knowledge and skills inhibited their ability to conduct, organise, utilise and appraise research literature; these were significant barriers to the translation process. Conversely, key stakeholder collaboration was highlighted as a crucial facilitator.

Translation of health research evidence to real-world environments is pivotal for the success of any clinical practice or implementation of health policy (Bahadori et al., 2016; Bayuo, 2017). However, the translation process is often wrought with multiple barriers, as reported in this review. Inadequacy of individuals' knowledge and skills to conduct, organise, utilise and appraise research literature are the main individual-related barriers to translating research findings into clinical practice. These results are congruent with previous studies, highlighting that the lack of sufficient skills to appraise, interpret and utilise research findings deepens the research-practice gap (Bahadori et al., 2016). Training and engaging professionals in research practice, including the identification and use of research evidence, may be essential to boost their knowledge and skills to participate in the research process and interpret the findings (Mickan et al., 2017).

Additional barriers to research translation identified in this study included a lack of research education leading to disinterest, motivational challenges and scepticism over the potential of research evidence to be translated into clinical practice (Bahadori et al., 2016; Curtis et al., 2017). Several healthcare systems have invested considerably to address these knowledge barriers over the last two decades (Grimshaw et al., 2012). For instance, practice guidelines were in part developed to reduce the time needed for sourcing and reading research papers. The investment was also focused on reaching electronic libraries of open access evidence sources to enhance access to research articles and training to enhance research literacy skills (Grimshaw et al., 2012). However, these investments have confronted additional challenges, such as staff shortages, which limit clinicians' capacity to engage with and review research, irrespective of its format (Bahadori et al., 2016).

A lack of interest in the research process is another significant individual-level barrier noted in this study. The factors driving engagement and interest in research are likely to be multifaceted, potentially ranging from individual perceptions of the value of research to wider systemic issues such as limited clinical academic career pathways that would further enhance one's research skills (Brandenburg & Ward, 2022). At the micro level, translational barriers may also be reduced by motivating healthcare professionals. Individual-level facilitators involve a clear understanding of the target population, who could benefit from the research findings, so that the research evidence can be customised and communicated in a practical way to enable easy translation. Our results are aligned with previous reports indicating that successful dissemination and utilisation of research evidence can be achieved following the identification of the right audience and tailoring of messages using appropriate mediums (Curtis et al., 2017; Kothari & Wathen, 2017; Van der Graaf et al., 2018).

At the organisational level, translating research into the clinical environment requires resources. A lack of resources, such as limited access to research databases and the requisite equipment for them (such as IT infrastructure), are the leading organisational barriers to applying research findings in the clinical setting (Curtis et al., 2017). Lack of resources also indirectly affect the professional development of staff. Time constraints, heavy workload and lack of an adequately skilled workforce to read and understand research processes limit the translation of research into clinical practice. Existing literature aligns with these findings (Bahadori et al., 2016; Edwards et al., 2019); time was reported as an essential factor for utilising research evidence in clinical environments (Bahadori et al., 2016). A study conducted in eastern Turkey underscored that ‘the lack of time to implement the new ideas in the workplaces’ and ‘the lack of sufficient time for reading the studies’ were the most common barriers to translating research findings into clinical practice (Tan et al., 2012).

Our observations indicate policymakers’ mistrust in the potential of translating research into clinical practice. This affects both health policy development and systematic public investments in research programmes. A sizeable proportion of policymakers’ mistrust stems from their lack of research skills. Early identification and partnering with all key stakeholders (policymakers, evidence producers and the beneficiaries of research, such as the community) may overcome this challenge. Similar models have been suggested in earlier studies (Ongolo-Zogo et al., 2018) with, for example, technology-driven interactive models providing all stakeholders and beneficiaries with constant engagement and updated information to enable them to support evidence-based models (Cairney & Oliver, 2017; Van der Graaf et al., 2018). Alongside inclusion and participation of key stakeholders, focusing on transparency and accessibility of timely research findings can ensure that policymakers access and engage with research findings (Donnelly et al., 2018).

Strengths and limitations

Using a deductive approach to ascertain the barriers and facilitators may not have allowed for new and emergent insights to arise from the literature. However, the framework approach provided a useful way of structuring and mapping the key findings and literature in this area. A strength of our study is in the identification and stratification of the barriers and facilitators at three levels, which can help with devising targeted strategies to overcome identified challenges.

Conclusions

This systematic review highlights the key barriers and facilitators for translating research to clinical practice at the individual and organisational levels. Research evidence is essential to improving healthcare practice, thereby also enhancing patient outcomes. Through adopting an approach that explored factors at the individual and organisational levels, our findings can guide the development of targeted interventions to overcome the identified barriers. Furthermore, wider efforts to foster effective collaboration and cooperation between all stakeholders could lead to improvements in the translation of health research findings into clinical practice.

Ethics approval

Ethical approval was not required for this study because it used data from published papers.

References

- Abu-Odah, H., Mikati, D., & Arawi, T. (2020). Deconstructing Palliative Care in Areas of Armed Conflict: Needs, Challenges, and Concerns. In I. Laher (Ed.), *Handbook of Healthcare in the Arab World* (pp. 1-17). Springer International Publishing.
https://doi.org/10.1007/978-3-319-74365-3_225-1
- Athanasakis, E. (2013). Nurses' research behavior and barriers to research utilization into clinical nursing practice: a closer look. *International Journal of Caring Sciences*, 6(1), 16-28.
<http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=104240445&site=ehost-live>
- Bahadori, M., Raadabadi, M., Ravangard, R., & Mahaki, B. (2016). The barriers to the application of the research findings from the nurses' perspective: A case study in a teaching hospital. *Journal of education and health promotion*, 5, 14-14. <https://doi.org/10.4103/2277-9531.184553>
- Barnett-Page, E., & Thomas, J. (2009). Methods for the synthesis of qualitative research: a critical review. *BMC Medical Research Methodology*, 9(1), 59. <https://doi.org/10.1186/1471-2288-9-59>
- Barratt, H., Shaw, J., Simpson, L., Bhatia, S., & Fulop, N. (2017). Health services research: building capacity to meet the needs of the health care system. *Journal of health services research & policy*, 22(4), 243-249. <https://doi.org/10.1177/1355819617714815>
- Bayuo, J. (2017). "Worlds apart": the case of evidence utilization in Ghanaian health facilities. *JB/Evidence Synthesis*, 15(9), 2225-2226.
- Brandenburg, C., & Ward, E. C. (2022). "There hasn't been a career structure to step into": a qualitative study on perceptions of allied health clinician researcher careers. *Health research policy and systems*, 20(1), 6. <https://doi.org/10.1186/s12961-021-00801-2>
- Brownson, R. C., Fielding, J. E., & Green, L. W. (2018). Building capacity for evidence-based public health: reconciling the pulls of practice and the push of research. *Annu Rev Public Health*, 39, 27-53. <https://doi.org/10.1146/annurev-publhealth-040617-014746>
- Cairney, P., & Oliver, K. (2017). Evidence-based policymaking is not like evidence-based medicine, so how far should you go to bridge the divide between evidence and policy? *Health research policy and systems*, 15(1), 35. <https://doi.org/10.1186/s12961-017-0192-x>
- Conalogue, D. M., Kinn, S., Mulligan, J.-A., & McNeil, M. (2017). International consultation on long-term global health research priorities, research capacity and research uptake in developing countries. *Health research policy and systems*, 15(1), 24-24.
<https://doi.org/10.1186/s12961-017-0181-0>

- Curtis, K., Fry, M., Shaban, R. Z., & Considine, J. (2017). Translating research findings to clinical nursing practice. *J Clin Nurs*, 26(5-6), 862-872. <https://doi.org/10.1111/jocn.13586>
- Derman, R. J., & Jaeger, F. J. (2018). Overcoming challenges to dissemination and implementation of research findings in under-resourced countries. *Reproductive Health*, 15(1), 121-126. <https://doi.org/10.1186/s12978-018-0538-z>
- Edwards, A., Zweigenthal, V., & Olivier, J. (2019). Evidence map of knowledge translation strategies, outcomes, facilitators and barriers in African health systems [Review]. *Health research policy and systems*, 17(1). <https://doi.org/10.1186/s12961-019-0419-0>
- Engebretsen, E., Sandset, T. J., & Ødemark, J. (2017). Expanding the knowledge translation metaphor. *Health research policy and systems*, 15(1), 19-19. <https://doi.org/10.1186/s12961-017-0184-x>
- Gervais, M.-J., Marion, C., Dagenais, C., Chiocchio, F., & Houffort, N. (2015). Dealing with the complexity of evaluating knowledge transfer strategies: Guiding principles for developing valid instruments. *Research Evaluation*, 25(1), 62-69. <https://doi.org/10.1093/reseval/rvv034>
- Grimshaw, J., Eccles, M. P., Lavis, J. N., Hill, S. J., & Squires, J. E. (2012). Knowledge translation of research findings. *Implement Sci*, 7, 50-50. <https://doi.org/10.1186/1748-5908-7-50>
- Kalassian, K. G., Dremsizov, T., & Angus, D. C. (2002). Translating research evidence into clinical practice: new challenges for critical care. *Critical care.*, 6(1), 11-14. <https://doi.org/10.1186/cc1446>
- Kothari, A., & Wathen, C. N. (2017). Integrated knowledge translation: digging deeper, moving forward. *J Epidemiol Community Health*, 71(6), 619-623. <https://doi.org/10.1136/jech-2016-208490>
- Kreindler, S. A. (2018). Advancing the evaluation of integrated knowledge translation. *Health research policy and systems*, 16(1), 104-104. <https://doi.org/10.1186/s12961-018-0383-0>
- La Torre, G., Backhaus, I., & Mannocci, A. (2015). Rating for narrative reviews: concept and development of the International Narrative Systematic Assessment tool. *Senses and Sciences*, 2(2), 31-35.
- Langlois, E. V., Becerril Montekio, V., Young, T., Song, K., Alcalde-Rabanal, J., & Tran, N. (2016). Enhancing evidence informed policymaking in complex health systems: lessons from multi-site collaborative approaches. *Health research policy and systems*, 14, 20-20. <https://doi.org/10.1186/s12961-016-0089-0>

- LaRocca, R., Yost, J., Dobbins, M., Ciliska, D., & Butt, M. (2012). The effectiveness of knowledge translation strategies used in public health: a systematic review. *BMC Public Health*, 12(1), 751. <https://doi.org/10.1186/1471-2458-12-751>
- Lawrence, L. M., Bishop, A., & Curran, J. (2019). Integrated Knowledge Translation with Public Health Policy Makers: A Scoping Review. *Healthcare Policy*, 14(3), 55-77. <http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=136113277&site=ehost-live>
- Légaré, F., Ratté, S., Gravel, K., & Graham, I. D. (2008). Barriers and facilitators to implementing shared decision-making in clinical practice: Update of a systematic review of health professionals' perceptions. *Patient Educ Couns*, 73(3), 526-535. <https://doi.org/https://doi.org/10.1016/j.pec.2008.07.018>
- Mickan, S., Wenke, R., Weir, K., Bialocerkowski, A., & Noble, C. (2017). Strategies for research engagement of clinicians in allied health (STRETCH): a mixed methods research protocol. *BMJ Open*, 7(9), e014876. <https://doi.org/10.1136/bmjopen-2016-014876>
- Milner, M., Estabrooks, C. A., & Myrick, F. (2006). Research utilization and clinical nurse educators: a systematic review. *Journal of Evaluation in Clinical Practice*, 12(6), 639-655. <https://doi.org/10.1111/j.1365-2753.2006.00632.x>
- Norström, A. V., Cvitanovic, C., Löf, M. F., West, S., Wyborn, C., Balvanera, P., Bednarek, A. T., Bennett, E. M., Biggs, R., de Bremond, A., Campbell, B. M., Canadell, J. G., Carpenter, S. R., Folke, C., Fulton, E. A., Gaffney, O., Gelcich, S., Jouffray, J.-B., Leach, M., Le Tissier, M., Martín-López, B., Louder, E., Loutre, M.-F., Meadow, A. M., Nagendra, H., Payne, D., Peterson, G. D., Reyers, B., Scholes, R., Speranza, C. I., Spierenburg, M., Stafford-Smith, M., Tengö, M., van der Hel, S., van Putten, I., & Österblom, H. (2020). Principles for knowledge co-production in sustainability research. *Nature Sustainability*, 3(3), 182-190. <https://doi.org/10.1038/s41893-019-0448-2>
- Oliver, K., Innvar, S., Lorenc, T., Woodman, J., & Thomas, J. (2014). A systematic review of barriers to and facilitators of the use of evidence by policymakers. *BMC Health Serv Res*, 14(1), 2. <https://doi.org/10.1186/1472-6963-14-2>
- Ongolo-Zogo, P., Lavis, J. N., Tomson, G., & Sewankambo, N. K. (2018). Assessing the influence of knowledge translation platforms on health system policy processes to achieve the health millennium development goals in Cameroon and Uganda: a comparative case study. *Health Policy Plan*, 33(4), 539-554. <https://doi.org/10.1093/heapol/czx194>
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J., Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S., McGuinness, L. A., Stewart, L. A., Thomas, J., Tricco, A. C., Welch, V. A., Whiting, P., & Moher, D. (2021). The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Bmj*, 372, n71. <https://doi.org/10.1136/bmj.n71>

- Poot, C. C., van der Kleij, R. M., Brakema, E. A., Vermond, D., Williams, S., Cragg, L., van den Broek, J. M., & Chavannes, N. H. (2018). From research to evidence-informed decision making: a systematic approach. *Journal of Public Health, 40*(suppl_1), i3-i12. <https://doi.org/10.1093/pubmed/fox153>
- Pope, C., Ziebland, S., & Mays, N. (2000). Qualitative research in health care. Analysing qualitative data. *BMJ., 320*(7227), 114-116. <https://doi.org/10.1136/bmj.320.7227.114>
- Ramón, C., Nievas-Soriano, B. J., García-González, J., Alarcón-Rodríguez, R., Requena-Mullor, M., & Lozano-Paniagua, D. (2022). Motivation and Barriers to Research among Nursing Professionals in Southeast Spain. *Healthcare (Basel), 10*(4). <https://doi.org/10.3390/healthcare10040675>
- Sadeghi-Bazargani, H., Tabrizi, J. S., & Azami-Aghdash, S. (2014). Barriers to evidence-based medicine: a systematic review. *Journal of Evaluation in Clinical Practice, 20*(6), 793-802. <https://doi.org/10.1111/jep.12222>
- Shayan, S. J., Kiwanuka, F., & Nakaye, Z. (2019). Barriers associated with evidence-based practice among nurses in low- and middle-income countries: a systematic review. *Worldviews on Evidence-Based Nursing, 16*(1), 12-20. <https://doi.org/10.1111/wvn.12337>
- Shea, B. J., Reeves, B. C., Wells, G., Thuku, M., Hamel, C., Moran, J., Moher, D., Tugwell, P., Welch, V., Kristjansson, E., & Henry, D. A. (2017). AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ., 358*, j4008. <https://doi.org/10.1136/bmj.j4008>
- Straus, S. E., Tetroe, J. M., & Graham, I. D. (2011). Knowledge translation is the use of knowledge in health care decision making. *Journal of Clinical Epidemiology, 64*(1), 6-10. <https://doi.org/https://doi.org/10.1016/j.jclinepi.2009.08.016>
- Tan, M., Akgün Sahin, Z., & Kardas Özdemir, F. (2012). Barriers of research utilization from the perspective of nurses in Eastern Turkey. *Nurs Outlook, 60*(1), 44-50. <https://doi.org/10.1016/j.outlook.2011.07.002>
- The Canadian Institutes of Health Research. (2016). *About us*. Retrieved 22. February from <https://cihr-irsc.gc.ca/e/29418.html>
- Van der Graaf, P., Francis, O., Doe, E., Barrett, E., O'Rourke, M., & Docherty, G. (2018). Structural approaches to knowledge exchange: comparing practices across five centres of excellence in public health. *Journal of Public Health, 40*(suppl_1), i31-i38. <https://doi.org/10.1093/pubmed/fox150> %J Journal of Public Health
- Wallis, L. (2012). Barriers to implementing evidence-based practice remain high for U.S. nurses: getting past "we've always done it this way" is crucial. *Am J Nurs, 112*(12), 15. <https://doi.org/10.1097/01.Naj.0000423491.98489.70>

Woolf, S. H., Purnell, J. Q., Simon, S. M., Zimmerman, E. B., Camberos, G. J., Haley, A., & Fields, R. P. (2015). Translating evidence into population health improvement: strategies and barriers. *Annu Rev Public Health, 36*, 463-482. <https://doi.org/10.1146/annurev-publhealth-082214-110901>

World Health Organization. (2016). Innovative care for chronic conditions: Building blocks for actions: global report. 2002.

Table 1. Main Results of the Included Papers

Author, Study Year(s), Number of the included paper	Paper aim	Study design & analysis	Population countries	Main results	
				Barriers	Facilitators
Edwards et al. (2019) 2019 62 articles	"To provides a systematic overview of the literature on knowledge translation strategies employed by health system researchers and policymakers in African countries"	A systematic review Narrative synthesis	Healthcare researchers and policymakers African countries	Micro-level Deficient skills to perform knowledge translation activities, Meso-level Insufficient time at the organisation A paucity of resources that help professionals in immersing in health research	Meso-level Capacity-building workshops Macro-level Providing the most relevant high-quality research
Shayan et al. (2019) 2019 16 articles	"To assess barriers to EBP among nurses in low- and middle-income countries".	A systematic review Narrative synthesis	Nurses LMICs	Micro-level Difficulty in appraising research findings Professionals do have not enough time to read, conduct research, Professional are not knowledgeable on how to translate the findings to practice Meso-level Scant of the equipment's to implement research Scant of needed materials to implement research Difficulties in accessing information Lack of workforce Inadequate facilities to conduct research Lack of teamwork, Availability of training research courses, Lack of communication between academic, Mismatching between research theory and practice.	'Not reported'
Oliver et al. (2014) 2014 145 articles	"To identify new barriers of and facilitators to the use of evidence by policymakers"	A systematic review Narrative synthesis	Policymakers -	Meso-level Insufficient time at an organisation that assist professionals to use research findings Macro-level Lack of research methods skills among policymakers'	Meso-level Dissemination of research findings, Existence of and access to relevant research. Macro-level Collaboration between users and policymakers

Author, Study Year(s), Number of the included paper	Paper aim	Study design & analysis	Population countries	Main results	
				Barriers	Facilitators
Sadeghi-Bazargani et al. (2014) 2014 106 articles	"To systematically review and carry out an analysis on the barriers to evidence-based medicine."	A systematic review Narrative synthesis	HCPs -	Micro-level Lack of skills and understanding of statistic Meso-level Availability of equipment's and facilities to support research translation Lack of organisational support	'Not reported'
Légaré et al. (2008) 2008 38 articles	"To update a systematic review on the barriers and facilitators to implementing shared decision-making in clinical practice as perceived by health professionals."	A systematic review Narrative synthesis	HCPs -	Micro-level Lack of applicability due to patient characteristics. Meso-level Time constraints The clinical situation Lack of reimbursement Lack of resources	Micro-level HCPs motivation Meso-level Positive impact on the clinical process.
Lawrence et al. (2019) 2019 26 articles	"To report how decision-maker involvement in public health integrated knowledge translation research has been described and operationalised and whether the process was evaluated"	A scoping review Narrative synthesis	Policymakers -	Micro-level Low involvement of knowledge users Lack of expertise to present significant contributions Lack of motivation to update knowledge, Meso-level The usefulness of research to the knowledge users organisation Limited resource use Time constraints Lack of information sharing Macro-level No aligning research and policy considerations Conflicting priorities Others The language barrier, articles are written in English	Meso-level Budgeting for research activities (face-to-face meetings) Excellent interaction for sharing of the work Macro-level Identifying the right stakeholders Development of trust, Development mutual learning.

Author, Study Year(s), Number of the included paper	Paper aim	Study design & analysis	Population countries	Main results	
				Barriers	Facilitators
Derman and Jaeger (2018) 2018	"To overcome challenges to dissemination and implementation of research findings in under-resourced countries"	Narrative review Narrative synthesis	LMICs	Micro-level (individual) Inadequate communication and dissemination	Micro-level Consideration in advance of the audience likely to be interested in study findings. Meso-level Dissemination of the knowledge gained through our research Macro-level Engagement of stakeholders in the early stage of research process.
Athanasakis (2013) 2013 Literature review	"To review of nurses' research behaviour and the barriers that nurses meet in order to utilise research evidence into clinical nursing practice"	Narrative review Narrative synthesis	Nurses -	Micro-level Unaware of HCPs about research Unable HCPs to recognise the quality of research Mistrust of HCPs of the contributions of research findings. HCPs do not see the value of research for practice. Meso-level Not enough time to realise research findings Inadequate facilities for implementation Others The language barrier, articles are written in English	'Not reported'
Straus et al. (2011) 2011 Narrative review	"To provide an overview of the science and practice of knowledge translation."	Narrative review Narrative synthesis	- Global	Micro-level Professionals do have not sufficient knowledge and skills to appraise research. Low patients' adherence to recommendations, Meso-level Financial disincentives, lack of equipment Standards of care are not aligned with recommended practice.	'Not reported'

Author, Study Year(s), Number of the included paper	Paper aim	Study design & analysis	Population countries	Main results	
				Barriers	Facilitators
Kalassian et al. (2002) 2002 Literature review	To understand the challenges of translating research evidence into clinical in critical care	Narrative review Narrative synthesis	Clinicians -	<p>Micro-level Professionals do have not sufficient knowledge and skills to appraise research. No guidelines that inform research Salary and reimbursement</p> <p>Meso-level Financial incentives to promote practice guidelines may be expensive Unaffordable practice guidelines either by hospital administration or by key members of the health care system. Insufficient resources for implementing evidence-based guidelines. Struggle how to disseminate health care policies</p> <p>Macro-level Policymakers may be suspicious that there is an unacceptable increase in costs although evidence-based guidelines may improve health care.</p>	<p>Macro-level Formulating evidence-based guidelines “Developing and funding specific regional policies.”</p>

EBP= evidence-based practice; HCP= healthcare professionals; LMICs= low- and middle-income countries