

This is a repository copy of *Second-hand tobacco smoke exposure and smoke-free environments in Ethiopia: A scoping review and narrative synthesis*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/188831/>

Version: Published Version

Article:

Argefa, Terefe, Abraham Kassa, Selam and Mdege, Noreen Dadirai orcid.org/0000-0003-3189-3473 (2022) Second-hand tobacco smoke exposure and smoke-free environments in Ethiopia: A scoping review and narrative synthesis. *International Journal of Environmental Research and Public Health*. 8404. ISSN 1660-4601

<https://doi.org/10.3390/ijerph19148404>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Review

Second-Hand Tobacco Smoke Exposure and Smoke-Free Environments in Ethiopia: A Scoping Review and Narrative Synthesis

Terefe Gelibo Argefa ^{1,2}, Selam Abraham Kassa ² and Noreen Dadirai Mdege ^{2,3,4,*}

¹ ICAP in Ethiopia, Mailman School of Public Health, Columbia University, Addis Ababa P.O. Box 5566, Ethiopia; targerfa@developmentgateway.org

² Development Gateway, Inc., 1110 Vermont Ave NW Suite 500, Washington, DC 20005, USA; skassa@developmentgateway.org

³ Department of Health Sciences, University of York, York YO10 5DD, UK

⁴ Centre for Research in Health and Development, Amos Drive, York YO42 2BS, UK

* Correspondence: noreen.mdege@york.ac.uk

Abstract: Ethiopia passed a law prohibiting tobacco smoking in all public places in 2019. We conducted a scoping review to identify gaps in the existing literature on second-hand smoke (SHS) exposure and smoke-free environments in Ethiopia that need to be prioritised for future research to support policy and practice. We conducted systematic searches in January 2022 in the following databases: Medline, EMBASE, and PsycInfo. Two reviewers independently screened the identified study reports for eligibility and extracted data from the eligible studies. The extracted data was descriptively analysed, and research recommendations were drawn. A stakeholder consultation workshop was held to identify research topics on SHS exposure and smoke-free environments in Ethiopia that they perceived to be priorities for primary research. Of the 388 research reports identified, only nine were included in the scoping review. The topics explored includes prevalence of SHS exposure (six studies); knowledge on SHS exposure (three studies); compliance to smoke-free environments legislation (two studies); and exposure to anti-smoking messages (one study). The stakeholders prioritised further research addressing compliance monitoring and enforcement of the smoke free laws in Ethiopia. There is a need for studies that test new methods for compliance monitoring and enforcement, evaluate strategies to increase knowledge on the harms of SHS exposure and the smoke-free legislation, and evaluate the current smoke-free legislation in Ethiopia.

Keywords: smoking; tobacco smoke exposure; second-hand smoke; smoke-free environments



Citation: Argefa, T.G.; Kassa, S.A.; Mdege, N.D. Second-Hand Tobacco Smoke Exposure and Smoke-Free Environments in Ethiopia: A Scoping Review and Narrative Synthesis. *Int. J. Environ. Res. Public Health* **2022**, *19*, 8404. <https://doi.org/10.3390/ijerph19148404>

Academic Editors: Paul B. Tchounwou and Carla Berg

Received: 10 June 2022

Accepted: 7 July 2022

Published: 9 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Overall, 3.7% of adults in Ethiopia (6.2% of males, and 1.2% of females) smoke tobacco [1]. Exposure to second-hand tobacco smoke (SHS) results in 1.2 million deaths worldwide each year [2], with 47% and 28% of these deaths occurring in women and children, respectively [3]. Most of the deaths associated with SHS exposure are in low-income and middle-income countries (LMICs) [3]. Among adults in Ethiopia, 29.3% (6.5 million) of those who worked indoors reported that they were exposed to SHS in their workplace, and 12.6% (8.4 million) were exposed at home in 2016 [4]. In the same year, 60.4% of people visiting bars and nightclubs reported being exposed to SHS in these settings; and this was 31.1% in restaurants, 19.7% in government buildings, 11.4% in public transportation and 7.0% in health-care facilities [4]. SHS exposure among children is also high. In the last Global Youth Tobacco Survey (GYTS) among students aged between 13 and 15 years in Addis Ababa, about 29% lived in homes where others smoked in their presence; and over 40% were exposed to tobacco smoke in public places [5]. In a recent study among 1673 adolescents aged 13–19 years in Hawasa and Jimma town, 17% of non-smokers were exposed to SHS in their homes, and 61% were exposed in public places [6].

Ethiopia's Tobacco Control Strategic Plan 2010–2012 Ethiopian Calendar (E.C). (2017/18–2019/20) recognised SHS exposure as a public health threat, and made a commitment to enact and enforce smoke-free public places and environments [7]. In 2019, Ethiopia joined a group of 69 countries worldwide that provide protection from SHS exposure at best-practice level, with all public places completely smoke-free [8]. The country passed a law prohibiting smoking in all public places, i.e., any part of all indoor workplaces, all indoor public places, all means of public transport, and all common areas within condominium housings (Proclamation No. 1112/2019) [9]. In addition, smoking in any outdoor part of healthcare facilities, government institutions, facilities including schools intended mainly for those under the age of 21, higher education institutions, youth centres, and amusement parks is prohibited. The law also bans smoking or tobacco use in any indoor and outdoor space within 10 metres of any doorway, operable window, or air-intake mechanism of any public place or workplace [9]. Enforcement of the smoke-free regulations has been scaled-up through awareness raising and compliance inspections across the country: more than 16,000 inspections of a variety of public places across the country have been reported since 2019 [8].

Reducing SHS remains a top priority for the Ethiopian government [10]. Tobacco control stakeholders in Ethiopia, including government ministries, civil society organisations and academic institutions agree that there are some glaring gaps in evidence on the topic to inform policy and practice [10]. However, there has not been an attempt to bring together the currently existing evidence on SHS exposure and smoke-free environments in Ethiopia in order to identify specific research questions that need to be prioritised. We therefore conducted a scoping review and narrative synthesis aimed at examining the extent, range, and nature of research activity; summarising the findings from the studies identified; identifying gaps in the existing literature on SHS exposure and smoke-free environments in Ethiopia; and identifying research topics on SHS exposure and smoke-free environments in Ethiopia that need to be prioritised for primary research intended to support policy and practice.

2. Methods

2.1. Study Design

We utilised a well-established scoping review methodology comprising identifying relevant studies, selecting studies to include in the review, charting the data, collating, and summarising and reporting the results [11]. We also conducted a consultation exercise where we invited tobacco control stakeholders in Ethiopia who were not part of the review team to provide their insights and validate findings from the scoping review.

2.2. Study Identification

We conducted systematic searches in the following electronic databases using a comprehensive search strategy that was developed iteratively through a literature review and reviewing the results of initial scoping searches: Medline (1946 to 18 January 2022), EMBASE (1974 to 18 January 2021) and PsycInfo (1806 to January Week 2 2022). Our search strategy combined SHS exposure/smoke-free environment terms with Ethiopia (Supplementary Table S1). The following keywords were used for the searches: second-hand smoke, passive smoke, environmental exposure, environmental tobacco smoke, cigarette smoke, smoke-free, and Ethiopia. In addition to searching electronic databases, we also looked through the reference lists of included studies and consulted stakeholders and researchers working on the topic in Ethiopia to identify any study reports that we might have missed including grey literature. The stakeholders consulted included representatives from the Ministry of Health and other governmental and non-governmental institutions working on this topic in order to identify any relevant government reports or those from other stakeholders. No date, study design, publication type or language limits were imposed on the searches.

We followed the Cochrane and Centre for Reviews and Dissemination guidelines in designing searches [12,13]. The search strategies were reviewed by two reviewers

who were not involved in building the search using the Peer Review of Electronic Search Strategies checklist [14]. The searches were reported as per the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) literature search extension [15].

2.3. Study Selection

We used the Rayyan software (Rayyan—Intelligent Systematic Review) to manage and screen the research reports retrieved from the searches. Each study was independently screened for eligibility by two reviewers using a standardised, study selection form based on the prespecified study eligibility criteria. The form was piloted on five studies before use. The screening was done in two stages: screening of titles and abstracts of all retrieved records after removing duplicates; and then screening of the full texts of those studies identified in the first stage as potentially eligible. Any disagreements regarding study inclusion were resolved through discussion between the two reviewers.

Studies were eligible if they were carried out in Ethiopia anytime up to 18 January 2022, and addressed any aspect of SHS exposure, i.e., exposure to “the smoke emitted from the burning end of a cigarette or from other tobacco products usually in combination with the smoke exhaled by the smoker” [16], or smoke-free environments, i.e., environments where the air is 100% free from tobacco smoke (including, but is not limited to, air in which tobacco smoke cannot be seen, smelled, sensed or measured) [16] in Ethiopia. Studies that were not eligible if they were only available as an abstract.

2.4. Data Charting

A data charting form developed in Microsoft Excel (Office 2019) specifically for the review and based on the data domains and variables listed in Table 1 was used to chart the data from the included studies. The data extraction tables were accompanied by instructions and decision rules for coding data in order to increase consistency, reduce bias from subjective judgement and improve the validity and reliability of the process [13,17]. One researcher read each included study in full, putting any data that is relevant to the review under the appropriate data domains. For studies that were reported in more than one publication, we charted the data from each publication separately and then combined the data. A second reviewer checked the charted data for accuracy and quality. We also extracted main study results as reported in each study. Disagreements between reviewers were resolved through discussion and consensus.

Table 1. Data extraction domains and variables.

Domain	Variable
General study characteristics	Author, year of publication, geographic location (i.e., administrative district), study setting (e.g., school, health care facility, bar, restaurant, hotel, etc.), study design, sample size
Principal focus	Topic (e.g., exposure, enforcement, compliance, awareness, etc.), research question(s), type(s) of intervention(s) if applicable
Study population characteristics	Age, gender, ethnicity, socio-economic status

2.5. Data Collating, Summarising and Reporting

We conducted content analysis to identify and categorise various data items, and produced tables, graphs and narrative summaries that mapped the extent, range, and nature of research activity according to the domains and variables described in Table 1. We used these summaries to identify the key gaps in the existing literature and make recommendations for future research. We also narratively synthesised study findings according to their study focus areas. The reporting of the scoping review followed the PRISMA extension for scoping reviews [18].

2.6. Stakeholder Consultations

We held an online, one and a half hour, stakeholder consultation workshop to share and validate our findings. Specifically, we utilised this stage to identify, from the perspective of the stakeholders, after they had been presented with the scoping review results, what the priority research topics on SHS exposure and smoke-free environments in Ethiopia are. First, we presented the objectives, methods and results of the scoping review, and a preliminary list of recommendations that we had drawn from this literature (15 min). The stakeholders were given a chance to ask questions and seek clarification on any aspect of the presentation; and to discuss the recommendations amongst themselves, adding and removing recommendations from the preliminary list as they saw fit (30 min). We did not prescribe specific criteria for the discussions, but requested them to include the following as part of their considerations: the importance of the topic for policy making in Ethiopia; alignment with the government priorities; and the feasibility of conducting research on the topic, for example, in light of the current political climate and other resource constraints. After finalising the research recommendations list, the stakeholders were given one to two minutes to independently vote for their first choice in real time through a poll created on Zoom. The invited stakeholder organisations included academic and research institutions, civil society organisations, the World Health Organisation country office, the Ethiopia Food and Drug Administration, and relevant departments in the Ministry of Health. The list is provided in Supplementary Table S2.

3. Results

3.1. Search Results

The searches yielded 384 records and 366 remained after duplicates were removed. Of the 366 articles, the five studies that were eligible for full-text review were all found to be eligible for inclusion. Three additional eligible studies were identified from the reference list searches, and one more study was identified through stakeholder consultations. Therefore, a total of nine studies were included in the scoping review (Figure 1) [6,19–26].




Figure 1. Flow of studies in the review process.

3.2. General Characteristics of Studies

All of the included studies were cross-sectional surveys (Table 2) [6,19–26]. Four studies were conducted in Southern Ethiopia [6,22–24], whilst there were three studies in Central Ethiopia [19,26], and one each for East [25], West [21] and South West Ethiopia [6]. One study used data from a nationwide survey covering all 11 administrative districts [20]. Of the nine studies, there were two studies each for Addis Ababa [19,26], Alata Wondo [23,24] and Hawasa [6,22], and one each for Jimma [6], Nekemte [21] and Kersa towns [25]. Four of the studies were household level surveys [20,23–25], whilst two were conducted in schools [6,21], and one was at a university [22]. The remaining study covered a number of settings: food and beverage providing facilities; health care facilities; schools; youth centres; and government offices [19].

Table 2. General study characteristics.

Author, Year	Geographical Region	City/Town	Study Setting	Sample Size
Ababulgu et al., 2016 [6]	Southern and South West	Hawassa and Jimma towns	Schools	1704
Bekalu et al., 2022 [20]	Nation wide	All the 11 administrative regions	Households	10,150
Bobo et al., 2018 [21]	Western	Nekemte town	Schools	1088
Ethiopia Food and Drug Authority, 2021 [19]	Central	Addis Ababa (Bole and Arada sub-cities)	Food and beverage providing facilities; health care facilities; schools; youth centres; and government offices	221 (103 Food and Beverage Service Providers, 50 health service providers, 30 schools, 12 Youth Centres and 26 government offices included)
Kassa and Deyno, 2014 [22]	Southern	Hawassa town	University	590
Petersen et al., 2016 [23]	Southern	Alata Wondo town	Households	372
Petersen et al., 2018 [24]	Southern	Alata Wondo town	Households	372
Reda et al., 2013 [25]	Eastern	Kersa town	Households	600
Tadesse and Zawdie, 2019 [26]	Central Area	Addis Ababa	Hospitals	354

Sample sizes ranged from 221 to 10,150. All of the included studies were published between 2013 and 2022. Eight of the nine studies used data that was collected sometime between 2010 and 2018 [6,20–26], which is before Ethiopia’s current federal tobacco control legislation (Proclamation 1112/2019) [9], and the 2021 Tobacco Control Directive No. 771/2021 [27]. The remaining study collected data in 2021 in Addis Ababa, but this was before launching an initiative to make the city smoke-free [19].

3.3. Study Focus

Overall, six of the nine studies reported prevalence of SHS exposure places or home [6,21–25] one explored exposure to anti-tobacco messages [20]; three reported on knowledge on SHS exposure [20,24,25]; and two reported on compliance with smoke-free environments laws/regulations [19,26].

3.4. Study Population Characteristics

Two of the studies were conducted among school adolescents aged between 13 and 19 years (Table 3) [6,21]. The rest were conducted in adults: two among those ≥ 15 years of age [6,21]; two among those from 18 to 55 years of age [23,24]; one among university students aged from 20 to 24 years [22]; and one among hospital staff [26]. The study among hospital staff also conducted observations of 15 buildings located in the hospital premises of five hospitals that had started the implementation of smoke-free legislation [26]. Two of the adult sample studies had all female samples [23,24]. One study collected data from

food and beverage providing premises, health service premises, schools, youth centres and government offices through observation [19].

Table 3. Study population characteristics.

Author, Year	Population Group	Age Range Mean Age (Standard Deviation)	Gender
Ababulgu et al., 2016 [6]	Adolescents enrolled in grade 9 to 12	13–19 years Not reported	47.7% male; 52.3% female
Bekalu et al., 2022 [20]	Adults	≥15 years 31.2 (14.8) years	49.9% male; 50.1% female
Bobo et al., 2018 [21]	Adolescents attending high school	14–19 years Not reported	57.9% male; 42.1% female
Ethiopia Food and Drug Authority, 2021 [19]	Food, beverage and health service providers, schools, youth centres and government offices	-	-
Kassa and Deyno, 2014 [22]	Adults (University students)	20–24 years 20.7 (1.49) years	81.7% male; 18.3% female
Petersen et al., 2016 [23]	Adults	18–55 years 29.4 (6.9) years	100% female
Petersen et al., 2018 [24]	Adults	18–55 years 29.4 (6.9) years	100% female
Reda et al., 2013 [25]	Adults	≥15 years 35 (15.0) years	75.1% male; 24.9% female
Tadesse and Zawdie, 2019 [26]	Adults (hospital staff)	29 years * (Interquartile range 26 to 35 years)	54.3% male; 45.7% female

* Median.

3.5. Summary of Study Findings

The study results are summarised below by study focus areas.

3.5.1. Prevalence of SHS Exposure

Exposure among young people: The prevalence of self-reported SHS exposure in public places among adolescents attending school was at least 60% [6,21]. Self-reported exposure at home was also high, ranging from 17% to 38% of participants [6,21]. In the study among university students, 69% of participants reported exposure within university campuses, with 17% reporting daily exposure [22]. Exposure was in almost all areas within the campuses, including in the university compound, dormitories, toilets, lounges, cafes and class rooms [22]. Petersen et al. found that in 5.1% of households young children were frequently exposed to SHS in the home [23].

In the study by Ababulgu et al., the following significantly increased the odds (three times) of being exposed to SHS at home: being female when compared to being male (Odds Ratio (OR) 3.46; 95% CI: 2.62–4.57), having parents (OR 3.34; 95% CI: 2.37–5.03) or closest friends (OR 3.61; 95% CI: 2.41–5.41) who smoke compared to those who do not [6]. The following significantly increased the odds of SHS exposure in public places: having a peer who smokes (OR 3.76; 95% CI: 2.49–5.65) compared to those who did not, and not being informed on the danger of health effects of tobacco smoke (OR 5.32; 95% CI 4.13–6.81) [6].

Exposure among adults: In two studies, 15% [23] and 40% [25] of households reported that smoking was allowed in the home. In one of these studies, an additional 12% of households allowed smoking in the home with exceptions, and 17% did not have rules on second hand smoke [25]. In the study by Petersen et al., 14.4% of participants reported that smoking took place daily in their homes [23,24], whilst this was 33% in the study

by Reda et al. [25]. Petersen et al. reported that being an urban residents increased the likelihood to report that smoking was allowed in the home, or that SHS occurred daily in their household [23]. In addition, having a house member who is a current tobacco user, recent exposure to point-of-sale advertising, and living in a household where smoking is allowed in the home increased the likelihood of reporting daily occurrence of SHS in the home [23].

3.5.2. Knowledge on SHS Exposure

Reda et al. [25] and Petersen et al. [24] both reported high proportions of respondents who thought/believed that SHS exposure was harmful (73% and 98%, respectively). Nevertheless, knowledge on specific health conditions was high among participants in the study by Reda et al. than Petersen et al.: 82% versus 8% for heart disease, 90% versus 20% for respiratory problems; and 84% versus 5% for lung cancer, respectively. These two studies were conducted in different towns, and one was 75% male [25], while the other was 100% female [24].

In the study by Bekalu et al. [20], although the risk perception was high overall, the proportion reporting that SHS exposure causes serious illness were higher in those who were more educated, wealthier, and those who were urban residents.

3.5.3. Compliance with Smoke-Free Environments Laws/Regulations

In the study by Tadesse and Zawdie, only 2.8% of the inspected places in the five hospitals had anti-smoking signs, and this was more likely to be at entrances than other places [26]. Cigarette butts were observed in all hospitals and these were more likely to be found at the entrances (24%) and toilets (15%) than other places [26]. Actual smoking on hospital premises was observed in three of the five hospitals. Overall, 42% of hospital staff were aware of the hospital smoke-free legislation, whilst 34% were not aware and 24% were not sure that there was a smoke-free legislation that prohibited smoking within hospital premises. A small proportion of hospital staff (10%) reported not being compliant with smoke free legislation—the levels of knowledge about smoke-free legislation was poor among 69% of these staff, and 83% had unfavourable attitudes towards the legislation. In contrast, 42% of compliant hospital staff had poor knowledge of the legislation, and 47% had unfavourable attitudes towards it. Additionally, 28% of the hospital staff indicated that they had been enforcing the smoke-free legislation [26].

In the report by the Ethiopia Food and Drug Authority of their compliance monitoring exercise conducted in 2021 [19], overall compliance for the Bole and Arada sub-cities was low: 33% (4/12) for youth centres, 31% (8/26) for government offices, 54% (27/50) for health facilities, 45% (13/30) for educational facilities and 25% (26/103) for food and beverage venues [23]. Compliance was much higher in Arada when compared to Bole for most settings: 46% versus 16% for government offices, 67% versus 42% for health facilities and 68% versus 20% for educational facilities. Compliance was similar in food and beverage venues, 23% for Arada and 26% for Bole sub-city; and higher for Bole for youth centres (40% versus 0%). A number of challenges with compliance were reported including fear of conflict with clients, lack of client awareness of the legislation and cigarette sales within 100 m for the establishments. In addition, for food and beverage venues, there was a fear of losing customers, with foreign customers being treated differently (i.e., being allowed to smoke) to local ones. For schools, there were reports of people smoking along the school fence and also the presence of shisha houses in surrounding areas causing problems for compliance. Designated smoking areas were present in a few of the food and beverage venues and youth centres [19].

3.5.4. Exposure to Anti-Smoking Messages

In the study by Bekalu and colleagues, the association between exposure to anti-smoking messages and risk perception of SHS exposure was not significant (OR 1.26; 95% CI: 0.64–2.46) [20].

3.6. Stakeholder Consultation Results

After analysing and synthesising the results, two reviewers developed a preliminary list of research recommendations to be used as a starting point for the discussions with the stakeholders (Table 4). The stakeholder consultation workshop was attended by 11 participants representing nine stakeholder organisations: two represented the Ministry of Health, one represented the Ethiopia Food and Drug Administration, three represented academic institutions (one each for the Addis Ababa University, Department of preventive medicine; Hawassa University, College of Medicine and Health Sciences; and Jimma University, College of Health Sciences & Medicine) and five participants were from CSOs (two for Mathewos Wondu Ethiopia Cancer Society, and one each for the Health Development and Anti Malaria Association, MeQuamia Community Development Organization, and Campaign for Tobacco Free Kids).

Table 4. Research recommendations drawn from the literature.

Topic	Potential Research Objectives/Questions
Compliance monitoring and enforcement	Evaluation of the feasibility, acceptability, effectiveness, cost-effectiveness of alternative methods of compliance monitoring and enforcement, e.g., objective measures with visual aids that can be used to provide real-time feedback on air quality in a venue, including the implications of the recorded levels to health.
Compliance challenges	Investigate the impact of implementing smoke-free policies in hospitality venues on revenue. Develop and test strategies to overcome the other challenges such as cigarette sales within 100 m for the establishments and smoking within the vicinity of educational establishments.
Knowledge of the harms of SHS exposure	Develop and test the effectiveness and cost-effectiveness of different strategies for increasing knowledge of the harms of SHS exposure.
Awareness of the smoke free law	Develop and test the effectiveness and cost-effectiveness of different strategies for increasing Awareness of the smoke free law.
Exposure in educational facilities	Develop and test the effectiveness and cost-effectiveness of different strategies for decreasing exposure in educational facilities.
Impact of the smoke-free law	Evaluation of the impact of smoke-free laws using evaluative designs such as before and after comparisons or natural experiments where the data allows.

The stakeholders added '*Reducing levels of SHS exposure within households and communities*' to the list of research recommendations. Eight out of the eleven stakeholders cast their votes for their preferred research priority in the poll. The recommendation with the majority of votes was compliance monitoring and enforcement, specifically evaluating the feasibility, acceptability, effectiveness, cost-effectiveness of alternative methods of compliance monitoring and enforcement (Figure 2). There were no votes for compliance challenges and developing and testing the effectiveness and cost-effectiveness of different strategies for awareness among those who should comply as well as the general population. These topics are therefore not represented in Figure 2.




Figure 2. Stakeholder poll results.

4. Discussion

This scoping review highlights the scarcity of research on SHS exposure and smoke-free environments in Ethiopia. Of the nine identified studies, most (six) report on exposure to SHS either in the home or at academic institutions (i.e., schools and universities), or knowledge of harms due to SHS (three). Exposure in the other indoor public places, indoor workplaces and outdoor public places that are covered by the Ethiopian legislation have not been investigated in detail, except through the Global Adult Tobacco Survey which was last conducted in 2016 [4] and the GYTS conducted in 2003 [5]. Reports on the levels of compliance to legislation are scarce: only two were identified. One of these studies covered a wide range of settings, i.e., youth centres, government offices, health facilities, educational institutions and food and beverage venues, albeit only in two sub-cities in Addis Ababa [19]. All of the identified studies used data that was collected either before Ethiopia's current federal tobacco control legislation (Proclamation 1112/2019) [9], and the 2021 Tobacco Control Directive No. 771/2021 [27], or before local implementation of smoke-free initiatives.

Studies included in this review reported high levels of SHS exposure in the home among adolescents and adults, and at universities among young people. Being female, having members of households who smoke, living in urban areas and poor knowledge of the dangers of SHS exposure significantly increased the likelihood of SHS exposure in these settings. In addition, living in a household where smoking inside the home was permitted increased the likelihood of SHS exposure within the home. These findings are in line with prior research that has shown that women are worst affected by the SHS exposure disease burden, with 47% of deaths from SHS exposure occurring in female adults [3]. A recent analysis of SHS exposure among adolescents aged 12–15 years across 68 LMICs also reported a positive association between SHS exposure and parental smoking, with the association being larger for maternal smoking than paternal smoking [28]. Another study in The Gambia also found that lower education attainment, lower household income, urban residence, and not being aware of smoke-free regulations increased the risk of SHS exposure [29].

The studies included in the review found low levels of compliance with smoke-free laws in different establishments. In all of the studies, lack of knowledge of the legislation among staff working in those establishments as well as their clients seemed to be one of the causes of non-compliance and negative attitudes towards the legislation. The presence of cigarette sales within the vicinity, fear of conflict with clients, and fear of losing clients and revenue (in hospitality venues) also seem to play a part in non-compliance. Sub-optimal levels of compliance with smoke-free policies within hospitality venues have also been observed in other African countries [30–32]. In Ghana and Uganda, poor knowledge of the smoke-free legislation among hospitality venue staff was also reported [31,33]. Nevertheless, in a number of countries, including in Nigeria and Ghana, the attitude towards smoke-free laws among hospitality staff has been found to be generally positive [31,34–36], although the fear of revenue reductions seems to be a consistent finding within these venues [35,36].

The studies suggest that there could be demographic as well as geographical differences in both knowledge of the harms of SHS exposure and compliance to legislation. The study by Bekalu et al. observed a social gradient on knowledge, with the proportion reporting that SHS exposure caused serious illness being higher in those who were more educated and wealthier [20]. This proportion was also higher in urban than rural residents [20]. Furthermore, the proportion of participants with good knowledge of specific conditions that are attributable to SHS exposure was much higher in the study by Reda et al., which was conducted in Kersa town in a sample with 75% men [25], when compared to those reported by Petersen et al. in a study conducted in Alato town in an all-female sample [23].

In addition to the identification of research priorities, the findings from our stakeholder consultation meetings also highlighted a spread on opinions on what the research priorities should be. This could be as a result of the different interests and focus areas among the stakeholders. We incorporated stakeholder discussion on research priorities in order to facilitate understanding of each other's perspectives, the consideration of those different perspectives, and broad acceptance of the process and outcomes [37].

Reducing SHS exposure and increasing compliance to smoke-free laws remain a priority for Ethiopia. Increasing the knowledge and awareness of the harms of SHS exposure and the smoke-free laws among the public and staff working at different venues that need to be smoke-free seems to be important for reducing SHS exposure as well as increasing compliance. It could potentially reduce conflicts between staff within establishments and their clients and improve attitudes towards smoke-free legislation. Bekalu et al. suggests that mass media communication as a potential strategy for increasing knowledge and awareness of these issues [20]. Nevertheless, such a strategy would need to account for patterns of mass media use in Ethiopia, for example the types of mass media platforms commonly used and any sociodemographic (e.g., gender and age) or geographic (e.g., urban versus rural) differences that might exist in the access and use of those platforms [38–40].

The impact of the current smoke-free legislation on SHS exposure and compliance needs to be evaluated, for example using before and after comparisons where data allows. The current phased implementation of the legislation also offers an opportunity for natural experiments on the impact of the legislation on both exposure and compliance. Continuous evaluation of tobacco control policies could identify the gaps and promote the development of new strategies for improving compliance with the laws [41]. There is also a need to develop and test the effectiveness and cost-effectiveness of different strategies of increasing knowledge of the harms of SHS exposure and on the smoke-free laws. In addition to the use of mass media, other strategies could include workplace-, educational facility-, and community-based interventions. For example, educational facility-based interventions targeting children and young people, including teaching them to negotiate for smoke-free environments, or community-based interventions for example via faith-based institutions hold promise, although most of studies exploring them have mainly been in the context of exposure within the home [42–44]. The concerns around loss of revenue in hospitality

settings as a result of implementing smoke-free policies, need to be explored further. In addition, research should develop and test strategies to overcome the other challenges raised by staff members at different establishments such as cigarette sales within 100 m of the establishments and smoking within the vicinity of educational establishments. Another potential area of research is the evaluation of the feasibility, acceptability, effectiveness, cost-effectiveness of alternative methods of compliance monitoring and enforcement.

Compliance monitoring in Ethiopia and other countries worldwide is currently mainly through visual inspection of a venue using standard compliance indicators such as observed smoking; presence of designated smoking areas, ashtrays, no smoking signs, cigarette butts or the smell of tobacco smoke at the venues [45]. This method is attractive because it is relatively simple and inexpensive [45]. However, some studies have highlighted the potential usefulness of low-cost air quality monitors, such as those that measure the concentration of airborne particulate matter less than 2.5 microns (PM_{2.5}) in diameter. For example, in a study in Uganda, although the occurrence of at least one observational indicators of smoking within the 222 hospitality venues was in less than 50% of the venues, the average PM_{2.5} for a subsample of 108 study sites was 171 µg/m³, representing an unhealthy air quality rating [30]. The average rating in indoor venues indicated hazardous levels (PM_{2.5} concentration of 267.64 µg/m³), whilst outdoor venues had an average concentration of 85.60 µg/m³ indicating unhealthy air quality [30]. The results from these monitors can also be translated into visual aids that can be used to provide real-time feedback on air quality in a venue, including the implications of the recorded levels to health: this has been shown to be an effective educational and behaviour change motivational tool for reducing SHS exposure [46]. The majority of tobacco control stakeholders in Ethiopia who attended our workshop in fact voted this topic as their top priority.

Limitations of the Study

We used a comprehensive list of search terms. The searches were conducted in three of the main bibliographic databases for health-related research. Although attempts were made to identify grey literature through contacting key stakeholders, it is still possible that some of this literature was missed. Almost all the included studies were conducted before the implementation of Tobacco Control Directive; therefore, the findings may not be indicative of the implementation of the provisions of the current smoke free laws in Ethiopia. We gave stakeholders up to two minutes to independently vote for their top research priorities in real time during the meeting. This might have limited their ability to reflect on the discussions and gather more information to enable them to fully consider their choices. This approach however enabled us to maximise the number of responses obtained.

5. Conclusions

Our scoping review highlighted the scarcity of research on SHS exposure and smoke-free environments in Ethiopia. As prioritised by the stakeholders, further research is needed to address compliance monitoring and enforcement, particularly evaluating the feasibility, acceptability, effectiveness, cost-effectiveness of alternative methods of compliance monitoring and enforcement (e.g., objective measures with visual aids that can be used to provide real-time feedback on air quality in a venue), including the implications of the recorded levels to health.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijerph19148404/s1>, Table S1: Search strategies. Table S2: List of invited stakeholder organizations.

Author Contributions: N.D.M., T.G.A. and S.A.K. conceptualized the study; N.D.M. and T.G.A. designed the study methodology; N.D.M., T.G.A. and S.A.K. collected and analysed the data; N.D.M. was responsible for oversight and supervision; T.G.A. and N.D.M. prepared the first draft of the manuscript; all authors reviewed and edited the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This publication is based on research funded by the Bill & Melinda Gates Foundation (grant number INV-009670). The findings and conclusions contained within are those of the authors and do not necessarily reflect positions or policies of the Bill & Melinda Gates Foundation.

Institutional Review Board Statement: Ethical approval is not applicable for this study as this is a review and doesn't involve humans or animals.

Informed Consent Statement: Informed consent is not applicable for this study as this is a review and doesn't involve humans or animals.

Data Availability Statement: The data supporting the findings of this review are available within the paper and its supplementary materials.

Acknowledgments: This scoping review was conducted as part of the Tobacco Control Data Initiative (TCDI). TCDI is a program that covers six African countries (the Democratic Republic of Congo, Ethiopia, Kenya, Nigeria, South Africa, and Zambia) and aims to understand their tobacco data needs, identify existing data, confirm gaps in available tobacco data, collect new data to fill those gaps, and develop tools to enable policymakers to use essential data more effectively to inform tobacco policy. The programme is led by Development Gateway: an IREX Venture (DG), a global non-profit organization that specializes in data for development, in partnership with the University of Cape Town's Research Unit on the Economics of Excisable Products (REEP). We would like to extend our gratitude to the following DG Ethiopia country team members for their guidance, administrative & review support throughout the study: Rachel Kitonyo-Devotsu, Winnie Awuor and Andrea Ulrich. In addition, we would like to express our sincere gratitude to all stakeholders who participated in our stakeholder consultation meeting and their organisations.

Conflicts of Interest: The Bill & Melinda Gates Foundation had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results. The authors declare that they have no conflicts of interest related to publication of the manuscript.

References

1. Mengesha, S.D.; Teklu, K.T.; Weldetinsae, A.; Serte, M.G.; Kenea, M.A.; Dinssa, D.A.; Woldegabriel, M.G.; Alemayehu, T.A.; Belay, W.M. Tobacco use prevalence and its determinate factor in Ethiopia- finding of the 2016 Ethiopian GATS. *BMC Public Health* **2022**, *22*, 555. [CrossRef] [PubMed]
2. World Health Organization (WHO). Health Topics: Tobacco. 2022. Available online: <https://www.who.int/news-room/fact-sheets/detail/tobacco> (accessed on 5 January 2022).
3. Öberg, M.; Jaakkola, M.S.; Woodward, A.; Peruga, A.; Prüss-Ustün, A. Worldwide burden of disease from exposure to second-hand smoke: A retrospective analysis of data from 192 countries. *Lancet* **2011**, *377*, 139–146. [CrossRef]
4. Ethiopia Public Health Institute (EPHI); U.S. Centers for Disease Control and Prevention (CDC); World Health Organization (WHO). Global Adult Tobacco Survey (GATS) Ethiopia: Executive Summary 2016. In *Adult, Global Survey, Tobacco Summary, Executive*; 2016. Available online: <http://repository.iifphc.org/handle/123456789/469> (accessed on 5 January 2022).
5. World Health Organization (WHO). Ethiopia—Addis Ababa (Ages 13–15) Global Youth Tobacco Survey (GYTS) Fact Sheet; Ethiopia, 2003; p. 1. Available online: <https://www.afro.who.int/sites/default/files/GYTS/ethiopia.pdf> (accessed on 5 January 2022).
6. Ababulgu, S.A.; Dereje, N.; Girma, A. Secondhand tobacco smoke exposure among adolescents in an Ethiopian school. *Health Low-Resour. Settings* **2016**, *4*, 1–4. [CrossRef]
7. Food Medicine and Health Care Administration Control Authority of Ethiopia (FMHACA). Ethiopia Tobacco Control Strategic Plan 2010–2012 Ethiopian Calendar (2017/18–2019/2020). 31 October 2017; p. 1–123. Available online: <http://www.fmhaca.gov.et/wp-content/uploads/2019/03/EthiopiaTobacco-Control-Strategic-Plan.pdf> (accessed on 5 January 2022).
8. World Health Organization (WHO). WHO Report on the Global Tobacco Epidemic 2021: Addressing New and Emerging Products. In *Health Promotion*. 2021; pp. 1–212. Available online: <https://www.who.int/publications/i/item/9789240032095> (accessed on 5 January 2022).

9. Federal Democratic Republic of Ethiopia (FDRE). Federal Nagarit Gazette Proclamation Number 1112/2019. 2019; pp. 2–174. Available online: <http://www.fmhaca.gov.et/wp-content/uploads/2020/06/Food-and-Medicine-Administration-Proclamation-1112.pdf> (accessed on 5 January 2022).
10. Development Gateway Inc (DG). *The Tobacco Control Data Initiative program: Ethiopia Assessment Report*; Development Gateway, Inc.: Washington, DC, USA, 2022.
11. Arksey, H.; O'Malley, L. Scoping studies: Towards a methodological framework. *Int. J. Soc. Res. Methodol.* **2005**, *8*, 19–32. [[CrossRef](#)]
12. Higgins, J.P.T.; Lasserson, T.; Chandler, J.; Tovey, D.; Churchill, R. *Methodological Expectations of Cochrane Intervention Reviews (MECIR) Standards for the Conduct and Reporting of New Cochrane Intervention Reviews, Reporting of Protocols and the Planning, Conduct and Reporting of Updates*; Trusted evidence; Informed decisions; Better health; Cochrane: London, UK, 2017.
13. Centre for Reviews and Dissemination(CRD). Systematic reviews: CRD's guidance for undertaking reviews in health care. In *CRD's Guidance for Undertaking Reviews in Health Care*; York Publishing Services Ltd.: York, UK, 2009; pp. 1–194.
14. McGowan, J.; Sampson, M.; Salzwedel, D.M.; Cogo, E.; Foerster, V.; Lefebvre, C. PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement. *J. Clin. Epidemiol.* **2016**, *75*, 40–46. [[CrossRef](#)] [[PubMed](#)]
15. Rethlefsen, M.L.; Kirtley, S.; Waffenschmidt, S.; Ayala, A.P.; Moher, D.; Page, M.J.; Koffel, J.B.; PRISMA-S Group. PRISMA-S: An extension to the PRISMA Statement for Reporting Literature Searches in Systematic Reviews. *Syst. Rev.* **2021**, *10*, 39. [[CrossRef](#)]
16. World Health Organization (WHO). Protection from exposure to second-hand tobacco smoke: Policy recommendations. In *Policy Recommendations*; World Health Organization: Geneva, Switzerland, 2007.
17. Higgins, J.P.T.; Altman, D.G.; Sterne, J.A.C.; The Cochrane Bias Methods Group. Chapter 8: Assessing Risk of Bias in Included Studies; Part 2: General Methods for Cochrane Reviews > 8 Assessing Risk of Bias in Included Studies. 2011. Available online: https://handbook-5-1.cochrane.org/chapter_8/8_assessing_risk_of_bias_in_included_studies.htm (accessed on 5 January 2022).
18. Tricco, A.C.; Lillie, E.; Zarin, W.; O'Brien, K.K.; Colquhoun, H.; Levac, D.; Moher, D.; Peters, M.D.J.; Horsley, T.; Weeks, L.; et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann. Intern. Med.* **2018**, *169*, 467–473. [[CrossRef](#)]
19. Ethiopia Food and Drug Authority (EFDA). *Addis Ababa Smoke Free Program Assessment in Bole and Arada Sub-Cities*; Ethiopia Food and Drug Authority (EFDA): Addis Ababa, Ethiopia, 2021.
20. Bekalu, M.A.; Gundersen, D.A.; Viswanath, K. Beyond Educating the Masses: The Role of Public Health Communication in Addressing Socioeconomic- and Residence-based Disparities in Tobacco Risk Perception. *Health Commun.* **2020**, *37*, 214–221. [[CrossRef](#)]
21. Bobo, F.; Thanasekaran, P.; Joice, A.J.R.; Yadecha, B.; Alebel, A. Susceptibility to cigarette smoking and associated factors among high school students in western Ethiopia. *BMC Res. Notes* **2018**, *11*, 626. [[CrossRef](#)]
22. Kassa, A.; Deyno, S. Prevalence and Determinants of Active and Passive Cigarette Smoking among undergraduate students at Hawassa University, Hawassa, Ethiopia. *J. Trop. Dis.* **2014**, *2*, 2. [[CrossRef](#)]
23. Petersen, A.B.; Thompson, L.M.; Dadi, G.B.; Tolcha, A.; Cataldo, J.K. Factors associated with secondhand tobacco smoke in the home: An exploratory cross-sectional study among women in Aleta Wondo, Ethiopia. *BMC Public Health* **2016**, *16*, 910. [[CrossRef](#)] [[PubMed](#)]
24. Petersen, A.B.; Thompson, L.M.; Dadi, G.B.; Tolcha, A.; Cataldo, J.K. An exploratory study of knowledge, attitudes, and beliefs related to tobacco use and secondhand smoke among women in Aleta Wondo, Ethiopia. *BMC Women's Health* **2018**, *18*, 154. [[CrossRef](#)] [[PubMed](#)]
25. Reda, A.A.; Kotz, D.; Biadgilign, S. Adult tobacco use practice and its correlates in eastern Ethiopia: A cross-sectional study. *Harm Reduct. J.* **2013**, *10*, 28. [[CrossRef](#)] [[PubMed](#)]
26. Tadesse, T.; Zawdie, B. Non-compliance and associated factors against smoke-free legislation among health care staffs in governmental hospitals in Addis Ababa, Ethiopia: An observational cross-sectional study. *BMC Public Health* **2019**, *19*, 91. [[CrossRef](#)] [[PubMed](#)]
27. Ethiopian Food Medicine and Healthcare Administration and Control Authority (FMHACA). Tobacco Control Directive Number 771/2021. 2021; pp. 1–21. Available online: <http://www.fmhaca.gov.et/wp-content/uploads/2021/04/Tobacco-Control-Directive-Number-771-2021.pdf> (accessed on 5 January 2022).
28. Xi, B.; Liang, Y.; Liu, Y.; Yan, Y.; Zhao, M.; Ma, C.; Bovet, P. Tobacco use and second-hand smoke exposure in young adolescents aged 12–15 years: Data from 68 low-income and middle-income countries. *Lancet Glob. Health* **2016**, *4*, e795–e805. [[CrossRef](#)]
29. Cham, B.; Mdege, N.; Bauld, L.; Britton, J.; D'Alessandro, U. Exposure to Second-Hand Smoke in Public Places and Barriers to the Implementation of Smoke-Free Regulations in The Gambia: A Population-Based Survey. *Int. J. Environ. Res. Public Health* **2021**, *18*, 6263. [[CrossRef](#)]
30. Gravely, S.; Nyamurungi, K.N.; Kabwama, S.N.; Okello, G.; Robertson, L.; Heng, K.K.C.; Ndikum, A.E.; Oginni, A.S.; Rusatira, J.C.; Kakoulides, S.; et al. Knowledge, opinions and compliance related to the 100% smoke-free law in hospitality venues in Kampala, Uganda: Cross-sectional results from the KOMPLY Project. *BMJ Open* **2018**, *8*, e017601. [[CrossRef](#)]
31. Singh, A.; Dobbie, F.; Kinnunen, T.I.; Okello, G.; Semple, S.; Okyere, P.B.; Logo, D.D.; Lartey, K.F.; McNeil, A.; Britton, J.; et al. Adherence to smoke-free policies in Ghana: Findings from a cross-sectional survey of hospitality venue owners and staff. *Tob. Prev. Cessat.* **2021**, *7*, 4. [[CrossRef](#)]

32. Karimi, K.J.; Ayah, R.; Olewe, T. Adherence to the Tobacco Control Act, 2007: Presence of a workplace policy on tobacco use in bars and restaurants in Nairobi, Kenya. *BMJ Open* **2016**, *6*, e012526. [[CrossRef](#)]
33. Robertson, L.; Nyamurungi, K.N.; Gravely, S.; Rusatira, J.C.; Oginni, A.; Kabwama, S.N.; Ndikum, A.E.; Bianco, E.; Yusuf, S.; Huffman, M.D. Implementation of 100% smoke-free law in Uganda: A qualitative study exploring civil society's perspective. *BMC Public Health* **2018**, *18*, 927. [[CrossRef](#)]
34. Aherrera, A.; Çarkoğlu, A.; Hayran, M.; Ergör, G.; Eirüder, T.; Kaplan, B.; Susan, J.; Zheng, L.; Cohen, J.E.; Navas-Acien, A. Factors that influence attitude and enforcement of the smoke-free law in Turkey: A survey of hospitality venue owners and employees. *Tob. Control* **2016**, *26*, 540–547. [[CrossRef](#)]
35. Alaaeddine, G.; Al Kuhaimi, T.; Al Assaad, R.; Dany, M.; Diab, R.; Hanna, E.; Hirmas, N.; Ismail, H.; Mahmassani, D.; Tellawi, R.S.; et al. Assessing knowledge and attitudes of owners or managers of hospitality venues regarding a policy banning indoor smoking. *Public Health* **2013**, *127*, 461–466. [[CrossRef](#)] [[PubMed](#)]
36. Odukoya, O.O.; Ohanusi, U.; Olokodana, B. Are hospitality venue employees aware and do they support the State-wide Regulation of Smoking Law in Lagos state Nigeria? *Tob. Prev. Cessat.* **2016**, *2*, 68. [[CrossRef](#)]
37. Grill, C. Involving stakeholders in research priority setting: A scoping review. *Res. Involv. Engag.* **2021**, *7*, 75. [[CrossRef](#)] [[PubMed](#)]
38. Achia, T.N.O. Tobacco Use and Mass Media Utilization in Sub-Saharan Africa. *PLoS ONE* **2015**, *10*, e0117219. [[CrossRef](#)]
39. Yesuf, K.A.; Liyew, A.D.; Bezabih, A.K. Impact of exposure to mass media on utilization modern contraceptive among adolescent married women in Ethiopia: Evidence from Ethiopia demographic health survey 2016. *Int. J. Sci. Rep.* **2021**, *7*, 434–442. [[CrossRef](#)]
40. Zikarge, T.H. Adolescents Exposure to Global Mass Media: Influences on Sexual Behavior and Sexuality in Hawassa City, Ethiopia. *Res. Humanit. Soc. Sci.* **2019**, *9*, 19–25.
41. Sureda, X.; Ballbè, M.; Martínez, C.; Fu, M.; Carabasa, E.; Saltó, E.; Martínez-Sánchez, J.M.; Fernández, E. Impact of tobacco control policies in hospitals: Evaluation of a national smoke-free campus ban in Spain. *Prev. Med. Rep.* **2014**, *1*, 56–61. [[CrossRef](#)]
42. Jabbour, S.; Fouad, F.M. Religion-based tobacco control interventions: How should WHO proceed? *Bull World Health Organ* **2004**, *82*, 923–927.
43. Marsh, J.; McNeill, A.; Lewis, S.; Coleman, T.; Bains, M.; Larwood, A.; Purdy, J.; Jones, L.L. Protecting children from secondhand smoke: A mixed-methods feasibility study of a novel smoke-free home intervention. *Pilot Feasibility Stud.* **2016**, *2*, 53. [[CrossRef](#)]
44. Shah, S.; Ainsworth, H.; Fairhurst, C.; Tilbrook, H.; Sheikh, A.; Amos, A.; Parrott, S.; Torgerson, D.; Thompson, H.; King, R.; et al. Muslim communities learning about second-hand smoke: A pilot cluster randomised controlled trial and cost-effectiveness analysis. *npj Prim. Care Respir. Med.* **2015**, *25*, 15052. [[CrossRef](#)] [[PubMed](#)]
45. International Union against Tuberculosis and Lung Disease(The Union); Campaign for Tobacco-Free Kids(CTFK); Johns Hopkins Bloomberg School of Public Health (JHSPH). Assessing Compliance with Smoke-Free Laws. In *A "How-to" Guide for Conducting Compliance Studies*; International Union against Tuberculosis and Lung Disease: Edinburgh, Scotland, 2014; pp. 1–36.
46. Hovell, M.F.; Bellettiere, J.; Liles, S.; Nguyen, B.; Berardi, V.; Johnson, C.; Matt, G.; Malone, J.; Boman-Davis, M.C.; Quintana, P.J.E.; et al. Randomised controlled trial of real-time feedback and brief coaching to reduce indoor smoking. *Tob. Control* **2020**, *29*, 183. [[CrossRef](#)] [[PubMed](#)]