



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/189053/>

Version: Accepted Version

Article:

Shashkova, Sviatlana, Nyström, Thomas and Leake, Mark C (2022) Copy Number Analysis of the Yeast Histone Deacetylase Complex Component Cti6 Directly in Living Cells. *Methods in Molecular Biology*. pp. 183-190. ISSN: 1064-3745

https://doi.org/10.1007/978-1-0716-2221-6_14

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Copy number analysis of the yeast histone deacetylase complex component

Cti6 directly in living cells

Sviatlana Shashkova^{1,2}, Thomas Nyström¹, Mark C Leake^{2,3}

¹ Department of Microbiology and Immunology, Institute of Biomedicine, Sahlgrenska Academy, University of Gothenburg, 405 30 Gothenburg, Sweden

²Department of Physics, University of York, York, YO10 5DD, UK

³Department of Biology, University of York, York, YO10 5DD, UK

Correspondence to: sviatlana.shashkova@gu.se

Running Head: Cti6 copy numbers

Abstract

Proteins are the one of the key components of cellular life that play a crucial role in most biological processes. Therefore, quantification of protein copy numbers is essential for revealing and better understanding of cellular behaviour and functions. Here we describe a single-molecule fluorescence-based method of protein copy numbers quantification directly in living cells. This enables quick and reliable estimations and comparison of protein of interest abundance without implementing large-scale studies.

Key words: histone deacetylase, single-molecule, copy number, living cells

1. Introduction

Proteins are one of the main functional components of cells. Protein abundance is the key in regulation of protein-protein and protein-nucleic acids interactions as well as enzymatic reactions which comprise most biological processes. This makes quantification of protein copy numbers essential for revealing and better understanding of cellular behaviour and functions.

Initially, methods for protein numbers estimation relied on gene expression measurements, such as microarray hybridisation (1). However, protein homeostasis resembles a balance between protein production, including mRNA processing and translation, and degradation.

Therefore, reliable quantification can be obtained only by counting protein numbers directly.

Multiple large-scale high-throughput studies on eukaryotic proteome, mainly that of the budding yeast *Saccharomyces cerevisiae*, have been performed using various methods. This

includes quantitative mass spectrometry, western blotting, fluorescence-based methods, such as wide field and confocal microscopy approaches as well as flow cytometry (2–8).

We report an easy manual method based on a single-molecule fluorescence data, which allows for a quick estimation of protein numbers in living eukaryotic cells. We have applied such an approach to the Cti6 protein of the chromatin remodelling system in the budding yeast *S. cerevisiae*. Cti6 is a component of the large histone deacetylase complex Rdp3L and participates in multiple cellular processes including iron tolerance and metabolism maintenance (9). Cti6 plays an important role in gene expression regulation via its binding to the SAGA activator complex (10). Moreover, Cti6 has been reported to convert one of the largest repression machineries, the Ssn6-Tup1 global co-repressor complex, into an activator (9, 11).

Here we report in detail all necessary steps to estimate molecular copy numbers of a protein of interest in living yeast *S. cerevisiae* cells based on the fluorescent signal. We describe endogenous genomically integrated labelling of the Cti6 protein with a monomeric form of GFP, mGFP, which contains the A206K point mutation (12). We used a super-resolution Slimfield microscopy set-up to visualise Cti6-mGFP directly in living cells. We then applied previously reported MATLAB code (13, 14) to track fluorescent complexes and estimate intensity of a single fluorescent molecule. We then show how these data can be used to estimate Cti6 copy numbers per cell.

2. Materials

2.1 Yeast strains

1. The BY4741 wild type yeast *S. cerevisiae* cells, pmGFPS plasmid (**14**) (*HIS3*, *GFPmut3 S65G, S72A, A206K*).
2. 40% (w/v) glucose solution.
3. Yeast Nitrogen Base (YNB) media (with complete amino acid supplement and -his): 1.7 g/l yeast nitrogen base without amino acids and $(\text{NH}_4)_2\text{SO}_4$, 5 g/l $(\text{NH}_4)_2\text{SO}_4$, complete amino acid supplement or -his amino acid supplement as indicated by manufacturer, pH 5.8-6.0, supplemented with 4% w/v glucose (added after autoclaving). To prepare YNB plates, add 20 g/l agar before autoclaving.
4. Yeast Peptone Dextrose (YPD) liquid medium (10 g/l yeast extract, 20 g/l bacto-peptone) supplemented with 4% w/v glucose (added after autoclaving). To prepare YPD plates, add 20 g/l agar before autoclaving.
5. Phusion high fidelity DNA polymerase 2U/ μl kit, dNTP mix.
6. Primers: [Fig 1 near here] for mGFP::his3 fragment amplification, Forward (Fw) and Reverse (Rv) primers contain ~16-20 bp of the 5' and 3' ends, respectively, of the mGFP::his3 fragment from pmGFPS plasmid plus ~50bp sequences up- (Fv) and downstream (Rv) of the Cti6 STOP codon (Fig. 1a). For the conformation PCR, Forward (Fconf) and Reverse (Rconf) primers are

sequences of ~16-20 bp located at the 3' end of the *CTI6* gene and 5' end of the *mGFP::his3* (Fig. 1b).

7. 0.8% agarose gel with Midori Green or any other nucleic acid stain in 0.5x TAE buffer.
8. 1kb DNA ladder, 6x loading buffer.

9. 1M LiAc water solution.
10. 100mM LiAc water solution.
11. 50% w/v PEG-4000 solution in water.

12. 10 mg/ml single strand DNA (we use salmon sperm DNA).

13. 0.02% SDS solution in water.

14. Standard epifluorescence microscope. We used Zeiss Axio Observer.Z1

inverted microscope with Apotome and AxioCam 506 345 camera, and a Plan-Apochromat 100x/1.40 Oil DIC M27 objective.




Fig. 1 Primer design for (a) amplification of a *mGFP::his3* fragment for further genomic integration to label the *Cti6* protein and (b) verification of a correct insertion of the fluorophore sequence.

2.2 Super-resolution microscopy

1. We utilised the super-resolution fluorescence Slimfield microscope, with ~ 6 W/cm² excitation intensity directed onto a sample, 50mW Obis 488 nm laser, 100x 1.49 TIRF 1.49 NA oil immersion objective, EMCCD camera (Prime 95B scientific CMOS; Teledyne Photometrics) with 50 nm/pixel magnification (14, 15).
2. 125 μ l Gene Frames.
3. Plasma-cleaned BK7 glass microscope coverslip (22 \times 50 mm).
4. 2x YNB complete liquid medium supplemented with 8% w/v glucose.
5. 2% agarose solution in Milli Q water.

2.3 For the computational analysis, we used a combination of MATLAB, ImageJ Fiji and Excel software.

3. Methods

3.1 Fluorescent labelling of the Cti6 protein

1. Streak the BY4741 wild type cells from a frozen stock, using a sterile pipette tip on a freshly prepared YPD plate, and incubate at 30 °C for at least 24 h.
2. Set a culture in a 14 ml tube by inoculating 3 ml of YPD with cells grown a YPD plate. Incubate at 30 °C, 180 rpm, overnight.

- Amplify the mGFP::his3 fragment from the pmGFPS plasmid (14) by using Fw and Rv primers, dNTP mix and Phusion high fidelity DNA polymerase 2U/ μ l kit as recommended by manufacturer. With following PCR programme:

Initiation	98.0°C	- 30 sec	
Denaturation	98.0°C	- 10 sec	} 30 cycles
Annealing	60.6°C	- 20 sec	
Elongation	72.0°C	- 60 sec	
Final elongation	72.0°C	- 5 min	

Keep at 4°C

Verify the size of the fragment (here ~1920 bp) on 0.8% agarose gel by mixing 5 μ l of a PCR reaction mix with 1 μ l of 6x Loading Dye and applying onto a gel (100V, 25 min in 0.5x TAE buffer). Use 1kb DNA ladder as a reference for the size of the amplified fragment. Store at -20°C until further use.

- Inoculate 50ml of YPD medium with the overnight culture to the final OD₆₀₀ ~0.2. Grow until mid-logarithmic phase, OD₆₀₀ ~0.7-1.0.
- Transform the culture with 70 μ l of the PCR mix containing mGFP::his3 fragment by standard LiAc protocol (16). In brief, mix 50 μ l of washed with water and then 100 mM LiAc cells with 240 μ l of 50% w/v PEG-4000, 36 μ l of 1M LiAc, 5 μ l of 10 mg/ml salmon sperm DNA and 70 μ l of the PCR mix

containing mGFP::his3 fragment. Incubate the entire mix at 30°C for 30 min, then heat-shock at 42°C for 25 min. Wash (*Note 1*) and plate the transformation mix onto YNB-his plates. Incubate at 30°C for 3 days.

6. Confirm the colonies for the green signal under the microscope. We used Zeiss Axio Observer.Z1 inverted microscope with Apotome and Axiocam 506 345 camera, and a Plan-Apochromat 100x/1.40 Oil DIC M27 objective.
7. Verify correct integration of the mGFP::his3 fragment by confirmation PCR (in detail below).
8. Streak correct clones on a new YNB-his plate (*Note 2*), incubate for 24 h at 30°C. Store at 4°C. For longer storing, collect grown cells, mix in a cryo tube with 1ml of YP+20% glycerol solution, put into -80°C freezer.

3.2 Confirmation PCR

1. Extract genomic DNA: take a bit of a transformant into 20 µl of 0.02% SDS, heat at 98°C for 3 min. Spin down at 6000 rpm, 10 sec. Mix 10 µl of the supernatant with 10 µl of MiliQ water.
2. Perform PCR using Phusion high fidelity DNA polymerase 2U/µl kit as recommended by manufacturer. With following the PCR programme as above but using 58.9°C as an annealing temperature. Verify the size of the

fragment (here ~175 bp) on 0.8% agarose gel (100V, 25 min in 0.5x TAE buffer).

3.3 Preparing cells for microscopy

1. Set overnight cultures of BY4741 wild type and Cti6-mGFP strains in two 14 ml tubes by inoculating 3 ml of YNB complete with cells grown on plates. Incubate at 30 °C, 180 rpm.
2. In the morning, subculture the cells into 2x 3ml of fresh YNB complete medium to the final OD₆₀₀ ~0.2. Grow for four hours at 30 °C, 180 rpm.
3. Spin cells down at 3000 rpm, 1 min, remove the supernatant. Resuspend cells in 0.5 ml of fresh YNB complete medium.

3.4 Preparing microscopy slides using 125 µl Gene Frame

1. Remove the larger plastic cover and apply the frame onto a standard microscopy glass slide.
2. Prepare 0.5 ml of 2x YNB complete medium supplemented with 8% w/v glucose in a 1.5 ml Eppendorf tube.

3. Add 0.5 ml of melted 2% agarose water solution, quickly mix by pipetting up and down, take 0.5 ml of obtained mixture and add onto the slide within the Gene Frame.
4. Quickly apply a plastic cover slip from the Gene Frame package to remove the access and evenly distribute the solution within the well. Let sit for a few min.
5. Carefully remove the plastic cover slip by sliding it off.
6. Remove the smaller plastic cover from the frame.
7. Add ca 5 μ l of the prepared cell in droplets across the entire pad. Leave to dry for 3-5 min.
8. Cover the pad with a plasma-cleaned BK7 glass microscope coverslip (22 \times 50 mm), avoid air bubbles.

3.5 Data acquisition

1. Place the sample under the Slimfield microscope and locate a cell, adjust the focus to the middle of the cell.
2. Acquire 2 brightfield images with 10ms exposure time.
3. Turn off the brightfield and acquire 1000 images at 5 ms exposure time with the 488nm laser at 20mW power to obtain images of Cti6-mGFP.

4. Repeat for at least 30 cells.
5. Repeat the whole procedure for the non-fluorescent parental strain BY4741, this will be used for autofluorescence correction during the copy number estimation.

3.6 Copy number analysis

1. Load the TIF file containing fluorescence data acquisition into MATLAB as $n \times m$ pixels by p frames and perform tracking of individual fluorescent spots as described previously (17).
2. We estimate the fluorescence intensity of an individual mGFP molecule (I_{single}) based on the intensity distribution of the Cti6-mGFP spots identified in the last 2/3 of the acquisition frames (Note 3). [Fig 2 near here] As at this point, single photobleaching events become more apparent and detectable,




Fig. 2. Distribution of all Cti6-mGFP spot intensities (a) and spots identified towards the end of photobleaching (b). Peak values are indicated.

therefore, the most common value (the peak) represents the intensity of a single fluorophore (Fig. 2).

3. Load TIF file containing fluorescence data acquisition into ImageJ Fiji software and open the brightest frame.
4. Choose the “oval” selection shape on the ImageJ tool bar and draw a circle around the cell. Keep the same size of the selection for all measurements.
5. To obtain numeric values of total fluorescence intensity of the cell by pressing ‘Ctrl’ + ‘M’ (Analyse → Measure). Record Raw Integrated Density number (RawIntDen) in Excel. Repeat for all files containing fluorescence data of BY4741 Cti6-mGFP cells.
6. Repeat steps 1-5 for the BY4741 wild type fluorescence files. Calculate the average autofluorescence.
7. Subtract the average autofluorescence value from each RawIntDen value recorded from the BY4741 Cti6-mGFP cells.
8. To estimate the copy numbers of Cti6-mGFP per cell, divide obtained values by I_{single} .




Fig. 3. Jitter plot of Cti6-mGFP copy numbers per cell across a population of 32 individual cells. Bar indicates the median.

Figure 3 [Fig 3 near here] shows distribution

of Cti6 numbers per cell across a population of 32 cells. We estimated the mean number of Cti6 molecules as 2368 ± 1029 .

4. Notes

1. To increase the transformation efficiency, prior plating, the cells can be washed and incubated in YPD medium supplemented with 4% w/v glucose for 1 h at 30°C, 180 rpm.
2. As the fragment is genomically integrated, it is very unlikely that it will be lost during cellular division. Therefore, once the correct clones were confirmed, it is not mandatory to have a selective medium.
3. *Isingle* can also be estimated by imaging purified mGFP immobilised on a cover slip.

Acknowledgements

This work was supported by the Royal Society Newton International Fellowship Alumni (AL\191025, AL\201039), Knut and Alice Wallenberg Foundation (KAW 2017-0091, KAW 2015.0272), Swedish Research Council (VR 2019-03937) and the Leverhulme Trust (grant RPG-2019-156).

References

1. Spellman PT, Sherlock G, Zhang MQ, et al (1998) Comprehensive identification of cell

cycle-regulated genes of the yeast *Saccharomyces cerevisiae* by microarray hybridization. *Mol Biol Cell* 9:3273–3297. doi:10.1091/mbc.9.12.3273

2. Ho B, Baryshnikova A, Brown GW (2018) Unification of Protein Abundance Datasets Yields a Quantitative *Saccharomyces cerevisiae* Proteome. *Cell Syst* 6:192-205.e3. doi:10.1016/j.cels.2017.12.004
3. Tkach JM, Yimit A, Lee AY, et al (2012) Dissecting DNA damage response pathways by analysing protein localization and abundance changes during DNA replication stress. *Nat Cell Biol* 14:966–976. doi:10.1038/ncb2549
4. Chong YT, Koh JLY, Friesen H, et al (2015) Yeast Proteome Dynamics from Single Cell Imaging and Automated Analysis. *Cell* 161:1413–1424. doi:http://dx.doi.org/10.1016/j.cell.2015.04.051
5. Davidson GS, Joe RM, Roy S, et al (2011) The proteomics of quiescent and nonquiescent cell differentiation in yeast stationary-phase cultures. *Mol Biol Cell* 22:988–998. doi:10.1091/mbc.E10-06-0499
6. De Godoy LMF, Olsen J V., Cox J, et al (2008) Comprehensive mass-spectrometry-based proteome quantification of haploid versus diploid yeast. *Nature* 455:1251–1254. doi:10.1038/nature07341
7. Déneraud N, Becker J, Delgado-Gonzalo R, et al (2013) A chemostat array enables the spatio-temporal analysis of the yeast proteome. *Proc Natl Acad Sci U S A* 110:15842–15847. doi:10.1073/pnas.1308265110

8. Ghaemmaghami S, Huh WK, Bower K, et al (2003) Global analysis of protein expression in yeast. *Nature* 425:737–741. doi:10.1038/nature02046
9. Puig S, Lau M, Thiele DJ (2004) Cti6 is an Rpd3-Sin3 histone deacetylase-associated protein required for growth under iron-limiting conditions in *Saccharomyces cerevisiae*. *J Biol Chem* 279:30298–30306. doi:10.1074/jbc.M313463200
10. Papamichos-Chronakis M, Petrakis T, Ktistaki E, et al (2002) Cti6, a PHD Domain Protein, Bridges the Cyc8-Tup1 Corepressor and the SAGA Coactivator to Overcome Repression at GAL1. *Mol Cell* 9:1297–1305. doi:10.1016/S1097-2765(02)00545-2
11. Hill SM, Hao X, Grönvall J, et al (2016) Asymmetric Inheritance of Aggregated Proteins and Age Reset in Yeast Are Regulated by Vac17-Dependent Vacuolar Functions. *Cell Rep* 16:826–838. doi:10.1016/j.celrep.2016.06.016
12. Zacharias DA, Violin JD, Newton AC, Tsien RY (2002) Partitioning of lipid-modified monomeric GFPs into membrane microdomains of live cells. *Science* (80-) 296:913–916. doi:10.1126/science.1068539
13. Wollman AJM, Leake MC (2016) Single molecule narrowfield microscopy of protein-DNA binding dynamics in glucose signal transduction of live yeast cells. *Methods Mol Biol* 1431:5–15. doi:10.1007/978-1-4939-3631-1_2
14. Wollman AJM, Shashkova S, Hedlund EG, et al (2017) Transcription factor clusters regulate genes in eukaryotic cells. *Elife* 6:e27451. doi:10.7554/eLife.27451
15. Laidlaw KME, Bisinski DD, Shashkova S, et al (2021) A glucose-starvation response

governs endocytic trafficking and eisosomal retention of surface cargoes in budding yeast. *J Cell Sci* 134:. doi:10.1242/jcs.257733

16. Gietz RD, Schiestl RH (2007) Frozen competent yeast cells that can be transformed with high efficiency using the LiAc/SS carrier DNA/PEG method. *Nat Protoc* 2:1–4. doi:10.1038/nprot.2007.17
17. Wollman AJM, Leake MC (2016) Single-Molecule Narrow-Field Microscopy of Protein--DNA Binding Dynamics in Glucose Signal Transduction of Live Yeast Cells. *Chromosom Archit Methods Protoc* 5–15. doi:10.1007/978-1-4939-3631-1_2