



This is a repository copy of *Systematic reviews in orthodontics: a fresh look to promote renewal and reduce redundancy*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/190002/>

Version: Accepted Version

Article:

Millett, D.T., Benson, P.E. orcid.org/0000-0003-0865-962X, Cunningham, S.J. et al. (4 more authors) (2022) Systematic reviews in orthodontics: a fresh look to promote renewal and reduce redundancy. *American Journal of Orthodontics and Dentofacial Orthopedics*, 162 (1). pp. 1-2. ISSN 0889-5406

<https://doi.org/10.1016/j.ajodo.2022.03.012>

Article available under the terms of the CC-BY-NC-ND licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Title:

Systematic Reviews in Orthodontics: A fresh look to promote renewal and reduce redundancy.

Authors:

Declan T. Millett, Professor of Orthodontics, Oral Health & Development, Cork University Dental School and Hospital, University College Cork, Ireland.

Philip E. Benson, Professor of Orthodontics, School of Clinical Dentistry, University of Sheffield, United Kingdom.

Susan J. Cunningham, Professor in Orthodontics, EDI Craniofacial & Development Sci, Eastman Dental Institute, University College London, United Kingdom.

Grant T. McIntyre, Consultant / Honorary Professor, Dundee Dental Hospital, University of Dundee Dundee, United Kingdom.

Padhraig S. Fleming, Professor of Orthodontics, Department of Orthodontics, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, United Kingdom.

Farhad B. Naini, Consultant Orthodontist, St. George's University Hospitals NHS Foundation Trust, Kingston Hospital NHS Foundation Trust, London, United Kingdom.

Aliki Tsihlaki, Senior Lecturer, Department of Orthodontics, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, United Kingdom.

Corresponding Author: Declan T. Millett, Professor of Orthodontics, Oral Health & Development, Cork University Dental School and Hospital, University College Cork, Ireland. d.millett@ucc.ie

Systematic Reviews in Orthodontics: A fresh look to promote renewal and reduce redundancy.

Systematic reviews are intended to collate and analyze the evidence regarding a specific aspect of clinical practice in a drive for evidence-based health care.¹ Unfortunately, many systematic reviews in medicine and dentistry^{2,3} conclude by stating that further evidence is required through well-designed randomized clinical trials.^{4,6} It has been mooted that within orthodontics, systematic reviews considerably outweigh the number of clinical trials⁷ with evidence that a substantial proportion of the former are limited or inconclusive. This is due in many cases to low quality studies preventing or limiting meta-analyses.⁸ When several systematic reviews exist on a topic, it is possible to evaluate and collate their findings in an overview or 'umbrella' review.⁹ Within overviews, however, considerable duplication and unnecessary overlap have been identified which represents significant research waste.¹⁰

To address these shortcomings, is there a need to revisit what a researcher should do when a systematic review is completed? Although the Cochrane Collaboration has adopted the concept of living systematic reviews (LSR)¹¹ with creation of a Living Evidence Network (LEN), this involves continual update of the review as new evidence becomes available, rather than providing a direct stimulus for the improvement of the primary evidence in a subject area. The limited yield from systematic reviews, meta-analyses and umbrella reviews has prompted us to make the following suggestions:

1] When a clinical systematic review is completed, aside from making recommendations as to how evidence should be improved, the authors should lead by example and ideally undertake a randomized controlled trial (RCT) but if not feasible or appropriate, another high quality study for example a prospective cohort study. This would hopefully lead authors to reflect on why they did the review in the first place, presumably to improve their clinical practice and that of others, rather than completion and publication of the review as the primary academic exercise.

2] All orthodontic systematic reviews should adopt the PRISMA guidelines for write-up and include a link to a peer-reviewed protocol for the "ideal" study in the area of interest. This would include a sample size calculation for the most clinically important outcome, timepoint at which the data are to be collected, an explicit list of secondary outcomes based on the orthodontic core outcome set¹² and detail the prescribed methodology as well as the optimal statistical analyses required. The pre-determined study protocol should be published on PROSPERO or in an open access journal such as Clinical Trials. It should also be made available on the authors' own webpages. We feel that this approach would have several advantages:

- It would facilitate the conduct of RCTs on a specific topic by having a peer-reviewed protocol to hand that may be used for local ethical applications.
- Several operators / centres could undertake the same trial contemporaneously or in succession in different settings to improve the generalizability of findings. Such multi-centre collaboration is to be encouraged.^{13,14}
- Most importantly, methodological variation would be reduced and hopefully confounders identified as well as accounted for in the sampling strategy or analyses.
- With the adoption of the same trial design, similar participants and outcomes being assessed preferably from the orthodontic core outcome set¹² at the same timepoints, meta-analyses would be performed to potentially increase the certainty of findings.

This would maximize yield from our clinical trials.¹⁵ For reviews incorporating comparisons of several interventions, defining a protocol for a subsequent trial is challenging. In those instances, focus would have to be directed at trialling what are regarded as the most clinically important outcomes.

It is essential to emphasize that we continue to recognize the importance of well-conducted systematic reviews with meta-analyses, as these are the pinnacle of the hierarchy of evidence. We also acknowledge that we are guilty in almost all cases of not having followed our own systematic reviews with related clinical trials and indeed the views portrayed here are not intended to be critical of investigators.

As has often been said "If you always do what you always did, you will always get what you always got" and it is our duty as researchers to generate meaningful quantitative and qualitative data with the aim

of improving clinical decision-making. Systematic reviews are a pivotal, complex and evolving field, and making progress will be challenging but ultimately rewarding. We can but try.

Acknowledgement

Claire Laide is thanked for her assistance.

References

1. Sackett DL, Rosenberg WMC, Gray JAM, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't? *BMJ* 1996;312-71.
2. Glennly A-M, Esposito M, Coulthard P, Worthington HV. The assessment of systematic reviews in dentistry. *Eur J Oral Sci* 2003;85-92.
3. Bassani R, Kahil G, Pereira R, Page MJ, Tricco AC, Moher D, Sarkis-Onofre R. Systematic reviews in dentistry: Current status, epidemiological and reporting characteristics. *J Dent* 2019;71-84.
4. Fleming PS, Seehra J, Polychronopoulou A, Fedorowicz Z, Pandis N. A PRISMA assessment of the reporting quality of systematic reviews in orthodontics. *Angle Orthod* 2013;158-163
5. Pandis, N., Fleming, PS, Worthington, H and Salanti, G. The quality of the evidence according to GRADE is predominantly low or very low in oral health systematic reviews. *PloS one* 2015;e0131644
6. Littlewood SJ, Millett DT, Doubleday B, Bearn DR, Worthington HV. Retention procedures for stabilising tooth position after treatment with orthodontic braces. *Cochrane Database of Systematic Reviews* 2016, DOI: 10.1002/14651858.CD002283.pub4.
7. Papageorgiou SN, Eliades T. Evidence-based orthodontics: Too many systematic reviews, too few trials. *J Orthod* 2019;9-12.
8. Koletsi, D., Fleming, P. S., Eliades, T., Pandis, N. The evidence from systematic reviews and meta-analyses published in orthodontic literature. Where do we stand?. *EJO* 2015;603-609.
9. Lewis R, Hendry M, Din N, Stanciu MA, Nafees S, Hendry A, Teoh ZH, Lloyd T, Parsonage R, Neal RD, Collier G, Huws DW. Pragmatic methods for reviewing exceptionally large bodies of evidence: systematic mapping review and overview of systematic reviews using lung cancer survival as an exemplar. *Syst Rev* 2019;8-171.
10. Lunny, C, Zhang, JH, Chen, A, Neelakant, T, Shinger, G, Stevens, A, Tasnim, S, Sadeghipouya, S, Adams, S, Zheng, YW and Lin, L. Topic duplication and research waste at the 'Overviews of systematic reviews' level: Survey of overlapping overviews, 29 June 2021, PREPRINT (Version 1) available at Research Square <https://doi.org/10.21203/rs.3.rs-665009/v1>
11. Elliott, JH, Synnot, A, Turner, T, Simmonds, M, Akl, EA, McDonald, S, Salanti, G, Meerpohl, J, MacLehose, H, Hilton, J and Tovey, D Living systematic review: 1. Introduction—the why, what, when, and how. *J Clin Epidemiol* 2017;23-30.
12. Tsihliaki A, O'Brien K, Benson PE, Marshman Z, Johal A, Colonio-Salazar FB, Harman NL, Fleming PS. Development of a core outcome set for use in routine orthodontic clinical trials. *Am J Orthod Dentofacial Orthop* 2020;650-660.
13. Bellomo R, Warrillow SJ, Reade MC. Why we should be wary of single-center trials. *Crit Care Med* 2009;3114-3119.
14. Bafeta A, Dechartres A, Trinquart L, Yavchitz A, Boutron I, Ravaud P. Impact of single centre status on estimates of intervention effects in trials with continuous outcomes: meta-epidemiological study. *BMJ* 2012;344.
15. Chalmers I, Glasziou P. Avoidable waste in the production and reporting of research evidence. *Lancet* 2009;86-89.