

This is a repository copy of *Measuring the digital divide among people with severe mental ill health using the essential digital skills framework*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/190398/>

Version: Published Version

---

**Article:**

Spanakis, P [orcid.org/0000-0002-2732-8300](https://orcid.org/0000-0002-2732-8300), Wadman, R [orcid.org/0000-0003-4205-0631](https://orcid.org/0000-0003-4205-0631), Walker, Lauren [orcid.org/0000-0003-2459-7860](https://orcid.org/0000-0003-2459-7860) et al. (6 more authors) (2022) Measuring the digital divide among people with severe mental ill health using the essential digital skills framework. *Perspectives in public health*. p. 17579139221106399. ISSN 1757-9147

<https://doi.org/10.1177/17579139221106399>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Measuring the digital divide among people with severe mental ill health using the essential digital skills framework

## Authors

**P Spanakis**   
 Department of Health Sciences, University of York, ARRC Building, Heslington, York YO10 5DD, UK  
 Department of Psychology, University of Crete, Rethymnon, Greece.  
 School of Psychology, Mediterranean College, Athens, Greece.  
 Email: panagiotis.spanakis@york.ac.uk

**R Wadman**  
 Department of Health Sciences, University of York, York, UK

**L Walker**   
 Department of Health Sciences, University of York, York, UK

**P Heron**  
 Department of Health Sciences, University of York, York, UK

**A Mathers**  
 The Good Things Foundation, Sheffield, UK; Royal Society of Arts, London, UK

**J Baker**  
 School of Healthcare, University of Leeds, Leeds, UK

**G Johnston**  
 Independent Peer Researcher, Clackmannan, UK

**S Gilbody**  
 Department of Health Sciences, University of York, York, UK

**E Peckham**  
 Department of Health Sciences, University of York, York, UK

**Corresponding author:**  
 Panagiotis Spanakis, as above

## Keywords

digital skills; digital divide; Internet; severe mental ill health; psychosis; bipolar

## Abstract

**Aims:** Amid the vast digitalisation of health and other services during the pandemic, people with no digital skills are at risk of digital exclusion. This risk might not abate by the end of the pandemic. This article seeks to understand whether people with severe mental ill health (SMI) have the necessary digital skills to adapt to these changes and avoid digital exclusion.

**Methods:** Two hundred and forty-nine adults with SMI across England completed a survey online or offline. They provided information on their digital skills based on the Essential Digital Skills (EDS) framework, sociodemographic information, and digital access. This is the first time that the EDS is benchmarked in people with SMI.

**Results:** 42.2% had no Foundation Skills, and 46.2% lacked skills for daily life (lacking Foundation or Life Skills). 23.0% of those working lacked skills for professional life (lacking Foundation or Work Skills). The most commonly missing skills were handling passwords and using the device settings (Foundation Skills) and online problem solving (Skills for Life). People were interested in learning more about approximately half of the skills they did not have. People were more likely to lack Foundation Skills if they were older, not in employment, had a psychosis-spectrum disorder, or had no Internet access at home.

**Conclusion:** A significant portion of people with SMI lacked Foundation Skills in this objective and benchmarked survey. This points to a high risk for digital exclusion and the need for focused policy and tailored health sector support to ensure people retain access to key services and develop digital skills and confidence. To our knowledge, this is the first time this has been described using the EDS framework. Services, including the National Health Service (NHS), need to be aware of and mitigate the risks.

## INTRODUCTION

During the COVID-19 pandemic, restrictions were applied to movement and social contact, to contain the spread of the virus.<sup>1,2</sup> Many daily activities (e.g. shopping, entertainment, and contacting friends and family) shifted from face-to-face to remote delivery, often via the Internet. The UK NHS experienced a similar shift, both in mental<sup>3–5</sup> and in general healthcare.<sup>6–8</sup>

These changes increased the demand for digital skills and led to more people using the Internet for longer, leading to estimations that in one year the UK made five years' worth of progress in digital engagement.<sup>9</sup> However, this

situation also highlighted the pre-existing digital divide (i.e. inequalities in digital access and engagement in the population) and the health inequalities that may derive from it.<sup>10</sup> People lacking digital skills might experience negative outcomes in relation to health, wellbeing, and social support, as they might struggle to access health services, Government and local authority services, and support network activities that are online.<sup>11</sup>

It remains unknown whether people with SMI (e.g. a diagnosis of bipolar or psychosis-spectrum disorder) have the digital skills to adapt to this new digitalisation of services.<sup>12</sup> Finding the

## Measuring the digital divide among people with severe mental ill health using the essential digital skills framework

Internet too difficult or too complex to understand has been commonly reported among people with SMI as a barrier to using the Internet, both before<sup>13–15</sup> and during the pandemic.<sup>16</sup> It is important to gain a deeper understanding of this issue, as digital exclusion in people with SMI might widen existing health and social inequalities. For example, people with SMI are at greater risk for long-term physical health conditions<sup>17</sup> and at greater need for regular healthcare appointments to monitor their conditions (e.g. the annual physical health check).<sup>18</sup> They might also be more likely to experience feelings of loneliness.<sup>19</sup> Lack of skills to use the Internet might obstruct access to important sources of support (e.g. video-calls with health professionals or friends and family, accessing health and community-related information online, and participating in online community activities). It is important to note that the risk of exclusion due to lack of skill will probably not abate at the end of the pandemic, considering that the pandemic accelerated plans for digital healthcare to become a mainstream option in the NHS.<sup>20</sup>

The EDS framework which has been adopted by the UK Department of Education defines the skills that people would need to have to fully benefit from using the Internet and digital devices,<sup>9</sup> and sets the standards for all formal digital skills training programmes in the country. The framework includes three domains: Foundation Skills, Skills for Life, and Skills for Work. The Foundation Skills domain concerns basic prerequisite knowledge (e.g. knowing how to use the settings in a device or how to connect to a secure Wi-Fi network), while the other two domains concern five skill types in everyday life and in work environments, respectively: (1) Being safe, legal and confident online; (2) Communicating; (3) Problem solving; (4) Transacting; and (5) Handling information and content. Although EDS is mapped with the general UK population annually through the Lloyds Bank UK Consumer Digital Index, it has never been benchmarked in people with SMI specifically.

This article reports on the Skills and Proficiency in Digital Essential

Requirements study (SPIDER) which assesses the digital skills of a sample of people with SMI amid the COVID-19 pandemic, based on the EDS framework. The primary objectives were to understand the extent to which people with SMI have EDS and how they compare to the general population, as well as to identify the areas of greatest deficit in skills and their association with key sociodemographic and health factors.

## METHODS

### Design and procedures

The SPIDER study recruited participants from a clinical cohort that participated in the Optimising Wellbeing in Self Isolation (OWLS) study. The cohort included people that were 18 years old or over and had a documented diagnosis of schizophrenia or delusional/psychotic illness (ICD 10 F20.X & F22.X or DSM equivalent) or bipolar disorder (ICD F31.X or DSM equivalent). The methods of recruitment to the OWLS study are outlined in the supplementary material (S1). To be eligible for SPIDER, participants had to have consented during OWLS to be contacted about future surveys. Participants were recruited to SPIDER from January 2021 to March 2021.

Interested participants were provided with an information sheet (read over the phone, or send by email, text message, or post), and those consenting to participate were given the option to complete the survey online via a link, or offline (over the phone with a researcher or in a hard copy survey sent by post.) Offline options were provided to ensure the inclusion of people that were not digitally engaged.

### Measures

All variables and analyses reported here have been preregistered<sup>21</sup> in Open Science Framework (OSF).

### Essential digital skills

Participants were asked whether they could complete a series of tasks related to using the Internet and digital devices, on their own if they ever need to (responses were yes or no). For those

answering 'no', they were asked if they would be interested in learning more about them (yes/no). The tasks were derived from the Lloyds UK Consumer Digital Index 2020<sup>22</sup> and the Scottish Council for Voluntary Organisations (SCVO) Essential Digital Skills Toolkit.<sup>23</sup> The complete list of tasks used in SPIDER is available at <https://mfr.osf.io/render?url=https://osf.io/keumd/?direct%26mode=render%26action=download%26mode=render>.

Participants were considered as having or not having a given skills domain according to the EDS framework.<sup>22</sup> For Foundation Skills, participants would need to report they could complete all tasks from a list of eight basic tasks (e.g. using a device setting to make its use easier, or connecting to a secure and safe Wi-Fi network).

For Skills for Life, we used a list of 30 tasks, grouped into five skill types: Communicating (e.g. I can set up an email account), Handling information (e.g. I can use search engines to find information and make use of search terms to generate better results), Transacting (e.g. I can safely buy things online), Problem solving (e.g. I can use the Internet to find how to do something online), and Being safe and legal online (I am careful with what I share online as I realise that the information I put online stays there and could be accessed in the future by other people). To be considered as having Skills for Life, participants should be able to complete at least one task from each skill type. Only participants who had Foundation Skills were assessed for Skills for Life.

For Skills for Work, we used a list of 14 tasks, grouped in the same skill types as in Skills for Life. The difference was that the tasks in each skill type were more relevant to a working environment (e.g. I can set up and manage an account on a professional online network/community, or I can organise, store and share work-related information on different computers, tablets or phones). To be considered as having Skills for Work participants should be able to complete at least one task in each skill type. Only participants who had Skills for Life and were in employment were assessed for Skills for Work.

## Measuring the digital divide among people with severe mental ill health using the essential digital skills framework

**Digital access**

In the initial OWLS survey, participants were asked if they owned a digital device (smartphones, computers, laptops, or tablets – yes/no), and whether they could access the Internet from home (yes/no). In the SPIDER study, we also asked whether people have used the Internet in the last 12 months (yes a lot, yes a little, or not at all) and if in the last 12 months they ever experienced not having enough data available to access the Internet as much as they would need (data poverty – yes/no).

**Sociodemographic and health variables**

Using data that were available in the initial OWLS dataset, we derived participants' age (grouped as 18–30, 31–45, 46–65, and 66+), gender (female, male or transgender), ethnicity (grouped as White background or Other than White), employment (grouped as being in paid employment or not) and care setting (grouped as primary or secondary mental healthcare). We also derived participants' socioeconomic deprivation index<sup>24</sup> according to their area of residency using their postcodes. Index scores range from 1 to 10 with higher scores indicating less deprivation and were grouped into very high deprivation (1–2), high deprivation (3–4), medium deprivation (5–6), low deprivation (7–8) and very low deprivation (9–10).

Participants' health records were inspected for consenting participants to obtain their SMI diagnosis, which was then categorised into psychosis-spectrum disorders (including schizophrenia, schizoaffective, or any other psychotic disorder), bipolar disorder, or other SMI (including participants who were eligible for OWLS on the basis of a psychosis or bipolar disorder diagnosis which was later changed in their health records to something different, as for example severe depressive disorder with psychotic features). For those not providing consent to access their records or insufficient identifiable information (e.g. name and date of birth), the diagnosis was coded as 'not recorded'. The 'not recorded' category is not reported in our preregistered plan but was added to retain in the analysis the 48 participants falling in this group.

**Analysis**

Descriptive statistics are provided for each skills domain (Foundation, Skills for Life, or Skills for Work). For participants lacking a skills domain, we present descriptive statistics for the most commonly missing skill types and for interest in learning more about these skill types.

Association of sociodemographic, health, and digital access variables (independent variables) with having Foundation Skills were examined with a univariable binary logistic regression model before adding these variables into a hierarchical multivariable binary logistic regression. In the multivariable model, all independent variables were inserted at once, with sociodemographic and health variables inserted in the first block and digital access variables in the second block. Statistical significance was set a  $p < .05$ , and analysis was conducted with IBM SPSS 26.

Some of the preregistered analysis was not possible due to very low counts in the sample. Among people with Foundation Skills, only nine people reported having no Skills for Life (93.0%) and four working people reported having no Skills for Work (92.0%). As a result, we do not present any descriptive statistics for missing skill domains and interest in learning more among those who reported having no Skills for Work. We also did not explore which factors are associated with having Skills for Life or Work.

To test for response bias, we examined whether participants who accepted or declined our invitation to the SPIDER study differed in age, gender, ethnicity, socioeconomic deprivation, care setting, and diagnosis. Differences were examined with a chi-square test (or the likelihood ratio if test assumptions were violated), apart from age where we used an independent samples t-test instead.

**RESULTS**  
**Sample**

We invited 315 adults with SMI to the SPIDER study and 249 (67.8%) participated. Those who accepted or declined the invitation did not differ in

terms of any of the examined sociodemographic characteristics (age:  $t(365) = -0.45$ ,  $p = .650$ ; gender: Likelihood Ratio (2) = 4.77,  $p = .092$ ; ethnicity:  $\chi^2(1) = 1.44$ ,  $p = .230$ ; deprivation:  $\chi^2(4) = 6.47$ ,  $p = .167$ ; care setting:  $\chi^2(1) = 0.63$ ,  $p = .429$ ; and diagnosis:  $\chi^2(3) = 6.07$ ,  $p = .108$ ).

Table 1 provides the sample characteristics. Participants had a mean age of 51.7 years old (range: 21–84), and the sample included 51.4% men, 46.6% women, 2% transgender, 15.6% people from other than White ethnic backgrounds and 44.6% who resided in high/very high deprivation areas. The primary diagnosis was psychosis-spectrum disorder (48.2%). The survey was completed online by 93 (37.3%) participants and over the phone or via the post by 156 (62.7%). Regarding digital access, 21.3% of the sample were non-users of the Internet, 12.4% did not own a digital device, 15.3% had no access to the Internet at home, and 11.2% had experienced data poverty.

**Essential digital skills**

Table 2 presents the EDS framework descriptive results. Out of the total sample ( $n = 249$ ), 105 participants (42.2%) reported no Foundation Skills. They were unable to complete on average 3.64 ( $\pm 2.50$ ) tasks out of eight in total but were interested in learning more about on average 2.22 ( $\pm 2.80$ ) of them. The tasks that people who reported no Foundation Skills were most often unable to perform were updating and changing passwords (68.6%) and using the device settings to improve its usability (e.g. changing font sizes, screen brightness, and screen contrasts) (61.9%). The least frequent skill deficit was not knowing how to turn on a device (17.1%) (Figure 1).

Among those who reported having Foundation Skills ( $n = 142$ , 57.0%), almost everyone ( $n = 132$ , 93.0%) reported having Skills for Life. Among the few people that reported not having Skills for Life ( $n = 9$ ), five (55.6%) reported having no problem-solving skills, and four (44.4%) no Transacting Skills. No deficiency was reported in Communication, Handling Information,

## Measuring the digital divide among people with severe mental ill health using the essential digital skills framework

Table 1	
Sample characteristics (n = 249)	
Variable (valid N)	N (%)
Age (249)	
18–30	28 (11.2)
31–45	64 (25.7)
46–65	100 (40.2)
66+	57 (22.9)
Gender (249)	
Female	116 (46.6)
Male	128 (51.4)
Transgender	5 (2.0)
Ethnicity (249)	
Asian	14 (5.6)
Black	4 (1.6)
Mixed	11 (4.4)
White British	200 (80.3)
White (other)	10 (4.0)
Other	10 (4.0)
Socioeconomic deprivation (241)	
Very high	60 (24.1)
High	51 (20.5)
Medium	49 (19.7)
Low	43 (17.3)
Very low	38 (15.3)
In paid employed (248)	
Yes	61 (24.5)
No	187 (75.1)
Diagnosis (249)	
Bipolar disorder	83 (33.3)
Psychosis-spectrum disorder	120 (48.2)

(Continued)

and Being safe and legal online (0% were missing these skill types). Participants that reported not having Skills for Life were unable to complete on average 14.6 ( $\pm$  6.5) tasks out of 30 overall but were interested in learning more about 7.7 ( $\pm$  7.4) on average. Among those who reported having Skills for Life and were in paid employment (n = 50), almost everyone (n = 46, 92.0%) reported having Skills for Work.

Overall, 114 participants from the total sample (45.7%) did not have the EDS for daily life, lacking either Foundation Skills or Skills for Life. Among the working population in our sample (n = 61), 14 participants (23.0%) did not have the EDS for professional life, lacking either Foundation Skills, Skills for Life, or Skills for Work.

### Sociodemographic associations with foundation skills

A multivariable binary logistic regression model (Table 3) demonstrated that reporting having Foundation Skills was eight and four times more likely among the 18–30 and 31–45 age groups, respectively (compared to 66+), three times more likely among those in paid employment and four times more likely in those with a bipolar disorder diagnosis (compared to psychosis-spectrum). It was also found that participants with no Internet access at home were seven times less likely to report they had Foundation Skills.

### DISCUSSION

We assessed digital skills in a sample of people with SMI using the EDS framework. To our knowledge, this is the first time this benchmark measure of knowledge and skills has been applied to a sample of people who use mental health services. It is concerning that 42% (n = 105) had no Foundation Skills, lacking prerequisite knowledge to interact with digital technologies and benefit from their use. The most problematic areas were handling passwords and using device settings to improve usability. Older people, those not currently in work, those with a psychosis-spectrum disorder, and those with no Internet access at home were at greater

## Measuring the digital divide among people with severe mental ill health using the essential digital skills framework

Table 1 (Continued)

Sample characteristics (n = 249)	
Variable (valid N)	N (%)
Other SMI	16 (6.4)
Not recorded	30 (12.0)
Care setting (247)	
Primary care	95 (38.2)
Secondary care	152 (61.0)
Using the Internet	
A lot	137 (55.0)
Just a little	58 (23.3)
Not at all	53 (21.3)
Owning any digital device (245)	
Yes	214 (85.9)
No	31 (12.4)
Access to the Internet at home (246)	
Yes	208 (83.5)
No	38 (15.3)
Ever felt not having enough data? (248)	
Yes	28 (11.2)
No	177 (71.1)
N/A (Not using the Internet)	43 (17.3)

SMI: severe mental ill health.

risk of lacking Foundation Skills. More positively, among people who had Foundation skills, almost everyone had Skills for Life and Skills for Work.

Our results reveal a digital divide between our SMI sample and the general population. People with SMI in this study were twice as likely to experience a deficit in either Foundation or Life Skills (n = 114, 46.2%) compared to the general population (22%).<sup>22</sup> During the pandemic, this might contribute to inequalities of access to digital services, including but not limited to healthcare services, due to a lack of familiarity or confidence with technology.<sup>25</sup> More worryingly, these

inequalities might remain beyond the pandemic,<sup>26</sup> given that transition to digital healthcare might become a more mainstream option.<sup>20,27</sup> Combined with the fact that most people in our study owned a digital device (n = 214, 85.9%), this implies that access to digital devices on its own may do not mitigate digital exclusion and that many people may use their devices only for very basic things. This suggests that digital skills training programmes are needed, that will be tailored to the needs of people with SMI, to help close this gap (see, for example, Recovery Colleges at <https://imroc.org/resources/1-recovery-colleges/>).

It was interesting that fewer of the working people in this study experienced a deficit in skills necessary for work, lacking either Foundation or Work Skills (n = 14, 23%), compared to the general population (52%).<sup>22</sup> This could be a sampling issue in our study and might not be representative of the total SMI population. However, it may also be explained by the fact that employment is less common in people with SMI,<sup>28–30</sup> as shown by the 75.1% (n = 187) rate of unemployment in this study, and has been associated with younger age and higher educational attainment,<sup>29,31,32</sup> as well as less severe SMI symptoms.<sup>29,32</sup> All of these might help people to acquire good digital skills, and this may be directly linked with this study's finding that employed participants were more likely to have Foundation Skills compared to those not in employment. Future studies should explore Skills for Work among people with SMI who are unemployed, as an indicative factor of their employment prospects and potential learning needs. Researchers should also seek to understand the contribution of digital skills deficits to the high unemployment rates in people with SMI (e.g. most jobs advertised online), especially in relation to other employment barriers related to SMI (e.g. ill health or stigma).

Looking further into skills deficits, we found that people with no Foundation Skills struggled the most with handling passwords and using device settings. As such, as important as it is to provide people with digital devices to tackle access issues (as happened a lot during the pandemic),<sup>33</sup> this should be combined with training and support for using these devices. Moreover, creating and maintaining a password is imperative to access several online services (e.g. GP services, streaming platforms, online banking, and Government online services, to name a few), so that deficits in that area may adversely affect not only people's confidence in using these services but also to the security of their information. As these skills seem to be often missing in the general population as well,<sup>22</sup> it is important for future studies to explore the specific barriers the SMI population might face, and how to

## Measuring the digital divide among people with severe mental ill health using the essential digital skills framework

Table 2	
Essential digital skills framework	
	N (%)
Skills Domains	
Foundation Skills <sup>a</sup>	
Yes	142 (57.0)
No	105 (42.2)
Skills for Life <sup>b</sup>	
Yes	132 (93.0)
No	9 (6.3)
Skills for Work <sup>c</sup>	
Yes	46 (92.0)
No	4 (8.0)
Overall	
Essential skills for daily life <sup>a</sup>	
Yes (having Foundation AND Life Skills)	132 (53.4)
No (Lacking Foundation OR Life Skills)	114 (46.2)
Essential skills for professional life <sup>d</sup>	
Yes (Having Foundation AND Work Skills)	46 (75.4)
No (Lacking Foundation OR Work Skills)	14 (23.0)
<sup>a</sup> Out of total sample n=249.	
<sup>b</sup> Out of those having Foundation Skills, n=142.	
<sup>c</sup> Out of those being in paid employment and having Skills for Life, n=50.	
<sup>d</sup> Out of those being in paid employment, n=61.	

address these from a person-centered design perspective. Among the few people who had Foundation Skills but no Skills for Life, the greatest area of deficit was problem-solving skills. Evidence suggests that lack of problem-solving skills in general, not just digital, is a common cognitive deficit found in people with schizophrenia.<sup>34</sup>

We also found that people with no Foundation Skills were interested in learning more about half of the tasks they could not perform. Given that lack of interest is a commonly reported barrier to digital engagement,<sup>9</sup> this suggests that people with SMI might have the

motivation to learn new skills if appropriate support becomes available. Efforts to reduce the digital divide in the SMI population could include tailored training in digital skills along with confidence-building measures and measures to increase the motivation to address digital exclusion.<sup>13</sup> Reducing the digital divide has the potential to reduce health inequalities. Future studies should seek to consult with those with lived experience to explore motivation in beginning and sustaining digital skills training, preference for training mode and setting, as well as barriers to engaging with web-based resources specifically

related to SMI conditions (e.g. fluctuating levels of wellness, low energy and motivation, paranoid thoughts, suspiciousness and concerns about privacy and difficulty processing information;),<sup>35</sup> in order to develop accessible skills training programmes tailored to the needs of this population.

Older people in this study were less likely to have Foundation Skills, as also found in the general population.<sup>22</sup> This is unsurprising, as age is traditionally associated with less digital engagement in SMI<sup>14-16</sup> and the general population<sup>36,37</sup> alike. People with psychosis-spectrum disorders were at greater risk for not having Foundation Skills compared to those with bipolar disorder, after adjusting for people's age or employment status. This supports our previous findings that during the pandemic restrictions people with bipolar disorder were more likely to use the Internet a lot for their daily activities compared to people with psychosis.<sup>16</sup> Cognitive and occupational dysfunctions that are common in schizophrenia<sup>38</sup> might not facilitate the development of digital skills. It might also be that specific SMI-related barriers to digital engagement (e.g. reduced concentration, hallucinations, or paranoid ideas<sup>13,39</sup>) are more common in people with psychosis than bipolar disorder. Digital exclusion is becoming increasingly concentrated in vulnerable populations (including older people with SMI who are likely to be heavier users of healthcare services) leading to worse health outcomes through both direct and indirect routes.<sup>40</sup> The increasing use of digital technology in the delivery of health services will directly affect those who are digitally excluded as they will not be able to access services that could benefit them. This could range from accessing appointment booking systems to health-promoting services or applications, along with accessing video appointments. Given the current drive towards comprehensive digital healthcare within the NHS,<sup>20</sup> the importance of digital inclusion for health and social care should, and to some extent is, being acknowledged.<sup>41</sup> The particular needs and barriers experienced by people with SMI should inform this wider policy context.

## Measuring the digital divide among people with severe mental ill health using the essential digital skills framework

Figure 1


**Prevalence of skills deficiency among participants with no Foundation Skills (n = 105)**

Table 3

**Association of having foundation skills with sociodemographic and digital access variables**

Variables (Valid N)	Have foundation skills N (%)	Univariable model		Multivariable model (n = 229)	
		OR	95% CI	AdjOR	95% CI
Age (247)					
18–30	24 (85.7%)	10.29**	3.14–33.72	7.56*	1.75–32.66
31–45	45 (70.3%)	4.06**	1.90–8.68	3.60*	1.22–10.57
46–65	52 (53.1%)	1.94	0.99–3.78	1.93	0.75–4.97
66+	21 (36.8%)	1		1	
Gender (242)					
Male	74 (58.3%)	1.07	0.65–1.79	1.78	0.89–3.53
Female	65 (56.5%)	1		1	
Ethnicity (247)					
Other than White	22 (56.4%)	1.05	0.53–2.10	1.05	0.43–2.60
White	120 (57.7%)	1		1	

(Continued)



## Measuring the digital divide among people with severe mental ill health using the essential digital skills framework

Table 3 (Continued)

<b>Association of having foundation skills with sociodemographic and digital access variables</b>					
<b>Variables (Valid N)</b>	<b>Have foundation skills N (%)</b>	<b>Univariable model</b>		<b>Multivariable model (n=229)</b>	
		<b>OR</b>	<b>95% CI</b>	<b>AdjOR</b>	<b>95% CI</b>
<b>Socioeconomic deprivation (239)</b>					
Very high	29 (49.2%)	0.63	0.28–1.44	0.65	0.22–1.97
High	26 (52.0%)	0.71	0.30–1.66	0.44	0.14–1.36
Medium	34 (69.4%)	1.48	0.61–3.60	1.37	0.44–4.23
Low	24 (55.8%)	0.82	0.34–2.00	0.56	0.18–1.73
Very Low	23 (60.5%)	1		1	
<b>In paid employment (247)</b>					
Yes	50 (83.3%)	5.16**	2.47–10.79	2.67*	1.07–6.66
No	92 (49.2%)	1		1	
<b>Diagnosis (247)</b>					
Not recorded	19 (63.3%)	1.94	0.85–4.44	1.40	0.48–4.07
Other SMI	9 (60.0%)	1.69	0.57–5.04	1.43	0.40–5.17
Bipolar	58 (69.9%)	2.61*	1.45–4.71	3.67*	1.56–8.64
Psychosis spectrum	56 (47.1%)	1		1	
<b>Care setting (245)</b>					
Secondary care	86 (57%)	0.94	0.56–1.58	1.15	0.55–2.38
Primary care	55 (58.5%)	1		1	
<b>Device ownership (243)</b>					
No	2 (6.5%)	0.04**	0.01–0.16	0.26	0.04–1.70
Yes	138 (65.1%)	1		1	
<b>Internet access at home (244)</b>					
No	3 (7.9%)	0.04**	0.01–0.15	0.14*	0.03–0.70
Yes	137 (66.5%)	1		1	

OR: odds ratio; CI: confidence interval; SMI: severe mental ill health.  
\*p < .05, \*\*p < .001.

The SPIDER study is not free of limitations. The sample size may have been underpowered for the estimate of the logistic regression, but this analysis was exploratory. We used a non-random sample that might lead to respondent

biases and limited generalisability, although the use of a sampling framework increases our confidence that we were recruiting across a good spread of the SMI population. Skills were self-evaluated by participants rather than

completing objective tests. Self-evaluation might be susceptible to other fluctuating factors (e.g. current mood), as well as response and social desirability bias (e.g. n = 134, 53.8% of the sample completed the survey over the phone

## Measuring the digital divide among people with severe mental ill health using the essential digital skills framework

with a researcher). It should be noted though that organising objective tests, especially in the context of the pandemic restrictions, would present several practical challenges. Participants were also assessed on the basis of predetermined tasks, rather than personalised tasks related to the specific activities they want/need to complete in their everyday lives. However, that would affect the comparability of results across different populations and existing datasets. Regarding digital access, we did not examine whether the devices that people owned were current enough or had to be shared with other members of the household, as often happened during the pandemic.<sup>42</sup> Despite the high rates of device ownership in our sample, real unobstructed access might be undermined by obsolete technologies or multiusership. Finally, in this study, we examined generic digital skills rather than skills and factors that may be more pertinent to this population and their use of digital health services (e.g. e-health literacy, online help-seeking and service use), which should be considered in future research.

## CONCLUSION

This study highlighted deficits in digital skills in the SMI population that are worse than in the general population.

This might suggest that an already socially disadvantaged population is at risk for further exclusions due to digital skills deficits, as many health services and social connections have moved to digital platforms during 2020/2021. Importantly and worryingly, this includes vital health services many of which might continue to operate digitally into the future. Services including the NHS need to be aware and have a responsibility to actively and immediately accommodate those with SMI who would prefer face-to-face rather than online contact, and whose health will likely suffer through the digital divide. In addition, further funding is needed for research into a widely available, person-centred, accessible training programme, co-produced with end-users to identify and remove barriers related to lack of digital skills.

## ACKNOWLEDGEMENTS

This report is independent research supported by the National Institute for Health Research Yorkshire and Humber Applied Research Collaboration. The views expressed in this publication are those of the author(s) and not necessarily those of the National Institute for Health Research or the Department of Health and Social Care. We thank the participants in the SPIDER study and NHS mental health staff for their support with this study.

## CONFLICT OF INTEREST

The author(s) declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.


## FUNDING

The author(s) disclosed receipt of the following financial support for the research, authorship and/or publication of this article: This work was supported by the Closing the Gap network which is funded by UK Research and Innovation, and their support is gratefully acknowledged (Grant reference: ES/S004459/).

## ETHICAL APPROVAL

Ethical approval for this study was granted by the Health Research Authority Northwest – Liverpool Central Research Ethics Committee (REC reference 20/NW/0276). Completing and returning the survey were considered implying consent to participate in the study.

## ORCID IDS

P Spanakis  <https://orcid.org/0000-0002-2732-8300>

L Walker  <https://orcid.org/0000-0003-2459-7860>

## SUPPLEMENTAL MATERIAL

Supplemental material for this article is available online.

## References

- Alfano V, Ercolano S. The efficacy of lockdown against COVID-19: a cross-country panel analysis. *Appl Heal Econ Heal Pol* 2020;**18**:509–17.
- May T. Lockdown-type measures look effective against covid-19. *BMJ* 2020;**370**:m2809.
- Chen S, Jones PB, Underwood BR *et al.* The early impact of COVID-19 on mental health and community physical health services and their patients' mortality in Cambridgeshire and Peterborough, UK. *J Psychiatr Res* 2020;**131**:244–54.
- Johns G, Tan J, Burhouse A *et al.* A visual step-by-step guide for clinicians to use video consultations in mental health services: NHS examples of real-time practice in times of normal and pandemic healthcare delivery. *BJPsych Bull* 2020;**44**:277–84.
- Johnson S, Dalton-Locke C, Vera San Juan N *et al.* Impact on mental health care and on mental health service users of the COVID-19 pandemic: a mixed methods survey of UK mental health care staff. *Soc Psychiatry Psychiatr Epidemiol* 2020;**56**:25–37.
- Mehta J, Yates T, Smith P *et al.* Rapid implementation of Microsoft Teams in response to COVID-19: one acute healthcare organisation's experience. *BMJ Health Care Inform* 2020;**27**(3):e100209.
- Nune A, Iyengar K, Ahmed A *et al.* Challenges in delivering rheumatology care during COVID-19 pandemic. *Clin Rheumatol* 2020;**39**(9):2817–21.
- Shah N, Thakrar A, Visvanathan S *et al.* Virtual online consultation platforms for secondary care: a review of the options. *BMJ Innov* 2021;**7**:135–40.
- Lloyds Bank. UK consumer digital index 2021, 2021. Available online at: <https://www.lloydsbank.com/banking-with-us/whats-happening/consumer-digital-index.html>
- Watts G. COVID-19 and the digital divide in the UK. *Lancet Digit Health* 2020;**2**(8):e395–6.
- Parliament Office of Science and Technology (POST). Developing essential digital skills, 2021. Available online at: <https://post.parliament.uk/research-briefings/post-pn-0643/>
- Spanakis P, Peckham E, Mathers A *et al.* The digital divide: amplifying health inequalities for people with severe mental illness in the time of COVID-19. *Br J Psychiatry* 2021;**219**(4):529–31.
- Greer B, Robotham D, Simblett S *et al.* Digital exclusion among mental health service users: qualitative investigation. *J Med Internet Res* 2019;**21**:e11696.
- Robotham D, Satkunanathan S, Doughty L *et al.* Do we still have a digital divide in mental health? A five-year survey follow-up. *JMIR* 2016;**18**:e309.
- Tobitt S, Percival R. Switched on or switched off? A survey of mobile, computer and Internet use in a community mental health rehabilitation sample. *J Ment Heal* 2017;**28**:4–10.
- Spanakis P, Heron P, Walker L *et al.* Use of the Internet and digital devices among people with severe mental ill health during the COVID-19 pandemic restrictions (Pre-print). *Medrxiv*. Epub 2021 June 21. DOI: 10.1101/2021.06.17.21259095.
- Dickerson F, Origoni A, Rowe K *et al.* Risk factors for natural cause mortality in a cohort of 1494 persons with serious mental illness. *Psychiatry Res* 2021;**298**:113755.
- Armitage R. Physical health checks for people with severe mental illness during COVID-19. *British J Gen Pract* 2021;**71**:171.
- Badcock JC, Adery LH, Park S. Loneliness in psychosis: a practical review and critique for

## Measuring the digital divide among people with severe mental ill health using the essential digital skills framework

- clinicians. *Clin Psych Sci Pract* 2020;**27**:e12345.
20. National Health Service (NHS). The NHS long term plan, 2019. Available online at: <https://www.longtermplan.nhs.uk/>
  21. Spanakis P, Heron P, Walker L *et al*. Skills and Proficiency in Digital Essential Requirements (SPIDER) – pre-registered analysis plan, 2021. DOI: 10.17605/OSF.IO/BS9TE.
  22. Lloyds Bank. UK Consumer Digital Index 2020, 2020. Available online at: [https://www.lloydsbank.com/assets/media/pdfs/banking\\_with\\_us/whats-happening/lb-consumer-digital-index-2020-report.pdf](https://www.lloydsbank.com/assets/media/pdfs/banking_with_us/whats-happening/lb-consumer-digital-index-2020-report.pdf)
  23. The Scottish Council for Voluntary Organisations (SCVO). Understanding and measuring essential digital skills, 2018. Available online at: <https://storage.googleapis.com/scvo-assets/test/digitalparticipation/documents/eds-measuring-understanding.pdf?version=0.0.29>
  24. Ministry of Housing Communities & Local Government. *English indices of deprivation*. London: Ministry of Housing Communities; 2019.
  25. Liberati E, Richards N, Parker J *et al*. Remote care for mental health: qualitative study with service users, carers and staff during the COVID-19 pandemic. *BMJ Open* 2021;**11**:e049210.
  26. Healthwatch England. Locked out: digitally excluded people's experiences of remote GP appointments, 2021. Available online at: <https://www.healthwatch.co.uk/report/2021-06-16/locked-out-digitally-excluded-peoples-experiences-remote-gp-appointments>
  27. Inkster B, O'Brien R, Selby E *et al*. Digital health management during and beyond the COVID-19 pandemic: opportunities, barriers, and recommendations. *JMIR Mental Health* 2021;**7**:e19246.
  28. Ajnakina O, Stubbs B, Francis E *et al*. Employment and relationship outcomes in first-episode psychosis: a systematic review and meta-analysis of longitudinal studies. *Schizophrenia Res* 2021;**231**:122–33.
  29. Holm M, Taipale H, Tanskanen A *et al*. Employment among people with schizophrenia or bipolar disorder: a population-based study using nationwide registers. *Acta Psychiatrica Scandinavica* 2020;**143**:61–171.
  30. Marwaha S, Durrani A, Singh S. Employment outcomes in people with bipolar disorder: a systematic review. *Acta Psychiatrica Scandinavica* 2013;**128**:179–93.
  31. Burke-Miller JK, Cook JA, Grey DD *et al*. Demographic characteristics and employment among people with severe mental illness in a multisite study. *Community Ment Health J* 2006;**42**(2):143–59.
  32. Fundora C, Cruz M, Barone K *et al*. Lifetime employment in schizophrenia: correlates of developing long term unemployment after being employed before. *Cogn Neuropsychiatry* 2021;**26**(2):95–106.
  33. All Parliamentary Group on Social Integration (APPG). Social connection in the COVID-19 crisis, 2020. Available online at: <https://ec.europa.eu/migrant-integration/sites/default/files/2020-06/Social-Connection-in-the-COVID-19-Crisis.pdf>
  34. Xia J, Li C. Problem solving skills for schizophrenia. *Cochr Datab Syst Rev* 2007;**2007**:CD006365.
  35. Arnold C, Williams A, Thomas N. Engaging with a web-based psychosocial intervention for psychosis: qualitative study of user experiences. *JMIR Mental Health* 2020;**7**:e16730.
  36. The Office of Communications. Adult's media use and attitudes, 2020. Available online at: <https://www.ofcom.org.uk/research-and-data/media-literacy-research/adults/adults-media-use-and-attitudes>
  37. National Health Service Digital (NHS Digital). Digital inclusion guide for health and social care, 2019. Available online at: <https://digital.nhs.uk/about-nhs-digital/our-work/digital-inclusion>
  38. Cowman M, Holleran L, Lonergan E *et al*. Cognitive predictors of social and occupational functioning in early psychosis: a systematic review and meta-analysis of cross-sectional and longitudinal data. *Schizophrenia Bull* 2021;**47**:1243–53.
  39. Schrank B, Sibitz I, Unger A *et al*. How patients with schizophrenia use the internet: qualitative study. *J Med Internet Res* 2010;**12**:e70.
  40. Health Education England. The topol review. Preparing the healthcare workforce to deliver the digital future, 2019. Available online at: <https://societyandethicsresearch.wellcomeconnectingscience.org/catalogue/the-topol-review-preparing-the-healthcare-workforce-to-deliver-the-digital-future-final>
  41. NHS Digital. Why digital inclusion matters to health and social care, 2019. Available online at: <https://digital.nhs.uk/about-nhs-digital/our-work/digital-inclusion/digital-inclusion-in-health-and-social-care#:~:text=In%20an%20increasingly%20digital%20world,the%20skills%20to%20use%20it>
  42. Child Poverty Action Group. The cost of learning in lockdown, 2021. Available online at: <https://cpag.org.uk/policy-and-campaigns/report/cost-learning-lockdown-family-experiences-school-closures>