



This is a repository copy of *The timing of the Svalbardian Orogeny in Svalbard: a review.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/190514/>

Version: Published Version

Article:

Koehl, J.-B.P., Marshall, J.E.A. and Lopes, G. orcid.org/0000-0002-6866-5127 (2022) The timing of the Svalbardian Orogeny in Svalbard: a review. *Solid Earth*, 13 (8). pp. 1353-1370. ISSN 1869-9510

<https://doi.org/10.5194/se-13-1353-2022>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



The timing of the Svalbardian Orogeny in Svalbard: a review

Jean-Baptiste P. Koehl¹, John E. A. Marshall², and Gilda Lopes^{3,4}

¹Department of Geosciences, University of Oslo, P.O. Box 1047 Blindern, NO-0316 Oslo, Norway

²School of Ocean and Earth Science, University of Southampton, National Oceanography Centre, European Way, SO143ZH Southampton, UK

³CIMA – Centre for Marine and Environmental Research, Universidade do Algarve, Campus de Gambelas, 8005-139 Faro, Portugal

⁴School of Biosciences, University of Sheffield, Western Bank, S102TN Sheffield, UK

Correspondence: Jean-Baptiste P. Koehl (jeanbaptiste.koehl@gmail.com)

Received: 1 February 2022 – Discussion started: 1 March 2022

Revised: 8 August 2022 – Accepted: 9 August 2022 – Published: 30 August 2022

Abstract. In the Late Devonian to earliest Mississippian, Svalbard was affected by a short-lived episode of deformation named the Svalbardian Orogeny. This event resulted in intense folding and thrusting in Devonian sedimentary successions. Deformation stopped prior to the deposition of Carboniferous to Permian sedimentary strata of the Billefjorden and Gipsdalen groups, which lie unconformably over folded Devonian strata. Later on, presumed Svalbardian structures were reworked during Eurekan tectonism in the early Cenozoic and partly eroded. At present, records of Svalbardian deformation are only preserved in narrow N–S-trending belts in central, northern, western, and southern Spitsbergen. Despite extensive field studies, the timing of the Svalbardian Orogeny is poorly constrained and remains a matter of debate in places because of conflicting ages and because of the complex tectonic history of Svalbard. The present contribution aims at reviewing and discussing all available age constraints for Svalbardian tectonism, including notably palynological, paleontological, and geochronological evidence. This has great implications for the plate tectonic reconstructions of Arctic regions and for the tectonic history of Svalbard. Palynological and paleontological evidence suggest that the Mimerdalen Subgroup is upper Givetian to lower Frasnian (ca. 385–380 Ma) in age and that the Billefjorden Group is mid-Famennian to Upper Mississippian (ca. 365–325 Ma) in age, constraining the Svalbardian event in central and northern Spitsbergen to 383–365 Ma if it ever occurred. Palynological ages indicate that the Adriabukta Formation in southern Spitsbergen is Middle Mississippian and therefore cannot have been involved in the Svalbardian event, thus suggest-

ing that all the deformation in southern Spitsbergen is early Cenozoic in age and that strain-partitioning processes had a major role in localizing deformation in weaker stratigraphic units. The few geochronological age constraints yielding Late Devonian–Mississippian ages in Svalbard may reflect either Svalbardian contraction or extensional processes and are therefore of no use to validate or invalidate the occurrence of the Svalbardian event. On the contrary, the contradicting lines of evidence used to support the occurrence of the Svalbardian event and new regional geophysical studies suggest that Svalbard was subjected to continuous extension from the late Silurian to early Permian times.

1 Introduction

The Svalbardian Orogeny, also known as the Innuitian or Ellesmerian Orogeny, refers to a short-lived episode of contraction and/or transpression that affected all levels of the crust and occurred in the Late Devonian (to earliest Mississippian?) when parts of the tectonic plates now constituting most of the Arctic (Laurentia and Baltica) collided with each other and deformed Proterozoic to mid-Paleozoic sedimentary basins and basement rocks in northeastern Russia (Malyshev et al., 2011; Luchitskaya et al., 2015), Canada (Thorsteinsson and Tozer, 1970; Trettin, 1973, 1991; Embry and Klovan, 1976; Embry, 1991; Harisson, 1995; Harisson and Brent, 2005; Piepjohn et al., 2008, 2013; Piepjohn and von Gosen, 2017), and Alaska (Grantz and May, 1984; Lane, 2007; Kumar et al., 2011); Proterozoic to Silurian metasedi-

mentary rocks in northern and northeastern Greenland (Higgins et al., 2000; Piepjohn et al., 2015); and Devonian collapse basins and Precambrian to lower Paleozoic basement in Norway (Roberts, 1983; Osmundsen et al., 1998) and Svalbard (Vogt, 1938; Harland et al., 1974; McCann, 2000; Piepjohn, 2000; Piepjohn et al., 2000; Fig. 1).

In Svalbard, Svalbardian contraction (transpression?) followed the Caledonian Orogeny (ca. 460–410 Ma; Horsfield, 1972; Dallmeyer et al., 1990; Johansson et al., 2004, 2005; Faehnrich et al., 2020) and subsequent deposition of thick upper Silurian–Devonian sedimentary successions during late- to post-orogenic collapse (Gee and Moody-Stuart, 1966; Friend et al., 1966; Friend and Moody-Stuart, 1972; Murascov and Mokin, 1979; Manby and Lyberis, 1992; Manby et al., 1994; Friend et al., 1997; McCann, 2000; Dallmann and Piepjohn, 2020) and led to the final accretion of Svalbard's three basement terranes (Harland and Wright, 1979; Ohta et al., 1989, 1995; Harland et al., 1992; Gee and Page, 1994). Although early accounts envisioned hundreds to thousands of kilometer-scale strike-slip movements along N–S-striking faults like the Billefjorden and Lomfjorden fault zones (e.g., Harland et al., 1974, 1992), more recent studies have shown that such large-scale strike-slip movements are unlikely (McCann, 2000; Michalski et al., 2012). In addition, new geochronological and structural work in northern Svalbard shows that collapse-related extension leading to the exhumation of the Bockfjorden Anticline as a core complex lasted from the late Silurian to the Late Devonian (Famenian at 368.42 ± 0.81 Ma; Braathen et al., 2018), i.e., possibly overlapping with Svalbardian contraction.

Evidence of Svalbardian tectonism includes dominantly west-verging folds and thrusts within several kilometer-thick, Devonian, late- to post-orogenic, collapse-related sedimentary rocks in central and northern Spitsbergen (Andrée Land Group, including the Mimerdalen Subgroup; Vogt, 1938; Harland et al., 1974; Manby and Lyberis, 1992; Manby et al., 1994; Friend et al., 1997; Piepjohn and Dallmann, 2014; Dallmann and Piepjohn, 2020) and dominantly east-verging folds and thrusts in Devonian (to Middle Mississippian?) sedimentary rocks in southern Spitsbergen (Marietoppen and Adriabukta formations; Dallmann, 1992; Bergh et al., 2011). The latter were interpreted to be unconformably covered by presumed undeformed, shale-rich, poorly exposed Triassic strata in Røkensåta (Dallmann, 1992). Important uncertainties around this interpretation are discussed for the first time in the present work and suggest that interpretation based on this outcrop should be given little to no credit.

Shortly (i.e., immediately up to a few million years) after the end of Svalbardian deformation, partly eroded Devonian sedimentary rocks in Spitsbergen were covered by fluvial, coal-rich deposits of the Billefjorden Group, possibly during widespread latest Devonian–Mississippian extension, and shallow marine strata of the Gipsdalen Group mostly deposited within narrow, kilometer-wide to tens of kilometers wide, N–S- to NW–SE-trending basins (Cutbill




Figure 1. Topographic and bathymetric map around Spitsbergen modified after Jakobsson et al. (2012). The location of exploration well 7816/12-1 is shown in white. Abbreviations are as follows: Ad stands for Adriabukta; Bi stands for Billefjorden, Bo stands for Blomstrandhalvøya, Br stands for Brøggerhalvøya, Fi stands for Fiskeknatten, Ga stands for Garmdalen, Hs stands for Hornsundneset, Hu stands for Hugindalen, Kg stands for Kongsfjorden, Kr stands for Krosspynten, Mi stands for Midterhuken, Py stands for Pyramiden, Re stands for Reindalspasset, Rø stands for Røkensåta, Tr stands for Triungen, and Yg stands for Yggdrasilkampen.

and Challinor, 1965; Maher, 1996; McCann and Dallmann, 1996; Braathen et al., 2011; Koehl and Muñoz-Barrera, 2018; see Fig. 2 for stratigraphy). Subsequently, Svalbardian structures were reworked by Eureka contraction and/or transpression during the opening of the Labrador Sea and Baf-




Figure 2. Late Paleozoic stratigraphic chart of the areas discussed in the text. The ages in the timescale are in Ma and are from Walker et al. (2018).

fin Bay between Canada and Greenland (Chalmers and Pulvertaft, 2001; Oakey and Chalmers, 2012), which resulted in the formation of the West Spitsbergen Fold-and-Thrust Belt between Kongsfjorden and Sørkapp (Harland, 1969; Lowell, 1972; Harland and Horsfield, 1974; Maher et al., 1986; Dallmann et al., 1988, 1993; Andresen et al., 1994; Bergh and Grogan, 2003; see location in Fig. 1) and of the Central Tertiary Basin in central Spitsbergen (Larsen, 1988; Petersen et al., 2016). As a result, Svalbardian structures were overprinted and reworked and now commonly display the same trends, plunges, strikes, dips, and kinematics as Eureka structures throughout the Arctic and, in many occurrences, coincide with (and are indistinguishable from) Eureka structures (e.g., Birkenmajer, 1964; Piepjohn et al., 2007, 2008, 2013, 2015; Bergh et al., 2011; Piepjohn and von Gosen, 2017; Dallmann and Piepjohn, 2020).

At present, original (unmodified) Svalbardian deformation is preserved only in a few narrow N–S-trending belts, including Dickson Land (Michaelsen et al., 1997; Piepjohn et al., 1997b; Michaelsen, 1998; Piepjohn, 2000), Andrée Land (Dallmann and Piepjohn, 2020), and Blomstrandhalvøya (Thiedig and Manby, 1992; Buggisch et al., 1994; Fig. 1). The best and most well-constrained example of Svalbardian tectonism is observed in central and northern Spitsbergen, where folded Lower to lowermost Upper Devonian sedimentary rocks of the Andrée Land Group and Mimerdalen Subgroup are unconformably overlain by apparently undeformed uppermost Devonian to lowermost Permian sedimentary strata of the Billefjorden and Gipsdalen groups (Vogt, 1938; Harland et al., 1974; Piepjohn, 2000; Piepjohn et al., 2000). The structures in the Dickson Land area are actually highly questionable and are addressed in two separate papers (Koehl et al., 2022b, c) and will therefore not be reviewed in detail in the present contribution.

Recent U–Th–Pb geochronology on monazite grains yielded 373–355 Ma (latest Devonian to earliest Mississippian) ages for amphibolite facies metamorphism along a gently west-dipping shear zone in Prins Karls Forland (location in Fig. 1) crosscutting Neoproterozoic basement rocks. These data provide evidence and time constraints for Svalbardian tectonism at depth of ca. 15 km (Faehnrich et al., 2017; Majka and Kościńska, 2017; Schneider et al., 2018; Kościńska et al., 2020). Potential Svalbardian (greenschist) facies metamorphism and mylonitization was also potentially identified in Oscar II Land (location in Fig. 1) and dated to 365–344 Ma through ^{40}Ar – ^{39}Ar and U–Th–Pb geochronology (Barnes et al., 2020).

Nonetheless, despite extensive previous works, Svalbardian tectonism at shallow crustal levels lacks accurate time constraints, and it is possible that, in certain places, structures ascribed to this event may have formed during the early Paleozoic Caledonian Orogeny or during the early Cenozoic Eureka tectonic event (e.g., Rippington et al., 2010). Distinguishing Svalbardian from Eureka structures is problematic. In Arctic Canada and Greenland, Ellesmerian structures

are thought to be overprinted almost everywhere by subsequent Eureka structures (e.g., Piepjohn et al., 2015). This is also the case to some extent in Svalbard, where Svalbardian and Eureka folds and thrusts are both believed to show dominantly east-verging geometries in the south but opposite vergence in the north where Svalbardian structures display mostly top-west attitudes (Dallmann and Piepjohn, 2020). Another issue arises from the complexity of the Eureka fold-and-thrust belt throughout Spitsbergen, which involves numerous décollements localized in shale-rich stratigraphic units, such as the Lower Triassic (Maher, 1984; Maher et al., 1986, 1989; Andresen et al., 1988; Bergh and Andresen, 1990; Haremo et al., 1990; Haremo and Andresen, 1992; Andresen et al., 1992; Dallmann et al., 1993; Bergh et al., 1997).

Furthermore, east- to northeast-plunging folds trending parallel to the inferred late- to post-orogenic extension direction in Middle Devonian collapse basins in western Norway were initially interpreted as Late Devonian to Mississippian, Svalbardian contractional and/or transpressional structures (Roberts, 1983). These are now known to have formed as transtensional folds during extensional collapse of the Caledonides (Chauvet and Séranne, 1994; Osmundsen and Andersen, 1994; Fossen et al., 2013). Thus, it is paramount to carefully constrain the timing of Svalbardian (and Ellesmerian) deformation throughout the Arctic to be able to evaluate the extent and impact of this tectonic event from a regional perspective and understand its interplay with potentially coeval collapse processes (e.g., Braathen et al., 2018; Maher et al., 2022).

Thus far, Svalbardian deformation is thought to have occurred during the Late Devonian to Early Mississippian, possibly initiating in the late Frasnian to Famennian (Vigran, 1964; Allen, 1965, 1973; Pčelina et al., 1986; Brinkmann, 1997; Schweitzer, 1999; Piepjohn et al., 2000; Fig. 2). The onset of deformation was presumably recorded by the deposition and syn-depositional deformation of coarse-grained sedimentary rocks of the Mimerdalen Subgroup in the late Famennian (Planteryggen and Plantekløfta formations in Fig. 2; Piepjohn and Dallmann, 2014). Deformation is believed to have stopped prior to the deposition of sedimentary rocks of the Billefjorden Group in the late Tournaisian (Vogt, 1938; Piepjohn, 2000).

The present contribution focuses on the debate around the timing of Svalbardian tectonism throughout Spitsbergen. In northern Spitsbergen, Svalbardian deformation was constrained to the late Famennian to earliest Mississippian by the identification of one specimen of *Retispora lepidophyta* in folded rocks of the Plantekløfta Formation (Schweitzer, 1999; Piepjohn et al., 2000). However, recent palynological and paleontological studies in northern and central Spitsbergen suggest slightly revised ages for the stratigraphic units used to constrain the timing of the Svalbardian Orogeny, including a Middle Devonian (minimum upper Givetian) age for rocks of the Tordalen Formation (Mimerdalen Subgroup; Berry and Marshall, 2015; Newman et al., 2019) and

a mid-Famennian age for the base of the Billefjorden Group (Scheibner et al., 2012; Lindemann et al., 2013; Marshall et al., 2015; Lopes, G., personal observation, 2019). In addition, the timing of Svalbardian deformation varies somewhat from north to south in Spitsbergen, and study of a palynological assemblage in the Adriabukta Formation in southern Spitsbergen constrained Svalbardian folding and faulting to the Viséan (Middle Mississippian; Birkenmajer and Turnau, 1962; Fig. 2). The present contribution reviews time constraints for Svalbardian tectonism in central, northern, southern, and western Svalbard and briefly discusses their implications.

Constraining the timing of the Svalbardian Orogeny in Svalbard with accuracy is of importance for paleogeographic and plate tectonics reconstructions in the Arctic. It is also important for the tectonic history of Svalbard, e.g., to evaluate potential interplay between late- to post-Caledonian extensional collapse, which resulted in the deposition of several kilometer-thick collapse basins (e.g., Murascov and Mokin, 1979; Manby and Lyberis, 1992; Friend et al., 1997; Braathen et al., 2018), and contractional tectonic processes that resulted in intense folding of these deposits (Vogt, 1938; Piepjohn, 2000; Dallmann and Piepjohn, 2020). Furthermore, the present study has implications for the methods used by geologists to interpret tectonic events worldwide, e.g., in pointing out that field studies based on long-distance observation of poorly exposed and inaccessible transects should be given little (if any) credit.

2 Review of age constraints in northern and central Spitsbergen

2.1 Age of the Mimerdalen Subgroup

The identification of one specimen of *Retispora lepidophyta* within strata of the Plantekløfta Formation by Brinkmann (1997, Table 14.3 therein) and Piepjohn et al. (2000; published in Schweitzer, 1999, Plate 6 in their Fig. 10 and Plate 7 in their Fig. 1) suggests a late Famennian age for the top of the Plantekløfta Formation and hence that Svalbardian tectonism terminated during the Famennian to Tournaisian in northern and central Spitsbergen.

Recent studies clearly demonstrated that the interpretation of *Retispora lepidophyta* by Brinkmann (1997), Schweitzer (1999), and Piepjohn et al. (2000) is erroneous. Notably, the lone figured specimen interpreted as *Retispora lepidophyta* by Schweitzer (1999) and Piepjohn et al. (2000) differs in size and shows significantly different morphological structures from typical *Retispora lepidophyta* (Playford, 1976; Berry and Marshall, 2015, Supplement DR3 therein; see also Supplement S1). In addition the fovea that characterize the spore's exoexine appear to be the result of damage by cubic diagenetic pyrite. Attempts have been made to locate the *Retispora lepidophyta* specimen figured

in Brinkmann (1997) and Schweitzer (1999) and used by Piepjohn et al. (2000) to date Svalbardian tectonism in central Spitsbergen for further analysis. These attempts were unfortunately unsuccessful (Marshall, J. E. A., personal observation, 2020).

Berry and Marshall (2015) re-evaluated the age of the Plantekløfta Formation to be early Frasnian based on fossils and miospores (ca. 383–380 Ma; see also their Supplements; Fig. 2). In addition, the paleontological study of Newman et al. (2019, 2020, 2021) recorded the presence of articulated fish in the Fiskekløfta Member of the Tordalen Formation (Fig. 2), i.e., undoubtedly in situ fossils, demonstrating a late to latest Givetian (ca. 385–383 Ma; Middle Devonian) age for this stratigraphic unit instead of late Famennian. If the relatively coarse grain size of the sedimentary deposits of the Planteryggen and Plantekløfta formations indeed reflects the onset of Svalbardian tectonism as suggested by Piepjohn and Dallmann (2014), the new paleontological and palynological ages constrain the initial phase of the Svalbardian Orogeny at 383–380 Ma.

A late Famennian age for the Plantekløfta Formation based on the lone specimen of *Retispora lepidophyta* in central Spitsbergen is the only contradictory evidence against a mid-Famennian age for the base of the Billefjorden Group and older age for the Mimerdalen Subgroup (Scheibner et al., 2012; Lindeman et al., 2013; Berry and Marshall, 2015; Marshall et al., 2015; Lopes et al., 2019; Newman et al., 2019; Lopes, G., personal observation, 2019).

2.2 Age of the Billefjorden Group

Recent palynological studies in central Spitsbergen dated the base of the Billefjorden Group in Triungen (see Fig. 1 for location) to the mid-Famennian (maximum ca. 365 Ma; Lindemann et al., 2013; Marshall et al., 2015; Lopes, G., personal observation, 2019; Fig. 2). At least 30 samples contained characteristic Famennian spore assemblages including *Cyrtospora cristifer*, *Cornispora monocornata*, *Cornispora bicornata*, *Cornispora tricornata*, *Lophozonotriletes lebedianensis*, *Knoxisporites dedaleus*, *Grandispora gracilis*, *Spelaeotriletes papulosus*, *Cristatisporites lupinovitchi*, *Lagenosisporites* sp., *Grandispora famensis*, and *Tergobulaspores immensus* (Marshall et al., 2015). Some samples from the lower part of the Billefjorden Group in Billefjorden also contained *Retispora lepidophyta* (Lopes, G., personal observation, 2019). These spore assemblages were also identified in sedimentary rocks in the lower part of the Billefjorden Group in northeastern Spitsbergen (Scheibner et al., 2012), thus strengthening a Famennian age for the base of this stratigraphic unit throughout northern and central Spitsbergen. Note that the base of the Billefjorden Group in Bjørnøya was also dated as Famennian based on palynology (Kaiser, 1970; Worsley and Edwards, 1976; Lopes et al., 2021). This strongly suggests that the Svalbardian deformation, which ended prior to the deposition of the Billefjorden

Group (Piepjohn, 2000), must have been terminated by the mid-Famennian in central and northern Spitsbergen. This implies a maximum duration of 18 Ma for this tectonic event.

Piepjohn and Dallmann (2014) proposed that the mid- to late-Famennian spores identified in the lower part of the Billefjorden Group were reworked based on their identification of one specimen of *Retispora lepidophyta* within the Plantekløfta Formation (Piepjohn et al., 2000). However, since this specimen clearly is a misidentification (Berry and Marshall, 2015, Supplement DR3 therein), the claim of reworking of mid- to late-Famennian spores found within the base of the Billefjorden Group in Triungen no longer has any supporting argument in Svalbard, and neither does the claim of Piepjohn et al. (2000) that the older Devonian spores found in the sample with the misidentified specimen of *Retispora lepidophyta* were reworked.

2.3 Other time constraints for deformation in central and northern Spitsbergen

At least some of the deformation in Lower to lowermost Upper Devonian strata of the Andrée Land Group and Mimerdalen Subgroup in central and northern Spitsbergen is early Cenozoic in age because uppermost Devonian to Mississippian strata of the Billefjorden Group, which overlie the Andrée Land Group and Mimerdalen Subgroup in the area, are intensely sheared top-west, e.g., in Pyramiden (Koehl, 2021) and Garmdalen (Koehl et al., 2020, 2022b; locations in Fig. 1). This is further supported by the interpretation of seismic data adjacent to nearshore portions of Billefjorden showing the presence of a bedding-parallel décollement between the Wood Bay Formation and the Gipsdalen Group (Koehl et al., 2020). These suggest a significant impact of strain partitioning during Eurekan deformation. Eurekan strain partitioning is further illustrated by tight plastic folding of Lower Devonian strata of the Andrée Land Group and brittle brecciation of the unconformity with Upper Pennsylvanian to Permian strata in Yggdrasilkampen (Manby et al., 1994, Fig. 11 therein) and by décollements within Middle Devonian deposits near the Billefjorden Fault Zone in Wijdefjorden (Marshall, J. E. A., personal observation, 2022; see Fig. 1 for location).

Another argument corroborating these data is the involvement in folding of Carboniferous picritic dikes dated at ca. 357 Ma (Evdokimov et al., 2006; monchiquite dikes in Gayer et al., 1966, and Manby and Lyberis, 1996) intruding Lower Devonian sedimentary rocks at Krosspynten (see location in Fig. 1).

In addition, part of the deformation recorded by Lower to lowermost Upper Devonian strata of the Andrée Land Group and Mimerdalen Subgroup is possibly related to extensional detachment folding in the Devonian (Chorowicz, 1992; Roy, 2007, 2009; Roy et al., 2022). This is also supported by recent field and geochronological studies in north-

western Spitsbergen (Braathen et al., 2018, 2020; Maher et al., 2022).

Thus, it is unclear how much (if any at all) of the deformation observed within Lower to lowermost Upper Devonian strata of the Andrée Land Group and Mimerdalen Subgroup in central and northern Spitsbergen actually reflects Ellesmerian tectonism.

3 Review of age constraints in southern Spitsbergen

3.1 Age of the Adriabukta Formation

In southern Spitsbergen, Lower to Middle Devonian sedimentary rocks of the Marietoppen Formation (time equivalent to the Pragian to Eifelian Wood Bay and Grey Hoek formations of the Andrée Land Group in central and northern Spitsbergen; Fig. 2) unconformably overlie Precambrian to early Paleozoic basement rocks and are overlain by sedimentary strata of the Adriabukta Formation that were deformed into tight east-verging folds presumably during Svalbardian tectonism (Birkenmajer, 1964). The age of the Adriabukta Formation was dated to the Middle Mississippian through analysis of palynomorphs from black shales at the base and within the formation (Birkenmajer and Turnau, 1962; Fig. 2). Dallmann et al. (1999) noted that because of the age discrepancy between the Middle Mississippian Adriabukta Formation and the Lower to lowermost Upper Devonian Andrée Land Group in central and northern Spitsbergen, the folding of the Adriabukta Formation could not be correlated to Svalbardian folding. Nevertheless, in 2011, Dallmann suggested that the Adriabukta Formation is actually Late Devonian in age based on structural correlation between presumed Svalbardian structures in the Adriabukta Formation and Svalbardian fold-and-thrust belts in central and northern Spitsbergen, thus generating a debate around the actual age of the formation. This is referenced as “Dallmann, W., personal communication (2009)” in Bergh et al. (2011).

The speculation and debate initiated by Dallmann around the age of the Adriabukta Formation is based on neither published material nor specific scientific evidence. By contrast, Birkenmajer and Turnau (1962) identified a count of 350 spore specimens from the Adriabukta Formation including specimens of *Lycospora*, *Tripartites*, and *Triquitrites*, which were then and are still characteristic of the Middle Mississippian (Hughes and Playford, 1961; Playford, 1962, 1963; Clayton 1996). Later palynological studies in Svalbard (Billefjorden; Lopes et al., 2019) and Europe (Clayton et al., 1977) corroborate the Middle Mississippian ages obtained by Birkenmajer and Turnau (1962) for the Adriabukta Formation. Thus, the debate around the Middle Mississippian ages obtained for the Adriabukta Formation by Birkenmajer and Turnau (1962) is not justified, and a Middle Mississippian age is entirely justified. The Adriabukta Formation in south-

ern Spitsbergen is therefore a time-equivalent example for the Billefjorden Group (e.g., Lopes et al., 2019).

The Middle Mississippian age of the Adriabukta Formation suggests that folding within this stratigraphic unit cannot be Late Devonian and is therefore not related to Svalbardian tectonism. A more likely origin for deformation within the Adriabukta Formation is the early Cenozoic Eureka tectonic event. The tightly folded character of the Adriabukta Formation was previously proposed to be related to the dominance of weak shale and to Cenozoic strain partitioning by Birkenmajer and Turnau (1962). This scenario is now the most likely explanation for differential deformation of shales of the Adriabukta Formation and for folding of the Marietoppen Formation in southern Spitsbergen.

3.2 Age of the Hornsundneset Formation

In Hornsundneset (Fig. 1), Siedlecki and Turnau (1964) analyzed eight samples from the Hornsundneset and Sergeijevfjellet formations of the Billefjorden Group. They proposed a Serpukhovian (Late Mississippian) age based on palynological results. However, a re-evaluation of their results showed that the Billefjorden Group in Hornsundneset (location in Fig. 1) is Middle Mississippian in age (Dallmann et al., 1999; Krajewski and Stempien-Salek, 2003), i.e., contemporaneous with the Adriabukta Formation (Fig. 2).

Interestingly, the Hornsundneset and Sergeijevfjellet formations are dominated by relatively hard, flat-lying beds of sandstone. Though located closer to the early Cenozoic collision zone with Greenland (i.e., within the West Spitsbergen Fold-and-Thrust Belt), these formations are relatively undeformed compared to the shale-dominated Adriabukta Formation (Siedlecki, 1960). This further supports a significant impact of strain partitioning on deformation patterns during the Eureka tectonic event in southern Spitsbergen.

3.3 Other time constraints

In Adriabukta (location in Fig. 1), the Adriabukta Formation is truncated by a major shear zone, the Mariekammen Shear Zone, which comprises hundreds of meter-long lenses of Cambrian metasedimentary basement rocks, shows a top-east reverse sense of shear and is unconformably overlain upwards by mildly folded Pennsylvanian strata of the Gipsdalen Group (Hyrnefjellet Formation), thus possibly reflecting Svalbardian tectonism (Birkenmajer and Turnau, 1962; Birkenmajer, 1964; Dallmann, 1992; Bergh et al., 2011). However, these previous studies did not account for the impact of Eureka tectonism in southern Spitsbergen. A simple restoration of the shear zone prior to Eureka deformation shows that, if this structure is indeed Mississippian in age, it must have formed as a normal fault and therefore cannot reflect Svalbardian contractional deformation (Supplement S2). It should be noted that other workers proposed that the Mariekammen Shear Zone formed as an early Cenozoic

structure (Dallmann, 1992; von Gosen and Piepjohn, 2001). The fact that the shear zone does not seem to crosscut the Hyrnefjellet Formation and instead abruptly dies out at the unconformity (see sketch in Fig. 5 in Bergh et al., 2011) instead supports a formation as a normal fault in the Mississippian (Supplement S2).

The Adriabukta Formation was intruded by two, thin, bedding-parallel, Early Cretaceous dolerite sills of the Diabasodden Suite (Senger et al., 2013) that are folded together with bedding surfaces (Birkenmajer and Morawski, 1960; Birkenmajer, 1964). If the Adriabukta Formation was already folded in the Early Cretaceous, the sills would have truncated both fold structures and bedding surfaces. For sills to intrude along bedding surfaces, these must have remained relatively undeformed, sub-planar, and sub-horizontal until the Early Cretaceous. The Early Cretaceous sills and Middle Mississippian sedimentary strata were then folded together during subsequent Eureka deformation. The two Early Cretaceous sills therefore further constrain the age of folding within the Adriabukta Formation and Marietoppen Formation to the early Cenozoic.

An early Cenozoic age for folding of shales of the Adriabukta Formation is further suggested by similar tight, east-verging fold geometries in Lower Triassic sedimentary strata incorporated as lenses into basement rocks in Fiskeknatten (location shown in Fig. 1; Birkenmajer, 1964).

Svalbardian deformation may be recorded in southernmost Spitsbergen (Røkensåta; Fig. 1 for location) where two outcrops of limited geographical extent (<1 km²) show poorly exposed, gently dipping, shale-rich, Lower Triassic sedimentary rocks over folded Middle Devonian strata (Dallmann, 1992). However, the two outcrops are of small size because they were extensively eroded, and the stratigraphic contact between Devonian and Triassic rocks is completely covered by loose material and located on steep mountain flanks (i.e., inaccessible for detailed inspection). In addition, Triassic successions in Spitsbergen dominantly consist of weak shale (Worsley and Mørk, 1978), which localized large amounts of Eureka deformation and displacement along décollement levels, and strain partitioning during early Cenozoic contraction is now known to have had a considerable influence on the deformation of shale units in southern Spitsbergen (e.g., tightly folded Middle Mississippian Adriabukta Formation versus undeformed Middle Mississippian Hornsundneset Formation; Siedlecki, 1960; Birkenmajer and Turnau, 1962). Furthermore, folds within Middle Devonian rocks in Røkensåta appear to die out upwards (see Fig. 4a in Dallmann, 1992). It is therefore possible that deformation in Røkensåta is also early Cenozoic in age.

Such heavily eroded and limited outcrops need to be interpreted with extreme caution. Lower Triassic strata throughout Spitsbergen are well known for hosting bedding-parallel Eureka décollements (Maher, 1984; Maher et al., 1986, 1989; Andresen et al., 1988; Bergh and Andresen, 1990; Haremo et al., 1990; Haremo and Andresen, 1992; Andresen

et al., 1992; Dallmann et al., 1993; Bergh et al., 1997). The most spectacular examples include the décollement in dark shales on the Midterhuken Peninsula (Maher, 1984; Dallmann et al., 1993; location shown in Fig. 1), the Berzeliustinden thrust in southern Spitsbergen (Dallmann, 1988), the Triassic décollement penetrated by the 7816/12-1 exploration well and well imaged on seismic data in Reindalspasset (Eide et al., 1991; Koehl, 2021, Fig. 5g therein; see Fig. 1 for location), and the “Lower Décollement Zone” in eastern Spitsbergen (Haremo et al., 1990; Andresen et al., 1992; Haremo and Andresen, 1992). A similar structure may very well have decoupled Eurekan deformation between folded Middle Devonian and overlying gently dipping Lower Triassic sedimentary strata in Røkensåta. Triassic shales are known to be much weaker than Devonian shales and to have preferentially localized Eurekan deformation at a much lower scale (e.g., décollements with kilometer-scale displacement in the Triassic shales versus open meso- to macro-scale folds with limited to no displacement within Devonian shales). This example stresses the importance of detailed inspection of extensively eroded outcrops, especially in glaciated Arctic areas, and highlights potential flaws in long-distance interpretation of kilometer-scale mountain flanks. Therefore, we propose that little to no weight should be given to any interpretation of these two poorly exposed and inaccessible outcrops until further inspection of the contact is made from very close range (e.g., using a drone).

4 Review of age constraints in western Spitsbergen

4.1 Conodont age in Blomstrandhalvøya

In western Spitsbergen, Thiedig and Manby (1992) and Kempe et al. (1997) showed that west-verging thrusts cross-cut Proterozoic and Devonian sedimentary rocks in Blomstrandhalvøya (location in Fig. 1). They used the westwards transport direction of these thrusts to suggest that they record Svalbardian tectonism because it is comparable to observations along inferred Svalbardian thrusts in Dickson Land and Andrée Land in central and northern Spitsbergen (Vogt, 1938; Harland et al., 1974; Piepjohn, 2000).

In addition, Kempe et al. (1997) also noted the presence of small NW-verging thrusts on Blomstrandhalvøya. Notably, they argued that the size of these thrust was different from that of Svalbardian structures and concluded that they must therefore be post-Devonian. Kempe et al. (1997) argued that, even though the NW-verging thrusts seemed to have formed in the early Cenozoic, NW-directed transport directions are not typical of early Cenozoic Eurekan tectonism, which produced NE-verging thrusts and folds in adjacent areas of Brøggerhalvøya (Bergh et al., 2000; Piepjohn et al., 2001; see location in Fig. 1). They therefore proposed that NW-verging thrusts on Blomstrandhalvøya formed during a discrete tectonic event in the Pennsylvanian to Cretaceous.

However, such a tectonic event is, thus far, unheard of in Spitsbergen. It is therefore more likely that the NW-verging thrusts in Blomstrandhalvøya formed in the early Cenozoic.

Previous works (e.g., Kempe et al., 1997) used the strike and vergence of structures in Blomstrandhalvøya to distinguish Eurekan from presumed Svalbardian structures. This argument is not valid because a single tectonic event may very well produce structures with varying vergence and strikes, e.g., the Eurekan in Svalbard, which resulted in the formation of east-verging structures in western and southwestern Spitsbergen (e.g., Maher et al., 1986; Dallmann et al., 1988, 1993; Andresen et al., 1994) and northeast-verging folds and thrusts in Brøggerhalvøya (e.g., Bergh et al., 2000; Piepjohn et al., 2001). Furthermore, recent regional studies have shown the occurrence of major, WNW–ESE-striking, several to tens of kilometers thick, thousands of kilometers long, inherited Timanian thrust systems extending from northwestern Russia to western Svalbard (Koehl, 2020; Koehl et al., 2021; Koehl et al., 2022a). One of these structures, the NNE-dipping Kongsfjorden–Cowanodden fault zone, extends into Kongsfjorden, where it was reactivated during the Caledonian and Eurekan events as a sinistral-reverse oblique-slip fault, thus partitioning deformation between northern and southern to western Svalbard during those two events and leading to oppositely verging Eurekan thrust across (e.g., west-verging in Andrée Land and Blomstrandhalvøya and east-verging in Røkensåta and Adriabukta and Hornsund) the fault and to bending Eurekan structures in the vicinity of the fault (e.g., in Brøggerhalvøya).

In western Blomstrandhalvøya, one sample in a presumably undeformed karst infill within a fissure in Proterozoic basement marbles that is a few meters wide yielded a Pennsylvanian to Permian age based on conodont fauna (Buggisch et al., 1994; Fig. 2). Since the karst infill was apparently not deformed, Buggisch et al. (1994) argued that the conodont fauna potentially constrained the formation of folds and west-verging thrusts on Blomstrandhalvøya to the Late Devonian (Svalbardian).

Nevertheless, several aspects of this feature call for caution regarding its bearing for Svalbardian tectonism. First, despite being located in an area deformed by Eurekan tectonism, e.g., Blomstrandhalvøya (e.g., NW-verging thrusts of Kempe et al., 1997) and Brøggerhalvøya; (Bergh et al., 2000; Piepjohn et al., 2001; Fig. 1), the Pennsylvanian to Permian cave seems to have escaped early Cenozoic deformation. This is possibly due to partitioning of Eurekan strain, which is known to have had a significant influence on deformation patterns in Brøggerhalvøya (e.g., Bergh et al., 2000). Thus, this small-scale karst feature is not an appropriate marker to discuss the timing of regional tectonic events in Blomstrandhalvøya.

Second, the cave is located within relatively undeformed Proterozoic marbles and away from presumed Svalbardian west-verging thrusts and associated deformed Lower Devonian sedimentary rocks on Blomstrandhalvøya. Hence, the

karst infill is inappropriate to constrain the timing of Svalbardian deformation in Blomstrandhalvøya. The deformation in basement marble (if there is any at all at the location of the karst) could very well be Caledonian as previously suggested by Michalski (2018).

Third, the karst is the only one of its kind yielding a Pennsylvanian to Permian age and is, moreover, based on only one sample with a poorly preserved conodont fauna (Buggisch et al., 1994). In their study, Buggisch et al. (1994) specified that the assignment to published species was difficult due to the poor preservation of the elements. Hence, further studies of caves and conodont fauna on Blomstrandhalvøya are therefore needed to further assess the reliability of the age obtained by Buggisch et al. (1994) and its implication (if any at all) for Svalbardian tectonism. Considering all pieces of evidence gathered thus far, the folds and thrusts in Proterozoic to Lower Devonian rocks in Blomstrandhalvøya may all be Caledonian and Eurekan in age since no appropriate constraints are available to date any potential Svalbardian deformation.

4.2 Amphibolite facies metamorphism in Prins Karls Forland

In Prins Karls Forland (see Fig. 1 for location), amphibolite facies metamorphism was dated to 373–355 Ma by ion microprobe and ^{40}Ar – ^{39}Ar geochronology and was postulated to be prograde and thus to record deep-crustal Svalbardian tectonism (ca. 15 km depth; Majka and Kościńska, 2017; Faehnrich et al., 2017; Schneider et al., 2018; Kościńska et al., 2020). This episode of deep-crustal metamorphism is coeval with shallow-crustal Svalbardian tectonism in central and northern Spitsbergen dated to ca. 383–365 Ma by recent paleontological and palynological studies (Scheibner et al., 2012; Lindemann et al., 2013; Marshall et al., 2015; Berry and Marshall, 2015; Newman et al., 2019; Lopes, G., personal observation, 2019). However, none of the ages in Prins Karls Forland are of any use in discussing the timing of the Svalbardian event since they could either reflect crustal thickening or late- to post-orogenic collapse.

Kinematic indicators along the dated west-dipping shear zone display top-SW to top-NW normal sense of shear (Schneider et al., 2018, Fig. 3b, e and f therein), which is incompatible with a formation during contractional (Svalbardian) tectonism. Instead, the shear sense suggests a close relationship with Devonian extensional collapse of the Caledonides. Notably, amphibolite facies metamorphism in Prins Karls Forland is also coeval with and occurred at comparable depth as deep-crustal, late Caledonian, high-pressure metamorphism along the conjugate eastern to northeastern Greenland margin (Gilotti et al., 2004; McClelland et al., 2006; Augland et al., 2010, 2011), which developed synchronously with the deposition of Devonian to Mississippian collapse basins along low-angle extensional detachments at the surface (Stemmerik et al., 1991, 1998, 2000;

Larsen and Bengaard, 1991; Strachan, 1994; Larsen et al., 2008). During late- to post-orogenic collapse, deep contractional tectonics occurring typically at greenschist to amphibolite facies conditions (Snoke, 1980; Lister and Davis, 1989; Krabbendam and Dewey, 1998) are commonly associated with near-surface extension (Platt, 1986; Rey et al., 2001, 2011; Teyssier et al., 2005).

Amphibolite facies metamorphism in Prins Karls Forland was also coeval with collapse-related core complex exhumation in northwestern Spitsbergen (latest movement at 368 Ma; Braathen et al., 2018). Hence, despite the postulated prograde character of amphibolite facies metamorphism in Prins Karls Forland, its timing appears to coincide with Late Devonian extensional events in nearby areas. If the postulated prograde character of amphibolite facies metamorphism in Prins Karls Forland is to be reconciled with the observed overall top-SW to top-NW normal sense of shear (Schneider et al., 2018, Fig. 3b, e and f therein) and with extensional tectonics in northwestern Spitsbergen (Braathen et al., 2018), then the shear zone and associated prograde metamorphism may reflect gradual burial linked to the deposition of thick collapse sediments and/or normal movements along the shear zone.

Since the Late Devonian to Mississippian (373–355 Ma) amphibolite facies metamorphism in basement rocks in Prins Karls Forland probably occurred at ca. 15 km depth, the timing and nature of metamorphism may not have any implications for the nature of paleostress and resulting deformation in shallow-crustal Devonian sedimentary rocks in Spitsbergen (e.g., coeval ultra-high-pressure metamorphism at depth and extensional collapse at the surface in Greenland in the Devonian to Mississippian; Strachan, 1994; Gilotti et al., 2004; McClelland et al., 2006).

Furthermore, the geochronological ages obtained by Kościńska et al. (2020) show broad ranges (430–336 Ma for monazite population I, 419–261 Ma for population II, and 443–226 Ma for population III) all ranging from the Silurian (Caledonian?) to the Carboniferous–Triassic. In addition, the ages obtained are associated with large σ_1 errors (12.4–20.2 Myr for population I, 19.6–49.9 Myr for population II, and 17.1–64.4 Myr for population III; see online supplement S1 in Kościńska et al., 2020). Since the length of Svalbardian tectonism in shallow-crustal Lower to lowermost Upper Devonian sedimentary rock in central and northern Spitsbergen is constrained to a maximum time span of 18 million years (383–365 Ma), i.e., a time span comparable with the σ_1 errors associated with the ages obtained by Kościńska et al. (2020), these ages are inappropriate to discuss the timing of Svalbardian tectonism in Svalbard (Schaltegger et al., 2015).

4.3 Greenschist facies metamorphism and thermal overprints in Oscar II Land

Geochronological ages in Oscar II Land (location in Fig. 1) are also useless in discussing the timing of the Svalbardian event since they may equally reflect extensional processes. Greenschist facies metamorphism that yielded 365–344 Ma ^{40}Ar – ^{39}Ar and U–Th–Pb ages (Barnes et al., 2020) were re-evaluated to ca. 410 Ma (Early Devonian) by Ziemniak et al. (2020), who obtained comparable ages for the same unit without the 365–344 Ma disturbance, which they attribute to fluid circulation. This is also partly supported by the poorer statistical reliability of the 365–344 Ma ages as documented by Barnes et al. (2020). The 365–344 Ma episode of low-grade metamorphism was coeval with the deposition of Lower Devonian sedimentary rocks in central and northern Spitsbergen in the Devonian Graben during late–post-orogenic collapse of the Caledonides and thus may also be related to extensional processes (Gee and Moody-Stuart, 1966; Friend et al., 1966; Friend and Moody-Stuart, 1972; Murasov and Mokin, 1979; Manby and Lyberis, 1992; Friend et al., 1997; McCann, 2000).

In addition, Michalski et al. (2017) provided evidence of two episodes of thermal overprints at 377–326 and ca. 300 Ma in pre-Caledonian rocks in Oscar II Land using ^{40}Ar – ^{39}Ar geochronology. The latter event is believed to be related to rifting. The former event at 377–326 Ma partly overlaps with the presumed timing of the Svalbardian Orogeny in central and northern Spitsbergen at ca. 383–365 Ma (Scheibner et al., 2012; Lindemann et al., 2013; Marshall et al., 2015; Berry and Marshall, 2015; Newman et al., 2019; Lopes, G., personal observation, 2019) and with the timing of 373–355 Ma amphibolite facies metamorphism in western Spitsbergen (Majka and Kościńska, 2017; Faehrich et al., 2017; Schneider et al., 2018; Kościńska et al., 2020). It is, however, not possible to infer tectonic stress orientation, and this event may very well be related to Svalbardian tectonism or to late Caledonian extensional processes in northeastern Greenland and Prins Karls Forland (Stemmerik et al., 1991, 1998, 2000; Larsen and Bengaard, 1991; Strachan, 1994; Larsen et al., 2008; Schneider et al., 2018; see also previous section) and in northern Spitsbergen (Chorowicz, 1992; Roy, 2007, 2009; Braathen et al., 2018, 2020; Roy et al., 2022; Maher et al., 2022).

5 Discussion and re-evaluation of the timing and extent of Svalbardian tectonism

The present brief review of age constraints in Spitsbergen shows a few noteworthy aspects of dating Svalbardian tectonism in Svalbard. In southern Spitsbergen, Middle Mississippian palynological ages for the tightly folded, shale-rich Adriabukta formation (Birkenmajer and Turnau, 1962) and its intrusion by two Early Cretaceous dolerite sills

that are folded together with bedding surfaces (Birkenmajer and Morawski, 1960; Birkenmajer, 1964) show that folding in this area may be exclusively and entirely early Cenozoic in age. Comparable Middle Mississippian palynological ages for the contemporaneous but undeformed, sandstone-dominated Hornsundneset Formation ca. 20 km to the southwest (Siedlecki, 1960; Siedlecki and Turnau, 1964) and mild folding of clastic-rich Pennsylvanian to Permian rocks in Adriabukta (Birkenmajer, 1964; Bergh et al., 2011) illustrate the strong impact of Eurekan strain partitioning on deformation patterns in southern Spitsbergen as previously considered by Birkenmajer and Turnau (1962) and Koehl (2020a).

The only potential record of Svalbardian tectonism in southern Spitsbergen occurs at Røkensåta. However, as previously discussed, the low quality of the only two exposures (stratigraphic contact covered by loose material), their very limited extent ($<1\text{ km}^2$), their inaccessibility for detailed inspection (located on steep mountain flanks), the significant impact of early Cenozoic strain partitioning in southern Spitsbergen (Birkenmajer and Turnau, 1962), and the geometry of folds within Middle Devonian rocks at this locality (dying out upwards; Dallmann, 1992) call for caution and further detailed investigation of structural and stratigraphic relationships at this locality. Nevertheless, if Eurekan tectonism alone produced the intense deformation in Adriabukta, it is possible that deformation in Røkensåta is exclusively early Cenozoic as well.

In central and northern Spitsbergen, Svalbardian tectonism was constrained to ca. 383–365 Ma (i.e., a maximum duration of 18 million years) by recent paleontological and palynological studies in sedimentary rocks of the Mimerdalen Subgroup (Berry and Marshall, 2015; Newman et al., 2019, 2020, 2021) and Billefjorden Group (Scheibner et al., 2012; Lindemann et al., 2013; Marshall et al., 2015; Lopes, G., personal observation, 2019). The only contradictory late Famennian age obtained by Piepjohn et al. (2000) via identification of one specimen of *Retispora lepidophyta* in one sample of the Plantekløfta Formation of the Mimerdalen Subgroup is now known to be a clear misidentification (Berry and Marshall, 2015, Supplement DR3 therein).

Despite the accurate and precise paleontological and palynological time constraints for Svalbardian tectonism in central and northern Spitsbergen, no geochronological constraints exist yet for discrete Svalbardian structures. In addition, the central and northern part of Spitsbergen was strongly affected by early Cenozoic Eurekan tectonism during which strain partitioning played an important role in localizing deformation in weak, shale-rich lithostratigraphic units like the Billefjorden Group (e.g., Koehl, 2021). Moreover, evidence for extensional detachment-related folding in northwestern (Braathen et al., 2018, 2020; Maher et al., 2022) and northern Spitsbergen (Chorowicz, 1992; Roy, 2007, 2009; Roy et al., 2022) in the Middle to Late Devonian may also have contributed to deformation patterns observed within Lower to lowermost Upper Devonian strata of

the Andrée Land Group and Mimerdalen Subgroup. Thus, it is unclear how much (if any at all) of the deformation observed within Lower to lowermost Upper Devonian strata in central and northern Spitsbergen actually reflects Svalbardian tectonism. Further studies are therefore clearly needed to quantify the impact of the Svalbardian Orogeny and to segregate discrete Svalbardian from Devonian extensional (detachment) faulting and folding and from early Cenozoic Eurekan folding and thrusting.

Another line of controversy is the incredibly rapid switch from extension-related normal faulting in the Early to Middle Devonian, to Svalbardian contraction in the Late Devonian, and back to dominantly extensional setting in the mid-Famennian in central and northern Spitsbergen. Notably, the Wood Bay Formation and Fiskekløfta Member of the Tordalen Formation are down-faulted by normal faults in southern Hugindalen and unconformably covered by the Planteryggen Formation (Hugindalen Phase in Piepjohn, 2000 and Dallmann and Piepjohn, 2020). The Fiskekløfta Member was dated to the latest Givetian (top of the unit at ca. 383 Ma) and the Plantekløfta Formation to the early Frasnian (383–380 Ma; Berry and Marshall, 2015; Newman et al., 2019, 2020, 2021). Since the conglomeratic beds of the Planteryggen and Plantekløfta formations are advocated by Piepjohn and Dallmann (2014) to reflect the onset of Svalbardian tectonism, this would therefore imply an abrupt switch in plate tectonic movements and stresses at exactly 383 Ma, i.e., completed within a million years at the maximum. In addition, mid-Famennian to Upper Mississippian sedimentary rocks of the Billefjorden Group and Pennsylvanian to lower Permian rocks of the Gipsdalen Group, which overlie the Andrée Land Group in central and northern Spitsbergen, are believed to have been deposited in extensional basins (Cutbill et al., 1976; Aakvik, 1981; Gjelberg, 1984; Braathen et al., 2011; Koehl and Muñoz-Barrera, 2018; Smyrak-Sikora et al., 2018). This implies another rapid reversal in regional plate tectonics movements from contraction to extension at ca. 365 Ma. Since regional plate tectonics reorganization and tectonic stress reorientation are known to be relatively slow and gradual processes, such abrupt switches are regarded as highly unlikely. Considering the extensional setting inferred in both the Early to Middle Devonian (Chorowicz, 1992; Piepjohn, 2000; Roy, 2007, 2009; Braathen et al., 2018, 2020; Dallmann and Piepjohn, 2020; Roy et al., 2022; Maher et al., 2022) and mid-Famennian to early Permian (Cutbill et al., 1976; Aakvik, 1981; Gjelberg, 1984; Braathen et al., 2011; Smyrak-Sikora et al., 2018), it is more likely that Svalbardian contraction never occurred and that the area was subjected to continuous extension throughout the Devonian to Carboniferous. This is also supported by late Silurian to Late Devonian extensional detachment faulting and folding at 430–368 Ma in northwestern Spitsbergen (Braathen et al., 2018) and in the Middle to Late Devonian in northern Spitsbergen (Chorowicz, 1992; Roy, 2007, 2009; Roy et al., 2022).

The 383–365 Ma estimate for tentative Svalbardian deformation in shallow-crustal Lower to lowermost Upper Devonian sedimentary rocks in central and northern Spitsbergen partly overlaps with the timing of deep-crustal, 373–355 Ma, amphibolite facies metamorphism in Prins Karls Forland (Majka and Košmińska, 2017; Faehnrich et al., 2017; Schneider et al., 2018; Košmińska et al., 2020) and thermal events in Oscar II Land at 377–326 Ma (Michalski et al., 2017). However, the 383–365 Ma estimate reflects the age of stratigraphy in central and northern Spitsbergen, not the age of any specific Svalbardian structure. In addition, due to conflicting lines of evidence (e.g., postulated prograde metamorphism associated with normal sense of shear), the nature of tectonic stresses during tectonothermal events in Prins Karls Forland and Oscar II Land remains debatable.

Paleomagnetic and ^{40}Ar – ^{39}Ar geochronological data from Michalski et al. (2017) do not support a pre-Caledonian link or proximity between the Pearya terrane and western Spitsbergen. As part of the same trend, detrital zircons in western and central Spitsbergen show affinities with northern Baltica rather than Laurentia in the Paleozoic (Gasser and Andresen, 2013). This suggests that western and central Spitsbergen were located away from the main Ellesmerian belt in northern Greenland and Arctic Canada and thus may have escaped Ellesmerian tectonism. This is further supported by the recent discovery of several kilometer-thick late Neoproterozoic thrust systems that are thousands of kilometers long and crosscut the whole Barents Sea and the Svalbard Archipelago, thus suggesting that the Svalbard Archipelago was already accreted and attached to Baltica in the late Neoproterozoic (Koehl, 2020b; Koehl et al., 2022a).

6 Conclusion

There should be no debate as to the age of the Mimerdalen Subgroup and Billefjorden Group. These are upper Givetian to lower Frasnian (ca. 385–380 Ma) and mid-Famennian to Upper Mississippian (ca. 365–325 Ma), respectively, and are robustly constrained by palynological and paleontological markers. The single palynomorph specimen that was not in line with these ages that was found in the Mimerdalen Subgroup is a clear misidentification of *Retispora lepidophyta*. Thus, the timing of Svalbardian tectonism in central and northern Spitsbergen is constrained to 383–365 Ma. Nonetheless, because of the strong impact of Eurekan strain partitioning and extensional detachment-related folding and faulting, much is left to do to quantify the impact, extent, and timing of Svalbardian tectonism in this area (if it ever occurred). Future studies should focus on geochronological dating of presumed Svalbardian thrusts.

There is also no debate about the age of the Adriabukta Formation in southern Spitsbergen. Palynological evidence confirms that this formation is Middle Mississippian in age and is therefore a time-equivalent example of the unde-

formed, sandstone-rich Hornsundeneset Formation. Hence, folding in the Adriabukta Formation is entirely and exclusively ascribed to Eurekan tectonism, and the tight character of folding is ascribed to strain partitioning in the early Cenozoic. Due to lack of robust minimum time constraints, the occurrence of Svalbardian tectonism in southern Spitsbergen is highly doubtful. Future studies could, if feasible, focus on establishing clear tectonic and stratigraphic relationships in Røkensåta.

Postulated prograde amphibolite facies metamorphism at 373–355 Ma in pre-Caledonian basement rocks in Prins Karls Forland occurred at a depth of ca. 15 km and, thus, has no bearings on the nature of tectonic stress and associated deformation in shallow-crustal Devonian to Mississippian sedimentary rocks. Top-SW to top-NW normal sense of shear along the dated shear zone suggests that this episode of postulated prograde metamorphism may actually be related to shallow-crustal, extensional collapse processes, possibly reflecting progressive burial and movements along the shear zone during the deposition of collapse sediments. Similar processes are well documented on the conjugate margin of Svalbard in northeastern Greenland and in northwestern Spitsbergen, and these processes involve deep, late Caledonian, high-pressure metamorphism and shallow-crustal extensional detachments.

Considering the dominantly extensional tectonic settings inferred for shallow-crustal rocks in late Silurian to early Permian times and the multiple inconsistencies and contradicting lines of evidence associated to the Svalbardian Orogeny throughout Svalbard, the accretion of Svalbard to Baltica as early as the late Neoproterozoic, and the two abrupt and rapid switches in tectonic stress orientation required in the Late Devonian to account for Svalbardian tectonism, it is much more likely that the whole archipelago was subjected to continuous extension from the late Silurian to early Permian times and escaped Svalbardian deformation.

Data availability. No data sets were used in this article.

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/se-13-1353-2022-supplement>.

Author contributions. JBPK wrote the manuscript and designed the figures (contribution: 50%). JEAM and GL provided critical input and corrections to the manuscript and figures (contribution: 25% each).

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. The authors thank the referees, Michael Newman and Keith Dewing, the editor, Silvia Gardin, and Chris Berry and Karsten Piepjohn for helpful comments and discussion on the manuscript.

Financial support. The present study was financed by the ARCEX (Research Centre for Arctic Petroleum Exploration; grant no. 228107), SEAMSTRESS (Centre for Earth Evolution and Dynamics; grant no. 287865), and CEED projects (grant no. 223272) funded by the Research Council of Norway, the Tromsø Research Foundation, and six industry partners.

Review statement. This paper was edited by Silvia Gardin and reviewed by Michael Newman and Keith Dewing.

References

- Aakvik, R.: Fasies analyse av Undre Karbonske kullførende sedimenter, Billefjorden, Spitsbergen, PhD Thesis, University of Bergen, Bergen, Norway, 219 p., 1981.
- Allen, K. C.: Lower and Middle Devonian spores of north and central Vestspitsbergen, *Palaeontology*, 8, 678–748, 1965.
- Allen, K. C.: Further information on the Lower and Middle Devonian spores from Dickson Land, Spitsbergen, *Norsk Polarinstittutt Årbok 1971*, 7–15, https://brage.npolar.no/npolar-xmlui/bitstream/handle/11250/172763/Aarbok_1971.pdf?sequence=1&isAllowed=y (last access: 25 August 2022), 1973.
- Andresen, A., Bergh, S. G., Hansen, H., Kløvjan, O., Kristensen, S. E., Livbjerg, F., Lund, T., Mair, B. F., Midbøe, P. and Nøttvedt, A.: Geometry and structural development of the Billefjorden and Lomfjorden Fault Zones in the Isfjorden-Sabine Land Area, Spitsbergen, Abstract 18, *Nordiske Geologiske Vintermøde, København*, 33–34, 1988.
- Andresen, A., Haremo, P., Swensson, E. and Bergh, S. G.: Structural geology around the southern termination of the Lomfjorden Fault Complex, Agardhdalen, east Spitsbergen, *Norsk Geol. Tidsskr.*, 72, 83–91, 1992.
- Andresen, A., Bergh, S. G. and Haremo, P.: Basin inversion and thin-skinned deformation associated with the Tertiary transpressional west Spitsbergen Orogen, in: *Proceedings of the International Conference on Arctic Margins*, edited by: Thurston, D. K. and Fujita, K., Anchorage, Alaska, USA, September 1992, 161–166, <https://www.boem.gov/sites/default/files/about-boem/BOEM-Regions/Alaska-Region/ICAM/1994-ICAM/ICAM92-161.pdf> (last access: 25 August 2022), 1994.
- Augland, L. E., Andresen, A. and Corfu, F.: Age, structural setting, and exhumation of the Liverpool Land eclogite terrane, East Greenland Caledonides, *Lithosphere*, 2, 267–286, 2010.

- Augland, L. E., Andresen, A. and Corfu, F.: Terrane transfer during the Caledonian orogeny: Baltican affinities of the Liverpool Land Eclogite Terrane in East Greenland, *Geol. Soc. London*, 168, 15–26, 2011.
- Barnes, C. J., Walczak, K., Janots, E., Schneider, D. and Majka, J.: Timing of Paleozoic Exhumation and Deformation of the High-Pressure Vestgötabreen Complex at the Motalafjella Nunatak, Svalbard, *Minerals*, 10, 125, <https://doi.org/10.3390/min10020125>, 2020.
- Bergh, S. G. and Andresen, A.: Structural development of the Tertiary fold-and-thrust belt in east Oscar II Land, Spitsbergen, *Polar Res.*, 8, 217–236, 1990.
- Bergh, S. G. and Grogan, P.: Tertiary structure of the Sørkapp–Hornsund Region, South Spitsbergen, and implications for the offshore southern continuation of the fold-thrust Belt, *Norsk Geol. Tidsskr.*, 83, 43–60, 2003.
- Bergh, S. G., Braathen, A., and Andresen, A.: Interaction of Basement-Involved and Thin-Skinned Tectonism in the Tertiary Fold-Thrust Belt of Central Spitsbergen, Svalbard, *AAPG Bulletin*, 81, 637–661, 1997.
- Bergh, S. G., Maher Jr., H. D., and Braathen, A.: Tertiary divergent thrust directions from partitioned transpression, Brøggerhalvøya, Spitsbergen, *Norsk Geol. Tidsskr.*, 80, 63–82, 2000.
- Bergh, S. G., Maher Jr., H. D. and Braathen, A.: Late Devonian transpressional tectonics in Spitsbergen, Svalbard, and implications for basement uplift of the Sørkapp–Hornsund High, *J. Geol. Soc. London*, 168, 441–456, 2011.
- Berry, C. M. and Marshall, J. E. A.: Lycopsid forests in the early Late Devonian paleoequatorial zone of Svalbard, *Geology*, 43, 1043–1046, 2015.
- Birkenmajer, K.: Devonian, Carboniferous and Permian formations of Hornsund, Vestspitsbergen, *Studia Geologica Polonica*, 11, 47–123, 1964.
- Birkenmajer, K. and Morawski, T.: Dolerite intrusions of Wedel-Jarlsberg Land Vestspitsbergen, *Stud. Geol. Polon.*, 4, 103–123, 1960.
- Birkenmayer, K. and Turnau, E.: Lower Carboniferous age of the so-called Wijde Bay Series in Hornsund, Vestspitsbergen, *Nor. Polarinst. Årb.*, 1961, 41–61, 1962.
- Braathen, A., Bælum, K., Maher Jr., H. D. and Buckley, S. J.: Growth of extensional faults and folds during deposition of an evaporite-dominated half-graben basin; the Carboniferous Billefjorden Trough, Svalbard, *Norsk Geol. Tidsskr.*, 91, 137–160, 2011.
- Braathen, A., Osmundsen, P. T., Maher Jr., H. D. and Ganerød, M.: The Keisarhjelmen detachment records Silurian–Devonian extensional collapse in Northern Svalbard, *Terra Nova*, 30, 34–39, 2018.
- Braathen, A., Ganerød, M., Maher Jr., H., Myhre, P. I., Osmundsen, P. T., Redfield, T. and Serck, C.: Devonian extensional tectonics in Svalbard; Raudfjorden's synclinal basin above the Keisarhjelmen detachment, 34th Nordic Geological Winter Meeting, 8–10 January 2020, Oslo, Norway, <https://www.geologi.no/konferanser/foredrag-abstracts/file/206-abstract-ngwm20> (last access: 25 August 2022), 2020.
- Brinkmann, L.: Geologie des östlichen zentralen Dickson Landes und Palynologie der Mimerdalen Formation (Devon), Spitzbergen. (Geology of eastern-central Dickson Land and palynology of the Mimerdalen Formation [Devonian], Spitsbergen.), unpublished Master's Thesis, University of Münster, Münster, Germany, 94 p., 1997.
- Buggisch, W., Piepjohn, K., Thiedig, F. and von Gosen, W.: A Middle Carboniferous Conodont Fauna from Blomstrandhalvøya (NW-Spitsbergen): Implications on the Age of Post-Devonian Karstification and the Svalbardian Deformation, *Polarforschung*, 62, 83–90, 1994.
- Chalmers, J. A. and Pulvertaft, T. C. R.: Development of the continental margins of the Labrador Sea: a review, in: *Non-Volcanic Rifting of Continental Margins: A Comparison of Evidence from Land and Sea*, edited by: Wilson, R. C. L., Taylor, R. B. and Froitzheim, N., *Geol. Soc. London, Spec. Publi.*, 187, 77–105, 2001.
- Chauvet, A. and Séranne, M.: Extension-parallel folding in the Scandinavian Caledonides: implications for late-orogenic processes, *Tectonophys.*, 238, 31–54, 1994.
- Chorowicz, J.: Gravity-induced detachment of Devonian basin sediments in northern Svalbard, *Norsk Geol. Tidsskr.*, 72, 21–25, 1992.
- Clayton, G.: Mississippian miospores, in: *Palynology: principles and applications*, edited by: Jansonius, J. and McGregor, D.C., *AASP Foundation*, Salt Lake City, Utah, USA, 2, 589–596, <http://www.worldcat.org/title/889720416> (last access: 25 August 2022), 1996.
- Clayton, G., Coquel, R., Doubinger, J., Gueinn, K. J., Loboziak, S., Owens, B. and Streel, M.: Carboniferous Miospores of Western Europe: illustration and zonation, *Meded. Rijks Geol. Dienst*, 29, 1–71, 1977.
- Cutbill, J. L. and Challinor, A.: Revision of the Stratigraphical Scheme for the Carboniferous and Permian of Spitsbergen and Bjørnøya, *Geol. Mag.*, 102, 418–439, 1965.
- Cutbill, J. L., Henderson, W. G., and Wright, N. J. R.: The Billefjorden Group (Early Carboniferous) of central Spitsbergen, *Norsk Polarinst. Skr.*, 164, 57–89, 1976.
- Dallmann, W. K.: The structure of the Berzeliustinden area: evidence for thrust wedge tectonics in the Tertiary fold-and-thrust belt of Spitsbergen, *Polar Res.*, 6, 141–154, 1988.
- Dallmann, W. K.: Multiphase tectonic evolution of the Sørkapp–Hornsund mobile zone (Devonian, Carboniferous, Tertiary), Svalbard, *Norsk Geol. Tidsskr.*, 72, 49–66, 1992.
- Dallmann, W. K. and Piepjohn, K.: The structure of the Old Red Sandstone and the Svalbardian Orogenic Event (Ellesmerian Orogeny) in Svalbard, *Norg. Geol. Unders. B.*, 15, 106 p., <https://www.ngu.no/publikasjon/special-publication-152020> (last access: 25 August 2022), 2020.
- Dallmann, W. K., Ohta, Y. and Andresen, A.: Tertiary Tectonics of Svalbard, *Norsk Polarinstittutt Rapportserie*, 46, 112 p., <https://brage.npolar.no/npolar-xmlui/handle/11250/173348> (last access: 25 August 2022), 1988.
- Dallmann, W. K., Andresen, A., Bergh, S. G., Maher Jr., H. D. and Ohta, Y.: Tertiary fold-and-thrust belt of Spitsbergen Svalbard, *Norsk Polarinstittutt Meddelelser*, 128, 51 p., <https://brage.npolar.no/npolar-xmlui/handle/11250/2396840> (last access: 25 August 2022), 1993.
- Dallmann, W. K., Dypvik, H., Gjelberg, J. G., Harland, W. B., Johannessen, E. P., Keilen, H. B., Larssen, G. B., Lønøy, A., Midbøe, P. S., Mørk, A., Nagy, J., Nilsson, I., Nøttvedt, A., Olaussen, S., Pcelina, T. M., Steel, R. J. and Worsley, D.: *Lithostratigraphic Lexicon of Svalbard*, edited by: Dallmann,

- W. K., Norwegian Polar Institute, Polar Environmental Centre, Tromsø, Norway, http://mmmtk.ginras.ru/pdf/Dallmann%20et%20al.,1999_Svalbard.pdf and https://timescalefoundation.org/resources/NW_Europe_Lex/litho/svalbard/index.htm (last access: 25 August 2022), 1999.
- Dallmeyer, R. D., Peucat, J. J., Hirajima, T. and Ohta, Y.: Tectonothermal chronology within a blueschist–eclogite complex, west-central Spitsbergen, Svalbard: Evidence from $^{40}\text{Ar}/^{39}\text{Ar}$ and Rb–Sr mineral ages, *Lithos*, 24, 291–304, 1990.
- Eide, J. R., Ree, R. and Rockman, P. O.: Final well report – 7816/12-1 July 1991, Norsk Hydro A. S., Harstad, Norway, available upon request to Equinor and Store Norske Spitsbergen Kulkompani (SNSK), 1991.
- Embry, A. F.: Middle-Upper Devonian Clastic Wedge of the Arctic islands, in: *Geology of the Inuitian Orogen and Arctic Platform of Canada and Greenland*, edited by: Trettin, H. P., Geological Survey of Canada, *Geology of Canada*, Geological Society of America, *The Geology of North America*, 3, <https://geoscan.nrcan.gc.ca/starweb/geoscan/servlet.starweb?path=geoscan/fulle.web&search1=R=133959> (last access: 25 August 2022), 1991.
- Embry, A. F. and Klovan, J. E.: The Middle-Upper Devonian Clastic Wedge of the Franklinian Geosyncline, *Bull. Can. Petrol. Geol.*, 24, 485–639, 1976.
- Evdokimov, A. N., Burnaeva, M. Yu., Radina, E. S. and Sirotkin, A. N.: The First Find of Kimberlitic Accessory Minerals in Mafic–Ultramafic Dikes in Spitsbergen, *Doklady Earth Sciences*, 407, 275–279, 2006.
- Faehrich, K., Schneider, D., Manecki, M., Czerny, J., Myhre, P. I., Majka, J., Kościńska, K., Barnes, C. and Maraszewska, M.: Eurekan deformation on Prins Karls Forland, Svalbard – new insights from $\text{Ar}^{40}/\text{Ar}^{39}$ muscovite dating, *Geophys. Res. Abstracts*, 19, EGU General Assembly 23–28 April 2017, Vienna, Austria, <https://meetingorganizer.copernicus.org/EGU2017/EGU2017-642.pdf> (last access: 25 August 2022), 2017.
- Faehrich, K., Majka, J., Schneider, D., Mazur, S., Manecki, M., Ziemniak, G., Wala, V. T. and Strauss, J. V.: Geochronological constraints on Caledonian strike-slip displacement in Svalbard, with implications for the evolution of the Arctic, *Terra Nova*, 32, 290–299, 2020.
- Fossen, H., Teyssier, C. and Whitney, D. L.: Transtensional folding, *J. Struct. Geol.*, 56, 89–102, 2013.
- Friend, P. F. and Moody-Stuart, M.: Sedimentation of the Wood Bay Formation (Devonian) of Spitsbergen: Regional analysis of a late orogenic basin, *Norsk Polarinst. Skr.*, 157, 80 p., <https://brage.npolar.no/npolar-xmlui/bitstream/handle/11250/173898/Skrifter157.pdf?sequence=1&isAllowed=y> (last access: 25 August 2022), 1972.
- Friend, P. F., Heintz, N. and Moody-Stuart, M.: New unit terms for the Devonian of Spitsbergen and a new stratigraphical scheme for the Wood Bay Formation, *Polarinst. Årbok*, 1965, 59–64, 1966.
- Friend, P. F., Harland, W. B., Rogers, D. A., Snape, I., and Thornley, R. S.: Late Silurian and Early Devonian stratigraphy and probable strike-slip tectonics in northwestern Spitsbergen, *Geol. Mag.*, 134, 459–479, 1997.
- Gasser, D. and Andresen, A.: Caledonian terrane amalgamation of Svalbard: detrital zircon provenance of Mesoproterozoic to Carboniferous strata from Oscar II Land, western Spitsbergen, *Geol. Mag.*, 150, 1103–1126, 2013.
- Gayer, R. A., Gee, D. G., Harland, W. B., Miller, J. A., Spall, H. R., Wallis, R. H. and Winsnes, T. S.: Radiometric age determinations on rocks from Spitsbergen, *Norsk Polarinstutt Skrifter*, 137, 43 p., <https://brage.npolar.no/npolar-xmlui/bitstream/handle/11250/173922/Skrifter137.pdf?sequence=1&isAllowed=y> (last access: 25 August 2022), 1966.
- Gee, D. G. and Moody-Stuart, M.: The base of the Old Red Sandstone in central north Haakon VII Land, Vestspitsbergen, *Polarinst. Årbok*, 1964, 57–68, 1966.
- Gee, D. G. and Page, L. M.: Caledonian terrane assembly on Svalbard: New evidence from $^{40}\text{Ar}/^{39}\text{Ar}$ dating in Ny Friesland, *Am. J. Sci.*, 294, 1166–1186, 1994.
- Gilotti, J. A., Nutman, A. P., and Brueckner, H. K.: Devonian to Carboniferous collision in the Greenland Caledonides: U–Pb zircon and Sm–Nd ages of high-pressure and ultrahigh-pressure metamorphism, *Contrib. Mineral Petrol.*, 148, 216–235, 2004.
- Gjelberg, J. G.: Early–Middle Carboniferous sedimentation on Svalbard. A study of ancient alluvial and coastal marine sedimentation in rift- and strike-slip basins, PhD Thesis, University of Bergen, Bergen, Norway, 306 p., <https://bora.uib.no/bora-xmlui/handle/1956/20981> (last access: 25 August 2022), 1984.
- Grantz, A. and May, S. D.: Summary Geologic Report for Barrow Arch Outer Continental Shelf (OCS) Planning Area, Chukchi Sea, Alaska, United States Department of the Interior, Geological Survey, 84–395, <https://pubs.er.usgs.gov/publication/ofr84395> (last access: 25 August 2022), 1984.
- Haremo, P. and Andresen, A.: Tertiary décollements thrusting and inversion structures along Billefjorden and Lomfjorden Fault Zones, East Central Spitsbergen, in: *Structural and Tectonic Modelling and its Application to Petroleum Geology*, edited by: Larsen, R. M., Brekke, H., Larsen, B. T. and Talleraas, E., Norwegian Petroleum Society (NPF) Special Publications, 1, 481–494, 1992.
- Haremo, P., Andresen, A., Dypvik, H., Nagy, J., Elverøi, A., Eike-land, T. A. and Johansen, H.: Structural development along the Billefjorden Fault Zone in the area between Kjellströmdalen and Adventdalen/Sassendalen, central Spitsbergen, *Polar Res.*, 8, 195–216, 1990.
- Harland, W. B.: Contribution of Spitsbergen to understanding of tectonic evolution of North Atlantic region, *AAPG Memoirs*, 12, 817–851, 1969.
- Harland, W. B. and Horsfield, W. T.: West Spitsbergen Orogen, in: *Mesozoic–Cenozoic orogenic belts*, edited by: Spencer, A. M., *Geol. Soc. London Spec. Publ.*, 4, 747–755, 1974.
- Harland, W. B. and Wright, N. J. R.: Alternative hypothesis for the pre-Carboniferous evolution of Svalbard, *Norsk Polarinst. Skr.*, 167, 89–117, 1979.
- Harland, W. B., Cutbill, L. J., Friend, P. F., Gobbett, D. J., Holliday, D. W., Maton, P. I., Parker, J. R. and Wallis, R. H.: The Billefjorden Fault Zone, Spitsbergen – the long history of a major tectonic lineament, *Norsk Polarinst. Skr.*, 161, 1–72, 1974.
- Harland, W. B., Scott, R. A., Auckland, K. A. and Snape, I.: The Ny Friesland Orogen, Spitsbergen, *Geol. Mag.*, 129, 679–708, 1992.
- Harrison, J. C.: Melville Island’s salt-based fold belt, Arctic Canada, *Geological Survey of Canada Bulletin*, 472, 344 p., <https://doi.org/10.4095/203576>, 1995.

- Harisson, J. C. and Brent, T. A.: Basins and fold belts of Prince Patrick Island and adjacent area, Canadian Arctic Islands, Geological Survey of Canada Bulletin, 560, 208 p., <https://doi.org/10.4095/220345>, 2005.
- Higgins, A. K., Soper, N. J. and Leslie, A. G.: The Ellesmerian and Caledonian Orogenic Belts of Greenland, *Polarforschung*, 68, 141–151, 2000.
- Horsfield, W. T.: Glauconite schists of Caledonian age from Spitsbergen, *Geol. Mag.*, 109, 29–36, 1972.
- Hughes, N. F. and Playford, G.: Palynological reconnaissance of the Lower Carboniferous of Spitsbergen, *Micropaleontology*, 7, 27–44, 1961.
- Jakobsson, M., Mayer, L., Coackley, B., Dowdeswell, J. A., Forbes, S., Fridman, B., Hodnesdal, H., Noormets, R., Pedersen, R., Rebesco, M., Schenke, H. W., Zarayskaya, Y., Accettella, D., Armstrong, A., Anderson, R. M., Bienhoff, P., Camerlenghi, A., Church, I., Edwards, M., Gardner, J. V., Hall, J. K., Hell, B., Hestvik, O., Kristoffersen, Y., Marcussen, C., Mohammad, R., Mosher, D., Nghiem, S. V., Pedrosa, M. T., Travaglini, P. G., and Weatherall, P.: The International Bathymetric Chart of the Arctic Ocean (IBCAO) Version 3.0, *Geophys. Res. Lett.*, 39, L12609, <https://doi.org/10.1029/2012GL052219>, 2012.
- Johansson, Å., Larionov, A. N., Gee, D. G., Ohta, Y., Tebenkov, A. M. and Sandelin, S.: Greenvillian and Caledonian tectonomagmatic activity in northeasternmost Svalbard, in: The Neoproterozoic Timanide Orogen of Eastern Baltica, edited by: Gee, D. G. and Pease, V., *Geol. Soc. London Memoirs*, 30, 207–232, 2004.
- Johansson, Å., Gee, D. G., Larionov, A. N., Ohta, Y. and Tebenkov, A. M.: Greenvillian and Caledonian evolution of eastern Svalbard – a tale of two orogenies, *Terra Nova*, 17, 317–325, 2005.
- Kaiser, H.: Die Oberdevon-Flora der Bäreninsel 3. Mikoflora des höheren Oberdevons und des Unterkarbons, *Palaeontographica Abt. B*, 129, 71–124, 1970.
- Kempe, M., Niehoff, U., Piepjohn, K. and Thiedig, F.: Kaledonische und svalbardische Entwicklung im Grundgebirge auf der Blomstrandhalvøya, NW-Spitzbergen, *Münster. Forsch. Geol. Paläont.*, 82, 121–128, 1997.
- Koehl, J.-B. P.: Devonian–Mississippian collapse and core complex exhumation, and partial decoupling and partitioning of Eurekan deformation as alternatives to the Ellesmerian Orogeny in Spitsbergen, *Solid Earth Discuss.* [preprint], <https://doi.org/10.5194/se-2019-200>, 2020a.
- Koehl, J.-B. P.: Impact of Timanian thrusts on the Phanerozoic tectonic history of Svalbard, Keynote lecture, EGU General Assembly, 3–8 May 2020, Vienna, Austria, <https://meetingorganizer.copernicus.org/EGU2020/displays/34748> (last access: 25 August 2022), 2020b.
- Koehl, J.-B. P.: Early Cenozoic Eurekan strain partitioning and decoupling in central Spitsbergen, Svalbard, *Solid Earth*, 12, 1025–1049, <https://doi.org/10.5194/se-12-1025-2021>, 2021.
- Koehl, J.-B. P. and Muñoz-Barrera, J. M.: From widespread Mississippian to localized Pennsylvanian extension in central Spitsbergen, Svalbard, *Solid Earth*, 9, 1535–1558, <https://doi.org/10.5194/se-9-1535-2018>, 2018.
- Koehl, J.-B. P., Collombin, M., Taule, C., Christophersen, G., and Allaart, L.: Influence of WNW–ESE-striking faults on Devonian–Permian sedimentary rocks in Billefjorden and implications for Ellesmerian and Eurekan tectonic events, <https://doi.org/10.13140/RG.2.2.35857.97129>, 2020.
- Koehl, J.-B. P., Cooke, F. A., and Plaza-Faverola, A. A.: Formation of a transform-parallel oceanic core complex along an inherited Timanian thrust, and impact on gas seepage in the Fram Strait, TSG Annual Meeting, 5–8 January, <https://doi.org/10.13140/RG.2.2.28315.49446>, 2021.
- Koehl, J.-B. P., Magee, C., and Anell, I. M.: Impact of Timanian thrust systems on the late Neoproterozoic–Phanerozoic tectonic evolution of the Barents Sea and Svalbard, *Solid Earth*, 13, 85–115, <https://doi.org/10.5194/se-13-85-2022>, 2022a.
- Koehl, J.-B. P., Stokmo, E. M. B., and Muñoz-Barrera, J. M.: On the Billefjorden Fault Zone in Garmndalen, central Spitsbergen, Carboniferous Basins in Svalbard, Canada and the Barents Sea, <https://doi.org/10.13140/RG.2.2.28031.33448>, 2022b.
- Koźmińska, K., Spear, F. S., Majka, J., Faehrich, K., Manecki, M., Piepjohn, K., and Dallmann, W. K.: Deciphering late Devonian–early Carboniferous $P - T - t$ path of mylonitized garnet-mica schists from Prins Karls Forland, Svalbard, *J. Metamorph., Geol.*, 38, 471–493, 2020.
- Krabbendam, M. and Dewey, J. F.: Exhumation of UHP rocks by transtension in the Western Gneiss Region, Scandinavian Caledonides, in: Continental Transpressional and Transtensional Tectonics, edited by: Holdsworth, R. E., Strachan, R. A. and Dewey, J. F., *Geol. Soc. London, Spec. Publi.*, 135, 159–181, 1998.
- Krajewski, K. P. and Stempien-Salek, M.: Overthrust Carboniferous strata (Sergejevffjellet Formation) at Lidfjellet, NW Sørkapp Land, Spitsbergen, *Polish Polar Res.*, 24, 61–72, 2003.
- Kumar, N., Granath, J. W., Emmet, P. A., Helwig, J. A., and Dinkelman, M. G.: Stratigraphic and Tectonic Framework of the US Chukchi Shelf: Exploration Insights from a New Regional Deep-seismic Reflection Survey, in: Arctic Petroleum Geology, edited by: Spencer, A. M., Gautier, D., Stoupakova, A., Embry, A. F. and Sørensen, K., *Geol. Soc. London Memoirs*, 35, 501–508, 2011.
- Lane, L. S.: Devonian–Carboniferous paleogeography and orogenesis, northern Yukon and adjacent Arctic Alaska, *Can. J. Earth Sci.*, 44, 679–694, 2007.
- Larsen, B. T.: Tertiary thrust tectonics in the east of Spitsbergen, and implications for the plate-tectonic development of the North-Atlantic, in: Tertiary Tectonics of Svalbard, edited by: Dallmann, W. K., Ohta, Y. and Andresen, A., *Norsk Polarinstittut Rapportserie*, 46, 85–88, 1988.
- Larsen, P.-H. and Bengaard, H.-J.: Devonian basin initiation in East Greenland: a result of sinistral wrench faulting and Caledonian extensional collapse, *J. Geol. Soc. London*, 148, 355–368, 1991.
- Larsen, P.-H., Olsen, H., and Clack, J. A.: The Devonian basin in East Greenland–Review of basin evolution and vertebrate assemblages, *GSA Bull. Mem.*, 202, 273–292, 2008.
- Lindemann, F.-J., Volohonsky, E., and Marshall, J. E.: A bonebed in the Hørbyreen Formation (Famennian–Viséan) on Spitsbergen, NGF Abstracts and Proceedings, 1, Winter Meeting, Oslo, 8–10 January, https://www.geologi.no/images/Konferanser/Vinterkonferansen2013_Abstracts.pdf (last access: 25 August 2022), 2013.
- Lister, G. S. and Davis, G. A.: The origin of metamorphic core complexes and detachment faults formed during Tertiary continental extension in the northern Colorado River region, USA, *J. Struct. Geol.*, 11, 65–94, 1989.

- Lopes, G., Mangerud, G., and Clayton, G.: The palynostratigraphy of the Mississippian Birger Johnsonfjellet section, Spitsbergen, Svalbard, *Palynology*, 43, 631–649, 2019.
- Lopes, G., Mangerud, G., Clayton, G., and Vigran, J. O.: Palynostratigraphic reassessment of the Late Devonian of Bjørnøya, Svalbard, *Rev. Palaeobot. Palynol.*, 286, 104376, <https://doi.org/10.1016/j.revpalbo.2020.104376>, 2021.
- Lowell, J. D.: Spitsbergen Tertiary Orogenic Belt and the Spitsbergen Fracture Zone, *Geol. Soc. Am. Bul.*, 83, 3091–3102, 1972.
- Luchitskaya, M. V., Sokolov, S. D., Kotov, A. B., Natapov, L. M., Belousova, E. A., and Katkov, S. M.: Late Paleozoic Granitic Rocks of the Chukchi Peninsula: Composition and Location in the Structure of the Russian Arctic, *Geotectonics*, 49, 243–266, 2015.
- Maher, H. D.: Structure and stratigraphy of the Midterhuken peninsula, Bellsund, west Spitsbergen, PhD Thesis, University of Wisconsin–Madison, Madison, USA, Norwegian Polar Institute, Saga Petroleum, 437, 1984.
- Maher Jr., H. D.: Atypical rifting during the Carboniferous of the NW Barents Shelf, Report for Saga Petroleum, November 1996, Saga Petroleum, 1996.
- Maher Jr., H. D., Craddock, C. and Maher, K.: Kinematics of Tertiary structures in upper Paleozoic and Mesozoic strata on Midterhuken, west Spitsbergen, *GSA Bulletin*, 97, 1411–1421, 1986.
- Maher Jr., H. D., Ringset, N. and Dallmann, W. K.: Tertiary structures in the platform cover strata of Nordenskiöld Land, Svalbard, *Polar Res.*, 7, 83–93, 1989.
- Maher, H., Braathen, A., Ganerod, M., Osmundsen, P. T., Redfield, T., Myhre, P. I., Serck, C. and Parcher, S.: Core complex fault rocks of the Silurian to Devonian Keisarhjelmen detachment in NW Spitsbergen, in: *New Developments in the Appalachians–Caledonian–Variscan Orogen*, edited by: Kuiper, Y. D., Murphy, J. B., Nance, R. D., Strachan, R. A. and Thompson, M. D., *GSA Spec. Paper*, 54, 265–286, 2022.
- Majka, J. and Kościńska, K.: Magmatic and metamorphic events recorded within the Southwestern basement province of Svalbard, *Arktos*, 3, 1–7, <https://doi.org/10.1007/s41063-017-0034-7>, 2017.
- Malyshev, N. A., Obmetko, V. V., Barinova, E. M. and Ikhsanov, B. I.: Tectonics of the sedimentary basins in the Russian sector of the Chuckchi Sea, in: *Proceedings of the International Conference on Arctic Margins VI*, edited by: Stone, D. B., Griukurov, G. E., Glough, J. G., Oakey, G. N. and Thurston, D. K., May 2011, Fairbanks, Alaska, USA, 203–209, 2011.
- Manby, G. M. and Lyberis, N.: Tectonic evolution of the Devonian Basin of northern Spitsbergen, *Norsk Geol. Tidsskr.*, 72, 7–19, 1992.
- Manby, G. M. and Lyberis, N.: State of stress and tectonic evolution of the West Spitsbergen Fold Belt, *Tectonophys.*, 267, 1–29, 1996.
- Manby, G. M., Lyberis, N., Chorowicz, J. and Thiedig, F.: Post-Caledonian tectonics along the Billefjorden fault zone, Svalbard, and implications for the Arctic region, *Geol. Soc. Am. Bul.*, 105, 201–216, 1994.
- Marshall, J., Lindemann, F. J., Finney, S. and Berry, C.: A Mid Famennian (Late Devonian) spore assemblage from Svalbard and its significance, CIMP Meeting, Bergen, Norway, 17–18 September, <https://cimp.weebly.com/uploads/6/4/0/>5/6405206/cimp2015_abstract_book.pdf (last access: 25 August 2022), 2015.
- McCann, A. J.: Deformation of the Old Red Sandstone of NW Spitsbergen; links to the Ellesmerian and Caledonian orogenies, in: *New Perspectives on the Old Red Sandstone*, edited by: Friends, P. F. and Williams, B. P. J., *Geol. Soc. London*, 180, 567–584, <https://doi.org/10.1144/GSL.SP.2000.180.01.30>, 2000.
- McCann, A. J. and Dallmann, W. K.: Reactivation of the long-lived Billefjorden Fault Zone in north central Spitsbergen, Svalbard, *Geol. Mag.*, 133, 63–84, 1996.
- McClelland, W. C., Power, S. E., Gilotti, J. A., Mazdab, F. K. and Wopenka, B.: U-Pb SHRIMP geochronology and trace-element geochemistry of coesite-bearing zircons, North-East Greenland Caledonides, *GSA, Spec. Paper*, 403, 23–43, 2006.
- Michaelsen, B.: Strukturgeologie des svalbardischen Überschiebungs- und Faltengürtels im zentralen, östlichen Dickson Land, Spitzbergen (Structural geology of the Svalbardian fold-and-thrust belt in central–eastern Dickson Land, Spitsbergen), Master's Thesis, University of Münster, Münster, Germany, available from the Norwegian Polar Institute, 134 p., 1998.
- Michaelsen, B., Piepjohn, K. and Brinkmann, L.: Struktur und Entwicklung der svalbardischen Mimerelva Syncline im zentralen Dickson Land, Spitzbergen, Münster. *Forsch. Geol. Paläont.*, 82, 203–214, 1997.
- Michalski, K.: Palaeomagnetism of metacarbonates and fracture fills of Kongsfjorden islands (western Spitsbergen): Towards a better understanding of late- to post-Caledonian tectonic rotations, *Polish Polar Res.*, 39, 51–75, 2018.
- Michalski, K., Lewandowski, M., and Manby, G.: New palaeomagnetic, petrographic and $^{40}\text{Ar}/^{39}\text{Ar}$ data to test palaeogeographic reconstructions of Caledonide Svalbard, *Geol. Mag.*, 149, 696–721, 2012.
- Michalski, K., Manby, G., Nejbort, K., Domanska-Siuda, J., and Burzynski, M.: Using palaeomagnetism and isotopic data to investigate late to post-Caledonian tectonothermal processes within the Western Terrane of Svalbard, *J. Geol. Soc., London*, 174, 572–590, 2017.
- Murascov, L. G. and Mokin, Ju. I.: Stratigraphic subdivision of the Devonian deposits of Spitsbergen, *Polarinst. Skr.*, 167, 249–261, 1979.
- Newman, M. J., Burrow, C. J., and den Blaauwen, J. L.: The Givetian vertebrate fauna from the Fiskekløfta Member (Mimerdalen Subgroup), Svalbard, Part I. Stratigraphic and faunal review, Part II. Acanthodii, *Norw. J. Geol.*, 99, 1–16, 2019.
- Newman, M. J., Burrow, C. J., and den Blaauwen, J. L.: A new species of ischnacanthiform acanthodian from the Givetian of Mimerdalen, Svalbard, *Norw. J. Geol.*, 99, 619–631, 2020.
- Newman, M. J., Burrow, C. J., den Blaauwen, J. L., and Giles, S.: A new actinopterygian *Cheirolepis jonesi* nov. sp. from the Givetian of Spitsbergen, Svalbard, *Norw. J. Geol.*, 101, 202103, <https://doi.org/10.17850/njg101-1-3>, 2021.
- Oakey, G. N. and Chalmers, J. A.: A new model for the Paleogene motion of Greenland relative to North America: Plate reconstructions of the Davis Strait and Nares Strait regions between Canada and Greenland, *J. of Geophys. Res.*, 117, B10401, <https://doi.org/10.1029/2011JB008942>, 2012.
- Ohta, Y., Dallmeyer, R. D., and Peucat, J. J.: Caledonian terranes in Svalbard, *GSA Spec. Paper*, 230, 1–15, 1989.

- Ohta, Y., Krasil'shchikov, A. A., Lepvrier, C., and Teben'kov, A. M.: Northern continuation of Caledonian high-pressure metamorphic rocks in central-western Spitsbergen, *Polar Res.*, 14, 303–315, 1995.
- Osmundsen, P. T. and Andersen, T. B.: Caledonian compressional and late-orogenic extensional deformation in the Staveneset area, Sunnfjord, Western Norway, *J. Struct. Geol.*, 16, 1385–1401, 1994.
- Osmundsen, P. T., Andersen, T. B., Markussen, S., and Svendby, A. K.: Tectonics and sedimentation in the hangingwall of a major extensional detachment: the Devonian Kvamshesten basin, western Norway, *Basin Res.*, 10, 213–234, 1998.
- Petersen, T. G., Thomsen, T. B., Olaussen, S., and Stemmerik, L.: Provenance shifts in an evolving Eurekan foreland basin: the Tertiary Central Basin, Spitsbergen, *J. Geol. Soc.*, 173, 634–648, 2016.
- Piepjohann, K.: The Svalbardian–Ellesmerian deformation of the Old Red Sandstone and the pre-Devonian basement in NW Spitsbergen (Svalbard), in: *New Perspectives on the Old Red Sandstone*, edited by: Friend, P. F. and Williams, B. P. J., *Geol. Soc. London Spec. Publi.*, 180, 585–601, <https://doi.org/10.1144/GSL.SP.2000.180.01.31>, 2000.
- Piepjohann, K. and Dallmann, W. K.: Stratigraphy of the uppermost Old Red Sandstone of Svalbard (Mimerdalen Subgroup), *Polar Res.*, 33, 19998, 2014.
- Piepjohann, K. and von Gosen, W.: Structural transect through Ellesmere Island (Canadian Arctic): superimposed Palaeozoic Ellesmerian and Cenozoic Eurekan deformation, in: *Circum-Arctic Lithosphere Evolution*, edited by: Pease, V. and Coackley, B., *Geol. Soc., London, Spec. Publi.*, 460, 33, <https://doi.org/10.1144/SP460.5>, 2017.
- Piepjohann, K., Brinkmann, L., Dissmann, B., Grewing, A., Michaelsen, B., and Kerp, H.: Geologische und strukturelle Entwicklung des Devon im zentralen Dickson Land, Spitzbergen, *Münster. Forsch. Geol. Paläont.*, 82, 175–202, 1997.
- Piepjohann, K., Brinkmann, L., Grewing, A., and Kerp, H.: New data on the age of the uppermost ORS and the lowermost post-ORS strata in Dickson Land (Spitsbergen) and implications for the age of the Svalbardian deformation, in: *New Perspectives on the Old Red Sandstone*, edited by: Friend, P. F. and Williams, B. P. J., *Geol. Soc. London Spec. Publi.*, 180, 603–609, 2000.
- Piepjohann, K., Thiedig, F., and Manby, G. M.: Nappe Stacking on Brøggerhalvøya, NW Spitsbergen, *Geol. Jb, B*, 91, 55–79, 2001.
- Piepjohann, K., von Gosen, W., Estrada, S., and Tessensohn, F.: Deciphering superimposed Ellesmerian and Eurekan deformation, Piper Pass area, northern Ellesmere Island (Nunavut), *Can. J. Earth Sci.*, 44, 1439–1452, 2007.
- Piepjohann, K., von Gosen, W., Tessensohn, F., and Saalman, K.: Ellesmerian fold-and-thrust belt (northeast Ellesmere Island, Nunavut) and its Eurekan overprint, in: *Geology of Northeast Ellesmere Island Adjacent to Kane Basin and Kennedy Channel, Nunavut*, edited by: Mayr, U., *Geological Soc. Can. Bull.*, 592, 285–303, 2008.
- Piepjohann, K., von Gosen, W., Läufer, A., McClelland, W. C., and Estrada, S.: Ellesmerian and Eurekan fault tectonics at the northern margin of Ellesmere Island (Canadian High Arctic), *Z. Dt. Ges. Geowiss.*, 164, 81–105, 2013.
- Piepjohann, K., von Gosen, W., Tessensohn, F., Reinhardt, L., McClelland, W. C., Dallmann, W. D., Gaedicke, C., and Harrison, J. C.: Tectonic map of the Ellesmerian and Eurekan deformation belts on Svalbard, North Greenland, and the Queen Elizabeth Islands (Canadian Arctic), *Arktos*, 1, 12, <https://doi.org/10.1007/s41063-015-0015-7>, 2015.
- Platt, J. P.: Dynamics of orogenic wedges and the uplift of high-pressure metamorphic rocks, *GSA Bull.*, 97, 1037–1053, 1986.
- Playford, G.: Lower Carboniferous microfloras of Spitsbergen, Part 1, *Paleontology*, 5, 550–618, 1962.
- Playford, G.: Lower Carboniferous microfloras of Spitsbergen, Part 2, *Paleontology*, 5, 4, 619–678, 1963.
- Playford, G.: Plant microfossils from the Upper Devonian and Lower Carboniferous of the Canning Basin, Western Australia, *Palaeontographica Abt. B*, 158, 1–71, 1976.
- Pčelina, T. M., Bogač, S. I., and Gavrilov, B. P.: Novye dannye po litostratigrafii devonskih otloženij rajona Mimerdalen arhipelaga Špicbergen (New data on the lithostratigraphy of the Devonian deposits of the region of Mimerdalen of the Svalbard Archipelago), in: *Geologija osadocnogo cehla arhipelaga Špicbergen (Geology of the sedimentary blanket of the archipelago of Spitsbergen)*, edited by: Krasil'shchikov, A. A. and Mirzaev, M. N., *Leningrad: Sevmorgeologija*, 7–19, 1986.
- Rey, P., Vanderhaeghe, O., and Teyssier, C.: Gravitational collapse of the continental crust: definition, regimes and modes, *Tectonophys.*, 342, 435–449, 2001.
- Rey, P., Teyssier, C., Kruckenberg, S. C., and Whitney, D. L.: Viscous collision in channel explains double domes in metamorphic core complexes, *Geology*, 39, 387–390, 2011.
- Ripington, S. J., Scott, R. A., Smyth, H., Bogolepova, O. K., and Gubanov, A. P.: The Ellesmerian Orogeny: Fact of Fiction?, *Geo-Canada 2010, Working with the Earth*, 10–14 May, Calgary, Alberta, Canada, https://geoconvention.com/wp-content/uploads/abstracts/2010/0411_GC2010_The_Ellesmerian_Orogeny.pdf (last access: 25 August 2022), 2010.
- Roberts, D.: Devonian Tectonic Deformation in the Norwegian Caledonides and Its Regional Perspectives, *Norg. Geol. Unders.*, 380, 85–96 1983.
- Roy, J.-C. L. G.: La géologie du fossé des Vieux Grès Rouges du Spitzberg (archipel du Svalbard, territoire de l'Arctique) – Synthèse stratigraphique, conséquences paléoenvironnementales et tectoniques synsédimentaires, *Mémoires des sciences de la Terre de l'Université Pierre et Marie Curie*, PhD Thesis, Pierre and Marie Curie University, Paris, France, 2007–15, 242 p., 2007.
- Roy, J.-C. L. G.: La saga des vieux grès rouges du Spitzberg (archipel du Svalbard, Arctique): Une histoire géologique et naturelle, *Charenton-le-pont: Auto-Edition Roy-Poulain*, 290 p., 2009.
- Roy, J.-C., Chorowicz, J., Deffontaines, B., Lepvrier, C., and Tardy, M.: Clues of gravity sliding tectonics at the Eifelien–Givetian boundary in the Old Red Sandstone of the [late Silurian?]-Devonian trough of Andrée Land (Spitsbergen), in: *La saga des vieux grès rouges du Spitzberg (archipel du Svalbard, Arctique): Une histoire géologique et naturelle*, edited by: Charenton-le-pont: Auto-Edition Roy-Poulain, *Norw. J. Geol.*, 2022, unpublished.
- Schaltegger, U., Schmidt, A. K. and Horstwood, M. S. A.: U–Th–Pb zircon geochronology by ID-TIMS, SIMS, and laser ablation ICP-MS: Recipes, interpretations, and opportunities, *Chem. Geol.*, 402, 89–110, 2015.

- Scheibner, C., Hartkopf-Fröder, C., Blomeier, D. and Forke, H.: The Mississippian (Lower Carboniferous) in northeast Spitsbergen (Svalbard) and a re-evaluation of the Billefjorden Group, *Z. Dt. Ges. Geowiss.*, 163/3, 293–308, 2012.
- Schneider, D. A., Faehrich, K., Majka, J., and Manecki, M.: $^{40}\text{Ar}/^{39}\text{Ar}$ geochronologic evidence of Eurekan deformation within the West Spitsbergen Fold and Thrust Belt, in: *Circum-Arctic Structural Events: Tectonic Evolution of the Arctic Margins and Trans-Arctic Links with Adjacent Orogens*, edited by: Piepjohn, K., Strauss, J. V., Reinhardt, L. and McClelland, W. C., GSA Special Paper, 541, 1–16, [https://doi.org/10.1130/2018.2541\(08\)](https://doi.org/10.1130/2018.2541(08)), 2018.
- Schweitzer, H.-J.: *Die Devonflora Spitzbergens (The Devonian flora of Spitsbergen)*, Palaeontographica Abteilung B Band, 252, Stuttgart, Schweizerbart Science Publishers, 1999.
- Senger, K., Roy, S., Braathen, A., Buckley, S., Bælum, K., Gernigon, L., Mjelde, R., Noormets, R., Ogata, K., Olaussen, S., Planke, S., Ruud, B. O., and Tveranger, J.: Geometries of doleritic intrusions in central Spitsbergen, Svalbard: an integrated study of an onshore-offshore magmatic province with applications to CO₂ sequestration, *Norw. J. Geol.*, 93, 143–166, 2013.
- Siedlecki, S.: *Culm Beds of the SW. Coast of Hornsund*, *Studia Geologica Polonica*, 4, 93–102, 1960.
- Siedlecki, S. and Turnau, E.: Palynological investigations of *Culm* in the area SW of Hornsund, Vestspitsbergen, *Studia Geologica Polonica*, 11, 125–138, 1964.
- Smyrak-Sikora, A. A., Johannessen, E. P., Olaussen, S., Sandal, G., and Braathen, A.: Sedimentary architecture during Carboniferous rift initiation – the arid Billefjorden Trough, Svalbard, *J. Geol. Soc. Lond.*, 176, 225–252, 2018.
- Snoke, A. W.: Transition from infrastructure to suprastructure in the northern Ruby Mountains, Nevada, in: *Cordilleran Metamorphic Core Complexes*, edited by: Crittenden Jr., M. D., Coney, P. J. and Davis, G. H., GSA Memoirs, 153, 287–333, <https://doi.org/10.1130/MEM153-p287>, 1980.
- Stemmerik, L., Vigran, J. O., and Piasecki, S.: Dating of late Paleozoic rifting events in the North Atlantic: New biostratigraphic data from the uppermost Devonian and Carboniferous of East Greenland, *Geology*, 19, 218–221, 1991.
- Stemmerik, L., Dalhoff, D., Larsen, B. D., Lyck, J., Mathiesen, A., and Nilsson, I.: Wandel Sea Basin, eastern North Greenland, *Geol. Greenland Bull.*, 180, 55–62, 1998.
- Stemmerik, L.: Late Palaeozoic evolution of the North Atlantic margin of Pangea, *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 161, 95–126, 2000.
- Strachan, R. A.: Evidence in North-East Greenland for Late Silurian–Early Devonian regional extension during the Caledonian orogeny, *Geology*, 22, 913–916, 1994.
- Teyssier, C., Ferré, E. C., Whitney, D. L., Norlander, B., Vandrahaeghe, O. and Parkinson, D.: Flow of partially molten crust and origin of detachments during collapse of the Cordilleran Orogen, in: *High-Strain Zones: Structure and Physical Properties*, edited by: Bruhn, D. and Burlini, L., Geol. Soc. London, Spec. Publ., 245, 39–64, 2005.
- Thiedig, F. and Manby, G.: Origins and deformation of post-Caledonian sediments on Blomstrandhalvøya and Lovénøyane, northwest Spitsbergen, *Norsk Geol. Tidsskr.*, 72, 27–33, 1992.
- Thorsteinsson, R. and Tozer, E. T.: Geology of the Arctic Archipelago, in: *Geology and Economic Minerals of Canada (5th edition)*, edited by: Douglass, R. J. W., Geological Survey of Canada, Economic Geology Report, 1, 547–590, <https://doi.org/10.4095/106153>, 1970.
- Trettin, H. P.: Early Paleozoic Evolution of Northern Parts of Canadian Arctic Archipelago, *AAPG Memoirs*, 19, 57–75, 1973.
- Trettin, H. P.: Geology of the Innuitian Orogen and Arctic Platform of Canada and Greenland, Geological Survey of Canada, *Geology of Canada*, 3, GSA, Geological Survey of Canada, *Geology of Canada Series no. 3*, Geological Society of America, *Geology of North America Series E*, Canada Communication Group – Publishing Ottawa, Canada KIA OS9, 569 p., <https://doi.org/10.4095/133959>, 1991.
- Vigran, J. O.: Spores from Devonian deposits, Mimerdalen, Spitsbergen, *Norsk Polarinstitutt Skrifter*, 132, 49 p., <https://brage.npolar.no/npolar-xmlui/handle/11250/173920> (last access: 25 August 2022), 1964.
- Vogt, T.: The stratigraphy and tectonics of the Old Red formations of Spitsbergen, *Abstracts of the Proceedings of the Geological Society London*, 1343, 88, 1938.
- Von Gosen, W. and Piepjohn, K.: Polyphase Deformation in the Eastern Hornsund Area, *Geol. Jb.*, B91, 291–312, 2001.
- Walker, J. D., Geissman, J. W., Bowring, S. A., and Babcock, L. E.: *Geologic Time Scale v. 5.0*, Geological Society of America, <https://doi.org/10.1130/2018.CTS005R3C>, 2018.
- Worsley, D. and Edwards, M. B.: The Upper Palaeozoic succession of Bjørnøya, *Norsk Polarinst. Årbok*, 1974, 17–34, 1976.
- Worsley, D. and Mørk, A.: The Triassic stratigraphy of southern Spitsbergen, *Polarinst. Årbok*, 1977, 43–60, 1978.
- Ziemniak, G., Majka, J., Manecki, M., Walczak, K., Jeanneret, P., Mazur, S., and Kościńska, K.: Early Devonian sinistral strike-slip in the Caledonian basement of Oscar II Land advocates for escape tectonics as a major mechanism for Svalbard terranes assembly, *Geophys. Res. Abstr.*, EGU General Assembly, 3–8 May 2020, Vienna, Austria, 2020. <https://doi.org/10.1130/2018.CTS005R3C>.