



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/190894/>

Version: Accepted Version

---

**Article:**

Curk, T, Dubacheva, GV, Brisson, AR et al. (2022) Controlling Superselectivity of Multivalent Interactions with Cofactors and Competitors. *Journal of the American Chemical Society*, 144 (38). pp. 17346-17350. ISSN: 0002-7863

<https://doi.org/10.1021/jacs.2c06942>

---

© 2022 The Authors. Published by American Chemical Society. "This is an author produced version of an article published in the *Journal of the American Chemical Society*. Uploaded in accordance with the publisher's self-archiving policy. "

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Controlling superselectivity of multivalent interactions with cofactors and competitors

Tine Curk,<sup>a,¶,\*</sup> Galina V. Dubacheva,<sup>b,¶</sup> Alain R. Brisson,<sup>c</sup> and Ralf P. Richter,<sup>d,\*</sup>

<sup>a</sup> Department of Materials Science and Engineering, Northwestern University, Evanston, IL 60208, USA

<sup>b</sup> Département de Chimie Moléculaire, Université Grenoble Alpes, CNRS UMR 5250, 38000 Grenoble, France

<sup>c</sup> UMR-CBMN, CNRS-University of Bordeaux-IPB, 33600 Pessac, France


<sup>d</sup> School of Biomedical Sciences, Faculty of Biological Sciences, School of Physics and Astronomy, Faculty of Engineering and Physical Sciences, Astbury Centre for Structural Molecular Biology, and Bragg Centre for Materials Research, University of Leeds, Leeds, LS2 9JT, United Kingdom

**KEYWORDS:** superselectivity, multivalent interactions, annexin, hyaluronan

**ABSTRACT:** Moieties that compete with multivalent interactions or act as cofactors are common in living systems, but their effect on multivalent binding remains poorly understood. We derive a theoretical model that shows how the superselectivity of multivalent interactions is modulated by the presence of cofactors or competitors. We find that the role of these participating moieties can be fully captured by a simple rescaling of the affinity constant of the individual ligand-receptor bonds. Theoretical predictions are supported by experimental data of the membrane repair protein annexin A5 binding to anionic lipid membranes in the presence of  $\text{Ca}^{2+}$  cofactors, and of the extracellular matrix polysaccharide hyaluronan (HA) binding to CD44 cell surface receptors in the presence of HA oligosaccharide competitors. The obtained findings should facilitate understanding of multivalent recognition in biological systems and open new routes for fine-tuning the selectivity of multivalent nanoprobes in medicinal chemistry.

Multivalent interactions involve the simultaneous formation of multiple supramolecular bonds, such as ligand-receptor binding<sup>1</sup> or host-guest complexation.<sup>2-3</sup> The combinatorial entropy of possible binding configurations gives rise to a supra-linear change in the number of bound multivalent probes as a function of receptor concentration.<sup>4-5</sup> This superselective behavior<sup>6</sup> allows specific targeting of surfaces displaying binding sites above a threshold surface concentration, while leaving surfaces with lower coverages virtually unaffected (Fig. 1A). The types of multivalent entities that display superselectivity vary widely, including proteins,<sup>7</sup> antibodies,<sup>4,8</sup> polymers,<sup>9-10</sup> viruses,<sup>11-13</sup> liposomes, and nanoparticles.<sup>14-15</sup> Resolving the mechanism of multivalent interactions is crucial both to understand the selectivity of biomolecular interactions and to facilitate the design of highly selective nanoprobes for diagnostics and therapies.<sup>16</sup>

Previous studies of superselectivity in synthetic and living systems have clarified the roles of the affinity of individual ligand-receptor bonds, probe valency, receptor surface density and in-plane mobility in multivalent binding.<sup>2, 14, 17</sup> In addition to these factors, biological systems commonly involve interacting moieties that modulate multivalent interactions. For example, many specific interactions in biochemistry require a cofactor (e.g., a multivalent ion or a small molecule) to form a bond and the strength of the interaction can be tuned by varying the concentration of cofactors.<sup>7, 18</sup> Likewise, competing interactions such as agonists vs. antagonists are common in biology. The effect of cofactors (Fig. 1B) or competitors (Fig. 1C) on multivalent binding remains largely unexplored, hampering the wider application of superselectivity concepts. Cofactors and competitors modulate the effective number of available receptors, and we hypothesize that a superselective response towards changes in receptor density naturally extends to modulations in cofactor or competitor concentrations (Fig. 1A, *bottom*).



**Fig. 1. Multivalent interactions in the presence of competitors and cofactors.** Superselectivity of multivalent probes to changes in receptor density (A, *top*) is modulated by the presence of cofactors (B) or competitors (C). Illustrative plot of probe density (solid black line) and corresponding selectivity parameters  $\alpha$  (dashed red line) vs. receptor density, and cofactor or competitor concentrations (A, *bottom*). Insets show the relevant reaction equilibria.

Here, we demonstrate based on simple theoretical arguments that cofactors and monovalent competitors impact superselective binding by effectively re-scaling the ligand-receptor affinity. We apply this insight to two important yet distinct examples of biomolecular interactions, namely, the  $\text{Ca}^{2+}$  dependent binding of the membrane repair protein annexin A5 (AnxA5) to anionic lipid membranes,<sup>7</sup> and the effect of competing oligosaccharides on the recognition of the extracellular matrix polysaccharide hyaluronan (HA) by CD44 cell surface receptors.<sup>19</sup>

The theory of multivalent binding<sup>2, 5-6</sup> predicts that the strength of the multivalent interaction, or avidity constant  $K_{av}$ , depends supra-linearly on the receptor density  $\Gamma_R$ , and the ligand–receptor dissociation constant  $K_d$  (Fig. 1A) as

$$K_{av} = a^3 N_A \left[ \left( 1 + \frac{n_R}{K_d v_{eff}} \right)^{n_L} - 1 \right], \quad [1]$$

where  $a$  and  $n_L$  are the size and valency of the multivalent probe, respectively,  $n_R = a^2 \Gamma_R$  the number of accessible receptors,  $N_A$  Avogadro’s number, and  $v_{eff}$  the effective free volume that each unbound ligand can explore (the ratio  $n_R/v_{eff}$  is also called ‘effective molarity’<sup>1, 20</sup>). When binding to a surface, this equation can be used as an input to the Langmuir isotherm, which predicts the surface density of adsorbed probes to be

$$\Gamma_p = \Gamma_{max} \frac{K_{av} c_p}{1 + K_{av} c_p}, \quad [2]$$

with the maximum possible density  $\Gamma_{max}$  and the concentration of unbound probes  $c_p$ . The binding is said to be superselective if the surface density increases faster than linearly with the receptor density, i.e., if the selectivity parameter

$$\alpha_R = \frac{d \log \Gamma_p}{d \log n_R} \quad [3]$$


is larger than unity (Fig. 1A). Here we extend this theory to fully capture the effect of cofactors and competitors, including selectivity with regard to cofactor concentration  $c_{cf}$  ( $\alpha_{cf} = d \log \Gamma_p / d \log c_{cf}$ ) and competitor concentration  $c_{mc}$  ( $\alpha_{mc} = -d \log \Gamma_p / d \log c_{mc}$ ; the minus sign ensures that  $\alpha_{mc} > 0$ , since binding generally decreases with  $c_{mc}$ ). The full theoretical derivation that considers the distribution of all possible binding states in equilibrium is provided in the Supporting Information, with only the main results being shown here.

**Cofactors.** We consider monovalent cofactors at (unbound) concentration  $c_{cf}$  that bind to ligands and receptors with the dissociation constants  $K_{d,L-cf}$  and  $K_{d,R-cf}$ , while the ligand–cofactor (or receptor–cofactor) complex binds to the receptors (or ligands) with constant  $K_{d,Lcf-R}$  (or  $K_{d,Rcf-L}$ ) (Fig. 1B). The effect of cofactors can be fully captured by using a generalized ligand–receptor ‘affinity’, with an effective dissociation constant

$$K_d^{(cf)} = \frac{K_{d,L-cf-R}}{c_{cf}} \left( 1 + \frac{c_{cf}}{K_{d,R-cf}} \right) \left( 1 + \frac{c_{cf}}{K_{d,L-cf}} \right), \quad [4]$$

where  $K_{d,L-cf-R} = K_{d,L-cf} K_{d,Lcf-R} = K_{d,R-cf} K_{d,Rcf-L}$  is the tripartite affinity constant.

At low cofactor concentrations,  $c_{cf} < K_{d,L-cf}$  and  $c_{cf} < K_{d,R-cf}$ , we can approximate  $K_d^{(cf)} \approx K_{d,L-cf-R} / c_{cf}$  and thus changing the cofactor concentration has the same effect as changing the receptor density  $n_R$  (Eq. [1]) and yields an equivalent superselective response ( $\alpha_{cf} \approx \alpha_R$ ; Fig. 1A, *bottom*). At intermediate concentrations,  $K_{d,R-cf} > c_{cf} > K_{d,L-cf}$  or  $K_{d,L-cf} > c_{cf} > K_{d,R-cf}$ , either the ligands or receptors are saturated with cofactors and changing the cofactor concentration has no effect:  $K_d^{(cf)} \approx \max[K_{d,Lcf-R}, K_{d,Rcf-L}]$ . Lastly, at very high concentrations,  $c_{cf} > K_{d,L-cf}$  and  $c_{cf} > K_{d,R-cf}$ , the oversaturation with cofactors weakens the effective binding:  $K_d^{(cf)} \approx c_{cf} K_{d,L-cf-R} / (K_{d,R-cf} K_{d,L-cf})$  and thus changing  $c_{cf}$  has the same effect as changing the inverse receptor density  $n_R^{-1}$  (Fig 2A). Often, however, only the low concentration regime is biologically relevant. These features can be employed to control the range of superselective receptor recognition by tuning the cofactor concentration (Fig. 2A). Thus, the influence of cofactors does not change the nature of multivalent binding, rather, it simply rescales the affinity constant according to Eq. [4].



**Fig. 2. Effect of cofactors.** (A) Example of the dependence of the selectivity parameter  $\alpha_R$  on the receptor surface density and cofactor concentration (Eqs. [1-4];  $n_L = 8$ ,  $c_p a^3 N_A = 0.001$ ,  $K_{d,R-cf} = 100 K_{d,L-cf}$ ). (B) Schematic of AnxA5 (PDB code 1AVR<sup>21</sup>) binding to supported lipid bilayers presenting PS lipids in a background of PC lipids. (C) Experimental dependence of AnxA5 (non-oligomerizing mutant at  $c_p = 0.56 \mu\text{M}$ ) binding on PS density at different  $\text{Ca}^{2+}$  concentrations (*symbols*; *error bars* represent experimental precision) is well reproduced by the theory (*solid lines in matching colors*) that explicitly models binding to the two types of lipids and membrane fluidity (see Supporting Information). (D) The sets of data at different  $\text{Ca}^{2+}$  concentration collapse onto a master curve when plotted as a function of  $f_{PS} \times [\text{Ca}^{2+}]$ . Slopes with  $\alpha$  values are included in (C) and (D) for reference.

A salient biological example of how cofactors influence multivalent interactions is AnxA5 binding to lipid membranes (Fig. 2B). AnxA5 functions as a cell membrane scaffolding and repair protein.<sup>22</sup> It preferentially binds anionic phospholipids, and requires  $\text{Ca}^{2+}$  as a cofactor for membrane binding.<sup>7</sup> In intact cells, anionic phospholipids reside in the inner (but not the outer) leaflet of the plasma membrane, whereas  $\text{Ca}^{2+}$  ions are virtually absent in the cytoplasm but present (in mM concentrations) outside the cell. AnxA5 thus binds to the cell membrane only upon membrane damage leading to influx of  $\text{Ca}^{2+}$  ions into the cell and possibly also to inter-leaflet lipid content mixing near the damage site.

Experimental data reveal superselective binding of AnxA5 to lipid membranes presenting anionic phosphatidyl serine (PS) in a background of zwitterionic phosphatidyl choline (PC) lipids,

and our theoretical model predicts well AnxA5 binding over four orders of magnitude of  $\text{Ca}^{2+}$  concentrations (Fig. 2C). Moreover, within the range of the investigated calcium concentrations, the binding of  $\text{Ca}^{2+}$  to both AnxA5 and PS lipids appears to be weak:  $c_{\text{cf}}/K_{\text{d,L-cf}} < 1$  and  $c_{\text{cf}}/K_{\text{d,R-cf}} < 1$ . Thus, Eq. [4] can be approximated as  $K_{\text{d}}^{(\text{cf})} = K_{\text{d,L-cf-R}}/c_{\text{cf}}$ , which implies that AnxA5 binding depends only on the product  $n_{\text{R}}c_{\text{cf}}$  (Eq. [1]), where  $n_{\text{R}} = f_{\text{PS}}(a/l)^2$ , with the protein cross-section  $a^2 = 25 \text{ nm}^2$ , the lipid cross-section  $l^2 = 0.7 \text{ nm}^2$ , and the PS lipid fraction  $f_{\text{PS}}$ . Indeed, when the AnxA5 binding data are plotted as a function of  $f_{\text{PS}}c_{\text{cf}}$ , all experimental data collapse onto a single master curve (Fig. 2D), thus validating our theory.

Our analysis identifies membrane recognition by AnxA5 as a striking example of superselective binding, demonstrating that binding is strongly superselective with respect to the cofactor  $\text{Ca}^{2+}$  as well as the receptor PS lipids, with maximal  $\alpha$  values  $\alpha_{\text{cf,max}} \approx \alpha_{\text{R,max}} \approx 4$  (Fig. 2D). This enables the protein to effectively respond to slight changes in the concentration of either of these two factors, which is crucial for its function as a membrane repair protein. We note that effective membrane repair additionally requires AnxA5 to organize into trimers and two-dimensional crystals on the membrane.<sup>22</sup> To probe superselective binding of the AnxA5 monomers, we have in Fig. 2 probed an AnxA5 mutant that does not oligomerize yet retains the membrane binding properties of the wild type protein. However, the superselective effects are retained, and even further accentuated, by the self-organization of the wild-type protein on the membrane (see Supporting Information).

**Competitors.** Similar to the theoretical treatment of cofactors, monovalent competitors are assumed to bind to surface receptors with the affinity constant  $K_{\text{d,R-mc}}$ . As shown in the full derivation of our analytical model, competitors at (unbound) concentration  $c_{\text{mc}}$  effectively rescale the ligand–receptor affinity  $K_{\text{d}}$  to

$$K_{\text{d}}^{(\text{mc})} = K_{\text{d}} \left( 1 + \frac{c_{\text{mc}}}{K_{\text{d,R-mc}}} \right). \quad [5]$$

The impact of this rescaling on superselective binding is illustrated in Fig. 3A and shows that increasing the competitor concentration pushes the range of superselective binding towards higher receptor densities. Equation [5] is well-known for monovalent interactions;<sup>29</sup> we here establish that it also applies to multivalent interactions and can be generalized to multiple competitor types (see Supporting Information).


We here test our simple model on data reported by Lesley et al.<sup>19</sup> on the inhibition of HA polysaccharide binding to CD44 cell surface receptors by HA oligosaccharides (Fig. 3B). That HA binding to cells depends sharply on receptor surface density is evident from previous work<sup>10, 23</sup>. Such superselective recognition is important for cell–extracellular matrix communication, and changes in HA presentation can dramatically affect recognition, e.g., inflammation entails degradation of large HA polysaccharides (MDa range) into small oligosaccharides. HA octasaccharides ( $\text{HA}_8$ ) just about fill the binding groove in a CD44 receptor,<sup>24</sup> and thus are effective monovalent competitors.

The simple analytical model (Eqs. [1-2]) with the re-scaled affinity  $K_{\text{d}}^{(\text{mc})}$  (Eq. [5]) reproduces the experimental data well (Fig. 3C), illustrating that it captures the salient features of the competition process. In the model, we fixed  $n_{\text{L}} = 500$  distinct sites for binding to CD44 receptors (consistent with an HA molecular mass of  $\sim 1 \text{ MDa}$  and a decasaccharide ‘footprint’ per receptor), a coil volume of  $a^3 = 4\pi R_{\text{g}}^3/3$  (with the radius of gyration,  $R_{\text{g}} \approx 90 \text{ nm}$ ,<sup>25</sup> and a concentration of  $c_{\text{P}} \approx 0.5 \text{ nM}$ ), and

$K_{\text{d,R-mc}} \approx 50 \mu\text{M}$  (within the broad range of reported values<sup>19, 24, 26</sup>). As the only fitting parameter, we determined  $n_{\text{R}}/(K_{\text{d}}v_{\text{eff}}) \approx 0.03$ , a value that is consistent with typical CD44 cell surface densities (see Supporting Information), i.e., the simple model makes reasonable quantitative predictions.

Importantly, we demonstrate that the binding response can be superselective with respect to the competitor concentration  $c_{\text{mc}}$  ( $\alpha_{\text{mc}} > 1$ ; Fig. 3C). The fact that the experimental dependence is less sharp than predicted theoretically is attributed to the relatively large polydispersity of HA polymers (ranging from 0.5 to 3 MDa) used in the experiments,<sup>19</sup> which is not considered by the analytical model.

The above re-analysis of data from the literature demonstrates the tangible benefits of superselectivity concepts. It is well known that small vs. large HA can exert opposing functional effects,<sup>27</sup> but the underpinning mechanisms have long remained elusive. With the theoretical tool presented here, we can rationalise how HA molecules of different sizes bind and compete with each other for receptors. Moreover, we can predict how changes in the presentation of HA (e.g., the effective mean size and size dispersity, which may be modulated by degradation or by cross-linking with soluble HA binding proteins) and its receptors (e.g., their affinity, surface density and clustering) modulate HA binding and downstream physiological processes.



**Fig. 3. Effect of monovalent competitors.** (A) Illustrative example of the dependence of the selectivity parameter  $\alpha_{\text{R}}$  on the receptor surface density and competitor concentration (Eqs. [1-3, 5];  $n_{\text{L}} = 8$ ,  $c_{\text{P}}\alpha^3 N_{\text{A}} = 0.001$ ). (B) Schematic of HA binding to CD44 obtained from a crystal structure.<sup>24</sup> (C) Competition of HA polysaccharides ( $\text{HA}_{\text{poly}}$ ) with octasaccharides ( $\text{HA}_8$ ) binding CD44 monovalently: experimental data from Ref. 19 (blue symbols), analytical fit (blue line) and the competitor selectivity  $\alpha_{\text{mc}}$  (red line).

In conclusion, we have developed new mechanistic understanding of multivalent recognition with cofactors and competitors. Rather than modifying the multivalent probe itself, the addition of monovalent binders as competitors or cofactors is a simple, and thus attractive, avenue to modulate superselective binding. This effect can be exploited, for example, to tune the

threshold receptor density  $I^*$  of a given probe (Fig. 3A), to target surfaces with low receptor density,<sup>28</sup> and for ‘superselective’ discrimination of cofactor concentrations (Fig. 2D). Our theory thus helps designing superselective probes for targeting and analytical purposes controlled by cofactors and competitors. Whilst the simple multivalent model (Eqs. [1-2]) assumes each ligand can bind to many receptors, the scaling expressions (Eqs. [4-5]) are general: they expand on similar and well-known expressions for monovalent interactions,<sup>29</sup> and also apply to systems with few receptors and many ligands (see Supporting Information).

## ASSOCIATED CONTENT

The supporting information contains theoretical derivations, and details of the AnxA5-to-membrane binding experiments and binding model. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## AUTHOR INFORMATION

### Corresponding Authors

\* Tine Curk – [tcurk@jhu.edu](mailto:tcurk@jhu.edu); Ralf Richter – [r.richter@leeds.ac.uk](mailto:r.richter@leeds.ac.uk)

### Author Contributions

† These authors contributed equally.

### Funding Sources

Marie Skłodowska-Curie Fellowship ‘ELNANO’ (845032) to TC, Marie Curie Career Integration Grant ‘CELLMULTIVINT’ (293803) to GVD, and European Research Council Starting Grant ‘JELLY’ (306435) and Biotechnology and Biological Sciences Research Council grant BB/T001631/1 to RPR.

## ACKNOWLEDGMENT

We thank Céline Gounou for her expert assistance in producing the AnxA5 protein.

## REFERENCES

1. Mammen, M.; Choi, S.-K.; Whitesides, G. M., Polyvalent interactions in biological systems: Implications for design and use of multivalent ligands and inhibitors. *Angew. Chem. Int. Ed.* **1998**, *37*, 2754-94.
2. Dubacheva, G. V.; Curk, T.; Auzély-Velty, R.; Frenkel, D.; Richter, R. P., Designing multivalent probes for tunable superselective targeting. *Proc Natl Acad Sci U S A* **2015**, *112*, 5579-84.
3. Dubacheva, G. V.; Curk, T.; Moggetti, B. M.; Auzély-Velty, R.; Frenkel, D.; Richter, R. P., Superselective targeting using multivalent polymers. *J Am Chem Soc* **2014**, *136*, 1722-5.
4. Carlson, C. B.; Mowery, P.; Owen, R. M.; Dykhuizen, E. C.; Kiessling, L. L., Selective tumor cell targeting using low-affinity, multivalent interactions. *ACS Chem Biol* **2007**, *2*, 119-27.
5. Curk, T.; Dobnikar, J.; Frenkel, D., Design principles for super selectivity using multivalent interactions. In *Multivalency: Concepts, research & applications*, Haag, R.; Huskens, J.; Prins, L.; Ravoo, B. J., Eds. John Wiley & Sons: Oxford, 2018.
6. Martinez-Veracoechea, F. J.; Frenkel, D., Designing super selectivity in multivalent nano-particle binding. *Proc Natl Acad Sci U S A* **2011**, *108*, 10963-8.
7. Richter, R. P.; Lai Kee Him, J.; Tessier, B.; Tessier, C.; Brisson, A., On the kinetics of adsorption and two-dimensional self-assembly of annexin A5 on supported lipid bilayers. *Biophys J* **2005**, *89*, 3372-85.
8. Nores, G. A.; Dohi, T.; Taniguchi, M.; Hakomori, S., Density-dependent recognition of cell surface gm3 by a certain anti-melanoma antibody, and gm3 lactone as a possible immunogen: Requirements for tumor-associated antigen and immunogen. *J Immunol* **1987**, *139*, 3171-6.

9. English, N. M.; Lesley, J. F.; Hyman, R., Site-specific de-n-glycosylation of cd44 can activate hyaluronan binding, and cd44 activation states show distinct threshold densities for hyaluronan binding. *Cancer Res.* **1998**, *59*, 3736-42.
10. Lawrance, W.; Banerji, S.; Day, A. J.; Bhattacharjee, S.; Jackson, D. G., Binding of hyaluronan to the native lymphatic vessel endothelial receptor lyve-1 is critically dependent on receptor clustering and hyaluronan organization. *J Biol Chem* **2016**, *291*, 8014-30.
11. Vachieri, S. G.; Xiong, X.; Collins, P. J.; Walker, P. A.; Martin, S. R.; Haire, L. F.; Zhang, Y.; McCauley, J. W.; Gamblin, S. J.; Skehel, J. J., Receptor binding by h10 influenza viruses. *Nature* **2014**, *511*, 475-7.
12. Overeem, N. J.; Hamming, P. H. E.; Grant, O. C.; Di Iorio, D.; Tieke, M.; Bertolino, M. C.; Li, Z.; Vos, G.; de Vries, R. P.; Woods, R. J.; Tito, N. B.; Boons, G. P. H.; van der Vries, E.; Huskens, J., Hierarchical multivalent effects control influenza host specificity. *ACS Cent Sci* **2020**, *6*, 2311-8.
13. Overeem, N. J.; Hamming, P. H. E.; Tieke, M.; van der Vries, E.; Huskens, J., Multivalent affinity profiling: Direct visualization of the superselective binding of influenza viruses. *ACS Nano* **2021**, *15*, 8525-36.
14. Scheepers, M. R. W.; van, I. L. J.; Prins, M. W. J., Multivalent weak interactions enhance selectivity of interparticle binding. *Proc Natl Acad Sci U S A* **2020**, *117*, 22690-7.
15. Linne, C.; Visco, D.; Angioletti-Uberti, S.; Laan, L.; Kraft, D. J., Direct visualization of superselective colloid-surface binding mediated by multivalent interactions. *Proc Natl Acad Sci U S A* **2021**, *118*, e210603611.
16. Schroeder, A.; Heller, D. A.; Winslow, M. M.; Dahlman, J. E.; Pratt, G. W.; Langer, R.; Jacks, T.; Anderson, D. G., Treating metastatic cancer with nanotechnology. *Nat Rev Cancer* **2011**, *12*, 39-50.
17. Dubacheva, G. V.; Curk, T.; Frenkel, D.; Richter, R. P., Multivalent recognition at fluid surfaces: The interplay of receptor clustering and superselectivity. *J Am Chem Soc* **2019**, *141*, 2577-88.
18. Jeppesen, B.; Smith, C.; Gibson, D. F.; Tait, J. F., Entropic and enthalpic contributions to annexin v-membrane binding: A comprehensive quantitative model. *J Biol Chem* **2008**, *283*, 6126-35.
19. Lesley, J.; Hascall, V. C.; Tammi, M.; Hyman, R., Hyaluronan binding by cell surface cd44. *J Biol Chem* **2000**, *275*, 26967-75.
20. Ercolani, G.; Schiaffino, L., Allosteric, chelate, and interannular cooperativity: A mise au point. *Angew Chem Int Ed Engl* **2011**, *50*, 1762-8.
21. Huber, R.; Berendes, R.; Burger, A.; Schneider, M.; Karshikov, A.; Luecke, H.; Romisch, J.; Paques, E., Crystal and molecular structure of human annexin v after refinement. Implications for structure, membrane binding and ion channel formation of the annexin family of proteins. *J Mol Biol* **1992**, *223*, 683-704.
22. Bouter, A.; Gounou, C.; Berat, R.; Tan, S.; Gallois, B.; Granier, T.; d'Estaintot, B. L.; Poschl, E.; Brachvogel, B.; Brisson, A. R., Annexin-A5 assembled into two-dimensional arrays promotes cell membrane repair. *Nat Commun* **2011**, *2*, 270.
23. Perschl, A.; Lesley, J.; English, N.; Trowbridge, I.; Hyman, R., Role of cd44 cytoplasmic domain in hyaluronan binding. *Eur J Immunol* **1995**, *25*, 495-501.
24. Banerji, S.; Wright, A. J.; Noble, M.; Mahoney, D. J.; Campbell, I. D.; Day, A. J.; Jackson, D. G., Structures of the cd44-hyaluronan complex provide insight into a fundamental carbohydrate-protein interaction. *Nat Struct Mol Biol* **2007**, *14*, 234-9.
25. Takahashi, R.; Al-Assaf, S.; Williams, P. A.; Kubota, K.; Okamoto, A.; Nishinari, K., Asymmetrical-flow field-flow fractionation with on-line multiangle light scattering detection. 1. Application to wormlike chain analysis of weakly stiff polymer chains. *Biomacromolecules* **2003**, *4*, 404-9.
26. Banerji, S.; Hide, B. R. S.; James, J. R.; Noble, M. E. M.; Jackson, D. G., Distinctive properties of the hyaluronan binding domain in the lymphatic endothelial receptor lyve-1 and their implications for receptor function. *J Biol Chem* **2010**, *285*, 10724-35.
27. Cyphert, J. M.; Trempus, C. S.; Garantzios, S., Size matters: Molecular weight specificity of hyaluronan effects in cell biology. *Int J Cell Biol* **2015**, *2015*, 563818.
28. Tito, N. B., Multivalent "attacker and guard" strategy for targeting surfaces with low receptor density. *J Chem Phys* **2019**, *150*, 184907.
29. Berg, J. M.; Tymoczko, J. L.; Gatto Jr., G. J.; Stryer, L., *Biochemistry*. Macmillan Learning: 2019.

---

For Table of Contents only.

