



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/192341/>

Version: Published Version

Article:

Tai, F.W.D., Ching, H.L., Sloan, M. et al. (2022) Comparison of patient tolerance and acceptability of magnet-controlled capsule endoscopy and flexible endoscopy in the investigation of dyspepsia. *Endoscopy International Open*, 10 (06). E735-E744. ISSN: 2364-3722

<https://doi.org/10.1055/a-1790-5996>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Comparison of patient tolerance and acceptability of magnet-controlled capsule endoscopy and flexible endoscopy in the investigation of dyspepsia



Authors

Foong Way David Tai^{1,2}, Hey Long Ching^{1,2}, Marion Sloan³, Reena Sidhu^{1,2}, Mark McAlindon²

Institutions

- 1 Academic Department of Gastroenterology and Hepatology, Sheffield Teaching Hospitals, Sheffield, United Kingdom
- 2 Department of Infection, Immunity and Cardiovascular Disease, University of Sheffield, Sheffield, United Kingdom
- 3 Sloan Medical Centre, Sheffield, United Kingdom

submitted 27.5.2021

accepted after revision 3.1.2022

Bibliography

Endosc Int Open 2022; 10: E735–E744

DOI 10.1055/a-1790-5996

ISSN 2364-3722

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

Corresponding author

Dr. Foong Way David Tai, Academic Department of Gastroenterology and Hepatology, Royal Hallamshire Hospital, Glossop Road, S10 2JF Sheffield, United Kingdom
Fax: +441142711901
david.tai@nhs.net

ABSTRACT

Background and study aims Oropharyngeal intubation during Esophagogastroduodenoscopy (EGD) is uncomfortable, associated with aerosol generation and transmission of airborne microbes. Less-invasive alternatives may be better tolerated. In this study, patient tolerance and acceptability of EGD and transnasal endoscopy (TNE) have been compared with magnet-controlled capsule endoscopy (MACE).

Patients and methods A comparison of MACE with EGD and TNE in the investigation of dyspepsia was performed. Factors affecting patient tolerance and acceptability were examined using the Endoscopy Concerns Scale (ECS) and Universal Patient Centeredness Questionnaire (UPC-Q).

Results Patients were significantly more distressed (scoring least to most distress: 1–10) by gagging (6 vs 1), choking (5 vs 1), bloating (2 vs 1), instrumentation (4 vs 1), discomfort during (5 vs 1) and after (2 vs 1) EGD compared to MACE (all $P < 0.0001$). Patients were more distressed by instrumentation (5 vs 1) and discomfort during (5 vs 1) TNE compared to MACE ($P = 0.001$). Patients were more accepting of MACE than EGD and TNE with a UPC-Q score (scoring least to most acceptable: 0–100) lower for EGD (50 vs 98, $P < 0.0001$) and TNE (75 vs 88, $P = 0.007$) than MACE, and a post-procedure ECS score (scoring most to least acceptable: 10–100) higher for EGD (34 vs 11, $P < 0.0001$) and TNE (25 vs 10.5, $P = 0.001$) than MACE. MACE would be preferred by 83% and 64% of patients even if EGD or TNE respectively was subsequently recommended to obtain biopsies in half of examinations.

Conclusions Gagging and choking during instrumentation, the main causes of patient distress during EGD, occurred less during TNE but tolerance, acceptability and patient experience favored MACE.

Introduction

Esophagogastroduodenoscopy (EGD) is performed in 3% of Medicare beneficiaries per year [1]. It is uncomfortable and many patients require intravenous sedation or general anesthesia. Adverse events occur in up to 1 in 200 EGDs and 60% are

cardiopulmonary events related to sedation and intubation [2, 3]. It is, therefore, understandable that patients may delay seeking medical advice for symptoms due to fear of investigation [4]. The COVID-19 pandemic has prompted consideration of the risks to endoscopy staff of aerosol-generating proce-

dures [5]. These concerns underlie a continuing search for less invasive upper gastrointestinal investigative tools which are effective, safe and acceptable to patients.

Transnasal endoscopy (TNE) using ultrathin instruments is better tolerated than EGD [6]. Capsule endoscopy does not involve intubation and the capsule can be moved and rotated in a stream of swallowed water to achieve gastric visualization using external magnetic control. Studies suggest that patients may prefer magnet-controlled capsule endoscopy (MACE) to EGD [7–10].

Our understanding of patients' upper gastrointestinal endoscopy experiences is mostly limited to procedural tolerance [11] and no studies have yet been designed to primarily examine patient experience of EGD and TNE by comparison to MACE. In addition to comparing tolerance for EGD, TNE, and MACE, we have also compared the broader experience with each using two patient-reported experience measures (PREMs): the Endoscopy Concerns Scale (ECS) and the Universal Patient Centeredness Questionnaire (UPC-Q) [12, 13].

Patients and methods

Patients

Patients between 18 and 80 years of age referred to Sheffield Teaching Hospitals for the endoscopic investigation of dyspepsia were invited to participate [14]. Those expressing interest in participating were consecutively screened for eligibility. Contraindications to MACE included a history of dysphagia, Crohn's disease, small bowel resection or previous abdominopelvic irradiation, long term (over six months) daily consumption of a nonsteroidal anti-inflammatory drugs and implanted metallic devices.

Interventions

Patients were offered the choice of EGD (with or without sedation) or TNE and invited to have MACE in the 2 weeks preceding their flexible endoscopy. Procedures were described and patients given standardized hospital information leaflets. Participants who agreed to have both MACE and flexible endoscopy were included in the study.

Robot-controlled MACE was performed using the NaviCam (AnX Robotica Corp, Texas, United States). The system comprises two joysticks which control an external magnet suspended on a robot arm above the patient recumbent on an examination couch. All examinations were performed by the same endoscopist. Prior to the examination, patients ingested 80 mg simethicone (Infacol, Teva UK) in 100 mL of water, followed by a series of position changes to wash the stomach and then consumed a further 500 to 1000 mL of water to distend the stomach before swallowing the capsule endoscope (AKEM-11SW; AnX Robotica Corp, Texas, United States). The procedure controls and maneuvers are described in detail elsewhere [15].

Flexible endoscopy was performed within 2 weeks of MACE by accredited endoscopists. All patients had oropharyngeal topical anesthesia and were offered benzodiazepine (midazolam) sedation prior to EGD (Olympus H260 or H290 or Pentax EC34-i10F gastroscope) or 5% lidocaine/0.5% phenylephrine nasal

spray (Alliance Healthcare Ltd, UK) 15 minutes prior to TNE (Olympus XP290N). TNE was performed by one endoscopist. Patients were excluded from the study if either MACE or flexible endoscopy were incomplete; if there was food in the stomach, the MACE procedure time was less than 10 minutes, or nasal intubation was not possible.

Data collection and analysis

Knowledge of what patients anticipate in advance of their investigation is necessary to understand their health seeking behavior and compliance. Prior to each examination patients were asked to score their anxiety on a visual analog scale (1–10: not at all to extremely), as well as 13 endoscopy-related concerns previously described by Condon et al. in their development of the ECS, namely: telling friends about, fasting and discomfort prior to, the test; intravenous cannulation, instrumentation (insertion of flexible endoscope or swallowing the capsule), expressions of emotions, the endoscopist seeing food in the stomach and feelings of gagging, choking, vomiting, bloating, discomfort during the test [13]. The instrumentation criterion in the ECS was adapted for MACE by asking the patient to score their experience of swallowing the capsule rather than insertion of the flexible endoscope. Summation of each of these 13 scores was used as a measure of pre-procedure acceptance (13–130: most to least acceptable).

Measures of patient acceptance were collected after the procedure.

1. An ECS questionnaire scoring 10 of 13 items described earlier quantified patients' actual experience. The three pre-procedural items (concerns related to telling friends about, fasting and discomfort prior to, the test) were not repeated. These 10 items were added together to provide a measure of acceptability in light of their actual experience (post-procedure ECS: 10–100 most to least acceptable).
2. Patients were asked to consider three scenarios: whether or not they would undergo the test again or advise a friend to do so in similar medical circumstances or have the test as a screen for cancer in 5 years. Patients were regarded as finding the test acceptable if they answered in the affirmative to all three questions.

The UPC-Q was used to assess and compare patient global experience with each form of endoscopy [12]. Each patient was asked to identify three aspects of the overall pathway that were most important to them. They were asked to rate their experience with each aspect (1–5: poor to excellent). They were then asked to rank the level of importance of the three aspects relative to each other by allocating a total of six points between the three aspects. Aspects of the care pathway listed as important to patients were categorized according to subject matter and reported. The overall UPC-Q score (of 0–100: least to most acceptable) was obtained using the following equation (where A_1 , A_2 and A_3 are the three most important aspects of the pathway chosen by each patient):

$$\text{UPC-Q score} = [(\text{Rate } A_1 \times \text{Rank } A_1/6) + (\text{Rate } A_2 \times \text{Rank } A_2/6) + (\text{Rate } A_3 \times \text{Rank } A_3/6) - 1] \times 25$$

3. Patients were asked to express a preference for one or the other test. EGD could be recommended following MACE if biopsies were needed. Therefore, patients were also asked to express a preference for the primary diagnostic test based on the probability of requiring a second procedure to obtain biopsies.

In a subset of patients, tandem recordings of MACE and flexible endoscopy were reviewed by a blinded expert endoscopist. Endoscopic findings and agreement between modalities are reported.

Outcome measures

The primary outcomes were to examine overall acceptability and experience with MACE compared to EGD or TNE. These were defined as pre-procedure patient anxiety and ECS scores, and post-procedure as the UPC-Q and post-procedure ECS scores. The secondary (tolerance) outcomes were to quantify the level of distress caused by each item of the ECS (13 pre-procedure and 10 post-procedure items).


Advice was sought from the Statistical Services Unit at the University of Sheffield. A single point (10%) difference in patient distress scores between groups was considered to be clinically significant. A sample of 44 per group would have a 90% power to detect a difference in distress scores of 1 between MACE and flexible endoscopy assuming a standard deviation of 2 using a paired t-test with a 0.05 two-sided significance level. A sample of 48 patients would allow for an a priori interim analysis to be performed after 16 transnasal gastroscopies which would have a 90% power to detect a difference in mean distress score of 2 and satisfy the requirements of full power.

Statistical methods

IBM SPSS Statistics for Macintosh, Version 24.0.0 (Armonk, New York, United States: IBM Corp) was used for statistical analysis. Continuous data was represented as median and interquartile range (IQR) respectively as the distribution among the majority of patient questionnaire data were non-parametric (Shapiro-Wilk test $P > 0.05$). Non-parametric paired differences in central tendencies were examined using Wilcoxon signed rank tests. Unpaired differences were examined using Mann Whitney U or Kruskal-Wallis H test. Categorical data is presented as number and percentages: n (%), and McNemar test was used to compare paired dichotomous variables. Statistical significance is defined as $P < 0.05$ and significance levels of multiple comparisons for primary and secondary outcomes were subjected to the Bonferroni correction. Construct validity of the UPC-Q was examined by Pearson's correlation of convergent and discriminant factors.

Ethics

This study was approved and conducted in accordance with the ethical standards of the South Central – Berkshire B Research Ethics Committee (16/SC/0606. ClinicalTrials.gov NCT03420729), and the 1964 declaration of Helsinki and its later amendments.



► Fig. 1 Participants in the trial.

Results

Participants

Of 111 patients who expressed an interest, 69 participated and 60 were included in the study (► Fig. 1). Characteristics of included patients undergoing MACE and EGD or TNE are reported in ► Table 1. MACE was complete in 61 (88%), EGD in 47 (100%), and TNE in 21 patients (95%). Nine patients were excluded. One converted to EGD following failed nasal intubation and eight had incomplete gastric MACE (► Fig. 1). Eighteen patients (40.9%) opted for sedation with EGD (median midazolam dose 2 mg (range 1.5–4); fentanyl 50 mcg (range 25–75)).

Patient anxiety, anticipatory concerns and views on acceptability of their proposed investigations

Before the procedure, patients were more anxious about having EGD than MACE, with median state anxiety scores of 5 and 2 ($P < 0.0001$) respectively, but not significantly more anxious about having TNE than MACE (► Table 2). Trait anxiety did not differ significantly between EGD and TNE cohorts (► Table 1). Median pre-procedure ECS scores were higher before EGD (39 vs 26, $P < 0.0001$) and TNE (42 and 32, $P = 0.04$) than MACE (► Table 2). Median distress scores were significantly more in anticipation of EGD compared to MACE with regard to procedure-related

► Table 1 Characteristics of included patients undergoing MACE followed by EGD or TNE. Age, gender, Hospital Anxiety and Depression scores (HADS), use of anxiolytics or antidepressants, body mass index (BMI) and previous experience of endoscopy reported in median (IQR) or n (%).

	MACE followed by		P
	EGD	TNE	
N (%)	44	16	
Age	53.0 (22)	52.5 (23)	0.77
Female gender, n (%)	27 (66)	6 (55)	0.50
HADS			
Anxiety score	5 (8)	2 (6)	0.81
Score >8, n (%)	12 (29)	3 (27)	0.90
Anxiolytic use, n (%)	0 (0)	0 (0)	–
Depression score	1 (5)	0 (4)	0.10
Score >8, n (%)	3 (7.3)	0 (0)	0.36
Anti-depressant use, n (%)	8 (18)	3 (19)	0.81
BMI kg/m ²	24 (10)	23 (9)	0.40
Previous EGD experience, n (%)	18 (41)	2 (12.5)	0.08
Previous TNE experience, n (%)	0 (0)	0 (0)	–

factors (gagging (5 vs 2, $P < 0.0001$) and procedural discomfort (5 vs 2, $P < 0.0001$). By comparison, discussing the procedure with friends, fasting pre-procedure, potential for intravenous cannulation, displays of emotions and the endoscopist seeing food in the stomach caused negligible distress. No statistically significant differences were seen in anticipatory concerns prior to TNE and MACE after Bonferroni's correction. There was a trend suggesting TNE might cause more distress than MACE when patients anticipated instrumentation (7.5 vs. 2.5, $P = 0.008$) and procedural discomfort (6.5 vs. 3, $P = 0.005$) (► **Table 2**).

Patient tolerance and factors causing distress during EGD, TNE and MACE

After their procedure, patients reported experiencing significantly more distress (median score) due to gagging (6 vs 1, $P < 0.0001$), choking (5 vs 1, $P < 0.0001$), instrumentation (4 vs 1, $P < 0.0001$), discomfort during (5 vs 1, $P < 0.0001$) and after (2 vs 1, $P < 0.0001$) EGD when compared to MACE (► **Table 3**). Similarly, patients reported significantly more distress due to instrumentation (4.5 vs 1, $P = 0.001$) and discomfort during (5 vs 1, $P = 0.001$) TNE compared to MACE (► **Table 3**). In those undergoing EGD with sedation, MACE was still significantly better tolerated (► **Table 4**).

No significant differences were found between pre- and post-EGD or TNE distress scores, suggesting patients correctly anticipated which procedural factors would cause them distress (► **Table 5**). However, factors related to tolerance of MACE (gagging, choking, abdominal bloating, instrumenta-

tion, discomfort during and after MACE) caused significantly less actual distress than anticipated.

Patient acceptance of EGD, TNE and MACE and effect of mucosal biopsies on patient preference MACE and flexible endoscopies

The UPC-Q was completed appropriately by 95% of participants ($n = 57/60$). For the purpose of data analysis, drinking the water, swallowing the capsule, procedural discomfort, and test duration were all categorized as being related to procedural tolerance; having sufficient information and being informed about test progress as being related to communication and having a comfortable environment and adverse effects as being related to procedural recovery and aftercare. Aspects identified by patients for EGD, TNE, and MACE (respectively) related to procedural tolerance in 67%, 54%, and 54% of patients; communication in 4%, 18%, and 8%; aftercare and recovery in 7%, 9%, and 5%; test results in 13%, 14%, and 18%; test accuracy in 2%, 4%, and 5%, and other concerns in 8%, 0%, and 10%.

Median ECS scores were significantly higher after EGD (34 vs 11, $P < 0.0001$) and TNE (25 vs 10.5, $P = 0.001$) compared to MACE. UPC-Q score was lower for EGD (50 vs 98, $P < 0.0001$) and TNE (75 vs 88, $P = 0.007$) than MACE. In the subgroup of 40.9% of patients who opted to have sedation with their EGD, MACE had a higher UPC-Q (100 vs 54, $P = 0.003$) and lower post-procedure ECS score (10.5 vs. 25, $P < 0.0001$), suggesting a greater acceptance of MACE than sedated EGD (► **Table 4**). As defined by affirmative answers to all three questions regarding preparedness to undergo the same test again or to recommend the test, 100%, 94% and 64% of patients found MACE, TNE and EGD respectively acceptable. However, all patients preferred MACE to EGD and 94% preferred MACE to TNE.

If tissue biopsies were necessary after MACE and the chance of requiring biopsies (requiring flexible endoscopy as a second test) was 1 in 20, 1 in 10, 1 in 5, 1 in 4 or 1 in 2, 100%, 100%, 94%, 94% and 83% would prefer MACE followed by EGD, rather than a direct to EGD strategy and 94%, 94%, 81%, 75% and 63% would prefer MACE followed by TNE rather than a direct to TNE strategy.

Performance characteristics of UPC-Q and ECS

The UPC-Q correlated with post-procedure ECS after both MACE ($r = -0.32$ $P = 0.01$) and gastroscopy ($r = -0.40$ $P = 0.002$) demonstrating convergent validity, but not with pre-procedure ECS ($P = 0.49$ and $P = 0.29$) nor state anxiety scores ($P = 0.25$ and $P = 0.26$) providing some evidence of discriminant validity.

Endoscopic findings

Twenty-three patients with tandem recording of EGD and MACE were included in the analysis of endoscopic findings. There was complete agreement in findings between MACE and EGD in 65% of patients (15/23) (► **Table 6**). Eight patients (35%) had normal examinations on both MACE and EGD. Thirty endoscopic findings were found in total: 66% of findings (20/30) were seen by both MACE and EGD, 16.7% (5/30) by MACE alone, and 16.7% (5/30) by EGD alone.

► Table 2 Comparison of patient pre-procedure anticipation when undergoing EGD or TNE vs MACE. Pairwise comparison of median (IQR) pre-procedure anxiety (1 – 10: Least to most) and pre-procedure endoscopic concern scale (ECS) score (most to least acceptable: 13–130). Components of the pre-procedure ECS distress scores (1 – 10: Least to most) are compared for description. Only p values for clinically significant differences (≥ 1 point difference) are reported.

	EGD (n=44)			TNE (n=16)		
	EGD	MACE	p	TNE	MACE	p
Pre procedure anxiety	5 (5)	2 (2)	<0.0001 ¹	4.5 (5)	4 (4)	0.57
Pre procedure ECS	39 (41)	26 (7)	<0.0001 ¹	42 (25)	32 (19)	0.04
Telling friends/colleagues about test	1 (0)	1 (3)	–	1 (1)	1 (2)	–
Fasting	1 (1)	1 (2)	–	1.5 (3)	1.5 (3)	–
Discomfort prior to procedure	1 (1)	1 (2)	–	1 (1)	1 (2)	–
Gagging	5 (4)	2 (5)	<0.0001*	3 (5)	3.5 (6)	–
Choking	5 (2)	2 (5)	0.05	3 (5)	3.5 (6)	–
Bloating	2 (2)	2 (4)	–	1 (1)	2 (2)	0.13
Vomiting	4 (2)	1 (6)	0.19	1.5 (1)	1 (3)	–
Doctor seeing food in stomach	1 (0)	1 (0)	–	1 (0)	1 (0)	–
Displaying emotions during the test	1 (1)	1 (3)	–	1 (1)	1 (3)	–
Instrumentation	5 (4)	3 (5)	0.03	7.5 (4)	2.5 (3)	0.008
Intravenous catheter	1 (1)	1 (3)	–	1 (1)	1 (2)	–
Discomfort during procedure	5 (3)	2 (6)	<0.0001 ²	6.5 (3)	3 (4)	0.005
Discomfort after procedure	2 (2)	2 (4)	–	4 (3)	3 (5)	0.40

¹ For primary outcomes, P<0.01 considered as statistically significant after Bonferroni correction for 4 comparisons.

² For descriptive outcomes, P<0.002 considered as statistically significant after Bonferroni correction for 26 comparisons

► Table 3 Comparison of patient experience when undergoing EGD or TNE vs MACE. Pairwise comparison of median (IQR) universal patient centredness questionnaire (UPC-Q) score (least to most acceptable: 0–100) and post-procedure endoscopic concern scale (ECS) score (most to least acceptable: 10–100). Components of the post-procedure ECS distress scores (1 – 10: Least to most) are also compared for description. Only p values for clinically significant differences (≥ 1 point difference) are reported.

	EGD (n=44)			TNE (n=16)		
	EGD	MACE	p	TNE	MACE	p
UPC-Q	50 (50)	98 (25)	<0.0001 ¹	75 (67)	88 (37)	0.007 ¹
Post procedure ECS	34 (32)	11 (1)	<0.0001 ¹	25 (15)	10.5 (5)	0.001 ¹
Gagging	6 (6)	1 (0)	<0.0001 ²	1.5 (2)	1 (0)	–
Choking	5 (6)	1 (0)	<0.0001 ²	1.5 (2)	1 (0)	–
Bloating	2 (4)	1 (0)	0.08	1 (3)	1 (1)	–
Vomiting	1 (3)	1 (0)	–	1 (0)	1 (0)	–
Doctor seeing food in stomach	1 (0)	1 (0)	–	1 (0)	1 (0)	–
Displaying emotions during the test	1 (4)	1 (0)	–	1 (1)	1 (0)	–
Instrumentation	4 (7)	1 (1)	<0.0001 ²	4.5 (4)	1 (0)	0.001 ²
Intravenous catheter	1 (1)	1 (0)	–	1 (0)	1 (0)	–
Discomfort during procedure	5 (5)	1 (0)	<0.0001 ²	5 (5)	1 (0)	0.001 ²
Discomfort after procedure	2 (4)	1 (0)	<0.0001 ²	2 (3)	1 (0)	0.01

¹ For primary outcomes, P<0.01 considered as statistically significant after Bonferroni correction for 4 comparisons.

² For descriptive outcomes P<0.002 considered as statistically significant after Bonferroni correction for 20 comparisons.

► **Table 4 Comparison of patient experience of subgroup of patients undergoing EGD with and without sedation compared to MACE.** Unpaired comparison of median (IQR) universal patient centredness questionnaire (UPC-Q) score (least to most acceptable: 0–100) and post-procedure endoscopic concern scale (ECS) score (most to least acceptable: 10–100). Components of the post-procedure ECS distress scores (1–10: Least to most) are also compared for description. Only p values for clinically significant differences (≥ 1 point difference) are reported.

	EGD (n = 44)					
	Sedated (n = 18)			Unsedated (n = 26)		
	EGD	MACE	p	EGD	MACE	p
UPC-Q	54 (52)	100 (19)	0.003 ¹	46 (50)	92 (25)	0.001 ¹
Post procedure ECS	25 (26)	10.5 (1.5)	<0.0001 ¹	34 (36)	11 (2)	<0.0001 ¹
Gagging	6 (6.3)	1 (0)	0.001 ²	7 (5)	1 (0)	<0.0001 ²
Choking	4 (5.3)	1 (0)	0.001 ²	5 (5)	1 (0)	<0.0001 ²
Bloating	1 (2)	1 (0)	–	4 (5)	1 (0)	0.001 ²
Vomiting	1 (1.5)	1 (0)	–	1 (4)	1 (0)	–
Doctor seeing food in stomach	1 (0)	1 (0)	–	1 (0)	1 (0)	–
Displaying emotions during the test	1 (4.3)	1 (0)	–	2 (4)	1 (0)	0.001 ²
Instrumentation	3 (4.5)	1 (1)	0.008	5 (7)	1 (1)	<0.0001 ²
Intravenous catheter	2 (3)	1 (0)	0.005	1 (0)	1 (0)	–
Discomfort during procedure	4 (6.3)	1 (0)	0.001 ²	6 (5)	1 (0)	<0.0001 ²
Discomfort after procedure	2 (3.3)	1 (0)	0.005	3 (4)	1 (0)	<0.0001 ²

¹ For primary outcomes, $P < 0.01$ considered as statistically significant after Bonferroni correction for 4 comparisons

² For descriptive outcomes $P < 0.002$ considered as statistically significant after Bonferroni correction for 20 comparisons.

► **Table 5 Comparison of pre-procedural anticipation and patient experience undergoing EGD, TNE and MACE.** Paired comparison of median (IQR) scores of distress cause by anticipation (before) and actual (after) endoscopy causing distress for each endoscopic modality (1–10: Least to most)

	EGD (n = 44)			TNE (n = 16)			MACE (n = 60)		
	Before	After	p	Before	After	p	Before	After	p
Gagging	5 (5)	6 (6)	0.22	3 (5.5)	1.5 (2)	0.10	3 (4)	1 (0)	<0.0001 ¹
Choking	5 (5)	5 (5.5)	–	3 (5.5)	1.5 (2)	0.05	2 (3)	1 (0)	<0.0001 ¹
Bloating	2 (4)	2 (3.5)	–	1 (1.8)	1 (2.8)	–	2 (2)	1 (1)	<0.0001 ¹
Vomiting	4 (6)	1 (3)	0.08	1.5 (2.8)	1 (0)	–	1 (2)	1 (0)	–
Doctor seeing food in stomach	1 (0)	1 (0)	–	1 (0)	1 (0)	–	1 (0)	1 (0)	–
Displaying emotions during the test	1 (2.5)	1 (4)	–	1 (2.8)	1 (1)	–	1 (1)	1 (0)	–
Instrumentation	5 (5)	4 (6.5)	0.16	7.5 (3)	4.5 (4)	0.04	3 (4)	1 (1)	<0.0001 ¹
Intravenous catheter	1 (3)	1 (1)	–	1 (1.8)	1 (0)	–	1 (1)	1 (0)	–
Discomfort during procedure	5 (6)	5 (5)	–	6.5 (3.8)	5 (4.5)	0.14	3 (2.5)	1 (0)	<0.0001 ¹
Discomfort after procedure	2 (4)	2 (4)	–	4 (3.8)	2 (2.8)	0.09	2 (3)	1 (0)	<0.0001 ¹

¹ For descriptive outcomes $P < 0.004$ considered as statistically significant after Bonferroni correction for 10 comparisons.

Discussion

The anticipation of EGD caused more distress in comparison to MACE due to concerns related to procedural tolerance (intubation and discomfort during EGD) and these were matched by

actual patient experience. This may explain why patients were much more anxious before EGD than MACE. Patients favored MACE over both EGD and TNE by some margin and by most tolerance measures, by acceptability, overall experience, and preference. The majority of patients would prefer MACE initially to

► **Table 6** Comparison of endoscopic findings of MACE and EGD

Case	MACE and EGD	MACE only	EGD only
1	Normal		
2		Fundal polyp	
3	Normal		
4	Antral angioectasia	Antral erosion	
5		(antral bulge) ¹	Esophagitis
6	Normal		
7	Normal		
8	2 prepyloric erosions		
9	D2 angioectasia		
10	Antral gastritis ²	Prepyloric erosion	
11	Antral erosion and a fundal polyp		
12			Pyloric erosion
13	Normal		
14	Normal		
15	Normal		
16	Oesophagitis ² and Fundal polyp		
17	Duodentitis		
18	Antral erosions		Duodentitis
19	Normal		
20	Esophagitis ²	Fundal polyp	Hiatus hernia, duodenitis
21	Fundal polyp x2, Antral gastritis		
22	Esophagitis ² , cluster of polyps on lower posterior body of stomach		
23	Antral gastritis, multiple > 20 flat polyps on body of stomach, hiatus hernia	Linear ulcer along lesser curve	

¹ Antral bulge not seen on EGD

² Not seen on live MACE examination, but seen on retrospective review of capsule video.

either EGD or TNE, even if there were a 50% chance of requiring further visits (and pre-procedure preparation) for a flexible endoscopy as a second test to obtain biopsies.

The aerosols generated by gagging and choking increase the risk of transmitting COVID-19 to endoscopy staff and patients [5]. This study suggests that compared to conventional EGD, risks could be mitigated by use of TNE, which induced less of an oropharyngeal reflex and consequent gagging and choking, or MACE, during which such a reflex was virtually absent. There are currently no published studies of aerosol generation in transnasal gastrointestinal endoscopy. A systematic review of nasal endoscopy report mixed findings with regard to aerosol generation during nasal procedures, but concluded that nasal anesthetics and decongestants are significantly aerosol-generating above baseline [16]. Furthermore, it was recently shown that joystick-controlled MACE could be performed on a patient in a separate room with audiovisual links to the endoscopist,

control station, and monitor, therefore eliminating the need for personal protective equipment [17].

EGD is the accepted gold standard in upper gastrointestinal investigation and TNE has comparable diagnostic accuracy in the esophagus [18, 19]. However, about 10% of early cancers are missed at initial EGD. [20] Proximal lesions identified during push enteroscopy [21] and capsule endoscopy [22] are presumed to have been missed by prior EGD. Recent studies of MACE using a handheld magnet suggest at least diagnostic equivalence with EGD (although both modalities missed pathology) [9, 10]. Moreover, in a 350 multicentre Chinese study the sensitivity, specificity, positive and negative predictive values of MACE compared to esophagogastroduodenoscopy (EGD) were 90.4% (84.7–96.1), 94.7% (91.9–97.5), 87.9% (81.7–94.0), and 95.9% (93.4–98.4), irrespective of site and size of focal gastric lesion [8]. MACE has been reported as taking between 14 and 26 minutes [15]. It has been shown that EGD last-

ing over 7 minutes has a threefold diagnostic yield of early cancer [23]. The relative benefit, therefore, of more responsive instrument control offered by EGD compared to a longer examination allowed by less-invasive or noninvasive tests needs further investigation.

Along with clinical effectiveness, patient experience is a further pillar in quality of care [24]. Studies of patient experience with endoscopy have focused on procedural tolerance and satisfaction [11,24]. However, this study shows that over one-third of the concerns volunteered by patients were unrelated to their procedural tolerance. That test results are important to patients is unsurprising, but many also expressed an interest in the overall investigative pathway, the detail (sensations experienced) and duration of the test. Patients valued a comfortable environment during recovery and information about potential adverse effects. A measure of satisfaction is unidimensional and conveys patient overall contentedness with their experience, but in contrast to PREMs, does not encompass all aspects of care nor discriminate which aspects are important [25].

Patient experience may affect compliance with investigation and participation in screening programs and services responsive to feedback can improve patient outcomes [25,26]. Therefore, we chose to use two PREMs, the ECS and UPC-Q. The ECS comprised aspects of patient concerns before, during, and after endoscopy, and the score derived was shown to demonstrate good construct validity and to correlate with patient acceptance of EGD [13]. The UPC-Q is a patient-generated index based on each individual's concerns, priorities, and experiences and serves to examine patient acceptability of a healthcare experience beyond the constraints set by an endoscopy paradigm. It performs reliably and correlates well against known measures of patient satisfaction in other inpatient and outpatient settings [12]. That it correlates with the ECS, an experience score designed for, and validated in, endoscopic practice, suggests that it could be used as a patient-related experience measure in this setting.

Our results are consistent with previous studies of tolerance and preference for capsule endoscopy over EGD [7–10,27,28] and TNE [29]. Patients are thought to formulate a notion of acceptability by comparing their expectation with actual experience [30]. Our patients accurately anticipated the unpleasant aspects of EGD, yet only 64% of patients regarded it as acceptable and it performed comparatively poorly in the UPC-Q. This contrasts with the 95% acceptability rate for EGD identified in the study by Condon et al [13]. However, acceptability is defined by the context in which it is assessed, and the experience of a less-invasive or noninvasive alternative in this study is likely to have adversely affected the acceptability and tolerability of conventional EGD, especially when patients had better MACE experience than they anticipated. This may, in part, also explain why transnasal endoscopy in this study may be more poorly tolerated than reported in previous studies in which patients were randomized to only one intervention. Finally, acceptability might also be affected by the cost of the investigative procedure in fee-for-service health care systems: a cost analysis of the different modalities is needed.

Conscious sedation for EGD had a disappointing impact. However, systematic review and meta-analysis showed that compared to no sedation, use of midazolam alone only improves patient satisfaction and willingness to repeat the procedure (not any aspect of tolerance nor overall experience) [31] at mean doses of 4.8 to 10.3 mg, far greater than would be used in current practice. That patients may not always feel “conscious” sedation is adequate may explain the move toward anesthetist-directed sedation using propofol [32]. Immediate serious adverse events of EGD are rare, occurring in between 0.32–0.39% of a recently described series of 1.38 million procedures [32]. Wang et al, however, showed that 1% of patients were treated for infection within 30 days of EGD [33] and therefore complications of EGD and sedation may be delayed and not always recognized.

The primary aim of the study was to compare patient experience with the three modalities. Patients were eligible if they needed investigation for dyspepsia, a cohort known to have a low yield of pathology [34]. Consistent with other studies comparing MACE and flexible endoscopy in recurrent iron deficiency [9] and upper gastrointestinal bleeding [10], both MACE and EGD miss lesions. No major adverse events have been reported following MACE, although experience with this technology is still limited. In the largest series of MACE procedures reported (n=3182), there were no reported cases of capsule retention [15]. Capsule aspiration may occur in 0.1% of patients at most, usually in men over 80 years of age. Devices were spontaneously expectorated by half the patients, the remainder needed bronchoscopic removal and no fatalities were reported [35]. Regarding TNE, minor epistaxis affects 5% of patients, but adverse cardiopulmonary events are significantly less than with EGD [36,37].

The ability to obtain biopsies remains an important advantage of flexible endoscopy. However, we previously found that while 84% of 500 patients having EGD to investigate dyspepsia had biopsies taken, they contributed to management in only 16% beyond empirical treatment with proton pump inhibitors or *Helicobacter pylori* “test and treat” strategies [38]. Even with a 50% chance that a second procedure would be required to obtain biopsies, most patients still preferred to have MACE as the initial investigation.

Conclusions

MACE is better tolerated and more acceptable than EGD and TNE. Although procedural tolerance was the major contributor to test acceptability, other factors related to the care pathway were identified by patients in one-third of cases. Patients expressed a preference for MACE followed by fiberoptic gastroscopy if biopsies were required.

Acknowledgments

The authors thank Øyvind Bjertnaes for his assistance in providing us with the Universal Patient Centeredness Questionnaire.

Competing interests

This study has been supported by AnX Robotica Corp (Texas, United States) in the provision of capsule endoscopes and travel expenses for Drs. Tai, Ching, and McAlindon to receive training in the use of the system; and the Dawkins and Strutt British Medical Association grant for salary support of Dr. Tai.

Funding

BMA Foundation for Medical Research

Clinical trial

Sheffield Teaching Hospitals NHS Foundation Trust
NCT03420729

TRIAL REGISTRATION: Interventional, parallel assignment, non-randomized, prospective at <https://www.clinicaltrials.gov/ct2/show/NCT03420729>

References

- [1] Pohl H, Robertson D, Welch HG. Repeated upper endoscopy in the Medicare population: a retrospective analysis. *Ann Intern Med* 2014; 160: 154
- [2] Ben-Menachem T, Decker GA, Early DS et al. Adverse events of upper gastrointestinal endoscopy. *Gastrointest Endosc* 2012; 76: 707–718
- [3] Levy I, Gralnek IM. Complications of diagnostic colonoscopy, upper endoscopy, and enteroscopy. *Best Pract Res Clin Gastroenterol* 2016; 30: 705–718
- [4] Macdonald S, Macleod U, Campbell NC et al. Systematic review of factors influencing patient and practitioner delay in diagnosis of upper gastrointestinal cancer. *In Br J Cancer* 2006: 1272–1280
- [5] Gralnek IM, Hassan C, Beilenhoff U et al. ESGE and ESGENA Position Statement on gastrointestinal endoscopy and the COVID-19 pandemic. *Endoscopy* 2020; 52: 483–490
- [6] Sami SS, Subramanian V, Ortiz-Fernández-Sordo J et al. Performance characteristics of unsedated ultrathin video endoscopy in the assessment of the upper gastrointestinal tract: systematic review and meta-analysis. *Gastrointest Endosc* 2015; 82: 782–792
- [7] Denzer UW, Rösch T, Hoytat B et al. Magnetically guided capsule versus conventional gastroscopy for upper abdominal complaints: a prospective blinded study. *J Clin Gastroenterol* 2015; 49: 101–107
- [8] Liao Z, Hou X, Lin-Hu EQ et al. Accuracy of magnetically controlled capsule endoscopy, compared with conventional gastroscopy, in detection of gastric diseases. *Clin Gastroenterol Hepatol* 2016; 14: 1266–1273.e1261
- [9] Ching HL, Hale MF, Kurien M et al. Diagnostic yield of magnetically assisted capsule endoscopy versus gastroscopy in recurrent and refractory iron deficiency anemia. *Endoscopy* 2018; 51: 409–418
- [10] Ching HL, Hale MF, Sidhu R et al. Magnetically assisted capsule endoscopy in suspected acute upper gastrointestinal bleeding versus esophagogastroduodenoscopy in detecting focal lesions. *Gastrointest Endosc* 2019; 90: 430–439
- [11] Brown S, Bevan R, Rubin G et al. Patient-derived measures of gastrointestinal endoscopy: a meta-narrative review of the literature. *Gastrointest Endosc* 2015; 81: 1130–1140.e1131–1139
- [12] Bjertnaes O, Iversen HH, Holmboe O et al. The Universal Patient Centeredness Questionnaire: reliability and validity of a one-page questionnaire following surveys in three patient populations. *Patient Relat Outcome Meas* 2016; 7: 55–62
- [13] Condon A, Graff L, Elliot L et al. Acceptance of colonoscopy requires more than test tolerance. *Can J Gastroenterol* 2008; 22: 41–47
- [14] NICE. Gastro-oesophageal reflux disease and dyspepsia in adults: investigation and management Clinical guideline [CG184]. 2014.
- [15] Zhao AJ, Qian YY, Sun H et al. Screening for gastric cancer with magnetically controlled capsule gastroscopy in asymptomatic individuals. *Gastrointest Endosc* 2018; 88: 466–474.e1
- [16] Matos S, Sharma A, Crosby D. Objective assessment of aerosolization during transnasal endoscopy: a systematic review. *Otolaryngol Head Neck Surg* 2021: doi:10.1177/01945998211050632
- [17] Pan J, Li Z, Liao Z. Noncontact endoscopy for infection-free gastric examination during the COVID-19 pandemic. *VideogastrointestinalE* 2020; 5: 402–403.e1
- [18] Shariff MK, Bird-Lieberman EL, O'Donovan M et al. Randomized crossover study comparing efficacy of transnasal endoscopy with that of standard endoscopy to detect Barrett's esophagus. *Gastrointest Endosc* 2012; 75: 954–961
- [19] Choe WH, Kim JH, Ko SY et al. Comparison of transnasal small-caliber vs. peroral conventional esophagogastroduodenoscopy for evaluating varices in unsedated cirrhotic patients. *Endoscopy* 2011; 43: 649–656
- [20] Menon S, Trudgill N. How commonly is upper gastrointestinal cancer missed at endoscopy? A meta-analysis. *Endosc Int Open* 2014; 2: E46–E50
- [21] Zaman A, Katon RM. Push enteroscopy for obscure gastrointestinal bleeding yields a high incidence of proximal lesions within reach of a standard endoscope. *Gastrointest Endosc* 1998; 47: 372–376
- [22] Sidhu R, Sanders DS, Kapur K et al. Capsule endoscopy changes patient management in routine clinical practice. *Dig Dis Sci* 2007; 52: 1382–1386
- [23] Teh JL, Tan JR, Lau LJ et al. Longer examination time improves detection of gastric cancer during diagnostic upper gastrointestinal endoscopy. *Clin Gastroenterol Hepatol* 2015; 13: 480–487.e482
- [24] Neilson LJ, Patterson J, Wagner Cv et al. Patient experience of gastrointestinal endoscopy: informing the development of the Newcastle ENDOPREM™. *Frontline Gastroenterol* 2020; 11: 209–217
- [25] Black N, Jenkinson C. Measuring patients' experiences and outcomes. *BMJ* 2009; 339: b2495
- [26] Tierney M, Bevan R, Rees CJ et al. What do patients want from their endoscopy experience? The importance of measuring and understanding patient attitudes to their care *Frontline Gastroenterol* 2016; 7: 191–198
- [27] Rey JF, Ogata H, Hosoe N et al. Blinded nonrandomized comparative study of gastric examination with a magnetically guided capsule endoscope and standard videoendoscope. *Gastrointest Endosc* 2012; 75: 373–381
- [28] Beg S, Card T, Warburton S et al. Diagnosis of Barrett's esophagus and esophageal varices using a magnetically assisted capsule endoscopy system. *Gastrointest Endosc* 2019; 4: 773–781.e1
- [29] Chak A, Alashkar BM, Isenberg GA et al. Comparative acceptability of transnasal esophagoscopy and esophageal capsule esophagoscopy: a randomized, controlled trial in veterans. *Gastrointest Endosc* 2014; 80: 774–782
- [30] Kravitz RL. Measuring patients' expectations and requests. *Ann Intern Med* 2001; 134: 881–888
- [31] McQuaid KR, Laine L. A systematic review and meta-analysis of randomized, controlled trials of moderate sedation for routine endoscopic procedures. *Gastrointest Endosc* 2008; 67: 910–923

- [32] Vargo JJ, Niklewski PJ, Williams JL et al. Patient safety during sedation by anesthesia professionals during routine upper endoscopy and colonoscopy: an analysis of 1.38 million procedures. *Gastrointest Endosc* 2017; 85: 101–108
- [33] Wang P, Xu T, Ngamruengphong S et al. Rates of infection after colonoscopy and esophagogastroduodenoscopy in ambulatory surgery centres in the USA. *Gut* 2018; 67: 1626–1636
- [34] Ford AC, Marwaha A, Lim A et al. What is the prevalence of clinically significant endoscopic findings in subjects with dyspepsia? Systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2010; 8: 830–837, 837.e831–832
- [35] Yung DE, Plevris JN, Koulaouzidis A. Short article: Aspiration of capsule endoscopes: a comprehensive review of the existing literature. *Eur J Gastroenterol Hepatol* 2017; 29: 428–434
- [36] Tanuma T, Morita Y, Doyama H. Current status of transnasal endoscopy worldwide using ultrathin videoscope for upper gastrointestinal tract. *Dig Endosc* 2016; 28: 25–31
- [37] Ai ZL, Lan CH, Fan LL et al. Unsedated transnasal upper gastrointestinal endoscopy has favorable diagnostic effectiveness, cardiopulmonary safety, and patient satisfaction compared with conventional or sedated endoscopy. *Surg Endosc* 2012; 26: 3565–3572
- [38] Ching H-L, Hale MF, Sidhu R et al. Reassessing the value of gastroscopy for the investigation of dyspepsia. *Frontline Gastroenterol* 2018; 9: 62–66