

This is a repository copy of *Methods of assessing value for money of UK-based early childhood public health interventions:A systematic literature review*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/193357/>

Version: Published Version

Article:

Murphy, Peter James, Hinde, Sebastian orcid.org/0000-0002-7117-4142, Fulbright, Helen Athena et al. (2 more authors) (2022) Methods of assessing value for money of UK-based early childhood public health interventions:A systematic literature review. British Medical Bulletin. Idac035. ISSN: 1471-8391

<https://doi.org/10.1093/bmb/ldac035>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Invited Review

Methods of assessing value for money of UK-based early childhood public health interventions: a systematic literature review

Peter Murphy^{1,*}, Sebastian Hinde¹, Helen Fulbright², Louise Padgett³, and Gerry Richardson¹

¹Centre for Health Economics, University of York, York, YO10 5DD, UK, ²Centre for Reviews and Dissemination, University of York, York, YO10 5DD, UK, and ³Department of Health Sciences, University of York, York, YO10 5DD, UK

*Correspondence address. Centre for Health Economics, University of York, York YO10 5DD, UK.

E-mail: peter.murphy@york.ac.uk

Received 14 June 2022; Revised 25 October 2022; Accepted 11 November 2022

Abstract

Introduction: Economic evaluation has an important role to play in the demonstration of value for money of early childhood public health interventions; however, concerns have been raised regarding their consistent application and relevance to commissioners. This systematic review of the literature therefore aims to collate the breadth of the existing economic evaluation evidence of these interventions and to identify the approaches adopted in the assessment of value.

Source of data: Recently published literature in Medline, EMBASE, Econ-Lit, Health Management Information Consortium, Cochrane CENTRAL, Cochrane Database of Systematic Reviews, Health Technology Assessment, NHS EED and Web of Science.

Areas of agreement: The importance of the early childhood period on future health and well-being as well as the potential to impact health inequalities making for a strong narrative case for expenditure in early childhood public health.

Areas of controversy: The most appropriate approaches to evaluating value for money of such preventative interventions relevant for UK decision-makers given the evident challenges.

Areas timely for developing research: Future resource allocations decisions regarding early childhood public health interventions may benefit from consistency in the evaluative frameworks and health outcomes captured, as well as consistency in approaches to incorporating non-health costs and outcomes, incorporating equity concerns and the use of appropriate time horizons.

Key words: economic evaluation, infant, child, public health

The importance of the early childhood period on future health and well-being is well established.^{1,2} Interventions in early life not only have the potential to impact diseases in adult life but also to impact health inequalities experienced throughout the life course.^{3–7} Policy recommendations in the UK and elsewhere have therefore reiterated the importance of increasing public or government expenditure by local and national decision-makers on those in early childhood^{6,8} as well as more broadly acknowledging the value of investing in preventative interventions.⁹

Despite the presence of a strong narrative case for prevention in early childhood, it is important that decisions to fund such interventions are based on systematic and robust assessments of clinical and economic evidence. Resources are limited and decisions to fund an intervention means the opportunity to fund alternative interventions are foregone. Economic evaluation provides a systematic and transparent framework to identify which interventions offer value for money and help inform the choice of competing claims on limited resources.¹⁰

population level, there remains less agreement on the most appropriate methods for conducting economic evaluation.^{12,13} This is particularly true for public health interventions targeting early childhood given the complex interplay between health, development, education, socioeconomic status and the family environment.^{14–16}

The range of methodological challenges of conducting economic evaluations of such early childhood interventions has been highlighted in the literature and includes: appropriate time horizons, measuring and valuing health outcomes, incorporating non-health costs and outcomes, and informing health equity concerns.^{17–19} These challenges may in-part explain why a number of large-scale early childhood preventative interventions have failed to show cost-effectiveness¹³ and why gaps exist between the evidence base and decision-making.²⁰

Previous literature reviews in a paediatric setting have focussed on economic evaluations of specific intervention categories, such as vaccinations,²¹ parenting interventions,^{22,23} health promotion,^{24–26} and oral health.²⁷ Furthermore, these reviews have not limited the results to a specific country or jurisdiction. We consider it important to identify the economic evidence relevant to UK public health decision-makers given the positive and normative reasons why results of economic evaluations may differ across jurisdictions.²⁸ A literature review of

public health intervention decisions made by NICE²⁹ focussed on those conducted in a UK context yet there was no limitation by age category and the review focussed on NICE guidance and not the wider evidence base.

This systematic review of the literature therefore aims to achieve a number of goals. First, to collate the breadth of the existing economic evaluation evidence of early childhood public health interventions conducted in a UK context. Second, to describe the methods and approaches adopted in the evidence base to highlight consistencies in the demonstration of value for money. Finally, to critically appraise the quality of the evidence base. By doing so, this review seeks to provide researchers and policymakers in the UK details of the relevant economic evaluation evidence, as well as highlighting the methodological challenges and deficiencies in conducting such analyses.

Methods

The protocol was registered with PROSPERO (CRD42021270751) and was conducted and reported in accordance with PRISMA guidelines.³⁰

Data sources and searches

An initial search strategy was designed in Ovid MEDLINE with the final strategy adapted with relevant subject headings (controlled vocabularies) and search syntax to each of the databases listed below. No language limits were applied, but papers were limited to 2000 onwards to ensure the relevance to the current research and policy deliberations. Details of the full search strategies are contained in Supplementary Material, [Supplementary Appendix 1](#).

The following databases were searched between August 16 and 23, 2021:

1. MEDLINE(R) ALL (Ovid): 1946 to August 13, 2021.
2. Embase (Ovid): 1974 to August 17, 2021.
3. Econlit (Ovid): 1886 to August 5, 2021.
4. Health Management Information Consortium (HMIC) (Ovid): 1979 to July 2021.

5. Cochrane Central Register of Controlled Trials (Wiley): 2021, Issue 8 in the Cochrane Library.
6. Cochrane Database of Systematic Reviews (Wiley): 2021, Issue 8 in the Cochrane Library.
7. Health Technology Assessment (CRD): Inception to March 2018.
8. Economic Evaluations Database (CRD): Inception to March 31, 2015;
9. Science Citation Index Expanded (Web of Science): 1900 to August 16, 2021.

Supplementary searches in the form of reference checking and backwards citation searching³¹ of systematic reviews identified in the primary database search were undertaken on October 15, 2021.

Study selection

Two review authors (P.M. and L.P.) independently conducted title and abstract screening of a random sample of 10% of the retrieved records. A kappa statistic for assessing inter-rater agreement³² was calculated. Upon the achievement of a kappa statistic of 0.8 or above, one reviewer screened the remaining titles and abstracts. Failure to achieve the required kappa statistic meant a further 10% would be screened by both reviewers until the required score was achieved. Reviewers screened 20% (two screening rounds) before the sufficient kappa statistic was achieved. This process was applied at both the title and abstract screening stage and the full text screening stage. Any discrepancies were resolved by discussion between the two review authors.

Records were included if they reported economic evaluations of public health interventions in the UK. Public health was defined using terms that broadly reflected interventions of health improvement and included terms for wider social determinants of health (Supplementary Material, [Supplementary Appendix 1](#)). Evaluations were limited to those of interventions for infants and children with a mean age of 5 years or under at baseline to reflect the infant, toddler and preschool years. Economic evaluations of interventions for ages above 5 years of age were considered for inclusion if

Sources of grey literature amongst the search results were included in Ovid's HMIC database, which includes literature on health management, health service policies, public health and social care with an emphasis on the UK and the NHS. Grey literature that was found through supplementary searches of previous systematic reviews was deemed eligible for inclusion.

Critical appraisal of the included studies was conducted through the use of the CHEERS checklist,¹⁰ which can be found in Supplementary Material [Supplementary Appendix 3](#).

Health protection programmes were the most common intervention category (32%; 22/69). The remaining interventions were newborn screening




Fig. 1 PRISMA diagram of the flow of included and excluded studies.

(16%; 11/69), parenting support (12%; 8/69), injury prevention (7%; 5/69), health promotion (4%; 3/69), oral health (9%; 6/69), childhood screening (9%; 6/69), breast feeding (6%; 4/69), reducing the risk of maltreatment (3%; 2/69) and finally, interventions that cover both parenting support and health promotion (3%; 2/69). The general characteristics of the identified evaluations can be seen in Table 1.

A little under half of the evaluations were QALY-based CEAs (46%; 32/69). The majority of evaluations were non-QALY-based CEAs (49%; 34/69), with the health outcomes used including life years gained/saved (29%; 10/34), oral health outcomes such as dental caries detected or number of teeth free

from decay (15%, 5/34) and cases of a specific disease or condition detected (12%, 4/34). See Table 2 for the list of outcomes. One evaluation (1%) was a CCA¹⁰⁴ and one (1%) a CCA alongside a CEA.⁹⁷ Of the studies reporting outcomes in monetary terms, the evaluations identified were SROI (4%; 3/69), CBA (3%; 2/69) with one of the CBAs being conducted alongside a CEA. Finally, four (4%) were not explicit about the type of evaluation used; however, detailed inspecting suggested two of them could be classified as CCA^{35,42,98} and two as ROI analysis.^{76,96}

The most commonly reported perspective was the NHS or NHS and personal social services (PSS) (55%, 38/69), which is consistent with the latest NICE methods guidance.¹¹ A considerable

Table 1 Summary of the evaluations

Author year	Intervention category	Intervention (comparator)	Population	Type of evaluation	Study type	Time horizon	Outcomes captured (quality of life instrument)	Primary result of the evaluation
Anokye 2020 ⁴⁰	Breast feeding	Nourishing Start for Health, NOSH (usual care)	Newborn	Non-QALY-based CEA	Trial based	1 year	Proportion baby breast fed at 6 weeks	£974 per additional breast-fed baby
Hoddinott 2012 ⁴¹	Breast feeding	FEeding Support Team, FEST (reactive telephone support)	Newborn	Non-QALY-based CEA	Trial based	6–8 weeks	Any breastfeeding; exclusive breastfeeding	£87 per additional woman any breastfeeding; £91 per additional woman exclusively breast feeding
Jacklin 2007 ³⁸ NICE 2008 ³⁹	Breast feeding	Breast feeding peer support (unclear)	Newborn	QALY-based CEA	Model based	Unclear	QALYs (unclear); premenopausal breast cancer averted; infant infections averted	No ICER reported
Pokhrel 2015 ⁴²	Breast feeding	Breast feeding support (no breast feeding support)	Newborn	CCA (not specified but reports costs and outcomes separately)	Model based	1 year for three acute conditions (GI, LRTI and AOM); lifetime for maternal BC; neonatal unit stay for NEC	Cost savings. Includes a cost derived using NMB assuming 20 000/QALYs for the breast cancer benefits.	Report outcomes using three different types of policies: Policies A, B and C (impacts on acute diseases (GI, LRTI and AOM)); Policy D (impacts NEC) and Policy E (impacts BC). Policy A2 saves £11.04 m; policy D2 saves £6.12 m and policy E2 saves £31.42 m (this includes QALYs gained)
Bamford 2007 ⁴³	Childhood screening	Alternative SES programmes (no SES)	4–5 years	QALY-based CEA	Model based	11 years	QALYs (HUI)	£2445 per QALY
Carlton 2008 ⁴⁴	Childhood screening	Amblyopia (and strabismus) screening (no screening)	3–5 years	QALY-based CEA	Model based	100 years	QALYs (utility values from the literature)	Screening at 3 years without autorefraction was the most cost-effective, ICER of £527 375 per QALY.
Craig 2011 ⁴⁵	Childhood screening	Grote strategy for short stature screening (UK strategy)	Under 3 years	QALY-based CEA	Model based	12 years	QALYs (utility values from literature and expert opinion)	£1144 per QALY
Fayter 2007 ⁴⁶	Childhood screening	Short stature screening (no monitoring)	5 years	QALY-based CEA	Model based	Lifetime	QALYs (utility values from literature)	£9500 per QALY gained
Fortnum 2016 ⁴⁷	Childhood screening	Hearing screening (no screening)	4–5 years	QALY-based CEA	Model based	4 years	QALYs (utility values from literature)	The SES programme is dominated
Grill 2006 ⁴⁸	Childhood screening	Hospital hearing screening (community)	Newborn	Non-QALY-based CEA	Model based	10 years	Quality weighted detected child months	£2423 per detected child; £25 per quality weighed detected child month
Barber 2015 ⁴⁹	Health promotion	Preschoolers in the Playground, PiP (usual care)	1–4 years	QALY-based CEA	Trial based	1 year	QALYs (EQ-5D and PedsQL)	£19 588 per QALY

(Continued)

Table 1 Continued.

Author year	Intervention category	Intervention (comparator)	Population	Type of evaluation	Study type	Time horizon	Outcomes captured (quality of life instrument)	Primary result of the evaluation
Hollingworth 2012 ⁵⁰	Health promotion	Obesity/overweight interventions (no/minimal intervention)	4–5 years	Non-QALY-based CEA	Model based	Lifetime	Life years gained	£66 567 per life year gained (BMI SD score reduction of 0.03); £13 589 per life year gained (0.13 BMI SD score reduction)
Renwick 2018 ⁵¹	Health promotion	Smoking home intervention (usual care)	Under 5 years	Non-QALY-based CEA	Trial based	12 weeks	Average 16–24 h levels of particulate matter of < 2.5 µm diameter (PM _{2.5}); the number of quitters	£131 per additional 10 µg/m ³ reduction of 16–24 h PM _{2.5} ; £71 per additional quitter
Atkins 2012 ⁵²	Health protection	RotaTeq (no vaccination)	Under 6 months	QALY-based CEA	Model based	50 years	QALYs (utility values from the literature)	Dynamic model: £27 133 per QALY. Static model: £34 728 per QALY. Other scenarios presented.
Baguelin 2015 ⁵³	Health protection	LAIV (no vaccination)	2–4 years	QALY-based CEA	Model based	10 years	QALYs (utility values from the literature)	£2613 per QALY
Beck 2021 ⁵⁴	Health protection	4CMenB vaccination (no vaccination)	Under 1 year	QALY-based CEA	Model based	100 years	QALYs (utility values from the literature)	£18 645 per QALY gained
Brisson 2003 ⁵⁵	Health protection	VZV vaccination, infant strategy (no vaccination)	12–15 months	QALY-based CEA	Model based	80 years	QALYs (HUI-2)	VZV infant vaccination strategy is dominated
Christensen 2013 ⁵⁶	Health protection	New 'MenB' vaccine (no vaccination)	2 months to 4 years	QALY-based CEA	Model based	100 years	QALYs (utility values from the literature)	Between £162 800 and £290 000 per QALY gained (cohort model); and between £91 800 and £97 600 per QALY (dynamic model)
Christensen 2014 ⁵⁷	Health protection	Bexsero (no vaccination)	2 months to 1 year	QALY-based CEA	Model based	100 years	QALYs (EQ-5DY)	Ranged from £163 100 to £221 000 per QALY.
Edmunds 2002 ⁵⁸	Health protection	Acellular pertussis booster (no vaccination)	4 years	Non-QALY-based CEA	Model based	lifetime	Life-years gained; general practitioner consultation; and hospitalization averted	Booster doses range from £8463 to £49 511 per life year gained from the health perspective; £2489 to £36 941 per life year gained from a societal perspective.
Hodgson 2020 ⁵⁹	Health protection	RSV vaccination, MAB (status quo)	Under 5 years	QALY-based CEA	Model based	10 years	QALYs (EQ-5D)	Results presented as the maximum purchasing price per course for programmes to be cost-effective. For: MAB-VHR-S (£4342.97); MAB-HR-S (£201.15); MAD; MAB-HR-S+ (£87.03); VAC-INF-S (£94.76).
Jit 2007 ⁶⁰	Health protection	Rotavirus vaccination (no vaccination)	2–4 months	QALY-based CEA and non-QALY-based CEA	Model based	Unclear	QALYs (HUI-2 and EQ-5D)	£79 905 per QALY gained; £525 per episode prevented; £3803 per hospitalization prevented (using RotaTeq). £60 928 per QALY gained; £391 per episode prevented; £3647 per hospitalization prevented (using Rotarix).
Jit 2009 ⁶¹	Health protection	Rotavirus vaccination	2–4 months	QALY-based CEA	Model based	5 years	QALYs (HUI-2 and EQ-5D)	The ICER is EUR110 000 per QALY gained (Rotarix vaccination programme) and EUR160 000 per QALY gained (RotaTeq vaccination programme)

(Continued)

Table 1 Continued.

Author year	Intervention category	Intervention (comparator)	Population	Type of evaluation	Study type	Time horizon	Outcomes captured (quality of life instrument)	Primary result of the evaluation
Jit 2010 ⁶²	Health protection	Rotavirus vaccination (current care)	2–4 months	QALY-based CEA	Model based	5 years	QALYs (HUI-2 and EQ-5D)	EUR110 000 per QALY gained (Rotarix vaccination) and EUR150 000 per QALY gained (RotaTeq vaccination)
Knerer 2012 ⁶³	Health protection	Pneumococcal vaccination (PCV-13)	Under 2 years	QALY-based CEA	Model based	94 years	QALYs (utility values from literature)	Pneumococcal conjugate vaccine dominates (positive QALYs, negative costs).
Lorgelly 2007 ⁶⁴	Health protection	Rotavirus vaccination programme (no vaccination)	Newborn	Non-QALY-based CEA	Model based	5 years	Gastroenteritis episode avoided; GP visit avoided; hospitalization visit avoided; life years saved	£60.41 per episode avoided; £177 212 per life year saved.
Martin 2009 ⁶⁵	Health protection	Rotarix (no vaccination)	Under 6 months	QALY-based CEA	Model based	Lifetime	QALYs (EQ-5D)	£23 298 per QALY
McIntosh 2003 ⁶⁶	Health protection	Pneumococcal vaccination (no vaccination)	Under 6 months	Non-QALY-based CEA	Model based	10 years	Life years saved	£31 512 per life year saved
Melegaro 2004 ⁶⁷	Health protection	Pneumococcal vaccination (no vaccination)	2 months to 2 years	Non-QALY-based CEA and QALY-based CEA	Model based	Lifetime	Life years gained; QALYs (utility values from literature)	£70 699 per life year gained; £31 021 per QALY
Pitman 2013 ⁶⁸	Health protection	Influenza vaccination (current policy)	2–4 years	QALY-based CEA	Model based	200 years	QALYs (utility decrement from literature)	TIV in 2–4-year olds is dominated. LAIV in 2–4-year olds is cost saving
Siddiqui 2011 ⁶⁹	Health protection	HBV programme (current vaccination practice)	Under 6 months	QALY-based CEA	Model based	99 years	QALYs (utility values from literature)	£263 000 per QALY (for universal infant vaccination programme); £90 000 per QALY (for the selective infant programme)
Thomas 2018 ⁷⁰	Health protection	RSV vaccination (no vaccination)	Under 2 years	CBA	Cohort study	Lifetime	Costs	B/C ratios: 7.726 for bronchopulmonary dysplasia; 0.694 for congenital heart disease; 1.391 for extreme immaturity; 1.426 for premature babies; 0.465 for all other RSV admissions. All results for year of 2012/2013.
Trotter 2002 ⁷¹	Health protection	Meningitis C vaccination (no vaccination)	Under 4 years	Non-QALY-based CEA	Model based	Lifetime	Life years saved	0–4-month programme: £14 630 per life year saved; 5–11-month programme: £9493 per life year saved; 1–4 year programme: £5826 per life year saved
Trotter 2006a ⁷²	Health protection	Meningococcal vaccination (no vaccination)	Under 1 year	Non-QALY-based CEA and QALY-based CEA	Model based	100 years	Life years gained and QALYs (utility values from literature)	2–4-month programme: £38 164 per life year saved and £31 152 per QALY.

(Continued)

Table 1 Continued.

Author year	Intervention category	Intervention (comparator)	Population	Type of evaluation	Study type	Time horizon	Outcomes captured (quality of life instrument)	Primary result of the evaluation
Trotter 2006b ⁷³	Health protection	Meningococcal vaccination (current schedule)	Under 2 years	Non-QALY-based CEA	Model based	75 years	Life years gained	Strategy 2: £4 498 000 per life year; Strategy 3a: (2, 4, 13 months) –£ 2000 per life year gained; Strategy 3ab: (3, 13 months) –£4 811 000 per life year gained; Strategy 4: –£16 419 000 per life year gained
Achana 2016 ⁷⁴	Injury prevention	Six intervention combinations of education, equipment, home inspection and fitting (usual care)	Under 4 years	QALY-based CEA and non-QALY-based CEA	Model based	100 years	QALYs (utility values from the literature) and numbers of poison cases avoided	Non-QALY-based CEA: lowest ICER was education at £2888 per poison avoided. QALY-based CEA: lowest ICER was education at £41 330 per QALY gained.
Kendrick 2017a ⁷⁵	Injury prevention	(i) Functional smoke alarm (usual care) (ii) Safe hot tap water temperature (usual care) (iii) Promoting safety gate possession and use (usual care) (iv) Promoting the safe storage of medicines (usual care) (v) Promoting the safe storage of household and other products (usual care)	Under 5 years	QALY-based CEA	Model based	100 years	QALYs (utility values from literature)	(i) Education + equipment is £34 200 per QALY gained. (ii) Education is £40 271 per QALY gained. (iii) Education is £284 068 per QALY gained. (iv) Education is £41 330 per QALY gained. (v) All interventions were more costly and less effective than usual care.
Kendrick 2017b ⁷⁵	Injury prevention	IPB with or without facilitation (usual care)	Under 3 years	Non-QALY-based CEA	Trial based	1 year	Probability of having a fire escape plan	Injury prevention briefing only: £1260 per additional fire escape plan, injury prevention briefing +£616.13 per additional fire escape plan
Phillips 2011 ⁷⁶	Injury prevention	Scald prevention (waiting list)	Under 5 years	Non-QALY-based CEA and ROI (not stated)	Trial based	1 year	Risk reduction (scalds)	Scald prevention intervention: net savings of £7273 per scald avoided (NHS perspective), £53 949 per scald avoided (societal perspective). The benefit per £1 spent is £1.41 for an NHS perspective and (£0.47) for a lifetime perspective.
Saramago 2014 ⁷⁷	Injury prevention	Fire injury prevention interventions (usual care)	Under 5 years	QALY-based CEA	Model based	100 years	QALYs (utility values from literature)	Non-dominated interventions: education plus low cost/free safety equipment, £34 200 per QALY gained; education plus low cost/free safety equipment plus fitting plus home inspection at £3 466 635 per QALY gained.
Bessey 2019 ⁷⁸	Newborn screening	SCID screening (no screening)	Newborn	QALY-based CEA	Model based	5 years	QALYs (EQ-5D-3L)	£18 222 per QALY gained

(Continued)

Table 1 Continued.

Author year	Intervention category	Intervention (comparator)	Population	Type of evaluation	Study type	Time horizon	Outcomes captured (quality of life instrument)	Primary result of the evaluation
Kay 2018 ⁹⁰	Oral health	Supervised tooth brushing (no intervention)	5 years	QALY-based CEA	Model based	3 years	QALYs (utility values from literature)	Spending <£55 per child on supervised tooth brushing is cost-effective; spending <£100 on varnish would be cost-effective over 3 years
Kowash 2006 ⁹¹	Oral health	Out-reach education programme (unclear)	Under 1 year	CBA and non-QALY-based CEA	Trial based	3 years	Monetary and decayed, missing or filled tooth or tooth surface	The B/C ratio is 5.6. Cost-effectiveness ratio is 1.8.
O'Neill 2017 ⁹²	Oral health	Caries prevention (advice only)	2–3 years	Non-QALY-based CEA	Trial based	3 years	Proportion caries free; number of carious surfaces; number of episodes of pain	£2092.59 per caries free person; £250.58 per carious surface; £259.07 per number of pain episodes
Tickle 2016 ⁹³	Oral health	NIC-PIP caries prevention (prevention advice alone)	2–3 years	Non-QALY-based CEA	Trial based	3 years	Caries-free person; carious surfaces; episodes of pain	£2092.59 per proportion caries free; £250.58 per number of carious surfaces; £259.07 per episode of pain
Barnardo's 2012a ⁹⁴	Parenting support	Barnardo's Children's Centre Service: Stay and Play (unclear)	Under 2 years	SROI	Cohort study	5 years	Monetary outcomes	Approximately £2 for every £1 invested
Barnardo's 2012b ⁹⁴	Parenting support	Barnardo's Children's Centre Service: Family Support Worker (unclear)	Under 5 years	SROI	Cohort study	5 years	Monetary outcomes	£4.50 for every £1 invested
Edwards 2007 ⁹⁵	Parenting support	The Webster-Stratton Incredible Years basic parenting programme (waiting list)	3–4 years	Non-QALY-based CEA	Trial based	1 year	ECBI-I	£71 per one point change in the ECBI-I score
Gardner 2017 ⁹⁶	Parenting support	IY Basic parenting programme (no intervention)	5 years	Non-QALY-based CEA and ROI	Model based	25 years	ECBI-I	A WTP of £109 per point improvement on the ECBI-I is 50% probability of being cost-effective. In the 'high-cost' scenario, the ROI is 'nearly fourfold'. Assumed to be an ROI of 4 for the results.
McAuley 2004 ⁹⁷	Parenting support	Home Start support (no home start support)	Under 5 years	Non-QALY-based CEA and CCA	Cohort study	1 year	PSI; EPDS; RSE; BITSEAS; MSSSI	The intervention was assumed to be dominated (no effect difference and increases costs in the Home Start arm)
Morell 2000a ³⁵ Morell 2000b ⁹⁸	Parenting support	Postnatal support from a community midwifery support worker (no support worker)	Newborn	CCA (not specified but reports costs and outcomes separately)	Trial based	6 months	SF-36; Duke functional social support; Edinburgh postnatal depression scale; number breastfeeding only; number formula milk feeding only	No evidence of differences in SF-36, Edinburgh postnatal depression scale, and Duke functional social support scale) and rates of breast feeding between the two groups. The difference in total NHS costs between the groups was £178.61.

(Continued)

Table 1 Continued.

Author year	Intervention category	Intervention (comparator)	Population	Type of evaluation	Study type	Time horizon	Outcomes captured (quality of life instrument)	Primary result of the evaluation
Simkiss 2013 ⁹⁹	Parenting support	The Family Links Nurturing Programme (no screening)	2–4 years	QALY-based CEA	Trial based	10 years	QALYs (SF-6D, PedsQL)	£34 913 per QALY over 5 years and £18 954 per QALY over 10 years
Tudor Edwards 2016 ¹⁰⁰	Parenting support	IY BASIC parenting programme (waiting list)	3–4 years	Non-QALY-based CEA	Trial based	6 months	SDQ; ECBI; APS	£1295 per one point improvement in SDQ; £237 per one point improvement in ECBI-I; £9477 per one point improvement in APS
Chance 2013 ¹⁰¹	Parenting support and health promotion	Cambridgeshire's Funded Two-year-old Childcare (unclear)	2 years	SROI	Cohort study	5 years	Monetary outcomes	£8.40 for every £1 invested.
Mujica Mota 2006 ¹⁰²	Parenting support and health promotion	Means-tested access to full-time or part-time day care at the Hackney Early Years Centre (childcare secured themselves)	6 months to 3.5 years	Non-QALY-based CEA	Trial based	18 months	Proportion of mothers in paid employment or education at 18 months	£38 550 per additional woman in paid employment; societal perspective shows it to be cost saving.
Barlow 2019 ¹⁰³	Reducing risk of abuse/maltreatment	Parents under Pressure, PuP (treatment as usual)	Under 2 years	QALY-based CEA	Trial based	1 year	QALYs (EQ-5D-5L)	£34 095 per QALY (NHS and PSS perspective); £56 269 per QALY (societal perspective)
Boyd 2016 ¹⁰⁴	Reducing risk of abuse/maltreatment	New Orleans-Glasgow model (existing Glasgow model)	Under 5 years	CCA	Cohort study	5 years	Probability of one and two episodes in care	Reduced probability of two episodes in care (incremental reduction of 0.41) and reduced mean cost per child in the model (incremental difference of £6820).

AOM, acute otitis media; APS, Arnold–O’Leary Parenting Scale; BC, breast cancer;; CHD, congenital heart defect; ECBI, Eyberg Child Behaviour Inventory; GI, Gastrointestinal infection; HUI, health utilities index; ICER, incremental cost-effectiveness ratio; LRTI, lower respiratory tract infection; NEC, necrotizing enterocolitis; PedsQL, paediatric quality of life inventory; QWB, quality of well-being; SDQ, strengths and difficulties questionnaire; SF-6D, short-form 36 health survey.

Table 2 Summary of approaches

	Category	Total	%
Type of evaluation [†]	QALY-based CEA	32/77	42
	Non-QALY-based CEA	34/77	44
	CCA	4/77	5
	CBA	2/77	3
	SROI	3/77	4
	ROI	2/77	1
Outcomes used in CEA	Life years	10/34	29
	Multiple oral health outcomes	5/34	15
	Cases detected	4/34	12
	ECBI-I	3/34	9
	Timely diagnosis	3/34	9
	Multiple breastfeeding outcomes	2/34	6
	Poison cases avoided	1/34	3
	Quality weighted detected child months	1/34	3
	Probability of having a fire escape plan	1/34	3
	Proportion of mothers in paid employment or education at 18 months	1/34	3
	Risk reduction (scalds)	1/34	3
	PM2.5 level and the number of quitters	1/34	3
	Unclear	1/34	3
Perspective	NHS or NHS and PSS	38/69	55
	Societal	9/69	13
	NHS (scenario analysis of societal)	9/69	13
	Public sector	6/69	9
	NHS, education services, patients and family	1/69	1
	Public payer	1/69	1
	Health and other public sector providers	1/69	1
	NHS and the family	1/69	1
	NHS and ‘other government departments’	1/69	1
	Children and their families	1/69	1
	NHS (includes scenario with lost labour costs to families of children)	1/69	1
Incorporation of equity considerations in analysis?	Yes	2/69	3
	No	67/69	97
Use of decision modelling?	Yes	46/69	67
	No	23/69	33

[†] For the purpose of the summary of results, the number of evaluations was considered to be 77 as eight evaluations presented the results of two types of evaluation.

number were defined as having a societal perspective in the base case (13%; 9/69) or presenting a societal perspective alongside an NHS or NHS and PSS perspective (13%; 9/69). The breadth of the incorporated costs and outcomes in the societal perspectives, however, differed across evaluations.

These included costs borne by the family/caregiver (72%; 13/18), lost wages (61%, 11/18), household expenditure (28%, 5/18), travel time (11%, 2/18), the local authority/council (22%, 4/18), legal costs (11%, 2/18) and education costs (17%; 3/18). The outcomes captured in the societal perspective

The overwhelming majority of evaluations did not formally incorporate equity considerations (97%; 67/69), see [Table 2](#). Two evaluations (3%)^{74,88} considered the cost-effectiveness results across two different social groups. This provided insight into the distribution of health-related outcomes and therefore the cost-effectiveness across social groups.

Figure 2 presents the reported results from each study, grouped by the framework used and the intervention group. For QALY-based CEAs, 26 evaluations reported an incremental cost per QALY

Finally, only 29 of the evaluations covered all of the issues of concern in the presentation and discussion of results. The results of the detailed CHEERS checklist are presented in Supplementary Material, [Supplementary Appendix 3](#).

The results show the breadth of UK-focussed economic evaluations of early childhood public health




Fig. 2 Results of the QALY-based CEAs, CBAs and SROI and ROI by intervention category.

interventions reported or discussed in the published literature. The methods adopted in the demonstration of value for money showed a lack of consistency across many aspects including the type of economic evaluation, the health outcomes captured and the perspective adopted. Fourteen papers reported on all aspects of the CHEERS Checklist meaning 55 (80%) were lacking elements required of a well-reported economic evaluation.

Many of the evaluated interventions were deemed value for money from the perspectives taken. Twelve (38%) of the QALY-based CEAs are cost-effective against the NICE policy threshold of £20 000 per QALY. However, the interpretation of some of the results may require particular consideration given the health, economic, political and social context of these studies may have changed between 2000 and present day.

Because of the evaluative framework chosen for many of the other studies a robust statement of value for money of the intervention is not always possible. For example, in non-QALY-based CEAs, an explicit statement of cost-effectiveness is challenging when the outcome is a metric other than a generic measure of health such as the QALY as it is not possible to compare across different health dimensions. Much has been made of the limitations and challenges of using QALYs for paediatric populations^{17,19} but their use does allow the comparison of interventions across diseases areas as well as the consideration of the displaced resources (or the 'opportunity cost').

All of the interventions evaluated using SROI, ROI or CBA frameworks could be considered value for money as they were deemed to generate more monetary benefits than the costs (having a ratio of >£1 of benefit per £1 of cost in the case of the SROI

Although only a minority, a number of evaluations attempted to incorporate the wider social value of the intervention beyond the value to the health care system. A total of 18 evaluations adopted a ‘societal perspective’ but the results identified a lack of consistency in the included aspects of value. The inclusion of lost productivity to the parent or caregiver (in the form of wages lost) featured heavily in the evaluations, as did incorporating costs falling on special education services and legal services, yet none featured consistently. The implication of such inconsistencies is that value judgements about what ‘should’ count are falling on the researchers rather than socially legitimate decision-makers.¹⁰⁷ Public health guidance issued by NICE¹⁰⁸ does allow for flexibility in the costs and outcomes considered in an economic evaluation, but the lack of explicit value judgements may facilitate such inconsistencies.

The focus of this review was to identify interventions relevant to UK decision-makers. However, there may be important information available in an international context to aid learnings around the use of methods and approaches relevant to the UK. Future research may consider describing the methods and approaches adopted in the global evidence base to highlight consistencies in the demonstration of value for money in those economic evaluations developed for an international context.

A limitation is that there may be relevant and uncaptured evaluations in the grey literature. This is evidenced through the identification of evaluations produced by NICE,³⁸ Social Value UK,¹⁰¹ Barnardo's⁹⁴ and the Joseph Rowntree Foundation,⁹⁷

which were not identified in the database search but rather through the literature reviews. The search strategy included the HMIC database and lists grey literature amongst its coverage, hence the decision to include relevant grey literature in the reference searches of systematic reviews. The inclusion of the grey literature identified through the supplementary reference searching does in part explain the high number of studies identified in this way (5 of the 13). We considered it important to include grey literature as it is included in the HEER tool,³³ yet a pragmatic decision was made for the purpose of this review. Future literature reviews may consider searching and identifying a wider range of grey literature sources.

A further limitation was the difficulty posed in defining ‘public health’ for the purpose of the search strategy. The review focussed on interventions that aimed to improve the health of the infant or child yet health improvement in early childhood may be dependent on lifestyle and environment not merely based on biology and genetics.⁶ It stands that a social model of health may have generated different results. A pragmatic decision was made to include health terms and terms to capture the wider determinants of health in the search strategy.

Conclusion

In addition to identifying the breadth of evidence available in the published literature, this review provides an overview of the inconsistent methodological approaches used. The lack of consistency identified in the methods has highlighted a number of issues that may require consideration in the future generation of economic evaluations of similar interventions to aid decision-making. It is hoped the results can provide a foundation to help improve decision-making and provide a starting point for methodological developments in the early childhood public health context.

Authors' contribution

PM, SH and GR discussed the direction of the research and the general approach to the systematic review. HF advised on and ran the search strategies. PM and LP screened identified records. PM drafted

the manuscript. All authors commented on the early drafts and the final version of the manuscript.

Funding

This paper presents independent research as part of a PhD programme funded by the National Institute for Health Research's (NIHR) Applied Research Collaboration Yorkshire and Humber. The views expressed are those of the author(s) and not necessarily those of the NIHR.

Conflicts of interest statement

All authors declare no conflict of interest.

Data availability

The data underlying this article are available in the article and in its online supplementary material.

References

1. Shonkoff JP, Richter L, van der Gaag J, et al. An integrated scientific framework for child survival and early childhood development. *Pediatrics* 2012;129:e460–72.
2. Richter LM, Daelmans B, Lombardi J, et al. Investing in the foundation of sustainable development: pathways to scale up for early childhood development. *Lancet* 2017;389:103–18.
3. Black RE, Taylor CE, Arole S, et al. Comprehensive review of the evidence regarding the effectiveness of community-based primary health care in improving maternal, neonatal and child health: 8. summary and recommendations of the Expert Panel. *J Glob Health* 2017;7:010908.
4. Daelmans B, Black MM, Lombardi J, et al. Effective interventions and strategies for improving early child development. *BMJ* 2015;351:h4029.
5. Clarke-Stewart KA. Interactions between mothers and their young children: characteristics and consequences. *Monogr Soc Res Child Dev* 1973;38:1–109.
6. Marmot M, Bell R. Fair society, healthy lives. *Public Health* 2012;126:S4–s10.
7. Marmot M. Health equity in England: the Marmot review 10 years on. *BMJ* 2020;368:m693.
8. Government HM. *The Best Start for Life: A Vision for the 1,001 Critical Days*. London: Crown copyright; 2021.

9. Strelitz J, Nolte E, Pitchforth E, et al. The economic case for a shift to prevention. In: Lemer C, Todd K, Cheung R, editors. *Annual Report of the Chief Medical Officer*. London: Department of Health; 2012. p. 47–88.
10. Drummond MF, Sculpher MJ, Claxton K, et al. *Methods for the Economic Evaluation of Health Care Programmes*. Oxford: Oxford University Press; 2015.
11. National Institute for Health and Care Excellence. *NICE Health Technology Evaluations: The Manual*. London: National Institute for Health and Care Excellence; 2022.
12. Mathes T, Antoine SL, Prengel P, et al. Health technology assessment of public health interventions: a synthesis of methodological guidance. *Int J Technol Assess Health Care* 2017;33:135–46.
13. Edwards RT, McIntosh E. *Applied Health Economics for Public Health Practice and Research*. Oxford: Oxford University Press; 2019.
14. Taulbut M, Walsh D. *Poverty, Parenting, and Poor Health: Comparing Early Years' Experiences in Scotland, England and Three City Regions*. Glasgow: Glasgow Centre for Population Health; 2013.
15. Graham H, Power C. Childhood disadvantage and health inequalities: a framework for policy based on lifecourse research. *Child Care Health Dev* 2004;30:671–8.
16. Kiernan KE, Huerta MC. Economic deprivation, maternal depression, parenting and children's cognitive and emotional development in early childhood. *Br J Sociol* 2008;59:783–806.
17. Petrou S. Methodological challenges surrounding QALY estimation for paediatric economic evaluation. *Cost Eff Resour Alloc* 2022;20:10.
18. Petrou S, Gray R. Methodological challenges posed by economic evaluations of early childhood intervention programmes. *Appl Health Econ Health Policy* 2005;4:175–81.
19. Rowen D, Rivero-Arias O, Devlin N, et al. Review of valuation methods of preference-based measures of health for economic evaluation in child and adolescent populations: where are we now and where are we going? *Pharmacoeconomics* 2020;38: 325–40.
20. Powell TG, Gheera M. *Early Intervention: A Background Paper*. London: House of Commons Library; 2021.
21. Boonacker CW, Broos PH, Sanders EA, et al. Cost effectiveness of pneumococcal conjugate vaccination against acute otitis media in children: a review. *Pharmacoeconomics* 2011;29:199–211.
22. Sampaio F, Nystrand C, Feldman I, Mihalopoulos C. Evidence for investing in parenting interventions aiming to improve child health: a systematic review of economic evaluations. *Eur Child Adolesc Psychiatry* 2022.
23. Charles JM, Bywater T, Edwards RT. Parenting interventions: a systematic review of the economic evidence. *Child Care Health Dev* 2011;37:462–74.
24. Jacklin P, Retse P, Kwan I, et al. *Rapid Economic Review of Public Health Interventions Designed to Improve the Nutrition of Children Aged 0-5 Years*. London: National Institute of Health and Care Excellence; 2006.
25. Batura N, Hill Z, Haghighparast-Bidgoli H, et al. Highlighting the evidence gap: how cost-effective are interventions to improve early childhood nutrition and development? *Health Policy Plan* 2015;30:813–21.
26. Döring N, Mayer S, Rasmussen F, et al. Economic evaluation of obesity prevention in early childhood: methods, limitations and recommendations. *Int J Environ Res Public Health* 2016;13:911.
27. Anopa Y, Conway DI. Exploring the cost-effectiveness of child dental caries prevention programmes. Are we comparing apples and oranges? *Evid Based Dent* 2020;21:5–7.
28. Drummond M, Barbieri M, Cook J, et al. Transferability of economic evaluations across jurisdictions: ISPOR Good Research Practices Task Force report. *Value Health* 2009;12:409–18.
29. Owen L, Morgan A, Fischer A, et al. The cost-effectiveness of public health interventions. *J Public Health* 2011;34:37–45.
30. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535.
31. Hinde S, Spackman E. Bidirectional citation searching to completion: an exploration of literature searching methods. *Pharmacoeconomics* 2015;33:5–11.
32. McHugh ML. Interrater reliability: the kappa statistic. *Biochem Med* 2012;22:276–82.
33. Public Health England. *Health Economics: Evidence Resource* 2019. <https://www.gov.uk/government/publications/health-economics-evidence-resource> (24 May 2022, date last accessed).
34. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
35. Morrell CJ, Spiby H, Stewart P, et al. Costs and benefits of community postnatal support workers: a randomised controlled trial. *Health Technol Assess* 2000;4:1–100.

36. Pandor A, Eastham J, Beverley C, et al. Clinical effectiveness and cost-effectiveness of neonatal screening for inborn errors of metabolism using tandem mass spectrometry: a systematic review. *Health Technol Assess* 2004;8:1–121.
37. Pandor A, Eastham J, Chilcott J, et al. Economics of tandem mass spectrometry screening of neonatal inherited disorders. *Int J Technol Assess Health Care* 2006;22:321–6.
38. Jacklin P, Retsa P, Dougherty M, et al. *Modelling the Cost Effectiveness of Interventions to Promote Breastfeeding*. London: National Institute for Health and Care Excellence; 2007.
39. National Institute for Health and Care Excellence. *Modelling the Cost-Effectiveness of Breastfeeding Support*. London: National Institute of Health and Care Excellence; 2007.
40. Anokye N, Coyle K, Relton C, et al. Cost-effectiveness of offering an area-level financial incentive on breast feeding: a within-cluster randomised controlled trial analysis. *Arch Dis Child* 2020;105:155–9.
41. Hoddinott P, Craig L, MacLennan G, et al. The FEeding Support Team (FEST) randomised, controlled feasibility trial of proactive and reactive telephone support for breastfeeding women living in disadvantaged areas. *BMJ Open* 2012;2:e000652.
42. Pokhrel S, Quigley MA, Fox-Rushby J, et al. Potential economic impacts from improving breastfeeding rates in the UK. *Arch Dis Child* 2015;100:334–40.
43. Bamford J, Fortnum H, Bristow K, et al. Current practice, accuracy, effectiveness and cost-effectiveness of the school entry hearing screen. *Health Technol Assess* 2007;11(32):1–168, iii–iv.
44. Carlton J, Karnon J, Czoski-Murray C, et al. The clinical effectiveness and cost-effectiveness of screening programmes for amblyopia and strabismus in children up to the age of 4–5 years: a systematic review and economic evaluation. In: *Health Technol Assess*, Vol. 12, 2008,iii,xi–194
45. Craig D, Fayer D, Stirk L, Crott R. Growth monitoring for short stature: update of a systematic review and economic model. *Health Technol Assess* 2011;15:11.
46. Fayer D, Nixon J, Hartley S, et al. A systematic review of the routine monitoring of growth in children of primary school age to identify growth-related conditions. *Health Technol Assess* 2007;11:xi–xii.
47. Fortnum H, Ukoumunne OC, Hyde C, et al. A programme of studies including assessment of diagnostic accuracy of school hearing screening tests and a cost-effectiveness model of school entry hearing screening programmes. *Health Technol Assess* 2016; 20(36):1–178.
48. Grill E, Uus K, Hessel F, et al. Neonatal hearing screening: modelling cost and effectiveness of hospital- and community-based screening. *BMC Health Serv Res* 2006;6:1–9.
49. Barber SE, Akhtar S, Jackson C, et al. Preschoolers in the playground: a pilot cluster randomised controlled trial of a physical activity intervention for children aged 18 months to 4 years. Southampton: NIHR Journals Library; 2015.
50. Hollingworth W, Hawkins J, Lawlor D, et al. Economic evaluation of lifestyle interventions to treat overweight or obesity in children. *Int J Obes* 2012;36:559–66.
51. Renwick C, Wu Q, Breton MO, et al. Cost-effectiveness of a complex intervention to reduce children's exposure to second-hand smoke in the home. *BMC Public Health* 2018;18:1252.
52. Atkins KE, Shim E, Carroll S, et al. The cost-effectiveness of pentavalent rotavirus vaccination in England and Wales. *Vaccine* 2012;30:6766–76.
53. Baguelin M, Camacho A, Flasche S, Edmunds WJ. Extending the elderly-and risk-group programme of vaccination against seasonal influenza in England and Wales: a cost-effectiveness study. *BMC Med* 2015;13:1–13.
54. Beck E, Klint J, Neine M, et al. Cost-effectiveness of 4CMenB infant vaccination in England: a comprehensive valuation considering the broad impact of serogroup B invasive meningococcal disease. *Value Health* 2021;24:91–104.
55. Brisson M, Edmunds W. Varicella vaccination in England and Wales: cost-utility analysis. *Arch Dis Child* 2003;88:862–9.
56. Christensen H, Hickman M, Edmunds WJ, et al. Introducing vaccination against serogroup B meningococcal disease: an economic and mathematical modelling study of potential impact. *Vaccine* 2013;31:2638–46.
57. Christensen H, Trotter CL, Hickman M, et al. Re-evaluating cost effectiveness of universal meningitis vaccination (Bexsero) in England: modelling study. *BMJ* 2014;349:5725.
58. Edmunds WJ, Brisson M, Melegaro A, et al. The potential cost-effectiveness of acellular pertussis booster vaccination in England and Wales. *Vaccine* 2002;20:1316–30.
59. Hodgson D, Pebody R, Panovska-Griffiths J, et al. Evaluating the next generation of RSV intervention strategies: a mathematical modelling study and cost-effectiveness analysis. *BMC Med* 2020;18:1–14.

60. Jit M, Edmunds W. Evaluating rotavirus vaccination in England and Wales: part II. The potential cost-effectiveness of vaccination. *Vaccine* 2007;25:3971–9.
61. Jit M, Bilcke J, Mangen M-JJ, et al. The cost-effectiveness of rotavirus vaccination: comparative analyses for five European countries and transferability in Europe. *Vaccine* 2009;27:6121–8.
62. Jit M, Mangen M-JJ, Melliez H, et al. An update to 'the cost-effectiveness of rotavirus vaccination: comparative analyses for five European countries and transferability in Europe'. *Vaccine* 2010;28:7457–9.
63. Knerer G, Ismaila A, Pearce D. Health and economic impact of PHiD-CV in Canada and the UK: a Markov modelling exercise. *J Med Econ* 2012;15:61–76.
64. Lorgelly PK, Joshi D, Iturriza Gómara M, et al. Exploring the cost effectiveness of an immunization programme for rotavirus gastroenteritis in the United Kingdom. *Epidemiol Infect* 2008;136:44–55.
65. Martin A, Batty A, Roberts JA, et al. Cost-effectiveness of infant vaccination with RIX4414 (Rotarix) in the UK. *Vaccine* 2009;27:4520–8.
66. McIntosh ED, Conway P, Willingham J, et al. The cost-burden of paediatric pneumococcal disease in the UK and the potential cost-effectiveness of prevention using 7-valent pneumococcal conjugate vaccine. *Vaccine* 2003;21:2564–72.
67. Melegaro A, Edmunds WJ. Cost-effectiveness analysis of pneumococcal conjugate vaccination in England and Wales. *Vaccine* 2004;22:4203–14.
68. Pitman RJ, Nagy LD, Sculpher MJ. Cost-effectiveness of childhood influenza vaccination in England and Wales: results from a dynamic transmission model. *Vaccine* 2013;31:927–42.
69. Siddiqui MR, Gay N, Edmunds WJ, et al. Economic evaluation of infant and adolescent hepatitis B vaccination in the UK. *Vaccine* 2011;29:466–75.
70. Thomas G. A cost-benefit analysis of the immunisation of children against respiratory syncytial virus (RSV) using the English Hospital Episode Statistics (HES) data set. *Eur J Health Econ* 2018;19:177–87.
71. Trotter CL, Edmunds WJ. Modelling cost effectiveness of meningococcal serogroup C conjugate vaccination campaign in England and Wales. *BMJ* 2002;324:809.
72. Trotter CL, Edmunds WJ, Ramsay ME, et al. Modeling future changes to the meningococcal serogroup C conjugate (MCC) vaccine program in England and Wales. *Hum Vaccin* 2006;2:68–73.
73. Trotter CL, Edmunds WJ. Reassessing the cost-effectiveness of meningococcal serogroup C conjugate (MCC) vaccines using a transmission dynamic model. *Med Decis Making* 2006;26:38–47.
74. Achana F, Sutton AJ, Kendrick D, et al. A decision analytic model to investigate the cost-effectiveness of poisoning prevention practices in households with young children. *BMC Public Health* 2016;16:705.
75. Kendrick D, Ablewhite J, Achana F, et al. Keeping children safe: a multicentre programme of research to increase the evidence base for preventing unintentional injuries in the home in the under-fives. Southampton: NIHR Journals Library; 2017. PMID: 28771290.
76. Phillips CJ, Humphreys I, Kendrick D, et al. Preventing bath water scalds: a cost-effectiveness analysis of introducing bath thermostatic mixer valves in social housing. *Inj Prev* 2011;17:238–43.
77. Saramago P, Cooper NJ, Sutton AJ, et al. Cost-effectiveness of interventions for increasing the possession of functioning smoke alarms in households with pre-school children: a modelling study. *BMC Public Health* 2014;14:459.
78. Bessey A, Chilcott J, Leaviss J, et al. A cost-effectiveness analysis of newborn screening for severe combined immunodeficiency in the UK. *Int J Neonatal Screen* 2019;5:28.
79. Bessey A, Chilcott JB, Leaviss J, et al. Economic impact of screening for X-linked Adrenoleukodystrophy within a newborn blood spot screening programme. *Orphanet J Rare Dis* 2018;13:1–11.
80. Burke MJ, Shenton RC, Taylor MJ. The economics of screening infants at risk of hearing impairment: an international analysis. *Int J Pediatr Otorhinolaryngol* 2012;76:212–8.
81. Davies S, Cronin E, Gill M, et al. Screening for sickle cell disease and thalassaemia: a systematic review with supplementary research. *Health Technol Assess* 2000;4:i–v.
82. Ewer A, Furmston A, Middleton L, et al. Pulse oximetry as a screening test for congenital heart defects in newborn infants: a test accuracy study with evaluation of acceptability and cost-effectiveness. *Health Technol Assess* 2012;16:1–184.
83. Griebisch I, Knowles RL, Brown J, et al. Comparing the clinical and economic effects of clinical examination, pulse oximetry, and echocardiography in newborn screening for congenital heart defects: a probabilistic cost-effectiveness model and value of information analysis. *Int J Technol Assess Health Care* 2007;23:192–204.
84. Knowles R, Griebisch I, Dezateux C, et al. Newborn screening for congenital heart defects: a systematic

- review and cost-effectiveness analysis. *Health Technol Assess* 2005;9:1–152.
85. Roberts TE, Barton PM, Auguste PE, et al. Pulse oximetry as a screening test for congenital heart defects in newborn infants: a cost-effectiveness analysis. *Arch Dis Child* 2012;97:221–6.
86. Simpson N, Anderson R, Sassi F, et al. The cost-effectiveness of neonatal screening for cystic fibrosis: an analysis of alternative scenarios using a decision model. *Cost Eff Resour Alloc* 2005;3:8.
87. Uus K, Bamford J, Taylor R. An analysis of the costs of implementing the National Newborn Hearing Screening Programme in England. *J Med Screen* 2006;13:14–9.
88. Davenport C, Elley K, Salas C, et al. The clinical effectiveness and cost-effectiveness of routine dental checks: a systematic review and economic evaluation. *Health Technol Assess* 2003;7:1–127.
89. Davies G, Worthington H, Ellwood R, et al. An assessment of the cost effectiveness of a postal toothpaste programme to prevent caries among five-year-old children in the North West of England. *Community Dent Health* 2003;20:207–10.
90. Kay E, Owen L, Taylor M, et al. The use of cost-utility analysis for the evaluation of caries prevention: an exploratory case study of two community-based public health interventions in a high-risk population in the UK. *Community Dent Health* 2018;35:30–6.
91. Kowash MB, Tumba KJ, Curzon ME. Cost-effectiveness of a long-term dental health education program for the prevention of early childhood caries. *Eur Arch Paediatr Dent* 2006;7:130–5.
92. O'Neill C, Worthington HV, Donaldson M, et al. Cost-effectiveness of caries prevention in practice: a randomized controlled trial. *J Dent Res* 2017;96:875–80.
93. Tickle M, O'Neill C, Donaldson M, et al. A randomised controlled trial to measure the effects and costs of a dental caries prevention regime for young children attending primary care dental services: the Northern Ireland Caries Prevention in Practice (NIC-PIP) trial. *Health Technol Assess* 2016;20:1–96.
94. Salisbury D, Blazey L, Dutchman D, et al. The Value of Early Intervention Identifying the Social Return of Barnardo's Children's. *Centre Services*. Birmingham: Barnardo's; 2012.
95. Edwards RT, C  illeachair A, Bywater T, et al. Parenting programme for parents of children at risk of developing conduct disorder: cost effectiveness analysis. *BMJ* 2007;334:682.
96. Gardner F, Leijten P, Mann J, et al. Could scale-up of parenting programmes improve child disruptive behaviour and reduce social inequalities: using individual participant data meta-analysis to establish for whom programmes are effective and cost-effective. *Public Health Res* 2017;5:1–144.
97. McAuley C, Knapp M, Beecham J, et al. *The Outcomes and Costs of Home-Start Support for Young Families Under Stress*. York: Joseph Rowntree Foundation; 2004.
98. Morrell CJ, Spiby H, Stewart P, et al. Costs and effectiveness of community postnatal support workers: randomised controlled trial. *BMJ* 2000;321:593–8.
99. Simkiss DE, Snooks HA, Stallard N, et al. Effectiveness and cost-effectiveness of a universal parenting skills programme in deprived communities: multicentre randomised controlled trial. *BMJ Open* 2013;3:e002851.
100. Edwards RT, Jones C, Berry V, et al. Incredible Years parenting programme: cost-effectiveness and implementation. *J Child Serv* 2016;11(1):54–72.
101. Chance T. *Cambridgeshire's Funded Two-Year-Old Childcare: Social Return on Investment Report*. Cambridge: Cambridgeshire Country Council; 2013.
102. Mujica Mota R, Lorgelly PK, Mugford M, et al. Out-of-home day care for families living in a disadvantaged area of London: economic evaluation alongside a RCT. *Child Care Health Dev* 2006;32:287–302.
103. Barlow J, Sembi S, Parsons H, et al. A randomized controlled trial and economic evaluation of the Parents Under Pressure program for parents in substance abuse treatment. *Drug Alcohol Depend* 2019;194:184–94.
104. Boyd KA, Balogun MO, Minnis H. Development of a radical foster care intervention in Glasgow, Scotland. *Health Promot Int* 2015;31:665–73.
105. Brousselle A, Benmarhnia T, Benhadj L. What are the benefits and risks of using return on investment to defend public health programs? *Prev Med Rep* 2016;3:135–8.
106. Hinde S, Horsfield L, Bojke L, et al. The relevant perspective of economic evaluations informing local decision makers: an exploration in weight loss services. *Appl Health Econ Health Policy* 2020;18:351–6.
107. Williams A. Is the QALY a technical solution to a political problem? Of course not! *Int J Health Serv* 1991;21:365–9.

108. NICE. *Developing NICE Guidelines: The Manual*. London: National Institute for Health and Care Excellence, 2014.
109. Ingleby FC, Woods LM, Atherton IM, et al. Describing socio-economic variation in life expectancy according to an individual's education, occupation and wage in England and Wales: an analysis of the ONS longitudinal study. *SSM Popul Health* 2021;14: 100815.
110. Cookson R, Mirelman AJ, Griffin S, et al. Using cost-effectiveness analysis to address health equity concerns. *Value Health* 2017;20:206–12.