



This is a repository copy of *Direct nitrogen, phosphorus and carbon exchanges between Mucoromycotina 'fine root endophyte' fungi and a flowering plant in novel monoxenic cultures*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/193804/>

Version: Published Version

Article:

Hoysted, G., Field, K. orcid.org/0000-0002-5196-2360, Sinanaj, B. et al. (3 more authors) (2023) Direct nitrogen, phosphorus and carbon exchanges between Mucoromycotina 'fine root endophyte' fungi and a flowering plant in novel monoxenic cultures. *New Phytologist*, 238 (1). pp. 70-79. ISSN: 0028-646X

<https://doi.org/10.1111/nph.18630>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Rapid report

Direct nitrogen, phosphorus and carbon exchanges between *Mucoromycotina* ‘fine root endophyte’ fungi and a flowering plant in novel monoxenic cultures

Authors for correspondence:

Silvia Pressel

Email: s.pressel@nhm.ac.uk

Katie Field

Email: k.j.field@sheffield.ac.uk

Received: 9 June 2022

Accepted: 15 November 2022

Grace A. Hoysted¹ , Katie J. Field¹ , Besiana Sinanaj¹ ,
Christopher A. Bell² , Martin I. Bidartondo^{3,4}  and Silvia Pressel⁵ 

¹Plants, Photosynthesis and Soil, School of Bioscience, University of Sheffield, Sheffield, S10 2TN, UK; ²Centre for Plant Sciences, University of Leeds, Leeds, LS2 9JT, UK; ³Department of Life Sciences, Imperial College London, London, SW7 2AZ, UK;

⁴Department of Ecosystem Stewardship, Royal Botanic Gardens, Kew, Richmond, TW9 3DS, UK; ⁵Department of Life Sciences, Natural History Museum, London, SW7 5BD, UK

New Phytologist (2023)

doi: 10.1111/nph.18630

Key words: arbuscular mycorrhizal fungi, carbon, clover, fine root endophytes, flowering plants, *Mucoromycotina*, soil nutrients, symbiosis.

Summary

- Most plants form mycorrhizal associations with mutualistic soil fungi. Through these partnerships, resources are exchanged including photosynthetically fixed carbon for fungal-acquired nutrients. Recently, it was shown that the diversity of associated fungi is greater than previously assumed, extending to *Mucoromycotina* fungi. These *Mucoromycotina* ‘fine root endophytes’ (MFRE) are widespread and generally co-colonise plant roots together with *Glomeromycotina* ‘coarse’ arbuscular mycorrhizal fungi (AMF). Until now, this co-occurrence has hindered the determination of the direct function of MFRE symbiosis.
- To overcome this major barrier, we developed new techniques for fungal isolation and culture and established the first monoxenic *in vitro* cultures of MFRE colonising a flowering plant, clover. Using radio- and stable-isotope tracers in these *in vitro* systems, we measured the transfer of ³³P, ¹⁵N and ¹⁴C between MFRE hyphae and the host plant.
- Our results provide the first unequivocal evidence that MFRE fungi are nutritional mutualists with a flowering plant by showing that clover gained both ¹⁵N and ³³P tracers directly from fungus in exchange for plant-fixed C in the absence of other micro-organisms.
- Our findings and methods pave the way for a new era in mycorrhizal research, firmly establishing MFRE as both mycorrhizal and functionally important in terrestrial ecosystems.

Introduction

Among Earth’s most important symbioses are the ancient plant–fungus partnerships known as mycorrhizas, or ‘mycorrhiza-like’ associations in plants without roots (Read *et al.*, 2000). These mutualisms, underpinned by the bidirectional exchange of plant-fixed carbon for fungal-acquired mineral nutrients (Raven & Allen, 2003), were instrumental in plant terrestrialisation > 500 million years ago (Morris *et al.*, 2018) by facilitating early, rootless plant access to mineral nutrients held within primeval soils (Pirozynski & Malloch, 1975; Field *et al.*, 2015a). Thus, fungi played a formative role in the development of Earth’s terrestrial ecosystems and climate through their contributions to global carbon and nutrient cycles (Taylor *et al.*, 2011; Mills *et al.*, 2018). Today, these associations are formed between most land plants and a diverse subset of soil fungi (Field & Pressel, 2018), including the

widespread arbuscular mycorrhizal fungi (AMF) in the subphylum *Glomeromycotina* (Brundrett & Tedersoo, 2018), estimated to occur in > 70% of plants (Smith & Read, 2008).

Until recently, AMF encompassed the globally distributed ‘fine root endophytes’ (FRE; *Glomus tenue*; Thippayrugs *et al.*, 1999), which are known to colonise several vascular plant families (Ali, 1969; Abbott, 1982; Thippayrugs *et al.*, 1999), but have been largely overlooked due to practical limitations in molecular detection and inability to study them apart from coexisting AMF (Orchard *et al.*, 2017a). Through improved molecular detection and identification (Bidartondo *et al.*, 2011), it is now clear that FRE are distinct from *Glomeromycotina* AMF, belonging instead to the Endogonales in the subphylum *Mucoromycotina* (Orchard *et al.*, 2017a), and recently renamed *Planticonsortium tenue* (Walker *et al.*, 2018). Thus, FRE previously reported in flowering plants (Ali, 1969; Abbott, 1982; Thippayrugs *et al.*, 1999) are likely

closely related to Endogonales fungal associates previously identified (Bidartondo *et al.*, 2011; Rimington *et al.*, 2015; Hoysted *et al.*, 2018, 2019) and shown to be mutualistic in terms of carbon-for-nutrient exchange in a range of nonflowering plants, albeit using nonsterile, soil-based experimental systems (Field *et al.*, 2015b, 2016, 2019; Hoysted *et al.*, 2019, 2020).

Colonisation by FRE is, like AMF, generally characterised by the presence of arbuscules and arbuscule-like structures (Orchard *et al.*, 2017b), while the small diameter of FRE hyphae ($< 1.5 \mu\text{m}$), with small swellings and 'fan-like' morphologies, is considered a distinctive trait that separates them from AMF (or 'coarse root endophytes') which consistently develop wider ($> 3 \mu\text{m}$ in diameter) hyphae and larger vesicles (Orchard *et al.*, 2017a). Morphological plasticity has been noted in transmission and scanning electron micrographs of the ultrastructure of Mucoromycotina FRE (MFRE) exclusively associating with liverworts (Field *et al.*, 2015b, 2016) and a vascular plant (Hoysted *et al.*, 2019). Most recently, cryo-SEM has confirmed uniformly thin hyphae and hyphal 'ropes' as potential diagnostic features of MFRE symbioses (Albornoz *et al.*, 2020). Differently from the strictly biotrophic AMF, MFRE are considered facultative saprotrophs as it has been possible to isolate them from host plants – both a nonvascular (Field *et al.*, 2015b) and vascular plant (as shown herein) – and to grow them axenically, that is without a host in culture.

Latest research indicates that MFRE play a nutritionally complementary role to AMF by facilitating plant nitrogen (N) assimilation alongside AMF-facilitated plant phosphorus (P) acquisition through co-colonisation of the same host (Field *et al.*, 2019). Such functional complementarity is further supported by the observation that MFRE transfer significant amounts of ^{15}N but relatively little ^{33}P isotope tracers to a host lycophyte in the first experimental demonstration of MFRE nutritional mutualism with a vascular plant (Hoysted *et al.*, 2019). The apparent ability of MFRE, but not AMF, to transfer N also from organic sources to host liverworts in nonsterile soil (Field *et al.*, 2019) together with their presumed facultative saprotrophic nature points to possible functional similarities with ectomycorrhizal fungi, an assumption in line with results from a recent network analysis of symbiotic fungal associations in liverworts (Rimington *et al.*, 2019). However, evidence for the precise role of MFRE, in the absence of other soil micro-organisms, remains equivocal.

To date, detailed research into plant–MFRE associations has been constrained by a lack of *in vitro* experimental systems that allow indisputable determination of the direct function of the MFRE symbiosis in isolation. Our recent knowledge of MFRE function has been derived largely from experiments using wild soil-based systems and wild-collected plants that naturally only associate with MFRE (Field *et al.*, 2015b, 2016, 2019; Hoysted *et al.*, 2019, 2020), or from soil culture-based experimental pots (Orchard *et al.*, 2017a; Albornoz *et al.*, 2020). Each of these methods has significantly enhanced our understanding of MFRE form and function, shedding new light on the importance of MFRE associations in nature, and remains useful in the studies of plants associating with mixed microbial communities.

However, the development of *in vitro* experimental systems capable of distinguishing between fungal symbionts in the absence of

other soil biota is now critical for the functional significance of MFRE associations to be fully defined (Sinanaj *et al.*, 2021). This is particularly important as evidence is increasingly pointing towards most plants forming simultaneous symbioses with AMF and MFRE (Field *et al.*, 2015a,b, 2016; Hoysted *et al.*, 2019, 2020) and there being complementarity in function between symbionts (Field *et al.*, 2019; Hoysted *et al.*, 2019). Recently, it was reported that a free-living Mucoromycotina, *Gongronella* sp. W5, utilises plant sucrose as a carbon source (Wang *et al.*, 2021); however, data showing mutualistic transfer of carbon-for-nutrients between MFRE and a plant host in the absence of other micro-organisms do not currently exist. The development of an *in vitro* experimental system is critical to achieve this and further understand function, development and signalling and to identify specific symbiotic structures and interfaces in MFRE, particularly for comparisons with model AM symbioses. Here, we resolve this research challenge by establishing experimentally tractable, monoxenic symbiotic cultures of MFRE and white clover (*Trifolium repens*), a flowering plant genus used in other recent studies of MFRE colonisation (e.g. Orchard *et al.*, 2017a; Albornoz *et al.*, 2020), albeit in nonsterile systems. Using radio- and stable-isotope tracers in our *in vitro* systems, we measured the transfer of ^{33}P , ^{15}N and ^{14}C between MFRE hyphae and the host plant to provide the first unequivocal evidence of mutualistic transfer of MFRE-assimilated nutrients for plant-fixed carbon with a flowering plant, in the absence of other microbes.

Materials and Methods

Isolation of MFRE symbionts

Lycopodiella inundata (L.) Holub gametophytes and young sporophytes (with protocorms; Hoysted *et al.*, 2019) were collected from Thursley Nature Reserve, Surrey, UK (SE 90081 39754) in September 2019 and processed immediately for fungal isolation or stored in their natural substrate in growth chambers at 20°C : 15°C day: night temperatures and a 16-h day length, $225 \mu\text{mol photons m}^{-2} \text{s}^{-1}$, for molecular analyses.

Gametophytes and young sporophytes were carefully cleaned of adhering substratum; rhizoids and (for sporophytes) leaves were removed before thorough rinsing in dH_2O by gentle shaking for 1 h, followed by surface-sterilisation for 1 min in 0.5% sodium hypochlorite. Sterile gametophytes were kept intact or halved, while the sporophytic protocorms were cut into *c.* 0.5-mm sections, placed onto fungal growth medium under sterile conditions and incubated in the dark at 27°C . The fungal medium was the same as that developed by Field *et al.* (2015b) except for a lower concentration of thiamine (thiamine HCl, $100 \mu\text{g}$). Once fungal outgrowth from plant fragments became visible (1–2 wk), hyphae were subcultured onto the same medium used for isolation and kept in the dark at 27°C (Fig. 1a–c).

Molecular identification of fungal symbionts

Molecular analyses of fungal symbionts of *Lycopodiella* were carried out within 1 wk of collection (Rimington *et al.*, 2015). Extraction and sequencing of DNA were performed using the method of Bidartondo




Fig. 1 *In vitro* isolation of Mucoromycotina fine root endophytes (MFRE) from *Lycopodiella inundata* (a–c) and colonisation of white clover (*Trifolium repens*) by isolated MFRE (d–i). (a) *L. inundata* gametophyte (*) with copious fungal outgrowth, magnified in (b) and imaged under a scanning electron microscope (c); note the fine hyphae with numerous swellings (*). (d, e) Monoxenic culture of *T. repens* and isolated MFRE after 12 wk of culture; (e) note the abundant mycelium extending from plugs of pure MFRE cultures (*) and enveloping the roots (arrowed), enlarged in (f) (see also Fig. 4). (g–i) Trypan blue/ink-stained roots of *Trifolium* showing fine, irregularly branching hyphae with larger vesicles or spores (g, arrowed), forming tightly wound intracellular coils with small intercalary and terminal swellings (h, arrowed); note also the subtending intracellular hyphal ropes (*). (i) Young, arbuscule-like structure (arrowed) forming inside a clover root cell. Bars: (d) 50 mm; (e) 2 mm; (a) 500 μ m; (b, f) 300 μ m; (g) 50 μ m; (c, h, i) 20 μ m.

et al. (2011). The universal fungal 18S primer combination NS1 (White *et al.*, 1990) and EF3 (Smit *et al.*, 1999) was used to amplify DNA which was cloned (Topota; Invitrogen) and sequenced using an Applied Biosystems Genetic Analyser 3730 (Waltham, MA, USA). Between four and eight clones were sequenced for each of eight samples and identified using NCBI BLAST (Altschul *et al.*, 1997). Sequence editing and assembly were performed in GENEIOUS v.5.6 (<http://geneious.com>). The alignment algorithms of MUSCLE were used within MEGA v.5.1 (Tamura *et al.*, 2013), with reference sequences from GenBank. Using UCHIME (Edgar *et al.*, 2011) within MOTHUR (<http://www.mothur.org>), confirmed sequences were not chimeric. Evolutionary models were tested in MEGA. Bayesian inference was carried out using MRBAYES (Huelsenbeck & Ronquist, 2001) and FIGTREE v.1.4 (<http://tree.bio.ed.ac.uk>) for

visualisation and editing. The same method was used for molecular identification of the fungus isolated from *Lycopodiella* and introduced in monoxenic microcosms with clover, as well as that of the fungus colonising the roots of clover in our monoxenic microcosms.

Axenic plants

Clover plants were cultured *in vitro* to establish monoxenic cultures with MFRE isolates. Seeds were surface-sterilised in 70% ethanol for 1 min, rinsed in water, followed by shaking in 5% commercial sodium hypochlorite for 30 min and then thorough rinsing in sterilised dH₂O. Seeds were placed in 140-mm triple-vented sterile Petri dishes containing modified Strullu–Romand (MSR) medium lacking vitamins and sucrose (Declerck *et al.*, 1998) solidified with 0.4%

Phytigel (Sigma-Aldrich), and adjusted to pH 5.5 before sterilisation, where they germinated in the dark, inverted at 27°C, after 3 d.

In vitro colonisation of clover by MFRE isolate

The same MSR medium used for seed germination was plated slanted in 140 mm sterile, triple-vented raised Petri dishes. Three-day-old axenic clover seedlings (one per dish) were placed with their root system adhering to the medium (and the shoot extending into the medium-free portion of the dish) in Petri dishes pre-inoculated (1 wk) with axenic MFRE hyphae. Control plants were placed in noninoculated dishes containing the same medium. The sterile monoxenic microcosms were sealed with Parafilm, and the root system was covered with aluminium foil to prevent photo-oxidation and placed vertically in a growth chamber at 27°C, with a 16-h photoperiod and a photosynthetic photon flux of 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Plates were undisturbed for 12 wk to allow time for the fungus to colonise plant roots. After 12 wk, fungal hyphae were growing in close association with clover roots (Figs 1d–f, 2a); roots were therefore harvested, cleared in 10% KOH and either stained with trypan blue (Hoysted *et al.*, 2019) or 5% ink-vinegar (Vierheilig *et al.*, 1998).

Quantification of fluxes of ^{33}P , ^{15}N and ^{14}C between MFRE and clover

In vitro clover cultures were split into three groups: monoxenic plates colonised by MFRE with hyphae intact and, as controls, monoxenic plates colonised by MFRE with hyphae severed and plates containing only axenic clover without fungus. One hundred microlitres of aqueous solution containing 0.5 MBq ^{33}P -orthophosphate (111 TBq mmol^{-1} SA, 0.15 ng ^{33}P supplied; Hartmann Analytics) and ^{15}N -ammonium chloride (1 mg ml^{-1} ; 0.1 mg ^{15}N added; Sigma-Aldrich) was introduced into a well in each of the plates (Fig. 2a, top-left detail panel). In the plates of clover colonised by MFRE that served as controls, a trench was cut in the medium using a sterile blade and subsequently backfilled with sterile medium to sever hyphae and prevent any direct fungal access to radio- and stable-isotope tracers (Fig. 2a, bottom-left detail panel). As a further control, we included isotope-containing wells in microcosms containing only clover to observe the direct isotope tracer uptake of the asymbiotic plant roots compared with plants that formed mycorrhizal associations with the MFRE isolate. By backfilling trenches with additional medium, diffusion of isotopes from the well was purposefully permitted in all microcosms, allowing us to account for the role of intact MFRE hyphae.




Fig. 2 Set-up of *in vitro* monoxenic experimental system used to quantify fluxes of nutrients exchanged between *Mucoromycotina* fine root endophytes (MFRE) and white clover (*Trifolium repens*). (a) Monoxenic cultures were developed in sterile conditions in 140 mm Petri dishes half-filled with modified Strullu–Romand (MSR) media, which lacked vitamins and sucrose, solidified with 0.4% Phytigel and poured at a slant. Three plugs of axenic MFRE culture were placed on the media and allowed to establish before the introduction of a *T. repens* seedling. Plates were sealed for 12 wk. One hundred microlitres of an aqueous solution containing ^{33}P or ^{15}N was added to a well in each plate (see top-left detail panel). In the control plates, a trench was cut and subsequently filled with sterile medium to sever MFRE and to ensure there was no direct fungal access to radio- and stable-isotope tracers (see bottom-left detail panel). Purple colour on hyphae denotes the flow of fungal-acquired isotope tracers. (b) The MSR media and its contents were sealed off from the aboveground plant tissue, and each plate was placed in a gas-tight chamber before ^{14}C -labelling. Lactic acid was added to ^{14}C -labelled sodium bicarbonate to release a 0.5 MBq pulse of $^{14}\text{CO}_2$ for the plant to fix. At the end of the 24-h labelling period, KOH was added to the system to capture any remaining headspace $^{14}\text{CO}_2$. Red circles in (a, b) denote regions that were sampled for the analysis of ^{33}P , ^{15}N and ^{14}C .

The plates were resealed with Parafilm in sterile conditions and placed in controlled environment chambers (Model no. MicroClima 1200; Snejders Labs, Tilburg, the Netherlands) with 16-h daytime (20°C and 70% humidity) and 8-h night-time (15°C and 70% humidity). Daytime photosynthetically active radiation (PAR), supplied by LED lighting, was 225 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. Atmospheric $[\text{CO}_2]$ was set at 440 $\mu\text{l l}^{-1}$, monitored using a sensor system (Vaisala, Birmingham, UK) and maintained through the addition of gaseous CO_2 . Assimilation of ^{33}P into the aboveground plant material was monitored using a hand-held Geiger counter held over the Petri dish.

For ^{14}C -labelling, the medium containing clover roots and MFRE hyphae was sealed off from the aboveground plant tissue and each plate placed in a gas-tight chamber (Fig. 2b). A 0.5-MBq pulse of $^{14}\text{CO}_2$ was then liberated into the headspace of sealed Petri dishes by adding 2 ml of 90% (w/v) lactic acid to 13.5 μl $\text{NaH}^{14}\text{CO}_3$ (specific activity 2.124 GBq mmol^{-1} ; Hartmann Analytics; Fig. 2b). Cultures inside sealed chambers were maintained under growth conditions as previously mentioned. Twenty-four hours after the addition of ^{33}P and ^{15}N tracers and liberating the pulse of $^{14}\text{CO}_2$, the plates were opened and root-and-shoot material was separated, freeze-dried, weighted and homogenised. For the analysis of ^{33}P , between 0.1 and 4 mg of shoot, root and fungal material was digested in 500 μl of concentrated sulphuric acid at 365°C for 15 min. Fifty microlitres of hydrogen peroxide was added to cooled samples and returned to the digest block (BT5D; Grant Instruments, Cambridge, UK). Cleared digest solutions were then diluted to 5 ml with dH_2O . ^{33}P -radioactivity of plant material was quantified through liquid scintillation (Tri-Carb® 3100TR; PerkinElmer, Beaconsfield, UK). One millilitre of each digest solution was added to 10 ml of Elmusify-safe scintillant, and ^{33}P content was calculated using Eqn 1.

$$M^{33}\text{P} = \left\{ \left[\frac{\text{cDPM}}{\text{SAct}} \right] M_{\text{wt}} \right\} \text{Df} \quad \text{Eqn 1}$$

where $M^{33}\text{P}$, mass of ^{33}P (mg); cDPM, counts as disintegrations per min; SAct, specific activity of the course (Bq mmol^{-1}); Df, dilution factor; and M_{wt} , molecular mass of P (Cameron *et al.*, 2007).

Between 0.1 and 4 mg of freeze-dried, homogenised root-and-shoot tissues were weighed into $6 \times 4 \text{ mm}^2$ tin capsules (Sercon, Crewe, UK), and ^{15}N abundance was determined using a continuous flow infrared mass spectrometry (IRMS; model no. PDZ 2020 IRMS; Sercon) using air as the reference standard; the IRMS detector was regularly calibrated to commercially available reference gases. The ^{15}N transferred from fungus to plant was then calculated using equations published by Field *et al.* (2016).

The ^{14}C activity of shoot, root and fungal samples was quantified through sample oxidation (307 Packard Sample Oxidation, Isotech) followed by liquid scintillation. Total C ($^{12}\text{C} + ^{14}\text{C}$) fixed by the plant and transferred to fungus was calculated as a function of the total volume and CO_2 content of the labelling chamber and the proportion of the supplied $^{14}\text{CO}_2$ label fixed by

plants. As severing the fungal hyphae only impacts the movement of $^{33}\text{P}/^{15}\text{N}$ from the well to the plant, the difference in total C between the values obtained for clover cultures with MFRE and those without fungus is considered equivalent to the total C transferred from plant to symbiotic fungus within the Phytigel for that microcosm, noting that a small proportion will be lost through respiration and accounting for plant-fixed C gained by MFRE via diffusion, root exudation and/or dark fixation. The total C budget for each microcosm was calculated using equations adapted from Cameron *et al.* (2007) (Eqn 2). Total per cent allocation of plant-fixed C to extraradical symbiotic fungal hyphae was calculated by subtracting the activity (in Becquerels) of clover cultures without fungus from that detected in monoxenic cultures with MFRE present, dividing this by the sum of activity detected in all components of each microcosm, then multiplying by 100.

$$M_c = \left(\left(\frac{A}{\text{SAct}} \right) M^{14}\text{C} \right) + (P_r \times M_{\text{wt}_c}) \quad \text{Eqn 2}$$

where M_c , mass of carbon transferred from plant to fungus; A , radioactivity of the tissue sample (Bq); SAct, specific activity of the source (Bq mol^{-1}); $M^{14}\text{C}$, atomic mass of ^{14}C ; P_r , proportion of the total ^{14}C label supplied present in the tissue; M_{wt_c} , mass of C in the CO_2 present in the labelling chamber (g) (from the ideal gas law; Eqn 3):

$$M_{\text{cd}} = M_{\text{cd}} \left(\frac{P V_{\text{cd}}}{RT} \right) \therefore m_{c=m_{\text{cd}}} \times 0.27292 \quad \text{Eqn 3}$$

where m_{cd} , mass of CO_2 (g); M_{cd} , molecular mass of CO_2 (44.01 g mol^{-1}); P , total pressure (kPa); V_{cd} , volume of CO_2 in the chamber (0.003 m^3); R , universal gas constant (J $\text{K}^{-1} \text{mol}^{-1}$); T , absolute temperature (K); m_c , mass of C in the CO_2 present in the labelling chamber (g), where 0.27292 is the proportion of C in CO_2 on a mass fraction basis.

Data analyses

Isotope tracing data were analysed in SPSS Statistics v.26 (IBM, Armonk, NY, USA). Data were tested for normality and homogeneity of variances using the Kolmogorov–Smirnov test for normality. Where assumptions for parametric tests were not met, data were transformed using \log_{10} . If assumptions for parametric tests were still not met, a nonparametric statistical test would be performed. The differences between plant assimilation of ^{33}P and ^{15}N were tested using Mann–Whitney U and Kruskal–Wallis analyses. Whiskers on box plots represent each of the data points (minimum to maximum) recorded during data collection.

Results

Molecular identification of the MFRE fungus

The molecular identification of the fungus colonising the roots of clover *in vitro* was confirmed as the same fungus introduced in our

monoxenic microcosms following isolation from wild gametophytes and young sporophytes of *L. inundata*, all of which matched (99.69%) *Endogonales* sp. GenBank [KJ952213](#) (Rimington *et al.*, 2015).

Axenic culture of plants and fungi and *in vitro* colonisation of clover by MFRE

Axenic MFRE isolates (Fig. 1a) comprised fine hyphal networks with small intercalary and terminal swellings < 20 μm in diameter (Fig. 1b,d). After *c.* 12 wk from the establishment of monoxenic microcosms containing axenic clover seedlings and MFRE hyphae, abundant mycelium extended from the original plugs of pure MFRE cultures (Fig. 1e) and enveloped the roots of clover (Fig. 1f). Root colonisation by MFRE was confirmed by molecular methods and staining (Fig. 1g–i), which revealed copious fungal colonisation with cytology typical of MFRE, consisting of fine (< 1.5 μm in diameter) irregularly branching hyphae with small intercalary and terminal swellings (Fig. 1g,h), sometimes forming tightly

wound coils and hyphal ropes (Fig. 1h), spores or vesicles (Fig. 1g) and fine arbuscule-like structures (Fig. 1i) in the root cortical cells of clover, with arbuscule-like structures confined to the inner cortex.

MFRE directly transfer ^{15}N and ^{33}P to clover

In the experimental microcosms where fungal hyphae remained intact (Fig. 2a), MFRE transferred an average of 48% ($\pm 18\%$) of the supplied ^{33}P tracer and an average of 43% ($\pm 21\%$) of the supplied ^{15}N tracer to the shoots of clover during the labelling period. We determined significant transfer of fungal-acquired ^{33}P to clover by comparing the amount of ^{33}P in shoot tissues where MFRE hyphae remained intact against amounts of ^{33}P in shoot tissue in microcosms where hyphae were severed (Fig. 3a; $P = 0.003$; Table 1) or where MFRE was not present (Fig. 3a; $P = 0.036$; Table 1) in terms of absolute quantities and when normalised to plant biomass (Fig. 3b; $P = 0.057$; Table 1). There was significantly more ^{15}N present in shoots where MFRE hyphae remained intact than in shoot tissue in microcosms where hyphae




Fig. 3 Nutrient fluxes between white clover (*Trifolium repens*) and Mucoromycotina fine root endophytes (MFRE). (a, b) Total shoot tissue phosphorus (^{33}P) in nanograms (ng) in sterile monoxenic cultures with intact fungi, trenched fungi and with no fungi present (a) and shoot concentration (ng g $^{-1}$) of fungal-acquired ^{33}P in *T. repens* (b). (c, d) Total shoot tissue (^{15}N) in nanograms (ng) in sterile monoxenic cultures with intact MFRE fungi (c) and shoot concentrations (ng g $^{-1}$) of fungal-acquired ^{15}N in *T. repens* (d). In (a, b), $n = 9$ for monoxenic cultures with intact fungi and $n = 5$ for those with trenched fungi and $n = 3$ for monoxenic cultures with no fungi present. In (c, d), $n = 16$ for monoxenic cultures with intact fungi and $n = 5$ for those with trenched fungi and $n = 6$ for monoxenic cultures with no fungi present (n indicates the number of biological replicates used during radio- and stable-isotope tracing experiments). Letters denote significant differences where $P < 0.05$, Mann–Whitney *U* and Kruskal–Wallis tests. Error bars represent the standard error of the mean, with all data points shown (minimum to maximum) collected during the experiments.

Table 1 Summary of statistical results of ^{15}N and ^{33}P isotope tracing experiments.

	Intact fungi vs Trenched fungi	Intact fungi vs No fungi	Trenched fungi vs No fungi
Absolute values ^{33}P (ng)			
Mann–Whitney U P -value	0.003	0.036	0.057
Kruskal–Wallis P -value	0.004		
Concentration values ^{33}P (ng g $^{-1}$)			
Mann–Whitney U P -value	0.001	0.009	0.036
Kruskal–Wallis P -value	0.001		
Absolute values ^{15}N (ng)			
Mann–Whitney U P -value	0.040	0.010	0.662
Kruskal–Wallis P -value	0.013		
Concentration values ^{15}N (ng g $^{-1}$)			
Mann–Whitney U P -value	0.349	0.056	1.000
Kruskal–Wallis P -value	0.156		

were severed (Fig. 3c; $P = 0.04$; Table 1) and axenic plates without MFRE (Fig. 3c; $P = 0.01$; Table 1) in terms of absolute quantities. However, when ^{15}N assimilation was normalised to plant biomass, the transfer was not significant (Fig. 3d, severed hyphae, $P = 0.349$; no MFRE present, $P = 0.056$; Table 1).

Plant-fixed carbon is detected in extraradical MFRE hyphae

Based on the drawdown and detection of ^{14}C in plant and fungal materials over the 24-h labelling period, we calculated a complete C budget for each microcosm (Supporting Information Fig. S1). There was significantly more plant-fixed carbon present in microcosms containing MFRE than in plant-only microcosms (Fig. 4a,b; $P = 0.047$). 606.29 ng (440.73 ng g $^{-1}$) of plant-fixed C was present in extraradical MFRE hyphae (Fig. 4a,b), equivalent to 0.87% of the total amount of C fixed by the clover during the 24-h labelling period.

Discussion

Our results provide the first unequivocal demonstration that symbiosis between a flowering plant, white clover, and a MFRE fungus is mutualistic, with the plant gaining both ^{15}N and ^{33}P tracers directly from the fungus while the fungus gains plant-fixed C, in the absence of other micro-organisms. While we focussed on a specific association, when considered together with those of previous studies (Field *et al.*, 2015b, 2016, 2019; Orchard *et al.*, 2017b; Hoysted *et al.*, 2019, 2020; Albornoz *et al.*, 2020, 2021), our findings indicate that MFRE symbioses are nutritionally mutualistic across diverse land plants.

Analysis of the fungus colonising the roots of clover, confirmed molecularly as an Endogonales (Mucoromycotina) isolate from a lycophyte (Hoysted *et al.*, 2019, 2020), revealed a morphology characterised by fine, irregularly branching hyphae with small intercalary and terminal swellings (Fig. 1g,h), forming vesicles or spores (Fig. 1g) as well as hyphal coils and hyphal ropes (Fig. 1h) alongside arbuscule-like structures (Fig. 1i). This morphology matches that described previously in a range of vascular (Orchard *et al.*, 2017b; Hoysted *et al.*, 2019, 2020; Albornoz *et al.*, 2020) and nonvascular plants (Field *et al.*, 2015b, 2016, 2019) colonised by MFRE.




Fig. 4 Total plant-derived carbon present in extraradical Mucoromycotina fine root endophytes (MFRE) fungal hyphae. (a) Total plant-derived carbon present in Phytigel with MFRE fungi present or plant-only microcosms where no MFRE were present after a 24-h labelling period (ng) and concentrations (ng g $^{-1}$) (b). For both (a) and (b), $n = 17$ for microcosms with MFRE present and $n = 7$ for microcosms with no MFRE present. Letters denote significant differences where $P < 0.05$, Mann–Whitney U and Kruskal–Wallis tests. Error bars represent the standard error of the mean, with all data points shown (minimum to maximum) collected during the experiments.

To date, MFRE research has been carried out using unpassured soil culture-based experimental systems (Orchard *et al.*, 2017a; Albornoz *et al.*, 2020) or wild-collected plants (Field

et al., 2015b, 2016, 2019; Hoysted *et al.*, 2020). As such, it is inevitable that these experiments included other soil micro-organisms alongside MFRE. As there is no information about how rhizosphere bacteria may influence MFRE metabolic characteristics and function, the inclusion of soil micro-organisms in previous studies made it impossible to determine the direct contribution of MFRE to host plant nutrition. Here, we show for the first time, using a novel *in vitro* monoxenic system, that MFRE directly assimilate and transfer both ^{33}P and ^{15}N to a flowering plant in the absence of other microbes.

Our results reveal significant transfer of fungal-acquired ^{33}P to clover; however, while we observed a clear trend of greater ^{15}N in plant shoots when MFRE hyphae are intact than where they are severed, the difference is not statistically significant. A possible explanation is that clover, a legume, is not heavily reliant on fungal symbionts for N assimilation, even in the absence of rhizobia. This could also be due to volatilisation and recapture of ammonium by the plant, and/or mass flow driven by plant transpiration in these microcosms. Nevertheless, and although the abundance of MFRE in our microcosms was not quantified, our results, when tentatively compared with those of previous studies on AMF (Thirkell *et al.*, 2020a,b), indicate that clover (without rhizobia) may assimilate more ^{15}N tracer via its MFRE symbiont per unit plant biomass than is typically assimilated by plants associated with AMF, albeit in nonsterile systems. This nutritional role, already indicated by studies of MFRE symbioses in nonflowering plants (Field *et al.*, 2019; Hoysted *et al.*, 2019), could help to explain the persistence of MFRE across most modern land plant lineages, facilitating plant access and assimilation of soil N (Howard *et al.*, 2022). The FRE have long been thought to enhance plant P, at least in soils with very low plant-available P (Crush, 1973; Rabatin *et al.*, 1993; Orchard *et al.*, 2017b; Alborno *et al.*, 2021); however, their potential role in plant N uptake has been overlooked. It is therefore important that N transfer and assimilation from MFRE to plant hosts are now investigated *in vitro* across a range of other flowering plants that do not associate with N-fixing bacteria. Parallel studies of N and P transfer by AMF are also required before meaningful comparisons between AMF and MFRE function can be made.

Because we used an inorganic N source, additional experiments are also needed to assess potential direct organic N utilisation by MFRE (Field *et al.*, 2019). A recent study on the ability of AMF to utilise N from organic sources (Rozmoš *et al.*, 2022) showed, using an *in vitro* monoxenic experimental system based on Ri T-DNA transformed chicory roots, that organic nitrogen utilisation by *Rhizophagus irregularis* was mediated by specific soil bacteria and accelerated by the presence of a protist. These findings, though not based on full plants, may explain the results of previous experiments using organic matter patches labelled with ^{15}N in soil-based microcosms, which showed successful transfer of the ^{15}N by AMF to host plants (Hodge *et al.*, 2000, 2001; Thirkell *et al.*, 2016). Thus, it is likely that AMF-associated and free-living rhizospheric bacteria as well as other soil fungi contained in the nonsterile fungal inoculum used in those experiments may have interacted with AMF (Vivas *et al.*, 2003; Frey-Klett *et al.*, 2007; Smith & Smith, 2011; Jiang *et al.*, 2021),

influencing the breakdown, mineralisation and assimilation of ^{15}N -labelled organic material. It is now important to determine whether similar processes may also explain organic N utilisation by MFRE (Field *et al.*, 2019) or whether these fungi, by virtue of their putative facultatively saprotrophic nature, can directly access and transfer N from organic sources. Since organic N represents a large proportion of total soil N, direct organic N utilisation by MFRE would have important implications for terrestrial N cycling (Hodge & Storer, 2015; Howard *et al.*, 2022).

Further research using monoxenic systems is also needed to compare the 'cost' in terms of plant-to-fungus transfer of C between AMF and MFRE symbioses. Our data show that symbiotic MFRE gain clover-fixed C (Fig. 4a,b), and previous experiments using soil culture-based systems suggested that the 'cost' of MFRE-vascular plant associations is at least on a par with, if not larger than, AMF-vascular plant associations (Hoysted *et al.*, 2020). However, this has only been compared in one vascular plant species; whether it holds true for *in vitro* monoxenic systems remains to be tested.

Previous research into the function of plant–MFRE symbioses raised fundamental questions about their persistence and ecological relevance in modern terrestrial ecosystems. We can now begin to address such questions with new experimental systems knowing that symbiotic MFRE are nutritionally mutualistic with flowering plants. Field *et al.* (2015b) demonstrated the ability of MFRE isolates to recolonise host liverworts *in vitro*; however, until now, this had not been achieved in vascular plants. Our *in vitro* system used a fungal isolate that originated from a wild-collected early-diverging vascular plant, *L. inundata*, and was introduced to white clover *in vitro*. This isolate was molecularly and cytologically confirmed to be colonising the roots of clover used in our experiment. This represents a novel, tractable *in vitro* experimental system designed to manipulate MFRE isolates and the resynthesis of their mycorrhizas with a flowering plant. It opens a realm of exciting possibilities for further research on MFRE mycorrhizal properties, including cytological, molecular and metabolomic comparisons with AMF where host plants are inoculated singly or co-colonised with both MFRE and AMF. Furthermore, a fundamental understanding of how MFRE distribution and function are affected by environmental factors such as temperature, water, light and atmospheric CO_2 , in addition to biotic factors such as interactions with other soil microbiota, can now be developed. Successful isolation from wild plants and axenic cultivation of an MFRE isolate offers exciting new opportunities to develop a model system for symbiotic MFRE and for omics in comparison with other fungi.

Acknowledgements

We gratefully acknowledge support from the European Research Council to KJF and SP (Consolidator Grant MYCOREV, 865225) and NERC to KJF, SP and MIB (NE/S009663/1; NE/N00941X/1). KJF is funded by a BBSRC Translational Fellowship (BB/M026825/1) and a Philip Leverhulme Prize (PLP-2017-079). We thank Juyeon Chae and Agatha Hamon-Pourquery for assistance with culture maintenance and Emily Magkourilou and

Emily Durant for technical assistance during the harvest of experiments.

Competing interests

None declared.

Author contributions

KJF, SP, MIB and GAH conceived and designed the investigation. SP collected the plant material, carried out the isolation of fungal symbionts and designed the monoxenic cultures with the help of GAH. KJF and CAB undertook the isotope tracing. GAH led the data analysis and writing. BS designed Figure 2 and contributed to the writing. All authors discussed the results and commented on the article. SP and KJF agree to serve as the authors responsible for contact and ensure communication.

ORCID

Christopher A. Bell  <https://orcid.org/0000-0002-7437-2793>
Martin I. Bidartondo  <https://orcid.org/0000-0003-3172-3036>
Katie J. Field  <https://orcid.org/0000-0002-5196-2360>
Grace A. Hoysted  <https://orcid.org/0000-0002-3469-2549>
Silvia Pressel  <https://orcid.org/0000-0001-9652-6338>
Besiana Sinanaj  <https://orcid.org/0000-0002-7124-7698>

Data availability

The data generated in this study are provided in the Main Manuscript file. Full datasets are available from the corresponding authors upon reasonable request. Sequence data have been deposited in GenBank with accession no. [KJ952213](https://doi.org/10.26434/chemrxiv-2023-kj952).

References

- Abbott LK. 1982. Comparative anatomy of vesicular arbuscular mycorrhizas formed on subterranean clover. *Australian Journal of Botany* 30: 485–499.
- Albornoz FE, Hayes PE, Orchard S, Clode PL, Nazeri NK, Standish RJ, Bending GD, Hilton S, Ryan MH. 2020. First cryo-scanning electron microscopy images and X-ray microanalyses of Mucoromycotina fine root endophytes in vascular plants. *Frontiers in Microbiology* 11: 2018.
- Albornoz FE, Orchard S, Standish RJ, Dickie IA, Bending G, Hilton S, Lardner T, Foster KJ, Gleeson DB, Bougoure J *et al.* 2021. Evidence for niche differentiation in the environmental responses of co-occurring Mucoromycotina fine root endophytes and Glomeromycotina arbuscular mycorrhizal fungi. *Microbial Ecology* 81: 864–873.
- Ali B. 1969. Occurrence and characteristics of the vesicular arbuscular endophytes of *Nardus stricta*. *Nova Hedwigia* 17: 409–425.
- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25: 3389–3402.
- Bidartondo MI, Read DJ, Trappe JM, Merckx V, Ligrone R, Duckett JG. 2011. The dawn of symbiosis between plants and fungi. *Biological Letters* 7: 574–577.
- Brundrett MC, Tedersoo L. 2018. Evolutionary history of mycorrhizal symbioses and global host plant diversity. *New Phytologist* 220: 1108–1115.
- Cameron DD, Johnson I, Leak JR, Read DJ. 2007. Mycorrhizal acquisition of inorganic phosphorus by the green-leaved terrestrial orchid *Goodyera repens*. *Annals of Botany* 99: 831–834.
- Crush JR. 1973. The effect of *Rhizopagus tenuis* mycorrhizas on ryegrass, cocksfoot and sweet vernal. *New Phytologist* 72: 965–973.
- Declerck S, Strullu DG, Planchette C. 1998. Monoxenic culture of the intraradical forms of *Glomus* sp. isolated from a tropical ecosystem: a proposed methodology for germplasm collection. *Mycologia* 90: 579–585.
- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27: 2194–2000.
- Field KJ, Bidartondo MI, Rimington WR, Hoysted GA, Beerling D, Cameron DD, Duckett JG, Leake JR, Pressel S. 2019. Functional complementarity of ancient plant–fungal mutualisms: contrasting nitrogen, phosphorus and carbon exchange between Mucoromycotina and Glomeromycotina fungal symbionts of liverworts. *New Phytologist* 223: 908–921.
- Field KJ, Pressel S. 2018. Unity in diversity: structural and functional insights into the ancient partnerships between plants and fungi. *New Phytologist* 220: 996–1011.
- Field KJ, Pressel S, Duckett JG, Rimington WR, Bidartondo MI. 2015a. Symbiotic options for the conquest of land. *Trends in Ecology & Evolution* 30: 477–486.
- Field KJ, Rimington WR, Bidartondo MI, Allinson KE, Beerling DJ, Cameron DD, Duckett JG, Leake JR, Pressel S. 2015b. First evidence of mutualism between ancient plant lineages (Haplomitriopsida liverworts) and Mucoromycotina fungi and its response to simulated Palaeozoic changes in atmospheric CO₂. *New Phytologist* 205: 743–756.
- Field KJ, Rimington WR, Bidartondo MI, Allinson KE, Beerling DJ, Cameron DD, Duckett JG, Leake JR, Pressel S. 2016. Functional analysis of liverworts in dual symbiosis with Glomeromycota and Mucoromycotina fungi under a simulated Palaeozoic CO₂ decline. *ISME Journal* 10: 1514–1526.
- Frey-Klett P, Garbaye J, Tarkka M. 2007. The mycorrhiza helper bacteria revisited. *New Phytologist* 176: 22–36.
- Hodge A, Campbell CD, Fitter AH. 2001. An arbuscular mycorrhizal fungus accelerates decomposition and acquires nitrogen directly from organic material. *Nature* 413: 297–299.
- Hodge A, Stewart J, Robinson D, Griffiths BS, Fitter AH. 2000. Competition between roots and soil microorganisms for nutrients from nitrogen-rich patches of varying complexity. *Journal of Ecology* 88: 150–164.
- Hodge A, Storer K. 2015. Arbuscular mycorrhiza and nitrogen: implications for individual plants through to ecosystems. *Plant and Soil* 386: 1–19.
- Howard N, Pressel N, Kaye RS, Daniell TJ, Field KJ. 2022. The potential role of Mucoromycotina ‘fine root endophytes’ in plant nitrogen nutrition. *Physiologia Plantarum* 174: e13715.
- Hoysted GA, Bidartondo MI, Duckett JG, Pressel S, Field KJ. 2020. Phenology and function in lycopod–Mucoromycotina symbiosis. *New Phytologist* 229: 2389–2394.
- Hoysted GA, Jacob A, Kowal J, Giesemann P, Bidartondo MI, Duckett JG, Gebauer G, Rimington WR, Schornack S, Pressel S *et al.* 2019. Mucoromycotina fine root endophyte fungi form nutritional mutualisms with vascular plants. *Plant Physiology* 181: 565–577.
- Hoysted GA, Kowal J, Jacob A, Rimington WR, Duckett JG, Pressel S, Orchard S, Ryan MH, Field KJ, Bidartondo MI. 2018. A mycorrhizal revolution. *Current Opinion in Plant Biology* 44: 1–6.
- Huelsenbeck JP, Ronquist F. 2001. MRBAYES: Bayesian inference and phylogenetic trees. *Bioinformatics* 17: 754–755.
- Jiang F, Zhang L, Zhou J, George TS, Feng G. 2021. Arbuscular mycorrhizal fungi enhance mineralisation of organic phosphorus by carrying bacteria along their extraradical hyphae. *New Phytologist* 230: 304–315.
- Mills BJ, Batterman SA, Field KJ. 2018. Nutrient acquisition by symbiotic fungi governs Palaeozoic climate transition. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 373: 20160503.
- Morris JL, Puttick MN, Clark JW, Edwards D, Kenrick P, Pressel S, Wellman CH, Yang Z, Schneider H, Donoghue PC. 2018. The timescale of early land plant evolution. *Proceedings of the National Academy of Sciences, USA* 115: 2274–2283.
- Orchard S, Standish RJ, Dickie IA, Renton M, Walker C, Moot D, Ryan MH. 2017a. Fine endophytes (*Glomus tenue*) are related to Mucoromycotina, not Glomeromycota. *New Phytologist* 213: 481–486.
- Orchard S, Standish RJ, Dickie IA, Renton M, Walker C, Moot D, Ryan MH. 2017b. Fine root endophytes under scrutiny: a review of the literature on

- arbuscule-producing fungi recently suggested to belong to the Mucoromycotina. *Mycorrhiza* 27: 619–638.
- Pirozynski KA, Malloch DW. 1975. The origin of land plants: a matter of mycotrophism. *Biosystems* 6: 153–164.
- Rabatin SC, Stinner BR, Paoletti MG. 1993. Vesicular-arbuscular mycorrhizal fungi, particularly *Glomus tenue*, in Venezuelan bromeliad epiphytes. *Mycorrhiza* 4: 17–20.
- Raven JA, Allen JF. 2003. Genomics and chloroplast evolution: what did cyanobacteria do for plants? *Genome Biology* 4: 209.
- Read DJ, Duckett JG, Francis R, Ligrone R, Russell A. 2000. Symbiotic fungal associations in 'lower' land plants. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 355: 815–831.
- Rimington WR, Pressel S, Duckett JG, Bidartondo MI. 2015. Fungal associations of basal vascular plants: reopening a closed book? *New Phytologist* 205: 1394–1398.
- Rimington WR, Pressel S, Duckett JG, Field KJ, Bidartondo MI. 2019. Evolution and networks in ancient and widespread symbioses between Mucoromycotina and liverworts. *Mycorrhiza* 29: 551–565.
- Rozmoš M, Bukovská P, Hršelová H, Kotianová M, Dudáš M, Gančarčíková K, Jansa J. 2022. Organic nitrogen utilisation by an arbuscular mycorrhizal fungus is mediated by specific soil bacteria and a protist. *ISME Journal* 16: 676–685.
- Sinanaj B, Hoysted G, Pressel S, Bidartondo MI, Field KJ. 2021. Critical research challenges facing Mucoromycotina 'fine root endophytes'. *New Phytologist Letter* 232: 1528–1534.
- Smit E, Leeflang P, Glandorf B, Van Elsas JD, Wernars K. 1999. Analysis of fungal diversity in the wheat rhizosphere by sequencing of cloned PCR-amplified genes encoding 18 S rRNA and temperature gradient gel electrophoresis. *Applied Environmental Microbiology* 65: 2614–2621.
- Smith SE, Read DJ. 2008. *Mycorrhizal symbiosis*, 3rd edn. London, UK: Academic Press.
- Smith SE, Smith FA. 2011. Roles of arbuscular mycorrhizas in plant nutrition and growth: new paradigms from cellular to ecosystem scales. *Annual Review of Plant Biology* 62: 227–250.
- Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S. 2013. MEGA6: molecular evolutionary genetics analysis v.6.0. *Molecular Biology and Evolution* 30: 2725–2729.
- Taylor L, Banwart S, Leake J, Beerling DJ. 2011. Modelling the evolutionary rise of ectomycorrhiza on sub-surface weathering environments and the geochemical carbon cycle. *American Journal of Science* 311: 369–403.
- Thippayrugs S, Bansal M, Abbott LK. 1999. Morphology and infectivity of fine endophyte in a Mediterranean environment. *Mycological Research* 103: 1369–1379.
- Thirkell TJ, Cameron DD, Hodge A. 2016. Resolving the 'nitrogen paradox' of arbuscular mycorrhizas: fertilisation with organic matter brings considerable benefits for plant nutrition and growth. *Plant, Cell & Environment* 39: 1683–1690.
- Thirkell TJ, Campbell M, Driver J, Pastok D, Merry B, Field KJ. 2020a. Cultivar-dependent increases in mycorrhizal nutrient acquisition by barley in response to elevated CO₂. *Plants, People, Planet* 3: 553–566.
- Thirkell TJ, Pastok D, Field KJ. 2020b. Carbon for nutrient exchange between arbuscular mycorrhizal fungi and wheat varies according to cultivar and changes in atmospheric carbon dioxide concentration. *Global Change Biology* 26: 1725–1738.
- Vierheilig H, Coughlan AP, Wyss URS, Piché Y. 1998. Ink and vinegar, a simple staining technique for arbuscular-mycorrhizal fungi. *Applied and Environmental Microbiology* 64: 5004–5007.
- Vivas A, Marulanda A, Ruiz-Lozano JM, Barea JM, Azcón R. 2003. Influence of a *Bacillus* sp. on physiological activities of two arbuscular mycorrhizal fungi and on plant responses to PEG-induced drought stress. *Mycorrhiza* 13: 249–256.
- Walker C, Gollotte A, Redecker D. 2018. A new genus, *Planticonsortium* (Mucoromycotina), and new combination (*P. tenue*), for the fine root endophyte, *Glomus tenue* (basonym *Rhizophagus tenuis*). *Mycorrhiza* 28: 213–219.
- Wang X, Fang J, Liu P, Liu J, Fang W, Fang Z, Xiao Y. 2021. Mucoromycotina fungi possess the ability to utilize plant sucrose as a carbon source: evidence from *Gongronella* sp. w5. *Frontiers in Microbiology* 11: 3322.
- White TJ, Bruns TD, Lee S, Taylor J. 1990. Analysis of phylogenetic relationships by amplification and direct sequencing of ribosomal RNA genes. In: Innis MA, Gelfand JJ, Sninsky JJ, White TJ, eds. *PCR protocols: a guide to methods and applications*. New York, NY, USA: Academic Press, 315–322.

Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

Fig. S1 Total carbon flux budget for monoxenic *in vitro* cultures of Mucoromycotina fine root endophyte hyphae (MFRE) colonising white clover (*Trifolium repens*).

Please note: Wiley is not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.