



This is a repository copy of *A deep dive into seagrass Rubisco catalytic properties uncovers a distinct evolutionary trajectory*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/195550/>

Version: Publishers draft (with formatting)

---

**Article:**

Degen, G.E. [orcid.org/0000-0002-0804-4169](https://orcid.org/0000-0002-0804-4169) (2022) A deep dive into seagrass Rubisco catalytic properties uncovers a distinct evolutionary trajectory. *Plant Physiology*. ISSN 0032-0889

<https://doi.org/10.1093/plphys/kiac546>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# A deep dive into seagrass Rubisco catalytic properties uncovers a distinct evolutionary trajectory

Gustaf E. Degen \*

School of Biosciences, University of Sheffield, S10 2TN, UK

\*Author for correspondence: [g.degen@sheffield.ac.uk](mailto:g.degen@sheffield.ac.uk)

Ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco) is the most abundant and arguably most important enzyme on Earth, as it is the entry point of carbon into the biosphere. CO<sub>2</sub> fixation by Rubisco onto ribulose-1,5 bisphosphate (RuBP) is the first step in the Calvin–Benson–Bassham cycle, resulting in two molecules of 3-phosphoglycerate (3PGA), which ultimately make up the biomass of photosynthetic organisms. Efficient CO<sub>2</sub> fixation is hindered by the inability of Rubisco to discriminate between CO<sub>2</sub> and O<sub>2</sub>. Oxygenation of RuBP leads to the production of toxic 2-phosphoglycolate, which is recycled back to 3PGA via photorespiration, consuming energy and releasing CO<sub>2</sub> in the process (Sage et al., 2012). In agriculturally important plants such as wheat (*Triticum aestivum*) or rice (*Oryza sativa*), photorespiration can cause losses greater than 25% of the CO<sub>2</sub> assimilated by photosynthesis (Walker et al., 2016), making Rubisco a key target for improving crop photosynthesis.


Rubisco activity and efficiency are measured by determining multiple parameters. The carboxylation velocity ( $V_c$ ) refers to the CO<sub>2</sub>-fixation activity per leaf area and is arguably the most important parameter. The turnover number ( $k_{cat}$ ) is the amount of product per active site per second. The Michaelis–Menten constants ( $K_m$ ) for CO<sub>2</sub> and O<sub>2</sub> ( $K_c$ ,  $K_o$ ) refer to the affinity of Rubisco to CO<sub>2</sub> and O<sub>2</sub>, respectively. A higher  $K_m$  means that more substrate is required to reach half of the maximum enzyme velocity, suggesting a lower affinity of the enzyme for the substrate. Finally, the specificity factor for CO<sub>2</sub> and O<sub>2</sub> ( $S_{C/O}$ ) is a measurement of the ability of Rubisco to discriminate between CO<sub>2</sub> and O<sub>2</sub>, where a higher value indicates that Rubisco binds more CO<sub>2</sub> relative to O<sub>2</sub>.

Surveying Rubisco kinetic diversity has received much attention in recent years. Analysis of Rubisco catalytic properties between the C3 plant wheat and its wild relatives uncovered a large diversity in Rubisco efficiencies (Prins et al., 2016). Comparing C3 plants to C4 plants, which evolved a carbon-concentrating mechanism (CCM) around 25 million years ago (mya; Sage et al., 2012), showed that  $k_{cat}^C$  in C4 plants was superior, whereas  $S_{C/O}$  was lower (Sage, 2002; Orr et al., 2016; Iñiguez et al., 2020). Due to the enrichment of CO<sub>2</sub> in C4 plants, the CO<sub>2</sub> specificity can be lower, allowing for higher rates of catalysis. This shows the impact CCMs have on the catalytic adaption of Rubisco.

To further the understanding of Rubisco evolution, marine environments are useful because of the slow diffusion of CO<sub>2</sub>. The limited supply of CO<sub>2</sub> to Rubisco in marine plants, such as seagrasses (*Posidonia*), has also resulted in the invention of CCMs, such as bicarbonate transporters. However, little is known about the evolutionary trajectory of catalytic properties in seagrasses, which colonized the oceans some 100 mya.

In this issue of *Plant Physiology*, Capó-Bauçà et al. (2022) investigated the kinetic properties of Rubisco in seagrass species from the Mediterranean and seas around Australia, revealing that these follow a different evolutionary trajectory compared to Rubiscos found in terrestrial angiosperms.

In the first set of experiments, the authors characterized Rubisco affinity for CO<sub>2</sub> and showed that  $K_c$  was much higher for the Mediterranean seagrass species, meaning that it had a lower affinity for CO<sub>2</sub> than the Australian species. Interestingly, the  $K_o$  of all seagrass species was much higher than those reported for other angiosperms, diverging from the linear relationship between  $K_c$  and  $K_o$  observed in the



**Figure 1** Australian and Mediterranean *posidonia* (seagrass) Ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco) catalytic properties. Australian and Mediterranean species diverged 68 mya. Analysis of the large Rubisco subunit (RbcL) and Rubisco properties revealed that the *Posidonia oceanica* species had two amino acid changes that contributed to a higher Michaelis–Menten constant for CO<sub>2</sub> ( $K_C$ ). Furthermore, only two species were found to have an effective CCM due to adaptation to their environment. Analyzing the relationship between  $K_C$  and turnover number for CO<sub>2</sub> ( $k_{cat}^C$ ) showed a distinct evolutionary trajectory of seagrasses, in contrast to land plants. Background image adapted from ‘*Posidonia oceanica* observed at La Ciotat’ by Frédéric Ducarme is licensed under CC BY-SA 4.0. Other figures adapted from Capó-Bauçà et al. (2022).

terrestrial plant Rubiscos. Surprisingly, the slower turnover rate in seagrasses was also accompanied by lower CO<sub>2</sub> affinity, in contrast to land plant Rubiscos. However, the O<sub>2</sub> affinities in seagrasses were also lower, which balances out the high  $K_C$ . Similar to other Rubiscos found in cyanobacteria or red algae (Shih et al., 2016), the authors concluded that seagrass Rubisco catalytic properties did not follow the same coevolutionary trend as terrestrial angiosperms.

This insight prompted the authors to investigate the amino acid sequence of the large Rubisco subunit (RbcL), which hosts the active site of Rubisco. They identified five amino acid changes in the Mediterranean species, which diverged

68 mya from the Australian relatives. Considering the location of these residues in the holoenzyme, the authors identified positions 279 and 449 as potentially accounting for the lower CO<sub>2</sub> affinity in the Mediterranean species. This also revealed that all Australian species shared the same RbcL sequence, suggesting that the observed low  $k_{cat}^C$  of one of these species is likely due to sequence differences in the small Rubisco subunit.

Not only did the seagrass species differ in their Rubisco catalytic properties but they also varied in the effectiveness of their CCMs. This was shown when using Tris-buffered seawater where photosynthetic activity depended on the efficiency of the CCM. In fact, only two out of the five species studied had an effective CCM, suggesting adaptation to the natural environment where light levels may influence the rate of energy production used to fuel CCMs. Thus, in some cases, an effective CCM is required, whereas for plants growing in deeper water, lower light, and higher CO<sub>2</sub> partial pressure, a less effective CCM is advantageous.

Overall, Capó-Bauçà et al. (2022) show that the evolution of seagrass Rubiscos in the ocean lead to different but advantageous changes in catalytic properties that are distinct from terrestrial angiosperm relatives (summarised in Figure 1). Their findings add to the growing body of work on Rubisco kinetic diversity, which is a valuable tool to improve Rubisco properties.

*Conflict of interest statement.* None declared.

## References

- Capó-Bauçà S, Whitney S, Iñiguez C, Serrano O, Rhodes T, Galmés J (2022) The trajectory in catalytic evolution of Rubisco in *Posidonia* seagrass species differs from terrestrial plants. *Plant Physiol* <http://doi.org/10.1093/plphys/kiac492>
- Iñiguez C, Capó-Bauçà S, Niinemets Ü, Stoll H, Aguiló-Nicolau P, Galmés J (2020) Evolutionary trends in RuBisCO kinetics and their co-evolution with CO<sub>2</sub> concentrating mechanisms. *Plant J* **101**(4): 897–918
- Orr DJ, Alcántara A, Kapralov MV, Andralojc PJ, Carmo-Silva E, Parry MAJ (2016) Surveying Rubisco diversity and temperature response to improve crop photosynthetic efficiency. *Plant Physiol* **172**(2): 707–717
- Prins A, Orr DJ, Andralojc PJ, Reynolds MP, Carmo-Silva E, Parry MAJ (2016) Rubisco catalytic properties of wild and domesticated relatives provide scope for improving wheat photosynthesis. *J Exp Bot* **67**(6): 1827–1838
- Sage RF (2002) Variation in the  $k_{cat}$  of Rubisco in C(3) and C(4) plants and some implications for photosynthetic performance at high and low temperature. *J Exp Bot* **53**(369): 609–620
- Sage RF, Sage TL, Kocacinar F (2012) Photorespiration and the evolution of C4 photosynthesis. *Ann Rev Plant Biol* **63**(1): 19–47
- Shih PM, Occhialini A, Cameron JC, Andralojc PJ, Parry MAJ, Kerkfeld CA (2016) Biochemical characterization of predicted Precambrian RuBisCO. *Nat Commun* **7**(1): 10382
- Walker BJ, VanLoocke A, Bernacchi CJ, Ort DR (2016) The costs of photorespiration to food production now and in the future. *Ann Rev Plant Biol* **67**(1): 107–129