



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/196567/>

Version: Accepted Version

Article:

Xiao, K-Q, Zhao, Y, Liang, C et al. (2023) Introducing the soil mineral carbon pump. Nature Reviews Earth & Environment, 4. pp. 135-136. ISSN: 2662-138X

<https://doi.org/10.1038/s43017-023-00396-y>

This article is protected by copyright. This is an author produced version of an article published in Nature Reviews Earth & Environment. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Introducing the soil mineral carbon pump

Ke-Qing Xiao^{1,†}, Yao Zhao², Chao Liang³, Mingyu Zhao⁴, Oliver W. Moore¹, Alba Otero-Fariña¹,
Yong-Guan Zhu^{5,6,7,†}, Karen Johnson⁸, Caroline L. Peacock¹

¹ School of Earth and Environment, University of Leeds, Leeds, UK

² State Key Laboratory of Environmental Criteria and Risk Assessment, Chinese Research Academy of Environmental Sciences, Beijing, China

³ Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang, China

⁴ Key Laboratory of Cenozoic Geology and Environment, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing, China

⁵ Key Laboratory of Urban Environment and Health, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen, China

⁶ State Key Laboratory of Environmental Chemistry and Ecotoxicology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, China

⁷ Zhejiang Key Laboratory of Urban Environmental Processes and Pollution Control, CAS Haixi Industrial Technology Innovation Center in Beilun, Ningbo, China

⁸ School of Engineering and Computing Sciences, Durham University, Durham, UK

†e-mail: k.q.xiao@leeds.ac.uk, ygzhu@rcees.ac.cn

Standfirst Microorganisms and minerals both contribute to organic carbon preservation and accumulation in soil. The soil microbial carbon pump describes the microbial processes, but a separate soil mineral carbon pump needs to be acknowledged and investigated.

[H1] Introduction

Soil impacts climate through the sequestration or release of carbon, which is impacted by soil organic matter formation¹. The role of microbes in soil organic carbon (OC) production and sequestration is described by the soil microbial carbon pump (MCP) concept². In this model, new organic compounds are produced through microbial anabolism, and are subsequently stabilized on mineral surfaces and within soil structures by the entombing effect. Although mineral-organic carbon associations are traditionally assumed to be protective, emerging evidence suggests these interactions are complicated and include numerous abiotic reactions not considered in the MCP. Here, we propose a distinct soil mineral carbon pump (MnCP) that works in parallel to the MCP.

[H1] The soil mineral carbon pump

The MnCP describes how soil minerals enhance the persistence and accumulation of OC. Soil minerals can transform plant or microorganism-derived labile OC into more stable forms through processes such as adsorption, occlusion, aggregation, redox reactions, and polymerization (Fig. 1). Adsorption, occlusion and aggregation can reduce the availability of OC by lowering its concentration in the dissolved pool, forming organo–minerals that are too large to be ingested by microbes and/or limiting the functioning of hydrolytic enzymes⁴. Clay minerals (kaolinite, montmorillonite) and metal (oxyhydr)oxides (iron oxides, birnessite) can drive polymerization, producing more recalcitrant OC (ref.³). Furthermore, redox reactions at mineral surfaces drive OC oxidation to CO₂ and can produce radicalized OC that can be complexed into larger molecules⁵.

The MnCP operates in various soil environments, potentially with a key role in OC sequestration in mineral soils. This pump is sustainable and can operate over long periods: there are abundant clays and metal (oxyhydr)oxides in soil to associate with labile OC, protecting it from microbial degradation⁴. Moreover, fluctuating redox conditions in soil can recycle reactive minerals that catalyze polymerization⁶, especially in environments like paddy soils and peatlands.

[H1] Relationship to the soil MCP

The initial adsorption of OC with minerals is usually followed by reactions that are distinct from those in the soil MCP; these reactions are included in the soil MnCP, as is mineral-catalyzed abiotic polymerization. However, the MnCP and MCP are not mutually exclusive. As exemplified by the

entombing effect in the MCP (ref.²), they work synergistically to preserve OC in soil. For example, redox reactions on OC-minerals interfaces can be either abiotic or biotic⁷, and the MnCP and MCP can be coupled through mineral–microbe interactions during minerals dissolution and electron transfer. Moreover, the extracellular polymeric substances excreted by microorganisms (as part of the MCP) can strengthen the aggregation of organic matter and minerals via a gluing effect⁸ (part of the MnCP) in soil.

[H1] Broader impacts

Minerals can be as important as microorganisms in increasing soil OC persistence and accumulation. The MnCP highlights this role and provides a framework for research into mineral–OC interactions, at fine scales and within the broader soil ecosystem. This work is needed to understand the complexity of soil OC (ref.⁹), and why some OC persists while other OC does not.

Acknowledging the suite of natural reactions that occur through the MnCP will be especially helpful in guiding analytical soil research. For example, the ability of minerals to catalyze polymerization of OC has been mostly found in laboratory settings³, but its occurrence in the soil and its broader impact on OC structure and persistence are unclear. Polymerization of OC can produce numerous new and larger molecules that are hard to identify analytically. Therefore, emerging new techniques (such as FT–ICR–MS, NEXAFS, and Nano–SIMS) should be used and combined to understand how the MnCP operates under different OC characteristics and mineralogy, and to find fingerprints of polymerized OC that might be used to identify its production and persistence in soils. Experimental work to demonstrate that soil mineral-catalyzed polymerization increases OC stability is also needed.

Interest in enhancing soil carbon sequestration to combat climate change is growing, so it is increasingly important to understand the mechanisms underlying OC preservation. Fostering and strengthening the MnCP in soil could be a part of this effort, analogously to enhanced silicate weathering methods¹⁰, providing another potential tool to stabilize carbon in the soil.

References

1. Davidson, E. A. & Janssens, I. A. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature*. **440**, 165–173 (2006).
2. Liang, C., Schimel, J. P. & Jastrow, J. D. The importance of anabolism in microbial control over soil carbon storage. *Nat. Microbiol.* **2**, 17105 (2017).
3. Kleber, M. et al. Dynamic interactions at the mineral–organic matter interface. *Nat. Rev. Earth Environ.* **2**, 402–421 (2021).
4. Keil, R. & Mayer, L. in *Treatise on geochemistry* (2014).
5. Johnson, K. et al. Towards a mechanistic understanding of carbon stabilization in manganese oxides. *Nat. Commun.* **6**, 7628 (2015).
6. Lovley, D. R. Dissimilatory Fe(III) and Mn(IV) reduction. *Microbiol. Rev.* **55**, 259 (1991).
7. Dong, H. et al. A critical review of mineral-microbe interaction and co-evolution: mechanisms and applications. *Natl. Sci. Rev.* **9**, nwac128(2022).
8. Zhang, M. et al. Selective retention of extracellular polymeric substances induced by adsorption to and coprecipitation with ferrihydrite. *Geochim. Cosmochim. Acta*. **299**, 15–34 (2021).
9. Lehmann, J. et al. Persistence of soil organic carbon caused by functional complexity. *Nat. Geosci.* **13**, 529–534 (2020).
10. Beerling, D. J. et al. Potential for large-scale CO₂ removal via enhanced rock weathering with croplands. *Nature*. **583**, 242–248 (2020).

Acknowledgments

This research project has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (Grant agreement No. 725613 MinOrg, C.L.P.) and the Strategic Priority Research Program of Chinese Academy of Sciences

107 (XDA28020103 and XDA28020101). M. Z is funded by the 100 Talents program of the Chinese
108 Academy of Sciences (E251520401).

109 **Competing interests**

110 The authors declare no competing interests.

111

112 **Fig. 1: The soil mineral and microbial carbon pumps.** **a|** Minerals enhance the persistence and
113 accumulation of organic carbon in soils in the soil mineral carbon pump (MnCP). Organic substrates
114 from plants and microorganisms are stabilized via processes such as adsorption, occlusion,
115 aggregation, redox reaction, and polymerization. **b|** The microbial carbon pump (MCP) is on the right
116 for comparison. Panel b is adapted from ref ², Springer Nature Ltd.

117