



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/197998/>

Version: Supplemental Material

Article:

Scott, N, Cairns, A, Prasad, P et al. (2023) Resection margin involvement after endoscopic excision of malignant colorectal polyps: definition of margin involvement and its impact upon tumour recurrence. *Histopathology*, 83 (1). pp. 80-90. ISSN: 0309-0167

<https://doi.org/10.1111/his.14903>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Table 2. Relationship between number of pathological risk factors present in the polypectomy specimen, performance of salvage surgery and residual tumour in surgical specimen.

No. of Pathological Risk Factors	Salvage Surgery	Tumour present in bowel wall	Lymph Node metastasis	Any Tumour in specimen
0	3/46 (6.5%)	0	0	0
1	25/75 (33.3%)	5 (20%)	2 (8%)	6 (24%)
2	22/33 (66.6%)	3 (13.6%)	5 (22.7%)	6 (27%)
3	4/4 (100%)	0	1 (25%)	1 (25%)
All Cases	54/158 (34.2%)	8 (14.8 %)	8 (14.8%)	13 (24.1%)