

Development, sensitivity and reliability of a French version of the Leeds Food Preference Questionnaire (LFPQ-fr) for the evaluation of food preferences and reward

Thivel D¹, Oustric P², Beaulieu K², Moore H¹, Bonjean L¹, Loglisci J^{3,4}, Georges M^{3,4}, Miyashita M⁵, Boirie Y⁶, Pereira B⁷, Finlayson G²

¹Laboratory of the Metabolic Adaptations to Exercise under Physiological and Pathological Conditions (AME2P), Clermont Auvergne University, Clermont-Ferrand, France.

²Appetite Control Energy Balance Group, School of Psychology, University of Leeds, Leeds LS2 9JT, UK.

³Department of Pulmonary Medicine and Intensive Care Unit, Constitutive Reference Center for Rare Pulmonary Diseases, University Hospital, Bourgogne-Franche-Comté, Burgundy University, Dijon, France

⁴Centre des Sciences du Goût et de l'Alimentation, AgroSup Dijon, CNRS UMR6265, INRAE UMR 1324, Université de Bourgogne Franche Comté, Dijon, France

⁵Faculty of Sport Sciences, Waseda University, Saitama 359-1192, Japan

⁶CSO-CALORIS, CHU Clermont-Ferrand, Clermont-Ferrand, France; Department of Human Nutrition, CHU Clermont-Ferrand, Clermont-Ferrand, France. Electronic address: yves.boirie@uca.fr

⁷Biostatistics Unit, DRCI, CHU Clermont-Ferrand, Clermont-Ferrand, France. Electronic address: bpereira@chu-clermontferrand.fr.

Corresponding authors

David Thivel, AME2P Laboratory, CRNH Auvergne, Clermont Auvergne University. Clermont-Ferrand, France.

David.thivel@uca.fr

Abstract

Background. There is a growing global interest in the evaluation of food reward, necessitating the adaptation of culturally appropriate instruments for use in empirical studies. This work presents the development and validation of a culturally adapted French version of the Leeds
40 Food Preference Questionnaire (LFPQ-fr).

Methods. The LFPQ-fr was developed and validated in healthy-weight adults using the following systematic approach: i) selection and validation of appropriate food pictures; ii) linguistic translation of liking and wanting constructs in the target population (n=430; 81% female; 42.2 ± 12.7 years); iii) validation of the sensitivity and reliability of the task performed
45 in a fasted state and in response to a standardized test meal (n=50; 50% female; 30.0 ± 8.4 years).

Results. During the first and second phases, the nutritional and perceptual validation of culturally appropriate food pictures and pertinent reward constructs, respectively, was demonstrated in a healthy-weight French sample. Findings from the third phase indicated that
50 all food reward components were sensitive to the test meal and showed moderate to high agreement in both fasted (Lin's CCC = .72-.94) and fed (Lin's CCC = .53-.80) appetitive states between visit 1 (V1) and visit (V2). Except for explicit liking fat bias, all primary outcomes were statistically consistent in fasted and fed states between V1 and V2. Changes in fat and taste biases in response to a standardized meal for all primary outcomes were also consistent
55 between V1 and V2 except for explicit liking fat bias (Lin's CCC = .49- .72).

Conclusion. The LFPQ-fr developed and tested in this study is a reproducible and reliable method to assess food reward in both the fasted and fed states in a healthy-weight French population.

60 **Key words.** Food reward, Food liking, Food wanting, Food preferences

Introduction

65 There has been an increasing interest in the study of appetite control over the last decade as
a potential vector for the regulation of body weight, and improvement of adherence to various
types of intervention. Whilst long considered an entirely physiological process, the regulation
of appetite is now understood to be determined by a complex integration of internal and
external multi-sensory signals processed centrally by homeostatic and hedonic systems of the
70 brain [1]. The homeostatic system has been extensively described and conceptualized as a
network of peripheral-central crosstalk conveying tonic and episodic signals modulating
hunger and satiety [2]. Simultaneously, the neural networks that operate as part of the
hedonic (i.e., reward-related) system are also sensitive to our external food environment and
endogenous processes [3]. Food reward may be characterized by two distinct yet integrated
75 neurobehavioral feedback mechanisms that constitute the expression of appetite: Liking and
Wanting [4]. Wanting, also referred to as Incentive Salience, represents the motivational
component of reward, typically measured as the potential to elicit/reinforce consummatory
behaviors. Conversely, Liking represents the affective component that is observed by the
pleasure experienced upon consumption of a palatable food [5]. With the prevalence of
80 obesity, eating disorders, and preoccupation with weight increasing in recent years, accurately
and reliably measuring food reward may contribute to the design of interventions that
improve appetite regulation [6].

While several methods have been proposed to assess an individual's food reward, the Leeds
Food Preference Questionnaire (LFPQ) has been proposed and developed as a valid,
85 systematic way to evaluate both liking and wanting for foods that vary on key nutritional and
sensory properties. Briefly, the LFPQ is a computer-based platform that operationalizes these
separable components through explicit assessment of liking and wanting for individual,
familiar foods, and implicit assessment of wanting during simulated food choices [7]. The LFPQ
has been shown to be sensitive to individual differences in eating behavior traits [8,9] and
90 states of hunger and satiety in response to a meal or preload [11,12]. Importantly, it has been
validated as a reliable predictor of actual food choice and intake in both laboratory and free-
living settings [4,10].

However, the LFPQ was developed in the context of British culinary culture, thus its utility in
assessing food reward and preferences among individuals from other cultures may be limited.

95 Cross-cultural studies have consistently shown that people indicate a stronger preference for
foods that are familiar to their own culture compared with foreign foods [13,14]. Indeed,
recent cross-cultural studies have clearly shown different neural, behavioral, perceptual and
sensory responses to culturally consistent and foreign meal types (e.g., classical western
versus non-western meals) [15-23]. While the construct validity of the LFPQ and its approach
100 to measuring food reward has been well established, cultural adaptations based on
population-specific food habits and identities may facilitate empirical research of appetite
control in a culturally diverse world.

To date, the LFPQ has been translated into about 20 languages (e.g., Japanese, Portuguese,
Chinese and Danish [24-27]), but relatively few iterations have followed a uniform and
105 systematic procedure to develop and implement cultural adaptations that would promote its
reliable application in empirical research worldwide [28].

Although studies have evaluated food reward in French adults and children with healthy-
weight, overweight or obesity using a French-language version of the original LFPQ [28], a
rigorous, systematic procedure of cultural adaptation with appropriate food imagery has yet
110 to be conducted for a French population. Therefore, the present work developed a cultural
adaptation of the LFPQ for a French population in accordance with the systematic procedure
outlined by Oustric and colleagues [28].

The reliability of the LFPQ-fr task would be supported by demonstration of consistent liking
and wanting for foods based on fat content (high versus low fat) and taste (sweet versus
115 savory) in the fasted state and after a fixed test meal during both lab visits (V1 and V2).
Moreover, validation of the task would be demonstrated by a general preference for low-fat
and savory foods in fasted, healthy-weight subjects and a shift towards bias for high fat and
sweet foods in response to a savory, low fat meal similar to results from studies using the
original British version [4].

120 **Methods**

The LFPQ-fr was developed and validated following a recommended procedure previously
detailed [28]. First, a culturally appropriate database of food images was selected and
validated based on familiarity, liking, intake frequency, and recognition of nutritional or
sensory properties by a sample of healthy-weight French adults. Second, a French translation

125 of the liking and wanting constructs was developed and tested for recognition in the same
population. Finally, the LFPQ-fr, comprised of these validated components, was then tested
for its sensitivity to different appetitive states (i.e., fasted versus after intake of a standardized
meal) and reliability over two lab visits on different test days in subset of the initial sample.
This study received an ethical agreement (CPP Ile de France III, 2019-A00853-54) and has been
130 registered as a clinical trial (clinicaltrial.gov, NCT04041830).

i) Food database creation

To create a food database appropriate for the French culture, 36 common French foods were
selected, prepared and photographed. The food database was pilot tested to contain: 1) ready
to eat and familiar foods within the French culture, 2) an acceptable number of food items to
135 complete the LFPQ tasks, 3) a typical portion size for each food, 4) foods appropriate to the
time of day of the measurements, 5) foods appropriate to the culture and habits of the French
population, 6) a diversity of foods within each category, and 7) food images of homogenous
quality and size.

Selected food images in the database were then allocated to an appropriate category based
140 on two nutrient- and sensory-based criteria: fat content and taste. Fat content was organized
dichotomously into high fat and low fat categories, with high fat foods deriving more than 40%
of their energy content from fat and low fat foods less than 20%. Foods were further
categorized based on a distinct sweet or savory taste, resulting in four eligible food categories:
High Fat Savory (HFSA), High Fat Sweet (HFSW), Low Fat Savory (LFSA), and Low Fat Sweet
145 (LFSW). To account for the effect of other macronutrients on appetite, protein and
carbohydrate levels were matched between high and low fat categories. Therefore, the savory
foods had high protein and low carbohydrate levels relative to the sweet foods within each
fat category (Table 1).

ii) Food database validation

150 An online questionnaire was created and disseminated by email to 3 areas in France (Dijon,
Clermont-Ferrand and Paris) to undertake the validation of the food database. These 3 areas
were either university or clinical sites but most of the respondents were working professionals
(e.g., researchers, doctors, technicians, nurses, or secretaries). Respondents that did not live
in France (n=7), were nutritionists/dieticians (n=3), or reported allergies to any of the foods

155 presented (n=10) were excluded from analyses. Among the 570 respondents that completed the questionnaire, 430 had responses to all the food questions, thus were included in analyses. The survey aimed to identify food pictures that were familiar, frequently eaten, liked, accurately perceived as sweet or savory, and low or high in fat, and consistent with lunchtime consumption. Indeed, food image familiarity, recognizability, and quality must be tested and
160 accounted for during validation due to the potential to confound liking and wanting scores [28]. To identify suitable food images, each image was presented independently, and the following eight questions were asked: Name: "Please name the food image?"; Frequency: "How often do you consume this food?" (5-point Likert scale, never to almost every day); Liking: "How pleasant does this food typically taste?" (100 mm-scale); Food categories: "Is this
165 food more sweet or savory?" (100 mm-scale); "Is this food low or high in fat?" (100 mm-scale); Time appropriateness: "How appropriate is it to consume this food in the morning/afternoon/evening?" (100 mm-scale).

The French translation of LFPQ task-specific instructions was conducted, discussed among various French native speakers, and comprehension thereof pilot tested using the online
170 survey. The food database was created in collaboration with researchers from Leeds, Dijon and Clermont universities.

Food images were considered appropriate if they satisfied the following conditions: 1) correctly recognized (defined by > 80% of the participants naming the food correctly); 2) habitually consumed (defined by a mean score greater than 2.5 out of 5, which was considered
175 to be representative of frequent contact with the food); 3) liked (defined by mean liking >70mm), 4) correctly recognized as sweet versus savory and low fat versus high fat (sweet and low fat if the mean value was < 40 mm and savory and high fat if > 60 mm, respectively); and 5) time appropriate (defined by mean >70mm for the specific time of day).

iii) Validation of the sensitivity and reliability of the LFPQ-fr

180 Population. A total of 50 healthy-weight adults (50% female) aged between 20-40 years old took part in this validation phase. Respondents first had to pass a medical screening performed by a nutritionist physician in order to confirm their eligibility and ability to complete the study. Mainly, the nutritionist confirmed the absence of eating disorders or chronic eating-related pathologies based on its interview and on the Dutch Eating Behavior

185 Questionnaire (DEBQ) [29] and Three Factors Eating Questionnaire (TFEQ) [31] and the Binge
Eating Scale (BES) [30]. Although not clear cut-offs are actually determined, the following have
been used for the TFEQ: restraint behaviors with a score > 10; disinhibition score > 8 and
susceptibility for hunger score > 7; for the DEBQ: Emotional, Restrained and External eating
scores >2.5 and for BES: score >10. Moreover, participants must have been weight-stable and
190 not undertaken any diet with an aim to lose weight within the last 6 months.

Design. Eligible participants completed anthropometric measurements and body composition
using the Tanita MC780 bio-impedance meter. They were then asked to visit the laboratory
on two occasions separated by at least 7 days. On both occasions, the participants were asked
to refrain from any structured physical activity over the preceding 48 hours and consume the
195 same prescribed breakfast at 08:00 am on each test day. The breakfast was individually
composed based on the participants eating habits but represented 9.5 to 10 kcal per kg of
body mass, respecting the appropriate nutritional recommendations (55% CHO, 30% lipids
and 15% protein). Importantly, all the participants were usual breakfast consumer. During
each test day, participants visited the laboratory at 11:00 am and compliance was checked by
200 an experimenter. At 11:30 am, they were asked to perform the computerized LFPQ-fr task,
which lasted about 15 minutes, before receiving a fixed test meal at 12:00pm. The test meal
consisted of a fixed quantity of Uncle Ben's tomato and olive oil rice and natural yogurt that
only varied based on the participant's gender (i.e., 900 kcal and 750 kcal for men and women,
respectively). The portion and presentation of the test meals were identical during both test
205 days. On both test days, meals were consumed alone over an allotted 30 minute period
without access to media or other social distractions. Participants were then asked to repeat
the LFPQ-fr task 15 minutes after consumption of the test meal. Additionally, appetite feelings
(hunger, fullness, prospective food consumption and desire to eat) were assessed using 150-
mm Visual Analogue Scales (VAS) [32] before and immediately after the test meal, as well as
210 30 minutes, 60 minutes, and 90 minutes after the meal.

Evaluation of food reward – Leeds Food Preference Questionnaire

The LFPQ consists of two distinct tasks. During the first task, explicit liking and wanting are
assessed for 16 distinct food images from all food categories (HFSA, HFSW, LFSA, LFSW)
individually using 100-mm VAS when presented with the following questions: i) "How pleasant
215 would it be to taste this food now?" (explicit liking) and; ii) "How much do you want to eat this

food now?" (explicit wanting). This was followed by a paired food task where the user is advised to choose between two food images from reciprocal food categories as rapidly and accurately as possible. In addition to quantifying choice frequency, an implicit wanting score is computed from reaction times covertly recorded during each selection that are adjusted for its selection frequency. During this task, there are a total 96 trials so that every possible food image pairing is presented and occurs in a random order. For each food reward component, two salient metrics are calculated: fat bias and taste bias. The fat bias score is calculated by subtracting mean scores of low fat foods from high fat foods, whilst taste bias score was obtained by subtracting mean scores of savory foods from sweet foods. Positive values for fat bias or taste bias indicates a preference for high-fat and sweet foods, respectively (for more details see Oustric et al. 2020 [28]).

Statistical analysis

The agreement of pre-meal and post-meal fat and taste biases for each salient reward component (i.e., explicit liking, explicit wanting, implicit wanting, and choice frequency) independently, between lab visits was analyzed using Lin's concordance correlation coefficients (CCC). To assess the reliability of scores in response to the test meal, delta scores were calculated by subtracting pre-meal from post-meal values and tested for agreement between visits using the same approach. To supplement these analyses, pre- and post-meal values were compared at each visit (V1 and V2) using random-effects models in order to analyze time (pre- and post-meal). Natural Logarithmic transformations of dependent variables were conducted when the assumption of normality was violated. Continuous variables were presented as means with standard deviation or median with interquartile range. The distribution of dependent variables and model residuals were tested for normality using Shapiro-Wilk test.

Statistical analyses were performed using Stata software (version 15, StataCorp, College Station, US). All statistical tests were carried out using a two-sided type I error rate at 5%. For the primary objective (i.e., to evaluate agreement), specific attention was paid to the magnitude of results reported with 95% limits of agreement and interpreted according to the conventions reported by Altman ([-0.7; -0.4] low agreement; [-0.4, 0.4] negligible agreement; [0.4, 0.7] mild agreement; and [>0.7] high agreement). No correction for multiple testing was

applied in the analysis of secondary objectives. Findings from these analyses were interpreted as exploratory.

250 **Results**

Food database validation

Participants involved in this stage of the validation process (N=430) ranged in age between 19 and 82 years (M=42.2, SD=12.7) and were predominately female (81%). 16 out of 36 food pictures matched the initial validation criteria and were selected for the experimental
255 validation of task reliability (third phase). Results from the food database validation and nutritional characteristics of the food are presented in Table 1 and Table S1 in the supplementary materials, respectively. All the food matched the selection criteria. However, it was difficult to find a diversity of foods within a category that are all liked and eaten at same time of day. This resulted in the LFSW category mainly being comprised of fruits, for example.
260 Moreover, it is worth noting that yogurt was perceived as high fat and sweets were not considered appropriate for lunchtime in the French sample, contrary to the British population on which the original LFPQ was validated. Based on the approach outlined by Oustric et al. 2020 [27], a hierarchical clustering was performed on R to graphically support and statistically validate the perception of each food image as belonging to one of the four distinct food
265 categories (HFSA, HFSW, LFSA and LFSW; see Figure 1). The 16 food images selected for the LFPQ-fr can be seen in Figure S1 (supplementary materials).

LFPQ-fr sensitivity and reliability

Population characteristics. This sample was 50% female and 30.0 ± 8.4 years old on average.
270 All participants were healthy weight with a mean BMI of 22.4 ± 2.4 kg/m², mean weight of 66.5 ± 11.4 kg, mean fat mass (FM) of $20.2 \pm 7.1\%$, and mean fat free mass (FFM) of 50.4 ± 11.2 kg. Regarding eating behavior traits, the sample had relatively low mean trait Binge Eating (BES; 6.4 ± 4.8). The TFEQ scores were: Cognitive Restraint = 1.71 ± 0.58 , Disinhibition: 2.04 ± 0.49 and Susceptibility to hunger: 2.04 ± 0.49 . The DEBQ scores were Emotional eating score:

275 1.86 ± 0.68; Restrained eating: 1.6 ± 0.4 and External eating: 1.8 ± 0.3. Altogether these scores indicated an absence of eating disorders and other deleterious eating patterns overall. No significant differences were observed between males and females. None of the volunteers were excluded based on these scores.

Task validity and sensitivity. Table 2 details the descriptive statistics for all LFPQ-fr values in 280 fasted and fed states on V1 and V2. The measures of central tendency of fat and taste biases for all primary outcomes were consistent with previous investigations and validations of the original British version and other cultural adaptations in both fasted and fed states and in healthy-weight samples [4, 24, 25, 28]. Indeed, similar to these previous studies, a preference for low-fat and savory foods was demonstrated in a fasted state and shifted towards stronger 285 reward for high-fat and sweet foods in response to intake of a low-fat, savory meal, indicative of sensory-specific satiety [9] (Table 3). Although significant changes in taste bias were observed for all dimensions during both visits, significant changes in fat bias were only observed for explicit wanting during both visits and explicit liking during V2.

Task reliability and agreement. Analyses of fasted and fed values revealed that all primary 290 reward components of the LFPQ-fr were statistically consistent except for explicit liking fat bias. That is, statistical differences were consistently detected between V1 and V2, indicative of good test re-test reliability. However, in the case of explicit liking fat bias, differences between visits in the fasted state were not significant ($p=.191$) whilst significant after the standardized test meal ($p=.019$). Importantly, Lin's CCC revealed moderate to high agreement 295 between the two visits for all the LFPQ-fr dimensions in fasted (CCC=.72-.94) and fed (CCC=.53-.80) appetitive states. The 95% limits of agreement obtained from the Bland & Altman analysis of pre- and post-meal values further support the moderate to high agreement between visits (Table 2).

In analyses of changes in response to the test meal, only the explicit liking fat bias 300 demonstrated inconsistent results between V1 and V2 as it was statistically significant on V2 ($p=.005$) but only approached significance on V1 ($p=.086$). The results for all other dimensions demonstrated consistency between V1 and V2. Moreover, delta values (i.e., changes from baseline to post-meal) for all LFPQ-fr dimensions were found to be reliable between V1 and V2 (CCC=.49-.72). The 95% limits of agreement obtained from the Bland & Altman analysis of 305 delta values can be seen in Table 3 and graphically depicted in Figure 2.

Appetite sensations. Trends in appetite sensations over the course of each test day were statistically similar between V1 and V2 (hunger: $p=0.095$; fullness: $p=0.707$; desire to eat: $p=0.740$ and prospective food consumption: $p=0.421$) as indicated by the non-significant time (pre-meal, post-meal, post+30min, post+60min, post+90min) x visit interactions. Figure 3 presents the evolution of hunger, fullness, desire to eat and prospective food consumption during V1 and V2.

Discussion

Although a French translation of the original British LFPQ has been used in several studies for the last 10 years approximately, the present work addressed a pressing need to validate a culturally adapted version of this instrument to reliably evaluate food reward and the hedonic control of appetite in a French population. Importantly, this cultural adaptation has been conducted following the rigorous and standardized set of procedures previously described [28]. Results from this study indicate that the food items and images selected during the developmental process and used to develop the LFPQ-fr were correctly identified and perceived by the population to be either low or high-fat and sweet or savory foods. Furthermore, the experimental evaluation of the newly developed LFPQ-fr demonstrated the reliability and reproducibility of salient food reward outcomes in both fasted and fed states.

In line with the methodological recommendations formulated for adapting the LFPQ to diverse cultures [28], the final set of 16 images that compose the LFPQ-fr was selected after an online survey conducted among 430 French adult participants who were asked to identify frequently consumed and liked food items that they could correctly recognize as sweet or savory and low or high fat, and typical of lunch time consumption. Specifically, the selected items were easily recognized by between 96.5% and 100% of the sample, which is highly satisfactory and in line with what was observed in similar previously published studies [24, 28]. Similarly, the sensory (sweet versus savoury) and nutritional (high versus low fat) qualities of the selected items were properly identified by the vast majority of the sample as displayed in Table 1 and the statistical validation displayed in Figure 1. Taken together, these results support the validity and cultural adaptation of the LFPQ-fr for a French population, and represent an important step since food choices have been shown to be affected by the familiarity with food images, which will inevitably vary across cultures [33].

The second step of the validation process consisted of asking healthy adults (half females and half males) to complete the LFPQ-fr before and after a standardized test meal on two separate occasions. The scores obtained in both fasted and fed states for all LFPQ-fr dimensions were in the ranges previously reported when using both the original [6, 28] and some recently
340 culturally adapted versions [24, 25], supporting the sensitivity to a test meal manipulation. Despite the high degree of experimental control by standardizing dietary and physical activity-related behaviors in the 2 days preceding each test day, some scores were found to be significantly different between the first and the second visit. This might be attribute to myriad environmental factors or other contextual reasons that appetite may be sensitive to and are
345 difficult to control for [1]. However, it important to note that the presence or absence of statistical differences remained consistent between pre- and post-meal values except for Explicit Liking fat bias. It should be further considered that the patterns in appetite feelings (hunger, fullness, prospective food consumption and desire to eat) from to baseline to 90 minutes post-meal were found to be nearly identical between V1 and V2. These observations
350 reinforce the condition and state-related consistency at the participant level between lab visits.

Importantly, due to the high inter-rater variability of appetite-related measures, the obtained concordance coefficients and the Bland & Altman analyses as represented graphically in Figure 2 and the LOA provide better evaluations of the test-re-test reliability of the LFPQ-fr.
355 Moreover, none of the changes in key LFPQ-fr dimension in response to the test meal was significantly different between the first and the second visit. Overall, the evaluation of the LFPQ-fr values across different appetitive states revealed a high reproducibility between the participants' two visits.

In line with previous cultural adaptations of the LFPQ conducted in other countries, the
360 present work produced a culturally appropriate and reliable version of the LFPQ suitable for French populations. While this validation study has been conducted in healthy men and women aged between 20 to 40 years, further studies should be conducted to assess its reliability in other French samples with different age ranges, weight status and health profiles. Additionally, it may be beneficial to conduct further studies in different settings or using
365 different test meals, for example, to elucidate whether the reliability of the LFPQ-fr extends to different research contexts.

References

- [1] H.R. Berthoud. Metabolic and hedonic drives in the neural control of appetite: Who is the boss? *Current Opinion in Neurobiology*. 21(6) (2011) 888–896.
375
- [2] J.E. Blundell, C. Gibbons, K. Beaulieu, N. Casanova, C. Duarte, G. Finlayson, R.J. Stubbs, M. Hopkins. The drive to eat in homo sapiens: Energy expenditure drives energy intake. *Physiol Behav*. 219 (2020) 112846.
- [3] H.R. Berthoud, H. Munzberg, & C. D. Morrison. Blaming the Brain for Obesity: Integration of Hedonic and Homeostatic Mechanisms. *Gastroenterology*, 152 (7) (2017) 1728–1738.
380
- [4] M. Dalton, & G. Finlayson. Psychobiological examination of liking and wanting for fat and sweet taste in trait binge eating females. *Physiology and Behavior*. 136 (2014) 128–134.
- [5] I. Morales, K.C. Berridge. 'Liking' and 'wanting' in eating and food reward: Brain mechanisms and clinical implications. *Physiol Behav*. 227 (2020) 113152.
- [6] G. Finlayson, & M. Dalton. Hedonics of food consumption: are food 'liking' and 'wanting' viable targets for appetite control in the obese? *Current Obesity Reports*. 1(1) (2012) 42–49.
385
- [7] G. Finlayson, N. King, & J.E. Blundell. Is it possible to dissociate 'liking' and 'wanting' for foods in humans? A novel experimental procedure. *Physiology and Behavior*. 90(1) (2007) 36–42.
- [8] M. Dalton, J.E. Blundell, & G. Finlayson. Effect of BMI and binge eating on food reward and energy intake: Further evidence for a binge eating subtype of obesity. *Obesity Facts*. 6(4) (2013) 348–359.
390
- [9] G. Finlayson, I. Bordes, S. Griffioen-Roose, C. de Graaf, J.E. Blundell. Susceptibility to overeating affects the impact of savory or sweet drinks on satiation, reward, and food intake in nonobese women. *Journal of Nutrition*. 142(1) (2012) 125–130
- [10] S.A. French, N.R. Mitchell, J. Wolfson, G. Finlayson, J.E. Blundell, R.W. Jeffery. Questionnaire and laboratory measures of eating behavior. Associations with energy intake and BMI in a community sample of working adults. *Appetite*. 72 (2014) 50–58.
395
- [11] S. Griffioen-Roose, G. Finlayson, M. Mars, J.E. Blundell, C. de Graaf. Measuring food reward and the transfer effect of sensory specific satiety. *Appetite*. 55(3) (2010) 648–655.
- [12] S. Griffioen-Roose, M. Mars, E. Siebelink, G. Finlayson, D. Tomé, C. de Graaf, C. Protein status elicits compensatory changes in food intake and food preferences. *The American Journal of Clinical Nutrition*. 95(1) (2011) 32–38
400

- [13] N. Li Food preferences of Chinese cruise tourists: an exploratory study. *J Foodserv Bus Res.* 25 (2011) 1–22.
- 405 [14] O. Wang, X. Gellynck, W. Verbeke. Perceptions of Chinese traditional food and European food among Chinese consumers. *Br Food J.* 118(2016) 2855–72.
- [15] X. Xu, J. Pu, A. Shaw, T. Jackson. Neural responsiveness to Chinese versus Western food images: An functional magnetic resonance imaging study of Chinese young adults. *Front Nutr.* 12 (9) (2022) 948039.
- 410 [16] M. Duerlund, B. Vad Andersen, K. Wang, R.C.K. Cha, D. V. Byrne. Post-Ingestive Sensations Driving Post-Ingestive Food Pleasure: A Cross-Cultural Consumer Study Comparing Denmark and China. *Foods.* 11;9 (5) (2020)617.
- [17] C. Sulmont-Rossé, R. Drabek, V.L. Almlí, H. Van Zyl, A. Patricia, M. Kern, J.A. Mcewan, G. Ares. A Cross-Cultural Perspective on Feeling Good in the Context of Foods and Beverages. *Food Res. Int.* 115 (2019) 292–301.
- 415 [18] G. Ares, A. Giménez, L. Vidal, Y. Zhou, A. Krystallis, G. Tsalis, R. Symoneaux, L.M. Cunha, A.P. de Moura, A. Claret., L. Guerrero, A.V. Cardello, A. Wright, L. Jefferies, M. Lloyd, D. Oliveira, R. Do We All Perceive Food-Related Wellbeing in the Same Way? Results from an Exploratory Cross-Cultural Study. *Food Qual. Prefer.* 52 (2016) 62–73.
- 420 [19] G. Ares, L. de Saldamando, A. Giménez, A. Claret, L.M. Cunha, L. Guerrero, A.P. de Mour, D.C.R. Oliveira, R. Symoneaux, R. Deliza. Consumers' Associations with Wellbeing in a Food-Related Context: A Cross-Cultural Study. *Food Qual. Prefer.* 40 (2015) 304–315.
- [20] H.J. Kim, S.J. Chung, K.O. Kim, B. Nielsen, R. Ishii, M.A. O'Mahony. Cross-Cultural Study of Acceptability and Food Pairing for Hot Sauces. *Appetite.* 123 (2018) 306–316.
- 425 [21] S.H. Kim, N. Petard, J.H. Hong. What Is Lost in Translation: A Cross-Cultural Study to Compare the Concept of Nuttiness and Its Perception in Soymilk among Korean, Chinese, and Western Groups. *Food Res. Int.* 105 (2018) 970–981.
- [22] H.S. Lee, K. Lopetcharat. Effect of Culture on Sensory and Consumer Research: Asian Perspectives. *Curr. Opin. Food Sci.* 15 (2017) 22–29.
- 430 [23] D.D. Torrico, S. Fuentes, C. Gonzalez Viejo, H. Ashman, F.R. Dunshea. Cross-Cultural Effects of Food Product Familiarity on Sensory Acceptability and Non-Invasive Physiological Responses of Consumers. *Food Res. Int.* 115 (2019) 439–450.
- 435 [24] A. Hiratsu, D. Thivel, K. Beaulieu, G. Finlayson, C. Nagayama, K. Kamemoto, S. Siripiyavatana, Y. Tataka, Y. Yamada, M. Miyashita. Development of the Leeds Food Preference Questionnaire in Japanese: Sensitivity and reproducibility of liking and wanting for food in fasted and fed states. *Food Quality and Preference.* 102, (2022) 104677.
- [25] J.P. Carvalho-Ferreira, G. Finlayson, D.T. da Cunha, G. Caldas, D. Bandoni, V.V. de Rosso. Adiposity and binge eating are related to liking and wanting for food in Brazil: A cultural adaptation of the Leeds food preference questionnaire. *Appetite.* 133 (2019) 174-183.
- 440 [26] H. Pedersen, J. Salling Quist, M. Møller Jensen, K.K. Bjerring Clemmensen, D. Vistisen, M.E. Jørgensen, K. Færch, G. Finlayson. Investigation of eye tracking, electrodermal activity and facial expressions as biometric signatures of food reward and intake in normal weight adults. *Food Quality and Preference.* 93 (2021) 104248.

- 445 [27] Y. Zhou, P. Oustric, X. Li, H. Zhu, G. Finlayson, C. Zhou. Neurobehavioral markers of food preference and reward in fasted and fed states and their association with eating behaviors in young Chinese adults. *Food Quality and Preference*. 103 (2023) 104689.
- [28] P. Oustric, D. Thivel, M. Dalton, K. Beaulieu, C. Gibbons, M. Hopkins, Blundell JE, G. Finlayson. Measuring food preference and reward: Application and cross-cultural adaptation of the Leeds Food Preference Questionnaire in human experimental research. *Food Quality and Preference*. 80 (2020) 103824
- 450 [29] P. Brunault, I. Rabemampianina, G. Apfeldorfer, N. Ballon, C. Couet, C. Réveillère, P. Gaillard, W. El-Hage. The Dutch Eating Behavior Questionnaire: Further psychometric validation and clinical implications of the French version in normal weight and obese persons. *Presse Med*. 44 (12) (2015) e363-72.
- 455 [30] P. Brunault, P. Gaillard, N. Ballon, C. Couet, P. Isnard, S. Cook, I. Delbachian, C.R. Courtois. [Validation of the French version of the Binge Eating Scale: Examination of its factor structure, internal consistency and construct validity in a non-clinical and a clinical population]. *Encephale*. 42 (5) (2016) 426-433.
- [31] A.J. Stunkard, S. Messick. The three-factor eating questionnaire to measure dietary restraint, disinhibition and hunger. *Journal of Psychosomatic Research*. 29 (1985) 71-83.
- 460 [32] Drapeau, Vicky, Neil King, Marion Hetherington, Eric Doucet, John Blundell, et Angelo Tremblay. 2007. « Appetite Sensations and Satiety Quotient: Predictors of Energy Intake and Weight Loss ». *Appetite* 48(2):159-66.
- 465 [33] C. H. C. Leenaars, J. C. Zant, A. Aussems, V. Faatz, D. Snackers, A Kalsbeek. The Leeds Food Preference Questionnaire after mild sleep restriction - A small feasibility study. *Physiology and Behavior*. 154 (2016) 28-33.

470

475

480

485

Figures legends

Figure 1. Scatter plot visualising the hierarchical clustering by taste and fat. To validate the classification in four food categories, a hierarchical clustering was made (see Oustric et al 2020). Mean results of the food validation survey for taste and fat have been scaled and the foods have been projected according to their new fat and taste coordinates. Positive ratings represent savoury taste or high-fat, respectively. Smaller points represent the foods and larger points depict the cluster's centre. The smaller the ellipse of the cluster, the more homogenous the cluster (e.g. LFSW). The further the food are from zero, the more separate are the clusters. This scatter plot attests of four distinct groups of food and allows to spot which food are closer to other clusters. Plot performed on R version 3.5.1 (R Core Team, 2013) using factoextra v1.0.5 package and enhanced hierarchical clustering (see Oustric et al. 2020).

Figure 2. Bland & Altman plots for food choice, explicit liking, explicit wanting and implicit wanting pre-post-meal deltas between visit 1 (V1) and visit 2 (V2) for both Taste and Fat. These graphs show the difference between the measures performed on V1 and V2 plotted against the means for each participant. The purple lines=Bias or mean differences between the 2 measures; red lines=95% limits of agreement of the mean difference. The critical difference is "two" times standard deviation of the difference between the 2 measures (half of the limits of agreement). The bias (purple line) should be close to zero and the limit of agreement narrow to support the reliability of the task. Participants should be evenly distributed along the means.

Figure 3. Evolution of Hunger (A), Desire to Eat (B), Fullness (C) and Prospective Food Consumption (D) during the first and second visits.

Figure S1. Food pictures selected in a French version of the Leeds Food Preference Questionnaire.

515

Table 1. Survey results on the selected food pictures (n=430)

Names	Recognition (%) ^a	Frequency (1-6)	Liking (mm)	Taste (mm)	Fat (mm)	Time of day (%) ^b
Brownie	99.30	3.24	82.76	4.73	85.56	82.41
Butter biscuits	100.00	2.84	73.00	9.88	82.88	74.40
Icecream	99.77	2.97	75.62	4.89	60.06	82.99
Raspberry tart	100.00	2.92	83.42	4.91	64.56	86.68
Apple puree	96.51	4.38	79.15	12.31	15.33	88.61
Banana	99.53	4.30	73.05	12.02	26.48	80.47
Strawberries salad	100.00	3.15	88.19	8.57	14.31	89.76
Fruits salad	100.00	4.29	84.13	12.26	9.77	89.68
Avocado with mayonnaise	99.77	3.14	72.17	80.73	74.66	76.96
Charcuterie	100.00	4.00	76.65	95.52	93.95	75.87
Comté	100.00	4.83	83.88	91.53	80.96	82.85
Quiche Lorraine	98.14	3.55	76.46	90.12	79.76	80.17
Chicken salad	99.30	4.00	71.03	85.68	28.48	82.94
Ratatouille	100.00	4.11	77.71	83.74	27.14	88.77
Shrimp skewer	100.00	3.27	79.28	85.81	29.07	77.94
Tomato salad	100.00	4.53	80.79	73.85	14.11	89.08
mean (sd)	99.52±0.9	3.72±0.65	78.58±5.00	47.28±40.19	49.19±30.97	83.10±5.21

^aRecognition is expressed as a percentage of the total number of the samples; ^bPercentage of participants who properly identified the appropriate time of day

Table 2. A French version of the Leeds Food Preference Questionnaire results and comparison between Visit 1 and Visit 2, pre- and post-meal.

	Visit 1						Visit 2						Pre vs post		Visit 1 vs Visit 2						
	Pre- meal			Post- meal			Pre- meal			Post- meal			Visit 1	Visit 2	Pre- meal			Post- meal			
	Mean	SD	p50[p25 ; p75]	Mean	SD	p50[p25 ; p75]	Mean	SD	p50[p25 ; p75]	Mean	SD	p50[p25 ; p75]	p	p	Lin [95%CI]	95% LOA (Bias)	p	Lin [95%CI]	95% LOA (Bias)	p	
Choice																					
<i>Fat</i>	-6,48	8,75	-7,00 [-11,00 ; -1,00]	-5,98	9,20	-5,50 [-13,01 ; 1,00]	-5,26	8,42	-5,00 [-10,00 ; -1,00]	-2,95	10,77	-3,00 [-9,00 ; 2,00]	0,59	0,204	0,94 [0,91 ; 0,97]	-6,01 to 4,35	0,044	0,66 [0,51 ; 0,81]	-18,11 to 12,86	0,015	
<i>Taste</i>	-16,00	12,12	-20,01 [-24,00 ; -8,00]	16,70	15,11	23,00 [8,00 ; 27,00]	-18,50	12,38	-21,00 [-26,00 ; -11,40]	18,87	13,16	23,50 [14,00 ; 27,50]	<0,0001	<0,0001	0,93 [0,89 ; 0,97]	-7,40 to 9,25	0,117	0,62 [0,45 ; 0,80]	-23,75 to 22,00	0,437	
Explicit Liking																					
<i>Fat</i>	-8,22	17,11	-5,56 [-18,25 ; 3,50]	-5,11	15,15	-1,31 [-11,5,0 ; 3,75]	-5,64	13,60	-6,00 [-11,75 ; 3,18]	-1,45	12,69	-1,87 [-8,62 ; 5,25]	0,086	0,005	0,80 [0,71 ; 0,90]	-19,42 to 16,35	0,191	0,80 [0,70 ; 0,90]	-18,39 to 13,32	0,019	
<i>Taste</i>	-17,82	22,16	-19,00 [-32,25 ; -4,37]	19,62	25,08	19,81 [5,25 ; 36,25]	-20,75	23,76	-21,31 [-36,31 ; -2,43]	22,41	18,74	19,75 [9,68 ; 39,43]	<0,0001	<0,0001	0,74 [0,61 ; 0,86]	-27,17 to 35,39	0,093	0,57 [0,38 ; 0,76]	-39,16 to 37,28	0,557	
Explicit Wanting																					
<i>Fat</i>	-9,01	16,04	-8,50 [-16,75 ; 0,50]	-4,09	14,20	-1,43 [-7,50 ; 2,12]	-7,57	13,08	-6,68 [-14,75 ; 0,37]	-1,26	11,65	-2,31 [-7,06 ; 1,31]	0,013	<0,0001	0,80 [0,70 ; 0,90]	-17,36 to 16,72	0,687	0,77 [0,66 ; 0,89]	-17,39 to 14,17	0,114	
<i>Taste</i>	-19,70	21,51	-18,81 [-37,12 ; -4,50]	18,62	25,05	17,81 [0,62 ; 34,62]	-19,60	22,66	-20,87 [-33,31 ; -6,12]	20,01	17,70	19,81 [5,81 ; 33,37]	<0,0001	<0,0001	0,89 [0,83 ; 0,95]	-17,94 to 21,10	0,366	0,63 [0,47 ; 0,79]	-34,00 to 35,01	0,967	
Implicit Wanting																					
<i>Fat</i>	-19,48	28,15	-22,29 [-34,28 ; -4,65]	-16,15	29,58	-11,91 [-36,08 ; 5,80]	-9,04	25,70	-10,67 [-21,99 ; 5,30]	-6,92	31,61	-5,86 [-23,54 ; 7,59]	0,358	0,52	0,76 [0,64 ; 0,87]	-40,65 to 23,54	<0,0001	0,64 [0,48 ; 0,81]	-56,30 to 40,37	0,017	
<i>Taste</i>	-41,03	42,61	-47,97 [-69,38 ; -17,02]	47,34	45,86	63,71 [13,83 ; 80,66]	-47,93	38,80	-52,19 [-76,16 ; -26,29]	47,21	42,03	62,93 [35,80 ; 74,44]	<0,0001	<0,0001	0,72 [0,58 ; 0,85]	-52,20 to 65,99	0,107	0,53 [0,32 ; 0,73]	-76,02 to 84,04	0,694	

Table 3. LFPQ-fr pre-post-meal deltas comparison between visit 1 and visit 2.

	V1 Delta Pre-Post-meal			V2 Delta Pre-Post-meal			p	V1 vs V2	
	Mean	SD	p50[p25 ; p75]	Mean	SD	p50[p25 ; p75]		Lin [95%CI]	95% LOA (Bias)
Choice									
<i>Fat</i>	0,57	7,10	1,00 [-3,00 ; 5,00]	1,31	8,98	1,00 [-4,00 ; 9,00]	0,509	0,59 [0,39 ; 0,78]	-15,01 to 13,55
<i>Taste</i>	31,91	17,48	32,00 [19,00 ; 47,00]	36,75	17,40	39,50 [25,00 ; 52,00]	0,204	0,71 [0,56 ; 0,87]	-26,55 to 23,43
Explicit Liking									
<i>Fat</i>	3,11	12,97	3,87 [-4,50 ; 13,12]	4,19	10,49	4,06 [-2,37 ; 8,93]	0,484	0,61 [0,44 ; 0,78]	-21,43 to 19,42
<i>Taste</i>	37,44	26,16	35,56 [16,00 ; 58,37]	43,17	30,18	41,43 [24,18 ; 56,68]	0,124	0,62 [0,45 ; 0,79]	-52,62 to 42,52
Explicit Wanting									
<i>Fat</i>	4,92	14,18	6,81 [-4,12 ; 13,37]	6,31	10,44	4,37 [0,56 ; 10,25]	0,424	0,55 [0,37 ; 0,74]	-24,41 to 21,82
<i>Taste</i>	39,67	27,43	37,06 [21,12 ; 59,50]	38,62	27,11	36,81 [21,06 ; 53,31]	0,877	0,71 [0,56 ; 0,87]	-41,13 to 38,82
Implicit Wanting									
<i>Fat</i>	3,33	25,88	0,05 [-12,53 ; 23,26]	2,11	23,06	-1,39 [-14,01 ; 21,13]	0,797	0,49 [0,27 ; 0,71]	-48,24 to 49,42
<i>Taste</i>	88,37	51,17	86,43 [52,27 ; 134,56]	95,14	45,50	87,58 [65,48 ; 134,10]	0,479	0,55 [0,35 ; 0,75]	-89,48 to 83,72

Table S1 (Supplementary). Nutritional composition of the selected food.

	Food	Kcal/100g	% Pro	% Carb	% Fat	Energy density (kcal/g)
HFSW	Brownie	440.00	5.27	48.58	53.18	4.40
	Icecream	175.59	8.70	49.58	52.22	1.76
	Raspberry tart	451.00	7.89	50.89	54.68	4.51
	Butter biscuits	503.00	4.45	49.20	50.10	5.03
mean±SD		392.4±147.13	6.58±2.04	49.56±0.97	52.54±1.92	3.92±1.47
LFSW	Banana	90.00	4.36	81.67	2.50	0.90
	Strawberries salad	52.24	4.74	76.06	2.95	0.52
	Apple puree	73.00	1.10	87.33	0.00	0.73
	Fruits salad	54.67	5.37	69.51	7.13	0.55
mean±SD		67.48±17.65	3.89±1.91	78.64±7.63	3.15±2.96	0.67±0.18
HFSA	Comté	418.00	25.55	0.00	74.50	4.18
	Avocado with mayonnaise	254.54	2.44	5.12	88.74	2.55
	Quiche Lorraine	274.00	12.99	22.17	70.29	2.74
	Charcuterie	338.48	29.78	1.59	68.59	3.38
mean±SD		321.26±73.8	17.69±12.41	7.22±10.19	75.53±9.15	3.21±0.74
LFSA	Chicken salad	156.00	54.03	3.95	19.95	1.56
	Shrimp skewer	109.00	69.72	15.62	9.58	1.09
	Ratatouille	58.00	8.28	18.75	18.62	0.58
	Salad with tomato	17.70	23.44	42.37	12.06	0.18
mean±SD		85.18±60.21	38.87±28.02	20.17±16.11	15.05±5.02	0.85±0.6
TOTAL						
mean±SD		216.58±167.04	16.76±19.79	38.9±29.95	36.57±30.31	2.17±1.67

HFSA: high-fat savoury, LFSA: low-fat savoury, HFSW: high-fat sweet, LFSW: low-fat sweet; % Pro: Percentage of total energy from protein, % Carb: percentage of total energy from carbohydrate, % Fat: percentage of total energy from fat.