



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/199092/>

Version: Published Version

Article:

McIntosh, Ian (2023) On the connected components of the moduli space of equivariant minimal surfaces in CH^2 . *Geometriae Dedicata*. 56. ISSN: 1572-9168

<https://doi.org/10.1007/s10711-023-00793-z>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



On the connected components of the moduli space of equivariant minimal surfaces in $\mathbb{C}\mathbb{H}^2$.

Ian McIntosh¹

Received: 20 August 2021 / Accepted: 25 March 2023 / Published online: 17 April 2023
© The Author(s) 2023

Abstract

An equivariant minimal surface in $\mathbb{C}\mathbb{H}^n$ is a minimal map of the Poincaré disc into $\mathbb{C}\mathbb{H}^n$ which intertwines two actions of the fundamental group of a closed surface Σ : a Fuchsian representation on the disc and an irreducible action by isometries on $\mathbb{C}\mathbb{H}^n$. The moduli space of these can be studied by relating it to the nilpotent cone in each moduli space of $PU(n, 1)$ -Higgs bundles over the conformal surface corresponding to the map. By providing a necessary condition for points on this nilpotent cone to be smooth this article shows that away from the points corresponding to branched minimal immersions or \pm -holomorphic immersions the moduli space is smooth. The argument is easily adapted to show that for $\mathbb{R}\mathbb{H}^n$ the full space of (unbranched) immersions is smooth. For $\mathbb{C}\mathbb{H}^2$ we show that the connected components of the moduli space of minimal immersions are indexed by the Toledo invariant and the Euler number of the normal bundle of the immersion. This is achieved by studying the limit points of the \mathbb{C}^\times -action on the nilpotent cone. It is shown that the limit points as $t \rightarrow 0$ lead only to branched minimal immersions or \pm -holomorphic immersions. In particular, the Euler number of the normal bundle can only jump by passing through branched minimal maps.

Keywords Minimal surface · Higgs bundle · Complex hyperbolic plane · Nilpotent cone

Mathematics Subject Classification (2020) 20H10 · 53C43 · 58E20

1 Introduction

Let Σ be a closed oriented surface of genus $g \geq 2$ and let N be a noncompact irreducible symmetric space. Denote by G the identity component of the isometry group of N . By an equivariant minimal surface we mean a minimal immersion $f : \mathcal{D} \rightarrow N$ of the Poincaré disc \mathcal{D} which intertwines the action of a Fuchsian representation $c : \pi_1 \Sigma \rightarrow \text{Aut}(\mathcal{D})$ with an irreducible representation $\rho : \pi_1 \Sigma \rightarrow G$. By a theorem of Corlette [5] (generalising a theorem of Donaldson for $N = \mathbb{R}\mathbb{H}^3$ [6]), the triple (f, c, ρ) is uniquely determined by the pair (c, ρ) , and if one considers that f is essentially unchanged by pre-composition or post-composition by isometries, then one only needs the conjugacy classes of c and ρ .

✉ Ian McIntosh
ian.mcintosh@york.ac.uk

¹ Department of Mathematics, University of York, York YO10 5DD, UK

Hence there is a natural way to assign a topology to the set $\mathcal{M}(\Sigma, N)$ of equivalence classes $[f, c, \rho]$ of equivariant minimal surfaces by embedding it in $\mathcal{T}_g \times \mathcal{R}(\pi_1 \Sigma, G)$ where \mathcal{T}_g is the Teichmüller space of Σ and $\mathcal{R}(\pi_1 \Sigma, G)$ is the character variety of G ., i.e., the moduli space of reductive representations of $\pi_1 \Sigma$ into G up to conjugacy.

This topology for $\mathcal{M}(\Sigma, N)$ was first proposed by Loftin and the author in [17] but prior to this, in [18], we had studied the set $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)$ and provided parametrisations for certain subsets which we called “components”, simply to mean “parts whose union is the whole”. We showed that each parametrisation equipped the component with the structure of a connected smooth complex manifold, most of which had the same dimension. These parametrisations exploited the nonabelian Hodge correspondence, through which one can identify an equivariant minimal surface with a $PU(2, 1)$ -Higgs bundle having certain properties. However, since in [18] we did not equip the total space $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)$ with a topology a priori, the question of its connected components had no framework in which to be addressed. One of the aims of this article is to answer this question now, given the topology described above (in a nutshell, the components identified in [18] are, with a few exceptions, the connected components). A key to understanding the connected components lies in having a criterion for points of the moduli space to be smooth, and this is provided below for $\mathcal{M}(\Sigma, N)$ when N is either $\mathbb{C}\mathbb{H}^n$ or $\mathbb{R}\mathbb{H}^n$. The second aim of this article is to show how these components fit together inside the larger set $\overline{\mathcal{M}}(\Sigma, \mathbb{C}\mathbb{H}^2)$ obtained by including all branched minimal surfaces. We show that this is not smooth, since singularities occur where the closures of the connected components of $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)$ meet. Note that $\overline{\mathcal{M}}(\Sigma, \mathbb{C}\mathbb{H}^2)$ was the space studied in [18], and the distinctive role of branched immersions explained here was not at all apparent during the writing of [18].

To summarise the results below, first recall that a $PU(n, 1)$ -Higgs bundle can be represented by a pair (E, Φ) where $E \rightarrow \Sigma_c$ is a holomorphic vector bundle of rank $n + 1$ over the Riemann surface $\Sigma_c \simeq \mathcal{D}/c$. This has a decomposition $E = V \oplus \underline{\mathbb{C}}$, where V is rank n and $\underline{\mathbb{C}}$ denotes the trivial line bundle. Then Φ is a holomorphic one-form with values in $\text{Hom}(\underline{\mathbb{C}}, V) \oplus \text{Hom}(V, \underline{\mathbb{C}})$. We write $\Phi = (\Phi_1, \Phi_2)$ to denote the two components. In general (E, Φ) corresponds to an equivariant, but possibly branched, minimal surface when $\text{tr}(\Phi^2) = 0$. Zeroes of Φ are branch points of the minimal surface and therefore we require Φ to have no zeroes (i.e., Φ_1, Φ_2 have no common zeroes) to obtain a point in $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^n)$. From [18] we know that one of Φ_1, Φ_2 is identically zero precisely when the minimal immersion is either holomorphic or anti-holomorphic. It is convenient to write $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^n)$ as a disjoint union $\mathcal{V} \cup \mathcal{W}$ where \mathcal{W} is the subvariety of \pm -holomorphic maps and \mathcal{V} is the complement. Our first main result concerns \mathcal{V} , and in fact applies also to $\mathcal{M}(\Sigma, \mathbb{R}\mathbb{H}^n) \subset \mathcal{V}$ using the totally geodesic embedding of $\mathbb{R}\mathbb{H}^n$ into $\mathbb{C}\mathbb{H}^n$.

Theorem 1.1 *Both $\mathcal{V} \subset \mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^n)$ and $\mathcal{M}(\Sigma, \mathbb{R}\mathbb{H}^n)$ are smooth manifolds. They have complex dimension $(g - 1) \dim(G)$ for $G = PU(n, 1)$ and $SO_0(n, 1)$ respectively.*

In fact what we prove is that in the nilpotent cone \mathcal{N}^c (i.e, the locus of $\text{tr}(\Phi^2) = 0$ in the Higgs bundle moduli space $\mathcal{H}(\Sigma_c, G)$ for $G = PU(n, 1)$ or $SO_0(n, 1)$) the regular points of the Hitchin function $\text{tr}(\Phi^2)$ are precisely the points for which Φ has no zeroes when neither Φ_j is identically zero.

Next we restrict our attention to $n = 2$. For $G = PU(2, 1)$ the connected components of $\mathcal{R}(\pi_1 \Sigma, G)$ are indexed by the Toledo invariant, τ . Using Toledo’s convention, as in [18], this satisfies $\tau \in \frac{2}{3}\mathbb{Z}$ and $|\tau| \leq 2(g - 1)$. We denote the subspace of triples $[f, c, \rho]$ for which ρ has Toledo invariant τ by $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)_\tau$ and it is clear that these are disconnected from each other. Minimal immersions which are not \pm -holomorphic have complex and anti-complex

points (points $p \in \Sigma$ for which $f_*T_p^{1,0}\Sigma$ is a subspace of either $T'\mathbb{C}\mathbb{H}^2$ or $T''\mathbb{C}\mathbb{H}^2$). These give effective divisors, D_2 and D_1 respectively. From [18] their degrees $d_j = \deg(D_j)$ satisfy

$$\tau = \frac{2}{3}(d_2 - d_1), \quad \chi(T\Sigma^\perp) = 2(g - 1) - d_1 - d_2, \tag{1.1}$$

$$0 \leq 2d_1 + d_2 < 6(g - 1), \quad 0 \leq d_1 + 2d_2 < 6(g - 1), \tag{1.2}$$

where $T\Sigma^\perp$ is the normal bundle of f , which means the quotient of $T\mathcal{D}^\perp \subset f^{-1}T\mathbb{C}\mathbb{H}^2$ by the natural action of $\pi_1\Sigma$. The inequalities are necessary and sufficient conditions for the existence of minimal immersions which are not \pm -holomorphic. In particular, the degrees d_1, d_2 are equivalent information to the pair τ and $\chi(T\Sigma^\perp)$. For holomorphic immersions the only topological invariant is the Toledo invariant, which determines $\chi(T\Sigma^\perp)$ through

$$\chi(T\Sigma^\perp) = -\frac{3}{2}\tau + 2(g - 1). \tag{1.3}$$

This follows by taking the quotient of the direct sum of holomorphic bundles $f^{-1}T'\mathbb{C}\mathbb{H}^2 = T^{1,0}\mathcal{D} \oplus T\mathcal{D}^\perp$ by the $\pi_1\Sigma$ -action.

Let $\mathcal{V}(d_1, d_2) \subset \mathcal{V}$ denote the subset of those immersions whose divisors have these degrees fixed. It was shown in [18] that each $\mathcal{V}(d_1, d_2)$ can be parametrised by a smooth connected complex manifold of dimension $8(g - 1)$ (actually in [18] we included the branched minimal immersions, which occur when $D_1 \cap D_2 \neq \emptyset$, but as these are described by a complex analytic subvariety removing these does not affect the connectedness of $\mathcal{V}(d_1, d_2)$). In Lemma 5.1 below we show that the parametrisation in [18] is smoothly compatible with the structure it inherits from Theorem 1.1. Since $\mathcal{V}(d_1, d_2)$ all have the same dimension they are therefore the connected components of \mathcal{V} . To establish the connected components of $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)$ it remains to see how these fit together with points of \mathcal{W} . Set $\mathcal{W}_\tau = \mathcal{W} \cap \mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)_\tau$ for each τ . We need only consider $\tau \geq 0$ since $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)$ carries a natural real involution $[f, c, \rho] \mapsto [\bar{f}, c, \bar{\rho}]$ for which $\tau(\bar{\rho}) = -\tau(\rho)$.

Theorem 1.2 *Let $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)_\tau$ denote the space of equivariant minimal immersions with Toledo invariant τ . Then $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)_0$ has connected components $\mathcal{V}(d_1, d_2)$ for $d_2 = d_1$. For $0 < \tau < 2(g - 1)$ the connected components of $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)_\tau$ are $\mathcal{V}(d_1, d_2)$ with $d_1 \neq 0$ and $\mathcal{V}(0, d_2) \cup \mathcal{W}_\tau$, while $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)_\tau = \mathcal{W}_\tau$ for $\tau = 2(g - 1)$.*

In particular, the connected components are indexed by τ and $\chi(T\Sigma^\perp)$: when $d_1 = 0$ the two equations (1.1) become the single equation (1.3).

This theorem is proved through a complete description of the limit points of the \mathbb{C}^\times -action $t \cdot (E, \Phi) = (E, t\Phi)$ on $PU(2, 1)$ -Higgs bundles, for both $t \rightarrow \infty$ and $t \rightarrow 0$. As well as proving the theorem, it gives us information about how the closures of the connected components fit together inside the space $\overline{\mathcal{M}}(\Sigma, \mathbb{C}\mathbb{H}^2)$ which includes all branched minimal immersions. On the Higgs bundle side this corresponds to the space of all (c, E, Φ) for which (E, Φ) lies in the nilpotent cone \mathcal{N}^c excluding the locus $\Phi = 0$ (since this corresponds to constant maps). Each \mathcal{N}^c is known to be stratified by the unstable manifolds of the downwards Morse flow for the Higgs field energy $\|\Phi\|_{L^2}^2$: for each connected component C of critical points these agree with

$$\mathcal{U}_C = \{(E, \Phi) : \lim_{t \rightarrow \infty} (E, t\Phi) \in C\}.$$

It was conjectured in [18], and is proven below, that except when C consists of local minima (there is precisely one such C for each τ and it equals \mathcal{W}_τ) these unstable manifolds

correspond to the fibres of the analytic family $\mathcal{V}(d_1, d_2)$ over \mathcal{T}_g . We understand how these fit together by examining the limit in the opposite direction, as $t \rightarrow 0$. It is well-known that when E is semi-stable (as a vector bundle) the limit $\lim_{t \rightarrow 0}(E, t\Phi)$ must have $\Phi = 0$ and therefore this can only occur when $\tau = 0$. When E is strictly unstable (and assuming $\tau \geq 0$ without loss of generality) we show that either (E, Φ_1) is a Hodge bundle, in which case this is limit, or the limit is determined by the maximal destabilizing line subbundle of V and always corresponds to a branched minimal immersion.

Thus we have the the following geometric picture of how these pieces fit together. Each fibre over c of the space $\overline{\mathcal{M}}(\Sigma, \mathbb{C}\mathbb{H}^2)_\tau$ is a union of these unstable manifolds, and their closures intersect either at branched minimal surfaces or at \pm -holomorphic surfaces. Removing these leaves a disjoint union of components $\mathcal{V}(d_1, d_2)$, one for each critical manifold of non-minima. In particular, one cannot pass from one component $\mathcal{V}(d_1, d_2)$ to another in $\overline{\mathcal{M}}(\Sigma, \mathbb{C}\mathbb{H}^2)$ without forcing the minimal immersion to branch somewhere. Another way of saying this is that $\chi(T\Sigma^\perp)$ can jump but only by passing through branched maps.

2 Definitions

Let $\mathbb{C}^{n,1}$ denote \mathbb{C}^{n+1} equipped with the pseudo-Hermitian inner product characterised by

$$\langle u, u \rangle = |u_1|^2 + \dots |u_n|^2 - |u_{n+1}|^2.$$

Let e_1, \dots, e_{n+1} denote the standard basis vectors for \mathbb{C}^{n+1} . We model $\mathbb{C}\mathbb{H}^n$ on the space of lines in $\mathbb{C}\mathbb{P}^n$ generated by a negative definite vector and fix a base point $o = [e_{n+1}]$. Throughout this article we set $G = PU(n, 1)$, the group of orientation preserving isometries of $\mathbb{C}\mathbb{H}^n$, and let $H \simeq U(n)$ denote the maximal compact subgroup which fixes the base point o . The Lie algebra $\mathfrak{g} = \mathfrak{su}(n, 1)$ of G has Cartan decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$ corresponding to the decomposition of elements

$$\begin{pmatrix} A & u \\ u^\dagger & a \end{pmatrix} = \begin{pmatrix} A & 0 \\ 0 & a \end{pmatrix} + \begin{pmatrix} 0 & u \\ u^\dagger & 0 \end{pmatrix} \tag{2.1}$$

where $A \in \mathfrak{u}(n)$, $a = -\text{tr}(A)$, $u \in \mathbb{C}^n$ and u^\dagger is the Hermitian transpose.

Let Σ be a closed oriented surface of genus $g \geq 2$ and let \mathcal{D} denote the Poincaré disc thought of as the universal cover of Σ (note that $\mathcal{D} \simeq \mathbb{C}\mathbb{H}^1$). A Fuchsian representation $c : \pi_1\Sigma \rightarrow PU(1, 1)$ equips Σ with a complex structure which will be denoted Σ_c . We will assume this complex structure is compatible with the orientation of Σ and hence the set of conjugacy classes of such Fuchsian representations can be identified with the Teichmüller space \mathcal{T}_g of Σ . An equivariant minimal surface in $\mathbb{C}\mathbb{H}^n$ is the equivalence class $[f, c, \rho]$ of a triple consisting of a Fuchsian representation c , an indecomposable representation $\rho : \pi_1\Sigma \rightarrow G$ and a minimal immersion $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^n$ which intertwines the actions of c and ρ , i.e.,

$$f \circ c(\delta) = \rho(\delta) \circ f, \quad \forall \delta \in \pi_1\mathcal{D}.$$

Equivalence is with respect to the action of $PU(1, 1) \times G$ by conjugation on (c, ρ) and the corresponding natural action on f . Indecomposability of ρ means exactly that its image does not lie in a proper subgroup of G . By the same argument as [17, Lemma 2.3] it is equivalent to the condition that f is linearly full (i.e., its image does not lie in a totally geodesic copy of $\mathbb{C}\mathbb{H}^k$ for $k < n$).

Remark 2.1 We could also say ρ is irreducible in its standard real representation on $\mathbb{C}^{n,1}$, although there is the potential for confusion with other meanings of irreducibility (say, for the adjoint representation of the complexification $G^{\mathbb{C}}$).

We use $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^n)$ to denote the set of equivariant minimal surfaces and give it the topology of its natural embedding

$$\mathcal{F} : \mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^n) \rightarrow \mathcal{T}_g \times \mathcal{R}(\pi_1 \Sigma, G); \quad [f, c, \rho] \rightarrow [c, \rho], \tag{2.2}$$

into the product of Teichmüller space with the character variety $\mathcal{R}(\pi_1 \Sigma, G)$ of reductive representations of $\pi_1 \Sigma$ into G modulo conjugacy. This map is injective by Corlette’s uniqueness theorem for equivariant harmonic maps under these conditions [5]. We will denote by $\mathcal{M}(\Sigma_c, \mathbb{C}\mathbb{H}^n)$ the fibre over fixed $c \in \mathcal{T}_g$.

Recall that, for each choice of c , non-abelian Hodge theory provides a homeomorphism between $\mathcal{R}(\pi_1 \Sigma, G)$ and the moduli space $\mathcal{H}(\Sigma_c, G)$ of polystable G -Higgs bundles over Σ_c (and this homeomorphism is analytic away from singularities). Indeed, for any simple noncompact G there is a homeomorphism between $\mathcal{T}_g \times \mathcal{R}(\pi_1 \Sigma, G)$ and the universal Higgs bundle moduli space $\mathcal{H}(\mathcal{C}_g, G)$ over \mathcal{T}_g [1, Thm 7.5] (here \mathcal{C}_g denotes the universal Teichmüller curve over \mathcal{T}_g). This is a complex analytic space for which the fibre over $c \in \mathcal{T}_g$ is biholomorphic to $\mathcal{H}(\Sigma_c, G)$. This homeomorphism is smooth about smooth points and therefore we obtain an embedding

$$\mathcal{F}' : \mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^n) \rightarrow \mathcal{H}(\mathcal{C}_g, G), \tag{2.3}$$

when $G = PU(n, 1)$. This equips $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^n)$ with a complex analytic structure.

We can easily characterise the image of \mathcal{F}' . Recall that in the projective equivalence class of the G -Higgs bundle (E, Φ) , E can be taken to have the form $V \oplus \underline{\mathbb{C}}$ where V is a holomorphic rank n bundle over Σ_c and $\underline{\mathbb{C}}$ denotes the trivial line bundle. The Higgs field Φ satisfies

$$\Phi \in (\text{Hom}(\underline{\mathbb{C}}, V) \oplus \text{Hom}(V, \underline{\mathbb{C}})) \otimes K,$$

and we will write $\Phi = (\Phi_1, \Phi_2)$ to indicate the two components for this decomposition. It is well-known that the harmonic map corresponding to (E, Φ) is weakly conformal (therefore branched minimal) when $\text{tr}(\Phi^2) = 0$. The branch points correspond exactly to the zeroes of Φ . Consequently we have the following elementary observation.

Lemma 2.1 *The pair (E, Φ) lies in the image of \mathcal{F}' if and only if it is stable (hence indecomposable) with $\text{tr}(\Phi^2) = 0$ and Φ having no zeroes.*

Each equivariant minimal surface (f, c, ρ) has associated to it the Toledo invariant τ of ρ and we denote the subset of triples with fixed τ by $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^n)_\tau$. Our convention will be that $\tau = -\frac{2}{n+1} \text{deg}(V)$ so that $|\tau| \leq 2(g - 1)$: this matches Toledo’s original definition and is the one used in [18].

We will also write $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^n)$ as a disjoint union $\mathcal{V} \cup \mathcal{W}$ where \mathcal{W} consists of all \pm -holomorphic immersions and \mathcal{V} is the complement. Each of these has its subsets $\mathcal{V}_\tau, \mathcal{W}_\tau$ containing those elements with fixed Toledo invariant. We note that under the non-abelian Hodge correspondence elements of \mathcal{W} correspond to Higgs bundles with $\Phi_2 = 0$ for $\tau > 0$ or $\Phi_1 = 0$ for $\tau < 0$. Since we always have $\Phi \neq 0$ we know $\mathcal{W}_0 = \emptyset$. Note that \mathcal{W}_τ corresponds to the length two Hodge bundles [18], which are the minima of the Higgs field energy $\|\Phi\|_{L^2}^2$ [4].

From [18] we know that when $n = 2$ the space \mathcal{V} is a disjoint union of subsets $\mathcal{V}(d_1, d_2)$, corresponding to $PU(2, 1)$ -Higgs bundles for which V and Φ are determined by an exact sequence [18]

$$0 \rightarrow K^{-1}(D_1) \xrightarrow{\Phi_1} V \xrightarrow{\Phi_2} K(-D_2) \rightarrow 0, \tag{2.4}$$

where $d_j = \text{deg}(D_j)$ satisfy the inequalities (1.2). There is a one-to-one correspondence between Higgs bundles of this type and data (c, D_1, D_2, ξ) where ξ is the extension class of this extension: this was used in [18] to give each $\mathcal{V}(d_1, d_2)$ the structure of a complex manifold of dimension $5(g - 1)$.

Finally, we will want to consider the strictly larger space $\overline{\mathcal{M}}(\Sigma, \mathbb{C}\mathbb{H}^2)$ which includes additionally the branched minimal immersions. Under \mathcal{F}' it maps to the locus $\text{tr}(\Phi^2) = 0$ excluding the Higgs bundles with $\Phi = 0$. We will write $\tilde{\mathcal{V}}$ and $\tilde{\mathcal{W}}$ when we extend \mathcal{V} and \mathcal{W} to include branched minimal immersions of the respective types. The total space is disconnected into Toledo invariant pieces $\overline{\mathcal{M}}(\Sigma, \mathbb{C}\mathbb{H}^2)_\tau$ and for $\tau \neq 0$ each of these is connected. This follows from the fact that for $\tau \neq 0$ each fibre over $c \in \mathcal{T}_g$ agrees with the component \mathcal{N}_τ^c of the nilpotent cone in Toledo invariant τ , and these are connected by the downwards Morse flow. For $\tau = 0$ this fibre is \mathcal{N}_0^c without the locus $\Phi = 0$. Since this locus is the critical manifold of minima we can no longer use the Morse flow argument, and an understanding of the connected components would require a different approach.

3 Smooth points of $\mathcal{M}(\Sigma, N)$

For $N = \mathbb{C}\mathbb{H}^n$ or $\mathbb{R}\mathbb{H}^n$ let $\mathcal{V} \subset \mathcal{M}(\Sigma, N)$ denote the subvariety of points where neither Φ_j is identically zero. For $\mathbb{R}\mathbb{H}^n$ this is the whole space while for $\mathbb{C}\mathbb{H}^n$ it consists of the equivariant minimal surfaces which are not \pm -holomorphic. Because we assume ρ is indecomposable there are no singular points of $\mathcal{T}_g \times \mathcal{R}(\pi_1, G)$ in \mathcal{V} .

Recall that \mathcal{C}_g denotes the universal Teichmüller curve over \mathcal{T}_g . If we denote by $H^0(\mathcal{C}_g, K^2)$ the bundle over \mathcal{T}_g with fibre $H^0(\Sigma_c, K^2)$ then the map

$$\mathcal{I} : \mathcal{H}(\mathcal{C}_g, G) \rightarrow H^0(\mathcal{C}_g, K^2); \quad (c, E, \Phi) \mapsto (c, \text{tr}(\Phi^2)),$$

is complex analytic. We will denote its zero locus (i.e., the preimage of the zero section of $H^0(\mathcal{C}_g, K^2)$) by \mathcal{N} : for real rank one groups like $PU(n, 1)$ and $SO_0(n, 1)$ this is the union $\cup_{c \in \mathcal{T}_g} \mathcal{N}^c$ of all nilpotent cones.

Let $\mathcal{N}^o \subset \mathcal{N}$ denote the open subvariety for which Φ vanishes nowhere and neither Φ_j is identically zero. Then \mathcal{N}^o is the image of \mathcal{V} in $\mathcal{H}(\mathcal{C}_g, G)$ and Theorem 1.1 is the statement that \mathcal{N}^o is smooth. We will prove this by showing that points of \mathcal{N}^o are regular points of \mathcal{I} . Since \mathcal{I} maps $\mathcal{H}(\Sigma_c, G)$ to $H^0(\Sigma_c, K^2)$ it suffices to show that each point of $(\mathcal{N}^c)^o = \mathcal{N}^o \cap \mathcal{N}^c$ is a regular point of \mathcal{I}_c , the restriction of \mathcal{I} to the fibre over c . We do this by generalising an argument used by Hitchin for $SL(2, \mathbb{C})$ -Higgs bundles [12] to compute the rank of $d\mathcal{I}_c$. For this we must recall the hypercohomology description of the tangent space to $\mathcal{H}(\Sigma_c, G)$ [3].

Let $P \rightarrow \Sigma_c$ denote the principal $H^{\mathbb{C}}$ -bundle whose associated bundle is E , and let $P(\mathfrak{h}^{\mathbb{C}}), \mathcal{P}(\mathfrak{m}^{\mathbb{C}})$ denote the holomorphic $\mathfrak{h}^{\mathbb{C}}$ and $\mathfrak{m}^{\mathbb{C}}$ bundles associated by the adjoint action of $H^{\mathbb{C}}$ on these subspaces of $\mathfrak{g}^{\mathbb{C}}$. One knows from [3] that when (E, Φ) is stable the tangent space at this point is isomorphic to the first hypercohomology $\mathbb{H}^1(\mathcal{A}^*)$ of the complex

$$\mathcal{A}^0 \xrightarrow{\text{ad } \Phi} \mathcal{A}^1 \rightarrow 0 \rightarrow \dots \tag{3.1}$$

where \mathcal{A}^0 is the sheaf of local sections of $P(\mathfrak{h}^{\mathbb{C}})$ and \mathcal{A}^1 is the sheaf of local sections of $P(\mathfrak{m}^{\mathbb{C}}) \otimes K$. As Hitchin [12] points out, by considering the “second” spectral sequence for the hypercohomology of \mathcal{A}^* (i.e., the spectral sequence for the filtration by the degree of Čech cochains) one obtains an isomorphism for $\mathbb{H}^1(\mathcal{A}^*)$ involving the kernel and cokernel sheaves

$$0 \rightarrow Ker \rightarrow \mathcal{A}^0 \xrightarrow{\text{ad } \Phi} \mathcal{A}^1 \rightarrow Coker \rightarrow 0, \tag{3.2}$$

namely,

$$\mathbb{H}^1(\mathcal{A}^*) \simeq H^1(Ker) \oplus H^0(Coker). \tag{3.3}$$

Now let B denote the Killing form on $\mathfrak{g}^{\mathbb{C}}$, specifically $B(\eta, \xi) = \text{tr}(\text{ad } \eta \text{ ad } \xi)$. By adjoint invariance $B(\Phi, \text{ad } \Phi(\eta)) = 0$ for all η , so that we have a well defined map of sheaves

$$B(\Phi, \cdot) : Coker \rightarrow K^2; \quad \eta \mapsto B(\Phi, \eta), \tag{3.4}$$

and an induced map from $H^0(Coker)$ to $H^0(K^2)$. When pre-composed with the projection of $\mathbb{H}^1(\mathcal{A}^*)$ onto $H^0(Coker)$ from (3.3), at a stable point (E, Φ) , this induced map agrees with $d\mathcal{I}_c$ [3, Remark 2.8(iv)]. Hence the rank of $d\mathcal{I}_c$ at (E, Φ) equals the rank of $B(\Phi, \cdot)$ on $H^0(Coker)$.

Lemma 3.1 *Suppose $\text{ad } \Phi$ has corank 1, i.e., $Coker$ is a rank 1 sheaf. Then $d\mathcal{I}_c$ has maximal rank if and only if Φ has no zeroes.*

Proof When $Coker$ is rank one $B(\Phi, \cdot)$ vanishes precisely at the zeroes of Φ with the same divisor of zeroes, D , and hence $Coker \simeq K^2(-D)$. Thus the induced map $H^0(Coker) \rightarrow H^0(K^2)$ is maximal rank, and hence $d\mathcal{I}$ has maximal rank, if and only if Φ has no zeroes. \square

Now our aim is to show that $\text{ad } \Phi$ has corank 1 for points of \mathcal{V} . First we do the case $N = \mathbb{C}\mathbb{H}^n$.

Lemma 3.2 *For $G = PU(n, 1)$, when neither Φ_j is identically zero the map $\text{ad } \Phi$ has corank one.*

Proof Following [18], since $\text{tr}(\Phi^2) = 0$ we have $\Phi_2 \circ \Phi_1 = 0$, and so we have a sequence

$$0 \rightarrow K^{-1}(D_1) \xrightarrow{\Phi_1} V \xrightarrow{\Phi_2} K(-D_2) \rightarrow 0, \tag{3.5}$$

where D_j is the divisor of zeroes of Φ_j . About each point $p \in \Sigma$ we can choose a local holomorphic chart (U, z) and a local holomorphic frame $\sigma_1, \dots, \sigma_n$ for V over U satisfying

$$\Phi(\sigma_0) = z^k \sigma_1 dz, \quad \Phi(\sigma_n) = z^l dz, \quad \Phi(\sigma_j) = 0 \text{ for } j \neq n.$$

Here k, l are, respectively, the degrees of D_1 and D_2 at the point p and σ_0 denotes a local trivialising section of the trivial bundle. With respect to this frame Φ is represented as a local section of $\text{End}(E) \otimes K$ by

$$\xi = \begin{pmatrix} 0 & \dots & 0 & z^k \\ 0 & \ddots & & 0 \\ \vdots & & & \vdots \\ 0 & \dots & z^l & 0 \end{pmatrix} dz = \begin{pmatrix} 0 & z^k e_1 \\ z^l e_n^t & 0 \end{pmatrix} dz, \tag{3.6}$$

where in the second expression the standard basis vectors e_1, \dots, e_n for \mathbb{C}^n have been used to write the matrix in block form. Let χ be a local holomorphic section of $P(\mathfrak{h}^{\mathbb{C}})$, identified with a locally holomorphic $\mathfrak{h}^{\mathbb{C}}$ -valued function. We compute

$$[\xi, \chi] = \left[\begin{pmatrix} 0 & z^k e_1 \\ z^l e_n^t & 0 \end{pmatrix} dz, \begin{pmatrix} A & 0 \\ 0 & a \end{pmatrix} \right] = \begin{pmatrix} 0 & z^k (aI_n - A)e_1 \\ z^l e_n^t (A - aI_n) & 0 \end{pmatrix} dz, \tag{3.7}$$

$a = -\text{tr}(A)$. Hence $[\xi, \chi] = 0$ holds over U when the $2n$ equations

$$(A + \text{tr}(A))e_1 = 0, \quad (A^t + \text{tr}(A))e_n = 0,$$

hold. In components these equations are

$$2A_{11} + \sum_{i=2}^n A_{ii} = 0, \quad A_{j1} = 0, \quad A_{nk} = 0, \quad 2A_{nn} + \sum_{i=1}^{n-1} A_{ii} = 0,$$

for $2 \leq j \leq n$ and $1 \leq k \leq n - 1$. We see that this gives $2n - 1$ independent equations since $A_{n1} = 0$ appears twice. Hence $\text{ad } \xi$ has rank $2n - 1 = \dim \mathfrak{m}^{\mathbb{C}} - 1$. \square

Now we deal with the case $N = \mathbb{R}\mathbb{H}^n$ and follow the description of the Higgs bundles for this case given in [17]. These have the form $E = V \oplus \mathbb{C}$ where V has $\det(V) = \mathbb{C}$ and carries a non-degenerate quadratic form Q_V . In this case $\Phi_2 = \Phi_1^T$ where the transpose is with respect to Q_V . It follows that for f to be an immersion both Φ_j vanish nowhere so both divisors D_1, D_2 are trivial. As a smooth bundle V splits into a direct sum $K^{-1} \oplus W \oplus K$ and with respect to this splitting we can write

$$Q_V = \begin{pmatrix} 0 & 0 & 1 \\ 0 & Q_W & 0 \\ 1 & 0 & 0 \end{pmatrix}. \tag{3.8}$$

Then Φ is represented by ξ in (3.6) with $k = l = 0$. For the Lie algebra we use

$$\mathfrak{so}(Q, 1) = \left\{ \begin{pmatrix} A & u \\ v^t & 0 \end{pmatrix} : A^t Q + Q A = 0, \quad Q u = v \right\} \simeq \mathfrak{so}(n + 1, \mathbb{C}),$$

where $A \in \text{End}(\mathbb{C}^n)$, $u, v \in \mathbb{C}^n$, Q is (3.8) with $Q_W = I_{n-2}$, and the transpose is the usual matrix transpose. The appropriate (complexified) symmetric space decomposition is obtained from the Cartan decomposition (2.1) above:

$$\mathfrak{so}(Q, 1) = \mathfrak{h}_Q^{\mathbb{C}} \oplus \mathfrak{m}_Q^{\mathbb{C}}, \quad \mathfrak{h}_Q^{\mathbb{C}} = \mathfrak{so}(Q, 1) \cap \mathfrak{h}^{\mathbb{C}}, \quad \mathfrak{m}_Q^{\mathbb{C}} = \mathfrak{so}(Q, 1) \cap \mathfrak{m}^{\mathbb{C}}.$$

Lemma 3.3 *For $G = SO_0(n, 1)$ the map $\text{ad } \Phi$ has corank one.*

Proof Following the same arguments as for $\mathbb{C}\mathbb{H}^n$ above but applied to $\mathfrak{so}(Q, 1)$ it suffices to show that the map $\text{ad } \xi : \mathfrak{h}_Q^{\mathbb{C}} \rightarrow \mathfrak{m}_Q^{\mathbb{C}}$ has corank 1. Using the equations (3.7) but with $\chi \in \mathfrak{m}_Q^{\mathbb{C}}$ we see that the image of $\text{ad } \xi$ is isomorphic to

$$\{Ae_1 : A^t = -QAQ\} \subset \mathbb{C}^n.$$

The symmetry $A^t = -QAQ$ implies that $A_{n1} = 0$ and this is the only condition Ae_1 must satisfy. Therefore $\text{im}(\text{ad } \xi)$ has codimension one in $\mathfrak{m}_Q^{\mathbb{C}}$. \square

To complete the proof of Theorem 1.1 we note that at regular points the dimension of $\mathcal{I}^{-1}(0)$ is

$$\begin{aligned} \dim \mathcal{H}(\mathcal{C}_g, G) - \dim H^0(\Sigma_c, K^2) &= \dim(\mathcal{T}_g) + \dim(\mathcal{R}(\pi_1 \Sigma, G)) - \dim H^0(\Sigma_c, K^2) \\ &= \dim(\mathcal{R}(\pi_1 \Sigma, G)). \end{aligned}$$

Here $\dim(\mathcal{R}(\pi_1 \Sigma, G))$ means the dimension of the smooth open subvariety of indecomposable representations. This has real dimension $(2g - 2) \dim(G)$.

Remark 3.1 Theorem 1.1 fills a small gap in the proof of Theorem 4.4 of [17], which gives a smooth parametrisation of the connected components of $\mathcal{M}(\Sigma, \mathbb{R}\mathbb{H}^4)$. The proof that the parametrisation is a local diffeomorphism needs the fact the $\mathcal{M}(\Sigma, \mathbb{R}\mathbb{H}^4)$ is non-singular.

4 Limit points of the \mathbb{C}^\times -action for $n = 2$

It is well-known that \mathbb{C}^\times acts on G -Higgs bundles by $t \cdot (E, \Phi) = (E, t\Phi)$, $t \in \mathbb{C}^\times$, and that the fixed points of this action are Hodge bundles, i.e., one can write $E = \bigoplus_{j=1}^m E_j$ so that $\Phi : E_j \rightarrow E_{j+1} \otimes K$ with $E_{m+1} = 0$. The integer m is called the *length* of the Hodge bundle. It is also well-known that these fixed points are exactly the critical points of the Higgs field energy $\mathfrak{E}(E, \Phi) = \|\Phi\|_{L^2}^2$. Given one of the connected components C of \mathbb{C}^\times -fixed points define

$$\mathcal{S}_C = \{(E, \Phi) : \lim_{t \rightarrow 0} (E, t\Phi) \in C\}, \quad \mathcal{U}_C = \{(E, \Phi) : \lim_{t \rightarrow \infty} (E, t\Phi) \in C\}.$$

By a theorem of Kirwan these agree with, respectively, the stable and unstable manifolds of C along the downwards Morse flow (gradient flow of $-\text{grad } \mathfrak{E}$). One knows that $\cup_C \mathcal{S}_C$ equals $\mathcal{H}(\Sigma_c, G)$ while $\cup_C \mathcal{U}_C$ equals the nilpotent cone \mathcal{N}^c .

Our aim in this section is to identify the limit points as $t \rightarrow \infty$ and $t \rightarrow 0$ inside $\overline{\mathcal{M}}(\Sigma, \mathbb{C}\mathbb{H}^2)$.

First we note that if $\Phi_j = 0$ for either $j = 1$ or $j = 2$, the pair (E, Φ) is plainly a Hodge bundle of length 2, hence a \mathbb{C}^\times -fixed point. So our interest is when neither Φ_j vanishes identically. In this case, using (2.4), as a C^∞ -bundle

$$V \simeq_{C^\infty} K^{-1}(D_1) \oplus K(-D_2). \tag{4.1}$$

The $\bar{\partial}$ -operator for the holomorphic structure of V can be written with respect to this smooth splitting as

$$\bar{\partial}_V = \begin{pmatrix} \bar{\partial}_1 & \beta \\ 0 & \bar{\partial}_2 \end{pmatrix}, \tag{4.2}$$

where β is a smooth $(0, 1)$ -form taking values in $\text{Hom}(K(-D_2), K^{-1}(D_1))$: its Dolbeault cohomology class $[\beta]$ corresponds to the extension class in $H^1(K^{-2}(D_1 + D_2))$ which determines the extension (2.4). When $[\beta] = 0$ it is easy to check that this gives a Hodge bundle [18], which we will denote by (E^∞, Φ^∞) . Here $E^\infty = V^\infty \oplus \underline{\mathbb{C}}$ where V^∞ is the holomorphically split bundle (4.1) and

$$\Phi_1^\infty : K^{-1}(D_1) \rightarrow V^\infty, \quad \Phi_2^\infty : V^\infty \rightarrow K(-D_2),$$

are the natural inclusion and projection respectively. Notice that $\Phi^\infty = \Phi$ as a map on smooth bundles.

Lemma 4.1 For $(E, \Phi) \in \mathcal{V}$, $\lim_{t \rightarrow \infty} (E, t\Phi) = (E^\infty, \Phi^\infty)$.

The proof is essentially the same as the proof of Prop. 4.9 in [17].

Proof We perform a gauge transformation. Set

$$g_t = \begin{pmatrix} t & 0 & 0 \\ 0 & t^{-1} & 0 \\ 0 & 0 & 1 \end{pmatrix}. \tag{4.3}$$

Then $g_t \Phi g_t^{-1} = t\Phi$ while

$$g_t^{-1} \bar{\partial}_E g_t = \begin{pmatrix} \bar{\partial}_1 & t^{-2} \beta & 0 \\ 0 & \bar{\partial}_2 & 0 \\ 0 & 0 & \bar{\partial} \end{pmatrix}. \tag{4.4}$$

Let E^t denote the holomorphic bundle for this $\bar{\partial}$ -operator. Then $(E, t\Phi) \simeq (E^t, \Phi)$ hence

$$\lim_{t \rightarrow \infty} (E, t\Phi) = \lim_{t \rightarrow \infty} (E^t, \Phi) = (E^\infty, \Phi^\infty).$$

□

Recall that in [18] the data $(c, D_1, D_2, [\beta])$ is used to parametrise the component $\mathcal{V}(d_1, d_2)$ by fixing the degrees of the divisors. Let $\bar{\mathcal{V}}(d_1, d_2)$ denote the closure of this in $\bar{\mathcal{V}}$, i.e., where D_1, D_2 can have points in common. By [18] this is a complex analytic family over Teichmüller space: let $\bar{\mathcal{V}}_c(d_1, d_2)$ denote the fibre over $c \in \mathcal{T}_g$. The previous lemma shows that, under the map from $\bar{\mathcal{V}}_c(d_1, d_2)$ to Higgs bundles over Σ_c , $(c, D_1, D_2, [t^{-2}\beta])$ maps to $(E, t\Phi)$ and therefore the limit as $t \rightarrow \infty$ is given by the trivial extension for V . Given that this parametrisation of Higgs bundles is smooth (which is proved in Lemma 5.1 below) we have the following corollary, which was conjectured in [18].

Corollary 4.2 $\bar{\mathcal{V}}_c(d_1, d_2)$ is diffeomorphic to the unstable manifold \mathcal{U}_C where C is the critical manifold of Hodge bundles for which $V = K^{-1}(D_1) \oplus K(-D_2)$ (i.e., the trivial extension).

Now we turn to limits as $t \rightarrow 0$. For a stable Higgs bundle (E, Φ) with Toledo invariant τ with $E = V \oplus \underline{\mathbb{C}}$ we note that $\tau = -\frac{2}{3} \deg(V)$ in our convention. Since the dual Higgs bundle has opposite Toledo invariant we may assume that $\tau \geq 0$.

Proposition 4.3 Let (E, Φ) be a stable Higgs bundle with $\tau \geq 0$ and let $(E^0, \Phi^0) = \lim_{t \rightarrow 0} (E, t\Phi)$. Suppose (E, Φ) is not a Hodge bundle and therefore is given by a non-trivial extension of the form (2.4) for effective divisors D_1, D_2 . Then exactly one of the following holds:

- (i) E is a semistable bundle, in which case $\tau = 0$ and $(E^0, \Phi^0) = (Gr(E), 0)$, where $Gr(E)$ is the associated graded bundle arising from a Jordan-Hölder filtration of E .
- (ii) $\tau > 0$ and every line subbundle L of V satisfies $\deg(L) < \frac{1}{3} \deg(V)$, in which case $(E^0, \Phi^0) = (E, \Phi_1)$ is a length two Hodge bundle.
- (iii) $\tau > 0$, and the maximal destabilizing line subbundle L of V has $\deg(L) = \frac{1}{3} \deg(V)$. In this case there is a positive divisor D for which $L \simeq K(-D_2 - D)$, $V/L \simeq K^{-1}(D_1 + D)$ and (E^0, Φ^0) is the polystable Higgs bundle

$$(K^{-1}(D_1 + D) \oplus \underline{\mathbb{C}}, \varphi) \oplus (K(-D_2 - D), 0), \tag{4.5}$$

where φ is the projection of $\Phi_1 : \underline{\mathbb{C}} \rightarrow V \otimes K$ to the quotient.

(iv) $\tau \geq 0$, V is unstable and its maximal destabilizing line subbundle L has $\deg(L) > \frac{1}{3} \deg(V)$. In this case there is a positive divisor D so that $L, V/L$ have the isomorphisms in (iii) but now (E^0, Φ^0) is a length three Hodge bundle of the form

$$(K^{-1}(D_1 + D) \oplus K(-D_2 - D) \oplus \mathbb{C}, \Phi^0), \tag{4.6}$$

where Φ_1^0 is the projection of Φ_1 to $(V/L) \otimes K$, and Φ_2^0 is the restriction of Φ_2 to L .

In case (i) the limit corresponds to a constant harmonic map. In all other cases except (ii) with $d_1 = 0$ (i.e., Φ_1 has no zeroes) the limit corresponds to a branched minimal surface. In case (iii) these are branched holomorphic maps to a totally geodesic $\mathbb{C}\mathbb{H}^1$ in $\mathbb{C}\mathbb{H}^2$.

Proof (i) If $E = V \oplus \mathbb{C}$ is a semistable bundle then $0 = \deg(\mathbb{C}) \leq \frac{1}{3} \deg(V)$. By assumption $\tau \geq 0$, hence $\tau = 0$. Now $(E, 0)$ is S -equivalent to $(Gr(E), 0)$ [20], which is polystable and hence the limit in moduli space. In particular, note that either V is polystable and $E = Gr(E)$, or V is strictly semistable and a non-trivial extension of its maximal degree line subbundle L of degree 0. Then a Jordan-Hölder filtration for E is $L \subset L \oplus \mathbb{C} \subset E$ and $Gr(E) = L \oplus \mathbb{C} \oplus V/L$. (ii) In this case write the holomorphic structure and Higgs field for $(E, t\Phi)$ in block decomposition with respect to $E = V \oplus \mathbb{C}$:

$$\bar{\partial}_E = \begin{pmatrix} \bar{\partial}_V & 0 \\ 0 & \bar{\partial} \end{pmatrix}, \quad \Phi = \begin{pmatrix} 0 & t\Phi_1 \\ t\Phi_2 & 0 \end{pmatrix}.$$

A simple computation shows that for

$$g_t = \begin{pmatrix} tI_V & 0 \\ 0 & 1 \end{pmatrix}$$

we have

$$\lim_{t \rightarrow 0} g_t^{-1} \bar{\partial}_E g_t = \bar{\partial}_E, \quad \lim_{t \rightarrow 0} g_t^{-1} t\Phi g_t = \begin{pmatrix} 0 & \Phi_1 \\ 0 & 0 \end{pmatrix}.$$

This represents the limit provided (E, Φ_1) is stable as a Higgs bundle. The Φ_1 -invariant subbundles of E are $\mathbb{C} \oplus \text{im } \Phi_1, V$ and any line subbundle of V . Since (E, Φ) is stable as a Higgs bundle and $\deg(V) < 0$ the first two already satisfy the slope inequality, as does $\text{im } \Phi_1$. So the additional condition is that every other line subbundle L of V satisfies $\deg(L) < \frac{1}{3} \deg(V)$. Note that the case $\tau = 0$ is covered by part (i).

(iii) First we note that $(V/L \oplus \mathbb{C}, \varphi)$ is a stable Higgs bundle, since V/L is the only φ -invariant proper subbundle and $\deg(V/L) = \frac{2}{3} \deg(V) < 0$ hence $\deg(V/L) < \frac{1}{2} \deg(V/L)$. Also $\frac{1}{2} \deg(V/L) = \deg(L)$ hence (4.5) is a polystable Higgs bundle. We also note that the restriction of Φ_2 to L is a holomorphic section of $L^{-1} \otimes K(-D_2)$ and this must have zeroes otherwise the extension (2.4) splits. Let $D > 0$ be the divisor of these zeroes, then $L \simeq K(-D_2 - D)$ and since $\det(V) \simeq \mathcal{O}(D_1 - D_2)$ it follows that $V/L \simeq K^{-1}(D_1 + D)$.

Now V can be written as an extension of the form

$$0 \rightarrow L \rightarrow V \rightarrow V/L \rightarrow 0. \tag{4.7}$$

This gives a C^∞ -isomorphism $E \simeq V/L \oplus L \oplus \mathbb{C}$. With respect to such a decomposition we can write

$$\bar{\partial}_E = \begin{pmatrix} \bar{\partial}_{V/L} & 0 & 0 \\ \beta & \bar{\partial}_L & 0 \\ 0 & 0 & \bar{\partial} \end{pmatrix}, \quad \Phi = \begin{pmatrix} 0 & 0 & \phi_{13} \\ 0 & 0 & \phi_{23} \\ \phi_{31} & \phi_{32} & 0 \end{pmatrix}.$$

Using the gauge transformation

$$g_t = \begin{pmatrix} t & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

a straightforward calculation gives

$$g_t^{-1} \bar{\partial}_E g_t = \begin{pmatrix} \bar{\partial}_{V/L} & 0 & 0 \\ t\beta & \bar{\partial}_L & 0 \\ 0 & 0 & \bar{\partial} \end{pmatrix}, \quad g_t^{-1}(t\Phi)g_t = \begin{pmatrix} 0 & 0 & \phi_{13} \\ 0 & 0 & t\phi_{23} \\ t^2\phi_{31} & t\phi_{32} & 0 \end{pmatrix}.$$

Setting $t = 0$ gives (4.5). Note that $\varphi = \varphi_{13}$ is the projection of $\Phi_1 : K^{-1}(D_1) \rightarrow V$ onto V/L and has divisor of zeroes $D_1 + D$ since it vanishes when either $\Phi_1 = 0$ or at the support of $\text{im } \Phi_1 \cap L$. But $\Phi_2 \circ \Phi_1 = 0$ so the support of D is all the points at which $\text{im } \Phi_1$ is not zero but lies in L .

(iv) As in the previous case we first show that the proposed limit (4.6) is a stable Higgs bundle. Since the only proper Φ^0 invariant subbundles are V/L and $V/L \oplus \underline{\mathbb{C}}$, and since $\text{deg}(V) \leq 0$, the stability condition is the single inequality $\frac{1}{2} \text{deg}(V/L) < \frac{1}{3} \text{deg}(V)$, i.e., $\text{deg}(L) > \frac{1}{3} \text{deg}(V)$. Just as in part (iii) we note that $L \simeq K(-D_2 - D)$ where $D > 0$ is the divisor of zeroes of Φ_2 restricted to L . Hence $V/L \simeq K^{-1}(D_1 + D)$.

Now to show this gives the limit we use the same argument as part (iii) but with

$$g_t = \begin{pmatrix} t^2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & t \end{pmatrix}.$$

In this case

$$g_t^{-1} \bar{\partial}_E g_t = \begin{pmatrix} \bar{\partial}_{V/L} & 0 & 0 \\ t^2\beta & \bar{\partial}_L & 0 \\ 0 & 0 & \bar{\partial} \end{pmatrix}, \quad g_t^{-1}(t\Phi)g_t = \begin{pmatrix} 0 & 0 & \phi_{13} \\ 0 & 0 & t^2\phi_{23} \\ t^2\phi_{31} & \phi_{32} & 0 \end{pmatrix}.$$

Setting $t = 0$ gives (4.6) with $\Phi_1^0 = \varphi_{13}$, the projection of Φ_1 onto $(V/L) \otimes K$, and $\Phi_2^0 = \varphi_{32}$, the restriction of Φ_2 to L . The latter has zeroes $D_2 + D$ by definition, and the former has zeroes $D_1 + D$ for the same reason as in part (iii). □

Note that this argument works perfectly well even when D_1, D_2 already have points in common (although D will not be the full branch point divisor in that case), and it tells us something about singularities of the nilpotent cone. When C consists of length three Hodge bundles \mathcal{U}_C is smooth and of dimension $5(g - 1)$: this follows either from the Morse index calculation of Gothen [9, Prop 3.2] or the dimension count in [18] for $\bar{V}_c(d_1, d_2)$, given Corollary 4.2. Any point on C which is a limit point as $t \rightarrow 0$ (i.e., from some $\mathcal{U}_{C'}$ for $C' \neq C$) lies on both the stable and unstable manifolds of C , which are transverse, and therefore the tangent space at this point has dimension greater than \mathcal{U}_C . Thus such limit points are singular points of the nilpotent cone. We can ask whether every length three Hodge bundle of the form (4.6) (i.e., with common divisor D) is such a singular point. The answer is no: by the following result there are many such Hodge bundles which are not limit

points. Recall from [18] that the necessary and sufficient conditions for (4.6) to give a stable Higgs bundle are the inequalities

$$0 \leq 2d'_1 + d'_2 < 6(g - 1), \quad 0 \leq d'_1 + 2d'_2 < 6(g - 1), \tag{4.8}$$

where $d'_j = \deg(D_j + D)$.

Proposition 4.4 *Let D_1, D_2, D be effective divisors, $D > 0$, set $D'_j = D_j + D$ and suppose their degrees d'_1, d'_2 satisfy $d'_1 \leq d'_2$ and $d'_1 + d'_2 < 2(g - 1)$. Then the Hodge bundle with V given by (4.6) does not lie at the limit as $t \rightarrow 0$ of a \mathbb{C}^\times -orbit in a different unstable manifold.*

Proof First, the two inequalities imply that (4.6) corresponds to a stable Higgs bundle with $\tau \geq 0$. By Prop. 4.3 such a limit can only be obtained from case (iv), where V has unique maximal destabilizing line subbundle $L \simeq K(-D'_2)$ and $V/L \simeq K^{-1}(D'_1)$. The existence of this requires

$$1 \leq \dim H^1(K^2(-D'_1 - D'_2)) = \dim H^0(K^{-1}(D'_1 + D'_2)),$$

using Serre duality. But the right hand dimension is zero whenever $\deg(K^{-1}(D'_1 + D'_2)) < 0$, i.e., when $d'_1 + d'_2 < 2(g - 1)$. □

5 Connected components of $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)$

Our aim is to prove Theorem 1.2. First we must recall some more facts from [18]. From [18] \mathcal{V} is a disjoint union $\cup_{(d_1, d_2)} \mathcal{V}(d_1, d_2)$ where the non-negative integers d_1, d_2 satisfy the inequalities (4.8). Points of $\mathcal{V}(d_1, d_2)$ are parametrised by the data (c, D_1, D_2, ξ) where D_1, D_2 are effective divisors of degree d_1, d_2 and ξ denotes the extension class for the extension (2.4). This class can be freely chosen in $H^1(\Sigma_c, K^{-2}(D_1 + D_2))$ and this parametrisation gives $\mathcal{V}(d_1, d_2)$ the structure of a complex analytic family over $c \in \mathcal{T}_g$. The fibre over $c \in \mathcal{T}_g$, which we will denote by $\mathcal{V}_c(d_1, d_2)$, is the holomorphic vector bundle over

$$\{(D_1, D_2) \in S^{d_1} \Sigma_c \times S^{d_2} \Sigma_c : D_1 \cap D_2 = \emptyset\}$$

with fibre $H^1(\Sigma_c, K^{-2}(D_1 + D_2))$.

We need to show first that this parametrisation is smooth with respect to the smooth structure of $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)$ given above, which uses its embedding in $\mathcal{T}_g \times \mathcal{R}(\pi_1 \Sigma, G)$. Since we are always dealing with points which are smooth in the latter we can use the nonabelian Hodge correspondence to identify the smooth locus of $\mathcal{R}(\pi_1, G)$ with the smooth locus of Higgs bundle moduli space $\mathcal{H}(\Sigma_c, G)$. It therefore suffices to prove the following lemma.

Lemma 5.1 *For each fixed $c \in \mathcal{T}_g$ we get a holomorphic embedding*

$$\mathcal{Z} : \overline{\mathcal{V}}_c(d_1, d_2) \rightarrow \mathcal{H}(\Sigma_c, G),$$

by assigning to (D_1, D_2, ξ) the Higgs bundle (E, Φ) for which $E = V \oplus \underline{\mathbb{C}}$ where V and Φ arise from the extension ξ using (2.4).

We know from [18] that \mathcal{Z} is one-to-one, so it suffices to show that it is a holomorphic immersion.

Before we begin the proof we need to fix a convention for which way to represent 1-cocycles with values in a vector bundle. For a vector bundle F with model fibre A and local transition relations $\psi_i = a_{ij} \psi_j$ between local trivialisations ψ_i over a Leray cover

$\{U_i\}$, let (ξ_{ij}, U_i, U_j) be a 1-cocycle for a class in $H^1(F)$, i.e., $\xi_{ij} \in \Gamma(U_i \cap U_j, F)$ satisfy $\xi_{ij} + \xi_{jk} = \xi_{ik}$. Then we choose to represent this by the local functions

$$c_{ij} = \psi_i(\xi_{ij}) : U_i \cap U_j \rightarrow A,$$

and the 1-cocycle conditions are equivalent to

$$c_{ij} + a_{ij}c_{jk} = c_{ik}.$$

Note that the opposite convention, to use $\psi_j(\xi_{ij})$ instead, is used in Gunning [11]. In what follows we need this for $H^1(\mathcal{A}^0)$ and $H^1(K^{-2}(D_1 + D_2))$.

Proof To describe the map \mathcal{Z} concretely about a point (D_1, D_2, ξ) , let P_1, \dots, P_d be the points in the support of $D_1 + D_2$ and let U_0, \dots, U_d be the Leray cover of Σ_c for which each U_j for $j \geq 1$ is an open disc about P_j , with $U_j \cap U_k = \emptyset$ for $j \neq k$, and $U_0 \cap U_j$ is an annulus excluding an open disc about P_j . Since V is determined by the extension class ξ , V has 1-cocycle $g = \{(g_{0j}, U_0, U_j) : 1 \leq j \leq d\}$ given by

$$g_{0j} = \begin{pmatrix} \alpha_{0j} & \lambda_{0j} \\ 0 & \beta_{0j} \end{pmatrix} \tag{5.1}$$

where

$$\alpha_{0j} = z^{-n_j} \frac{dz_0}{dz_j}, \quad \beta_{0j} = z_j^{m_j} \frac{dz_j}{dz_0}, \quad \lambda_{0j} = \beta_{0j} \xi_{0j} dz_0^2.$$

Here, for $j \geq 1$, z_j is a local parameter in U_j centred at P_j and n_j is the degree of D_1 at P_j while m_j is the degree of D_2 at P_j . By dz_0 we simply mean a non-vanishing holomorphic 1-form on U_0 . By ξ_{0j} we mean the local section of $K^{-2}(D_1 + D_2)$ over $U_0 \cap U_j$ which comprises a 1-cocycle representing the extension class ξ .

The 1-cocycle g determines a trivialisation χ_j of V , and dual χ_j^* of V^* , over U_j . When we think of the Higgs field components as $\Phi_1 \in H^0(V \otimes K)$ and $\Phi_2 \in H^0(V^* \otimes K)$ these satisfy

$$\chi_0(\Phi_1) = \begin{pmatrix} dz_0 \\ 0 \end{pmatrix}, \quad \chi_0^*(\Phi_2) = (0 \ dz_0),$$

and for $j \geq 1$

$$\chi_j(\Phi_1) = \begin{pmatrix} z_j^{n_j} dz_j \\ 0 \end{pmatrix}, \quad \chi_j^*(\Phi_2) = (0 \ z_j^{m_j} dz_j). \tag{5.2}$$

An open neighbourhood around $D_1 \in S^{d_1} \Sigma_c$ (respectively $D_2 \in S^{d_2} \Sigma_c$) is determined by collection of monic polynomials $z_j^{n_j} + u_j(z_j)$ (respectively $z_j^{m_j} + v_j(z_j)$) whose zeroes lie in U_j . Note that if P_j is not in the support of D_1 then $n_j = 0$ and u_j is the zero polynomial (and likewise for m_j, v_j when P_j is not in the support of D_2). The coefficients of the polynomials $u_j(z_j), v_j(z_j)$ provide a local chart about (D_1, D_2) . At a pair (D'_1, D'_2) in this neighbourhood of (D_1, D_2) any $\xi' \in H^1(K^{-2}(D'_1 + D'_2))$ can be represented by a 1-cocycle $\{(\xi'_{0j}, U_0, U_j)\}$. The map \mathcal{Z} maps (D'_1, D'_2, ξ') to the Higgs bundle constructed as above but with $z_j^{n_j}$ replaced by $z_j^{n_j} + u_j(z_j)$, $z_k^{m_k}$ replaced by $z_k^{m_k} + v_k(z_k)$, and ξ_{0j} replaced by ξ'_{0j} . It is easy to see that smooth variations of these parameters result in smooth variations of the Higgs bundle.

Our aim now is to show that $d\mathcal{Z}$ has trivial kernel at each point. First we describe the tangent space to $\overline{V}_c(d_1, d_2)$ at a given point. Since $\overline{V}_c(d_1, d_2)$ is a vector bundle over $S^{d_1} \Sigma_c \times S^{d_2} \Sigma_c$

we can fix a local trivialisation over a neighbourhood of (D_1, D_2, ξ) and identify the tangent space at that point with

$$T_{D_1}S^{d_1}\Sigma_c \oplus T_{D_2}S^{d_2} \oplus H^1(K^{-2}(D_1 + D_2)).$$

We will write elements of this in the form (w, y, η) . Further, the vector w (and similarly y) can be described as follows. It is well-known that

$$T_{D_1}S^{d_1}\Sigma_c = \bigoplus_j (\mathfrak{m}_{P_j}/\mathfrak{m}_{P_j}^{n_j+1}),$$

where $\mathfrak{m}_{P_j} \subset \mathcal{O}_{P_j}$ is the maximal ideal of locally holomorphic functions about P_j which vanish at P_j . We will identify

$$\mathfrak{m}_{P_j}/\mathfrak{m}_{P_j}^{n_j+1} \simeq \{z_j w_j(z_j) : w_j \in \mathbb{C}[z_j], \deg(w_j) \leq n_j - 1\},$$

and therefore write $w \in T_{D_1}S^{d_1}\Sigma_c$ as a tuple $w = (w_1, \dots, w_k)$ of polynomials. This is tangent to the curve $D_1(t)$ on $S^{d_1}\Sigma_c$ for which $D_1(t)$ is the divisor of zeroes given by the locally defined polynomial $z_j^{n_j} + t w_j(z_j)$ in U_j (for t sufficiently close to zero).

Given such a tangent vector (w, y, ξ) , let $(D_1(t), D_2(t), \xi(t))$ be the curve it is tangent to, obtained as above using $u_j = t w_j, v_j = t y_j$ and $\xi(t) = \xi + t \eta$ (using the local trivialisation of $\mathcal{V}_c(d_1, d_2)$). Let $\mathcal{Z}(t) = (E(t), \Phi(t))$ denote the image curve, with $E(t) = V(t) \oplus \mathbb{C}$, and let $g_{0j}(t)$ denote the transition functions for $V(t)$, in the form (5.1). These have entries

$$\alpha_{0j}(t) = \frac{1}{z^{n_j} + t w_j} \frac{dz_0}{dz_j}, \quad \beta_{0j}(t) = (z^{m_j} + t y_j) \frac{dz_j}{dz_0}, \quad \lambda_{0j}(t) = \beta_{0j} \xi(t)_{0j} dz_0^2.$$

In particular, $\chi_0(\Phi_1), \chi_0^*(\Phi_2)$ are time-independent. The derivative $\mathcal{Z}'(0)$ is represented by a hypercohomology class in

$$\mathbb{H}^1(\mathcal{A}^*) \simeq \frac{\{(A, B) \in \mathcal{C}^1(\mathcal{A}^0) \oplus \mathcal{C}^0(\mathcal{A}^1) : \delta A = 0, \delta B = [\Phi, A]\}}{\{(\delta C, [\Phi, C]) : C \in \mathcal{C}^0(\mathcal{A}^0)\}},$$

following [3]. Specifically, it is the class of a pair (A, B) for which A is represented by the 1-cocycle

$$g'(0)g^{-1} = \begin{pmatrix} \alpha'(0)\alpha^{-1} & -\lambda\alpha'(0)\alpha^{-1}\beta^{-1} + \lambda'(0)\beta^{-1} \\ 0 & \beta'(0)\beta^{-1} \end{pmatrix} \tag{5.3}$$

where we have dropped the Čech cocycle subscripts for notational simplicity. If we write $B = (B_1, B_2)$ where $B_j = \Phi'_j(0)$ then explicit calculation gives

$$\begin{aligned} \chi_0(B_1) &= \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad \chi_0^*(B_2) = (0 \ 0), \\ \chi_j(B_1) &= \begin{pmatrix} w_j dz_j \\ 0 \end{pmatrix}, \quad \chi_j^*(B_2) = (0 \ y_j dz_j). \end{aligned} \tag{5.4}$$

If $\mathcal{Z}'(0) = 0$ there must exist $C \in \mathcal{C}^0(\mathcal{A}^0)$ for which $A = \delta C$ and $B = [\Phi, C]$. We may assume C is represented by an upper triangular 0-cocycle of the form

$$\begin{pmatrix} a_j & b_j \\ 0 & c_j \end{pmatrix},$$

where the entries are holomorphic functions in U_j . The Lie bracket $[\Phi, C]$ corresponds to the pair $(-C\Phi_1, \Phi_2C)$ and calculation gives

$$\chi_j(C\Phi_1) = \begin{pmatrix} a_j z^{n_j} dz_j \\ 0 \end{pmatrix}, \quad \chi_j^*(\Phi_2C) = (0 \ c_j z^{m_j} dz_j).$$

Comparing this with (5.4), since $\deg(w_j) < n_j$ and $\deg(y_j) < m_j$ it follows that $B = [\Phi, C]$ if and only if both sides are zero, i.e., $a_j = 0 = c_j$ and w_j, y_j are both identically zero. This in turn means that the 1-cocycle in (5.3), which represents A , is

$$\begin{pmatrix} 0 & \eta_{0j} dz_0^2 \\ 0 & 0 \end{pmatrix}.$$

Now $A = \delta C$ is the condition that the 1-cocycle $\eta \in H^1(K^{-2}(D_1 + D_2))$ is trivial. Thus the kernel of dZ is trivial. □

It follows that each $\mathcal{V}(d_1, d_2)$ is smooth and connected as a submanifold of \mathcal{V} , and each is open since they have the same dimension. Thus we conclude:

Lemma 5.2 *The smooth submanifold $\mathcal{V} \subset \mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)$ is a disjoint union of connected components $\mathcal{V}(d_1, d_2)$.*

To complete the proof of Theorem 1.2 we need two more lemmas. Both concern $\tau > 0$. The first says that $\mathcal{V}(d_1, d_2)$ is disconnected from \mathcal{W}_τ when $d_1 \neq 0$. The second says $\mathcal{V}(0, d_2)$ is connected to \mathcal{W}_τ .

Lemma 5.3 *For $\tau > 0$, $\overline{\mathcal{V}}(d_1, d_2) \cap \mathcal{W}_\tau = \emptyset$ when $d_1 \neq 0$.*

Proof It suffices to prove that this is true for each fixed conformal structure c . So fix c and $\tau > 0$, and let C be the critical manifold of minima for the Higgs field energy $\mathfrak{E} = \|\Phi\|_{L^2}^2$ for this value of τ . Let $C_0 \subset C$ be the open subset of Higgs bundles for which Φ has no zeroes. Let C' be the critical manifold of \mathfrak{E} for which $\overline{\mathcal{V}}_c(d_1, d_2)$ corresponds to $\overline{U}_{C'}$, the closure of the unstable manifold of the downward gradient flow of \mathfrak{E} (equally, the unstable manifold for the \mathbb{C}^\times -action). The assertion is that $\mathcal{U}_{C'} \cap C_0 = \emptyset$ whenever $d_1 \neq 0$. Suppose $(E, \Phi) \in \overline{U}_{C'} \cap C_0$, then it is the limit of a sequence (E_k, Φ_k) of Higgs bundles in $\mathcal{U}_{C'}$. We may assume without loss of generality that each $\|\Phi_k\|_{L^2}^2$ lies below the penultimate critical value of \mathfrak{E} , and therefore all (E_k, Φ_k) lie in the stable manifold \mathcal{S}_C . Taking the limit as $t \rightarrow 0$ is a continuous map (since $\pi : \mathcal{S}_C \rightarrow C$ is a vector bundle) and therefore

$$(E, \Phi) = \lim_{k \rightarrow \infty} (E_k^0, \Phi_k^0),$$

where $(E_k^0, \Phi_k^0) = \lim_{t \rightarrow 0} (E_k, t\Phi_k)$. By Prop. 4.3 each Φ_k^0 has zeroes unless $d_1 = 0$. Therefore (E, Φ) lies in the closed subset $C \setminus C_0$. □

Lemma 5.4 *For every choice of $0 < d_2 < 3(g-1)$ there is a Higgs bundle (E, Φ) in $\mathcal{V}(0, d_2)$ for which $\lim_{t \rightarrow 0} (E, t\Phi)$ lies in \mathcal{W}_τ for $\tau = \frac{2}{3}d_2$.*

Proof Since $d_1 = 0$ the Higgs bundle (E, Φ) must correspond to an extension of the form

$$0 \rightarrow K^{-1} \xrightarrow{\Phi_1} V \xrightarrow{\Phi_2} K(-D_2) \rightarrow 0,$$

where $\text{deg}(D_2) = d_2$. By Prop. 4.3(ii) it suffices to show we can find such an extension for which V is a stable bundle. This is equivalent to the existence of an extension

$$0 \rightarrow \underline{\mathbb{C}} \rightarrow \lambda \rightarrow K^2(-D_2) \rightarrow 0,$$

for which $s(\lambda) = \text{deg}(\lambda) - 2 \max \text{deg}(L) \geq 1$, where the maximum is taken over all line subbundles of λ . Note that such extensions are parametrised by the space $H^1(K^{-2}(D_2)) \simeq H^0(K^3(-D_2))^*$. Such a situation is covered by [15, Prop 1.1]. Specifically, define $d = \text{deg}(K^2(-D_2)) = 4(g - 1) - d_2$ and let $s = 1$ when d is odd and $s = 2$ when d is even. This satisfies the conditions $4 - d \leq s \leq d$ required to apply [15, Prop 1.1] (in particular note that the critical case $4 - d = d$ can only occur for $d = 2$, in which case $4 - d = s = d$). Then by [15, Prop 1.1] $s(\lambda) \geq s$ if and only if the secant variety

$$\text{Sec}_{\frac{1}{2}(d+s-2)}(\Sigma_c) \subseteq \mathbb{P}H^0(K^3(-D_2))^*,$$

is a proper subvariety. Since $\text{deg}(K^3(-D_2)) > 2(g - 1)$ the Riemann-Roch theorem gives the dimension of $\mathbb{P}H^0(K^3(-D_2))^*$ as $5(g - 1) - d_2 - 1 = d + g - 2$. By [13] this secant variety has dimension $d + s - 3 \leq d - 1$ is therefore a proper subvariety for $g \geq 2$. \square

6 Open questions

Now that we understand the overall structure of $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)$ there are some important questions which deserve attention. I want to focus mainly on those that come from considering the natural projection

$$\mathcal{F}_2 : \mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2) \rightarrow \mathcal{R}(\pi_1 \Sigma, G); \quad [f, c, \rho] \mapsto \rho. \tag{6.1}$$

The domain and codomain have the same ‘‘typical’’ dimension. Goldman and Wentworth [8] proved that the image contains all convex cocompact representations, and for $PU(2, 1)$ convex cocompact is equivalent to (complex hyperbolic) quasi-Fuchsian (equally, quasi-isometric embedding; equally, Anosov embedding [10]). These representations are interesting because they act properly discontinuously on $\mathbb{C}\mathbb{H}^2$, so that in our context taking the quotient by this action produces a compact minimal surface in a locally symmetric space.

In general the set of Anosov representations is open [14, Thm 2.1], but very little is known about the structure of the set quasi-Fuchsian representations for $PU(2, 1)$, aside from Toledo’s proof that all representations with maximal invariant are actually \mathbb{C} -Fuchsian (they factor through $PU(1, 1) \rightarrow PU(2, 1)$). For non-maximal Toledo invariant:

- (i) The Hodge bundles with $d_1 = 0 = d_2$ correspond to totally geodesic Lagrangian embeddings: these lie at the ‘‘top’’ of $\mathcal{V}(0, 0)$ with respect to the downwards Morse flow. Under \mathcal{F}_2 they map to so-called \mathbb{R} -Fuchsian representations, those which factor through $SO(2, 1) \subset PU(2, 1)$. In an open tubular neighbourhood of these lie the almost \mathbb{R} -Fuchsian minimal surfaces of [18] whose representations were shown to be quasi-Fuchsian. The map \mathcal{F}_2 is invertible on this open neighbourhood. Parker & Platis [19] also constructed an open neighbourhood of quasi-Fuchsian representations around the \mathbb{R} -Fuchsian representations using a different construction.
- (ii) Goldman et al. [7] constructed quasi-Fuchsian representations for every even value of the Toledo invariant by glueing together disc bundles to construct the quotient spaces of $\mathbb{C}\mathbb{H}^2$. Anan’ in et al. [2] constructed different hyperbolic disc bundles for every non-maximal value of the Toledo invariant, which they conjecture possess holomorphic

(therefore minimal) sections. The corresponding representations are discrete and faithful.

With these facts in mind, the main questions are:

- (i) What properties of an equivariant minimal immersion ensure that the representation is quasi-Fuchsian? One expects that it must be a proper embedding, but is this sufficient?
- (ii) When does \mathcal{F}_2 fail to be injective, and what characterises the condition $d\mathcal{F}_2 = 0$?
- (iii) For the hyperbolic disc bundles in [2, 7] we are told their Toledo invariant and Euler number. Each admits at least one equivariant minimal immersion: how is $\chi(T\Sigma^\perp)$ related to that Euler number? One expects them to be equal.
- (iv) Does the image of every connected component of $\mathcal{M}(\Sigma, \mathbb{C}\mathbb{H}^2)$ contain a quasi-Fuchsian representation?
- (v) Are there any quasi-Fuchsian representations which only admit branched minimal surfaces?

Acknowledgements The initial stages of this research were carried out while the author was a visitor at the Simons Center for Geometry and Physics, Stony Brook University, in April 2019 during the programme “The Geometry and Physics of Hitchin Systems”, and then subsequently at the MFO Oberwolfach workshop “The Geometry and Physics of Higgs Bundles” in May 2019. I gratefully acknowledge the support of these institutions, and thank the organisers for the opportunity to participate. I also thank Andy Sanders and John Loftin for the stimulating discussions during the SCGP programme which raised some of the questions answered here.

Data availability Data sharing is not applicable to this article as no datasets were generated or analysed for this research.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Alessandrini, D., Collier, B.: The geometry of maximal components of the $PSp(4, \mathbb{R})$ character variety. *Geom. Topol.* **23**(3), 1251–1337 (2019)
2. Anan’in, S., Grossi, C., Gusevskii, N.: Complex hyperbolic structures on disc bundles over surfaces. *Int. Math. Res. Not. IMRN* **19**, 4295–4375 (2011)
3. Biswas, I., Ramanan, S.: An infinitesimal study of the moduli of Hitchin pairs. *J. Lond. Math. Soc.* **49**, 219–231 (1994)
4. Bradlow, S.B., Garcia-Prada, O., Gothen, P.B.: Surface group representations and $U(p, q)$ -Higgs bundles. *J. Differ. Geom.* **64**, 111–170 (2003)
5. Corlette, K.: Flat G -bundles with canonical metrics. *J. Differ. Geom.* **28**, 361–382 (1988)
6. Donaldson, S.K.: Twisted harmonic maps and the self-duality equations. *Proc. Lond. Math. Soc.* **3**, 127–131 (1987)
7. Goldman, W., Kapovich, M., Leeb, B.: Complex hyperbolic manifolds homotopy equivalent to a Riemann surface. *Commun. Anal. Geom.* **9**(1), 61–95 (2001)
8. Goldman, W., Wentworth, R.: Energy of twisted harmonic maps of Riemann surfaces. *Contemp. Math.* **432**, 45 (2007)
9. Gothen, P.B.: Topology of $U(2, 1)$ representation spaces. *Bull. Lond. Math. Soc.* **34**, 729–738 (2002)
10. Guichard, O., Wienhard, A.: Anosov representations: domains of discontinuity and applications. *Invent. Math.* **190**(2), 357–438 (2012)

11. Gunning, R.: Lectures on Vector Bundles over Riemann Surfaces. Princeton University Press, Princeton, New Jersey (1967)
12. Hitchin, N.: Critical loci for Higgs bundles. *Commun. Math. Phys.* **366**, 841–864 (2019)
13. Lange, H.: Higher secant varieties of curves and the theorem of Nagata on ruled surfaces. *Manuscripta Math.* **47**, 263–269 (1984)
14. Labourie, F.: Anosov flows, surface groups and curves in projective space. *Invent. Math.* **165**(1), 51–114 (2006)
15. Lange, H., Narasimhan, M.S.: Maximal subbundles of rank two vector bundles on curves. *Math. Ann.* **266**, 55–72 (1983)
16. Loftin, J., McIntosh, I.: Minimal Lagrangian surfaces in $\mathbb{C}\mathbb{H}^2$ and representations of surface groups into $SU(2, 1)$. *Geom. Dedic.* **162**, 67–93 (2013)
17. Loftin, J., McIntosh, I.: The moduli spaces of equivariant minimal surfaces in $\mathbb{R}\mathbb{H}^3$ and $\mathbb{R}\mathbb{H}^4$ via Higgs bundles. *Geom. Dedic.* **201**, 325–351 (2018)
18. Loftin, J., McIntosh, I.: Equivariant minimal surfaces in $\mathbb{C}\mathbb{H}^2$ and their Higgs bundles. *Asian J. Math.* **23**, 71–106 (2019)
19. Parker, J.R., Platis, I.: Open sets of maximal dimension in complex hyperbolic Quasi-Fuchsian space. *J. Differ. Geom.* **73**, 319–350 (2006)
20. Seshadri, C.S.: Space of unitary vector bundles on a compact Riemann surface. *Ann. Math.* **85**, 303–336 (1967)
21. Toledo, D.: Representations of surface groups in complex hyperbolic space. *J. Differ. Geom.* **29**, 125–133 (1989)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.