



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/199441/>

Version: Published Version

Article:

Hock, E.S., Franklin, M., Baxter, S. et al. (2023) Covariates of success in quitting smoking: a systematic review of studies from 2008 to 2021 conducted to inform the statistical analyses of quitting outcomes of a hospital-based tobacco dependence treatment service in the United Kingdom. *NIHR Open Research*, 3. 28. ISSN: 2633-4402

<https://doi.org/10.3310/nihropenres.13427.1>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>


Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



SYSTEMATIC REVIEW

Covariates of success in quitting smoking: a systematic review of studies from 2008 to 2021 conducted to inform the statistical analyses of quitting outcomes of a hospital-based tobacco dependence treatment service in the United Kingdom [version 1; peer review: awaiting peer review]

Emma S. Hock, Matthew Franklin, Susan Baxter, Mark Clowes, James Chilcott, Duncan Gillespie 

School of Health and Related Research, The University of Sheffield, Sheffield, England, UK

V1 **First published:** 19 May 2023, 3:28
<https://doi.org/10.3310/nihropenres.13427.1>
Latest published: 19 May 2023, 3:28
<https://doi.org/10.3310/nihropenres.13427.1>

Abstract

Background: Smoking cessation interventions are being introduced into routine secondary care in the United Kingdom (UK), but there are person and setting-related factors that could moderate their success in quitting smoking. This review was conducted as part of an evaluation of the QUIT hospital-based tobacco dependence treatment service (<https://sybics-quit.co.uk>). The aim of the review was to identify a comprehensive set of variables associated with quitting success among tobacco smokers contacting secondary healthcare services in the UK who are offered support to quit smoking and subsequently set a quit date. The results would then be used to inform the development of a statistical analysis plan to investigate quitting outcomes.

Methods: Systematic literature review of five electronic databases. Studies eligible for inclusion investigated quitting success in one of three contexts: (a) the general population in the UK; (b) people with a mental health condition; (c) quit attempts initiated within a secondary care setting. The outcome measures were parameters from statistical analysis showing the effects of covariates on quitting success with a statistically significant (i.e., p -value <0.05) association.

Results: The review identified 29 relevant studies and 14 covariates of quitting success, which we grouped into four categories: demographics (age; sex; ethnicity; socio-economic conditions; relationship status, cohabitation and social network), individual health status and healthcare setting (physical health, mental health), tobacco smoking variables (current tobacco consumption, smoking history, nicotine dependence; motivation to quit; quitting history), and

Open Peer Review

Approval Status *AWAITING PEER REVIEW*

Any reports and responses or comments on the article can be found at the end of the article.

intervention characteristics (reduction in amount smoked prior to quitting, the nature of behavioural support, tobacco dependence treatment duration, pharmacological aids).

Conclusions: In total, 14 data fields were identified that should be considered for inclusion in datasets and statistical analysis plans for evaluating the quitting outcomes of smoking cessation interventions initiated in secondary care contexts in the UK.

PROSPERO registration: CRD42021254551 (13/05/2021)

Keywords

smoking cessation, hospital, tobacco dependence, service evaluation

Corresponding author: Duncan Gillespie (duncan.gillespie@sheffield.ac.uk)

Author roles: **Hock ES:** Investigation, Writing – Original Draft Preparation, Writing – Review & Editing; **Franklin M:** Conceptualization, Funding Acquisition, Investigation, Project Administration, Supervision, Validation, Writing – Review & Editing; **Baxter S:** Funding Acquisition, Investigation, Writing – Review & Editing; **Clowes M:** Methodology; **Chilcott J:** Funding Acquisition, Validation, Writing – Review & Editing; **Gillespie D:** Funding Acquisition, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This study was supported by charity funding from Yorkshire Cancer Research as part of a commissioned evaluation of the QUIT hospital-based tobacco dependence treatment service (<https://sybics-quit.co.uk/>) (SA/R117), and by funds from Research England to generate knowledge to enhance the impact of this work [QR-Policy Support Fund]. Article processing charges were paid by the University of Sheffield Institutional Open Access Fund. Those funding the study had no involvement in its design, interpretation or the decision to submit this manuscript for publication.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2023 Hock ES *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Hock ES, Franklin M, Baxter S *et al.* **Covariates of success in quitting smoking: a systematic review of studies from 2008 to 2021 conducted to inform the statistical analyses of quitting outcomes of a hospital-based tobacco dependence treatment service in the United Kingdom [version 1; peer review: awaiting peer review]** NIHR Open Research 2023, 3:28 <https://doi.org/10.3310/nihropenres.13427.1>

First published: 19 May 2023, 3:28 <https://doi.org/10.3310/nihropenres.13427.1>

Plain english summary

Stop smoking interventions are being incorporated as a systematic and opt-out component of secondary care services in the UK's National Health Service (NHS), driven by the NHS's Long Term Plan. This review was conducted as part of an evaluation of the QUIT hospital-based tobacco dependence treatment service (<https://sybics-quit.co.uk>). To support the development of statistical analyses to find out what affects smokers' success in quitting smoking after contacting the service, research was needed to identify what characteristics of the individual smokers and the healthcare setting might be important for success in quitting. The main purpose of the review was to support the development of a statistical analysis plan of quitting outcomes. We looked at academic papers published between 2008 and 2021 that estimated the influence of different factors on success in quitting smoking. The results of the review summarise the list of factors that previous studies have found to have an influence on quitting outcomes. The list of factors was used to inform discussions about what data fields it would be important for the service to collect because that data could be important for helping the service to understand variation in quitting outcomes.

Introduction

Stop smoking interventions are increasingly being incorporated as a systematic and opt-out component of secondary healthcare services in the United Kingdom's (UK's) National Health Service (NHS), driven by a commitment to do so in the NHS's Long Term Plan¹⁻³. The general specification of the service pathway in acute inpatient settings is: (i) on admission, determine if the patient smokes; (ii) provide advice and treatment to support patient smokers not to smoke whilst in hospital; (iii) provide follow-up support after discharge from hospital to support the patient to quit smoking completely. This service pathway is based on the "Ottawa Model", following the early implementation of a hospital based tobacco dependence treatment service in Ottawa, Canada⁴, and subsequent implementation in the UK by the CURE service in Greater Manchester⁵. An evaluation framework for hospital based smoking cessation services in the UK was developed by consensus among UK stakeholders in acute and mental health NHS hospital Trusts⁶, and provides a guide to the key data fields to collect for service monitoring and evaluation. However, there is no specific guidance on what data fields might be important when undertaking "deep dives" into the data to investigate factors that might influence quitting success, which in this review we generically group under the term 'covariates' of quitting success. Without a comprehensive list of potentially influential covariates, there is a risk that important data fields might be omitted from the routine collection of service data or from statistical analyses that aim to investigate quitting outcomes.

The current best evidence on the covariates of tobacco smoking quit success comes from a systematic review by Vangeli *et al.*⁷, which examined worldwide evidence among the adult general population. The evidence presented by Vangeli *et al.* highlighted decreased quit success among smokers with higher nicotine dependence, smokers who smoked

more cigarettes each day, smokers who had made a previously unsuccessful quit attempt, and smokers who had not previously gone without smoking for a week or more. Older age and higher socio-economic status or income were also found by the review to be associated with higher quit success. However, there could also be factors specific to patient health, healthcare setting, and the features of smoking cessation interventions initiated in secondary care settings that Vangeli *et al.*'s review of factors in the general population did not include. For example, in the British Thoracic Society's national audits of smoking and smoking cessation intervention activities in acute NHS hospital Trusts⁸⁻¹⁰, the key characteristics that were used to describe variation in whether current smokers received care for their tobacco dependence were gender, age, consultant speciality, and the patients' route of contact with the secondary care service (elective / emergency).

This review was designed to support the evaluation of smoking cessation services in secondary care settings in the UK by identifying covariates worth considering in plans for the statistical analysis of quit success following contact with a hospital-based stop smoking advisor. The review was instigated by the need to identify key variables to include in the statistical analysis of quitting outcomes as part of an evaluation of the QUIT hospital-based tobacco dependence treatment service (<https://sybics-quit.co.uk>). The review was based on the question: 'What patient-, service- and setting-related factors influence the success of a quit attempt, including when initiated in a secondary care setting?' The populations of most interest were the UK and Canada, given that the Canadian Ottawa model is the exemplar for UK services. The review question and population restrictions aimed to capture covariates of quitting success relevant to the UK general population, relevant to people with a mental health condition in any setting and in any country, and relevant to care for tobacco dependence initiated within a secondary acute or mental health service in any country. Within each study identified, the sign of the statistical coefficient for each variable investigated was taken as a measure of the direction of its association with quitting success, and the statistical significance of that coefficient at the 95% level was used to indicate if the association was potentially identified by chance or not.

Methods

Patient and Public Involvement

Patients and the public were not involved in this review.

Study design

We undertook a systematic review of studies that used a statistical model to explore what covariates are associated with quitting success. We followed a systematic review approach but the review did make compromises as it was conducted as part of the process of the evaluation of a particular service and needed to fit into the time and resources available. These compromises were guided by the rapid review approach recommended by Tricco *et al.*^{11,12}: searching more than one database in one iteration, published literature, searches limited by date and language, research scope specified by two

researchers and a health librarian, and study selection and data abstraction by one reviewer and one verifier. Quality appraisal of studies was based on whether the reporting of statistical analysis was sufficient to provide estimates of the coefficient for each variable investigated and its statistical significance at the 95% level. The review approach taken thus aimed to produce a synthesis of available knowledge that was sufficient to meet the review's aim more quickly, ensuring logistical feasibility alongside restricted timelines, while minimising risk of bias^{11,13}. The protocol was registered on PROSPERO CRD42021254551 on 13th May 2021. Reporting follows PRISMA principles (<http://www.prisma-statement.org/>) (see *Extended data*¹⁴).

Definition of covariates, effect size, and statistical significance

We defined a covariate of quitting success (that we term a 'factor') as any independent variable that can strengthen, diminish, negate, or otherwise alter the association between independent and dependent variables (in this study, the dependent variables quantify success in quitting smoking)¹⁵.

As the dependent variable is binary (i.e., quit achieved or not by a particular time after initiating the quit attempt), we assumed that the most common statistical analysis conducted would be a form of logistic regression with effect sizes presented as odds-ratios (ORs) or unconverted beta coefficients. For descriptive purposes, when discussing effect sizes we use the following terminology whereby the binary 'outcome' is quitting success¹⁶:

- 'Equal odds' when $OR=1$; i.e., exposure does not affect odds of outcome
- 'Higher odds' when $OR>1$; i.e., exposure associated with higher odds of outcome
- 'Lower odds' when $OR<1$; i.e., exposure associated with lower odds of outcome

In keeping with the review's aim to identify a list of potentially important covariates of quitting success, we focused on identifying which covariates have been estimated to have a statistically significant relationship with quitting success (with statistical significance defined as $p < 0.05$) rather than focussing on effect size magnitude. We define 'no relationship' as meaning that a covariate did not have a statistically significant relationship with quit success (i.e., $p \geq 0.05$). We did not consider whether a relationship is causal or not, as we were interested only in association. If a study presented both univariate and multivariate analyses, we based the identification of important covariates on the multivariate analysis as this adjusts for the associations of other variables with quitting success.

Eligibility criteria

Inclusion was restricted to studies published in peer-reviewed journals, in English, and dating from 2008, the year of the National Institute of Health and Care Excellence Guidance PH10 (for England and Wales), in which Recommendation 8

stated that smoking cessation advice and support should be available in secondary care settings for everyone who smokes. Reviews were not included, but we checked references for any relevant studies. We included studies that presented statistical estimates of the effects of covariates on the success of a quit attempt.

We searched for studies statistically assessing quit attempts in three contexts: (a) the general population instigated in any setting within the UK; (b) people with a mental health condition instigated in any setting and in any country; (c) initiated within a secondary acute or mental health service in any country. The scope of (a) was limited to the UK for relevance and feasibility given the large number of studies worldwide.

Information sources

Searches were conducted in April 2021. A focused search strategy combining free-text terms with subject headings (e.g., MeSH) was run and translated for optimal effectiveness across the following databases: MEDLINE (including In-Process and Epub ahead of print); EMBASE; PsycINFO (all via Ovid); CINAHL (via EBSCO) and the Cochrane Library.

Search process

The search strategy was constructed around the facets of: Smoking cessation AND quitting success AND (UK OR mental health OR hospital setting). Due to the time-constrained nature of this review, searches prioritised specificity over sensitivity, but to mitigate the risk of missing relevant papers the strategy was validated against six studies already known by the authors to be potentially relevant: Le Grande *et al.*¹⁷, Lubitz *et al.*¹⁸, Ussher *et al.*¹⁹, Smit *et al.*²⁰, Vangeli *et al.*⁷, and Zhou *et al.*²¹. All six studies were retrieved by the search (see the *Extended data*¹⁴). Database search results were extracted directly to reference management software.

Study selection

Screening for studies relevant to each of our three contexts (a–c) was performed simultaneously, with included studies marked for relevance to each. Titles and abstracts were screened by one of three reviewers (EH, MF or SB); 70% of abstracts were checked by another reviewer (EH or MF). Full texts were assessed for inclusion by one reviewer and checked by another reviewer (EH or MF). Disagreements were resolved through discussion, with no need to involve a third reviewer.

Data extraction and synthesis

EH and MF designed and tested a spreadsheet for data extraction. Data were extracted and charted by EH and checked in regular meetings with MF and DG. The following data items were extracted: Reference information (first author and date), study type, country, setting (e.g., hospital type/department/ward), participant baseline characteristics (e.g., age, sex, socio-economic status, reason for admission, cigarettes/day smoked, number of previous quit attempts, nicotine dependence), measure of quit success (point prevalence abstinence or continuous abstinence, any time point but recorded separately per time-point). Relevant characteristics of the analysis were noted. For example, method of data collection, sample size,

time horizon, cessation time-point, measure of abstinence, whether ORs and model coefficients were captured, the model type, and whether a univariate or multivariate model. Detailed statistical results were also extracted: the whole model, where reported, including intercept and other coefficients, dependent and independent variable, any reported *p*-values, and goodness of fit statistics, if reported.

During the data extraction process, we began to develop an organisational framework by categorising studies according

to our three contexts, the covariates investigated and their effects on quit success. The organisational framework was then revised as results synthesis progressed. Covariates were grouped according to our final organisational framework.

Results

From 2,499 retrieved records, 29 studies were included in the synthesis (Figure 1), representing 21 studies relating to the UK general population context, six studies relating to mental health in the UK or Canada, and two studies relating to




Figure 1. PRISMA flow diagram of study inclusion.

secondary care in the UK or Canada. A list of excluded studies with reasons is available in Table S1 in Supplement 2 of the *Extended data*¹⁴.

Description of included studies

The characteristics of the included studies and participants’ characteristics are summarised in Table S2 and Table S3 in Supplement 2 of the *Extended data*¹⁴. Most studies had prospective, cross-sectional or retrospective designs; three studies were randomised controlled trials (RCTs).

Methodological differences between studies

Methodological differences are reported in Table S4 in Supplement 2 of the *Extended data*¹⁴. Smoking cessation was assessed in a variety of different ways across studies. The time horizon for reporting smoking abstinence following a quit attempt ranged from 2 weeks to 1 year. Abstinence was assessed as both point-prevalent and continuous, both by self-report (most frequently used for continuous abstinence) and validated by expired air carbon monoxide (CO; most frequently used to verify 7-day or 2-week point-prevalent abstinence, at ≤10 or ≤8 ppm). If a study conducted separate analyses for different durations of abstinence following a quit attempt, we reported the findings from each analysis independently. All

studies reported odds-ratios from a logistic regression, and two studies reported beta coefficients.

In terms of sample, the majority of UK studies were of the general population (15 studies) or community smoking cessation services (four studies), with three studies examining samples with specific characteristics (i.e., pregnant women, people aged 25–59 years, and English residents of Bangladeshi origin; see Table S3 in Supplement 2 of the *Extended data*¹⁴). Mental health population studies were from Canada and sampled from people attending community mental health services (four studies) or from the general population (two studies). The two secondary care studies recruited participants from a Canadian hospital-based smoking cessation clinic or UK cardiac rehabilitation setting.

Covariates of success in quitting tobacco smoking

Figure 2 summarises the covariates that had a statistically significant relationship with quit attempt success. Table S5 in Supplement 2 of the *Extended data*¹⁴ summarises the relationships between covariates and quit success. Table S6 in Supplement 2 of the *Extended data*¹⁴ provides a full description of the size and direction of covariate effects and the corresponding statistical significance.




Figure 2. List of covariates found to have a statistically significant association with quitting success in at least one study. Table S6 in Supplement 2 of the *Extended data*¹⁴ provides a full description of the size and direction of covariate effects and the corresponding statistical significance.

Demographics. Overall, 16 studies included demographic covariates; the factors related to quit outcome were age, sex, ethnicity, socioeconomic characteristics, smoker's relationship status, cohabitation and social network situation (Table S5 and Table S6 in Supplement 2 of the *Extended data*¹⁴).

Age. All studies showed higher odds of quit success with increasing age²²⁻²⁶. Six analyses reported in five papers found no relationship between age and quit success in the UK general population²⁷⁻³¹, two studies found no relationship for age in people with mental health conditions^{32,33}, and two studies found no relationship in a secondary care setting^{34,35}.

Sex. There were inconsistent findings for sex: in the UK general population, three studies reported higher odds of quitting success for males^{22,24,29} and two studies reported higher odds of quitting success for females^{25,27}. Two studies in an outpatient setting (cardiology and mental health services) found higher odds of quitting success in males^{34,36}. Six studies found no relationship between sex and quitting success in the UK general population^{23,26,28,30,31,37}, and two studies found no relationship in people with mental health conditions^{32,33}.

Ethnicity. One study reported higher odds of quitting success for Black ethnicity vs. White British ethnicity²⁴. One study reported no relationship between ethnicity and quitting success in the UK general population³⁰.

Socioeconomic characteristics. There was a varied definition of socioeconomic characteristics in the studies identified. Higher odds of quitting success were reported for people: with higher social grades^{24,28-30,38}; living in less deprived areas²⁶; higher income^{37,39}; higher occupational grades^{22,39}; more education^{27,39}; who paid for prescriptions vs. were exempt^{22,23}; had a higher reading level³⁰; people whose mothers worked in higher grade occupations during their childhood³⁹; and people who did not live in social housing⁴⁰. In the UK general population, one study reported no relationship between quitting success and the geographic Index of Multiple Deprivation (IMD) score for the location of the smoking cessation service²², five studies reported no relationship between quitting success and education^{25,27,30,37,39}, one study for prescription exemption status²⁵, and one study for employment status²⁵. In a secondary care setting, two studies reported no relationship between quitting success and the employment status of patients^{34,35}.

Relationship status, cohabitation and social network. A study in the UK general population found higher odds of quitting success for people who were single, divorced or separated vs. were married or living with a partner²⁵. However, a study of patients in care for cardiac rehabilitation found higher odds of quitting success for people who were married vs. single³⁵. In the UK general population, studies reported finding no relationship between quitting success and marital status³⁰, cohabitation status³⁹, or number of household smokers^{24,25}. One study of people with severe and persistent mental illness reported higher odds of quitting success for people with more social support for quitting from family/friends³².

Health and healthcare setting. There were eight studies that investigated the association between quitting success and the smoker's health or the healthcare setting in which the quit attempt was instigated; five reported covariates that had statistically significant relationships to quitting success (Table S5 and Table S6 in Supplement 2 of the *Extended data*¹⁴): level of cardiovascular risk; number of comorbidities; having a mental health diagnosis; having a history of depression; having a history of substance abuse.

Physical health. One study in an outpatient setting reported higher odds of quitting success for patients with low (vs. moderate or high) cardiovascular risk and patients with fewer comorbidities³⁵. However, no relationship was found between quitting success and moderate (vs. high) cardiovascular risk³⁵. Another study found no relationship between quitting success and the number of comorbidities that a patient had³⁴. One study reported no relationship between the clinical setting in which the patient was located at the time that they were referred to stop smoking support (Cardiology services/clinics vs. Respiriology services/clinics vs. other hospital services/clinics)³⁴.

Mental health. Lower odds of quitting success were reported for people with: a primary diagnosis of anxiety disorder vs. no disorder³³; recurrent, current or recent depression vs. no history of depression⁴¹; history of opiate abuse vs. history of alcohol abuse³³; history of alcohol abuse, opiate abuse and marijuana abuse vs. no history of substance abuse⁴². No relationship with quitting success was reported in three studies that investigated primary mental health diagnosis^{32,36,42}, two studies of PHQ-9 score^{32,43}, one study of having a history of substance abuse³², one study of HADS anxiety score and HADS depression score³⁵, and one study of history of psychiatric disorder and history of co-occurring substance use and psychiatric disorder³⁴.

Tobacco smoking variables. There were 17 studies in this category; 14 reported factors significantly related to quitting success (Table S5 and Table S6 in Supplement 2 of the *Extended data*¹⁴): daily cigarette consumption, carbon monoxide (CO) level at baseline, level of nicotine dependence, the most difficult situation not to smoke, determination / motivation to quit, and the history of previous attempts to quit smoking.

Current and previous cigarette consumption. Higher odds of pregnant women quitting smoking successfully were reported among women with lower pre-pregnancy cigarette consumption³⁹. No relationship between quitting success and the daily cigarette consumption prior to quitting was identified in one study in the UK general population²⁹, two studies of people with a mental health condition^{32,33} and one study in a secondary care setting³⁴. No relationship between quitting success and the age at which someone started to smoke regularly (age at smoking initiation) was reported by one study in the UK general population³⁰, two studies in people with a mental health condition^{32,33}, and one study in a secondary care setting³⁴.

Carbon monoxide (CO) level. The single study to find a relationship between quitting success and CO level prior to quitting was of a tailored smoking cessation programme for individuals with substance use disorders and mental illness; lower CO levels when the quit attempt began had higher odds of quitting success⁴². No relationship between quitting success and CO level was found by one study in people with a mental health condition³³, and one study in a secondary care setting³⁴.

Level of nicotine dependence. The 11 studies which identified statistically significant associations between quitting success and nicotine dependence prior to the quit attempt found mixed results: higher odds of quitting in smokers with lower nicotine dependence was found by nine studies in the UK general population^{22,25–30,37,44} and two studies of smoking cessation delivered in an outpatient setting^{32,33}. No relationship between quitting success and nicotine dependence was found by one study in the UK general population²⁴, two studies in people with a mental health condition^{36,42}, and one study in a secondary care setting³⁴. One study in the UK general population found higher odds of quitting success in smokers whose most difficult situation not to smoke was when feeling the urge to smoke, but the same study found no relationship with quitting success for when socialising, first thing in the morning, when angry or frustrated, when relaxing, and for ‘any other reason’³⁰. One study found no relationship between quitting success and the reported enjoyment of smoking²⁸.

Motivation to quit. Two studies in the UK general population found higher odds of quitting successfully for smokers who reported a determination to quit²⁴ or being motivated to quit³⁷. No relationships between quitting success and reported readiness to quit were found in one study in the UK general population³⁰, one study in people with a mental health condition³², and one study in a secondary care setting³⁴. One UK general population study found no relationship between quitting success and the reported reasons for quitting, main advantage of quitting, or main disadvantage of quitting³⁰.

Quitting characteristics. In terms of previous quit attempts, three studies in the UK general population^{27,29,30} and one study in a mental health setting³³ found higher odds of quitting successfully among smokers who had made more previous quit attempts or had previously been abstinent for longer periods. Specifically, higher odds of quitting successfully were found among those who had previously quit smoking for 3 months or more³⁰, made ≥ 2 quit attempts in the past 6 months²⁹, and had a longer duration of abstinence at the last attempt to quit^{27,33}. Three studies in the UK general population reported no relationship between quitting success and the number or duration of previous quit attempts^{25,29,45}, as did one study in people with a mental health condition³², and one study in an outpatient setting³⁴. One study in a UK general population reported no relationship between success in the current quit attempt and the time since the start of the last unsuccessful quit attempt²⁹.

Intervention characteristics. There were 21 studies that investigated the influence on quitting success of characteristics of the attempt to quit smoking; 17 studies reported factors significantly related to the success of quit attempts (Table S5 and Table S6 in Supplement 2 of the *Extended data*¹⁴). Factors related to the behaviour and choices of the individual smokers were whether smokers reduced or temporarily abstained from smoking before making a quit attempt, and various descriptors of the nature of support for the quit attempt. Pharmacological characteristics of the quit attempt were the type of pharmacological aid use, whether this was used alongside behavioural support, and the degree of compliance of the smoker making the quit attempt with the recommended guidelines for use of the pharmacotherapy chosen.

Reduction in amount smoked and/or temporary abstinence before quitting. Two studies found higher odds of quitting successfully for smokers who reduced the amount they smoked before attempting to quit smoking^{29,46}, including if this was with the support of pharmacotherapy⁴⁶. One study found no relationship between quitting success and whether the quit attempt was spontaneous, i.e. initiated as soon as the decision to quit has been made (compared with not making a spontaneous quit attempt)²⁹, and one study found no relationship between quitting success and whether the smoker reduced the amount smoked prior to quitting (compared with quitting without first reducing the amount smoked)²⁷.

Behavioural support type, setting and mode of contact. For the UK general population, higher odds of quitting were found for smokers who used a smoking cessation clinic and websites (compared with no support)^{40,47}, for smokers who used pharmacotherapy alongside help from a health professional or specialist smoking cessation advisor (compared with no support)⁴⁷, and for smokers who received support in specialist clinics^{22,45}, in the community (compared with other settings)^{25,26}, and with group support (compared with one-to-one or other support)^{22,23}. Lower odds of quitting were reported for smokers who used drop-in support (compared with one-to-one support)⁴⁵, and telephone support (compared with no support)⁴⁰. Other studies found no relationships between quitting success and the receipt of in-person behavioural support⁴⁰, the use of self-help materials⁴⁰, having one-to-one support⁴⁸, the setting of support for smoking cessation^{22,23,26,45}, having group therapy, or receiving support from a doctor or other health professional⁴⁷.

Tobacco dependence treatment duration and number of contacts. Higher odds of quitting success were associated with the number of contacts that a smoker had with a stop smoking advisor in the UK general population²⁴, and in studies of people with a mental health condition^{33,36,42}. Other studies found no relationship between quitting success and treatment duration or number of contacts^{22,32,34}.

Pharmacological aids. In the UK general population, higher odds of quitting success were found for smokers who used NRT (compared with no NRT/no cessation aids)^{22,40,45}, combination

NRT (compared with single NRT)³¹, varenicline (compared with no varenicline, no medication, or NRT)^{22,26,40,45}, bupropion (compared with no medication and NRT)^{22,25}, and for the use of any pharmacotherapy in general^{47,49}. There were also higher odds of quitting success with the use of e-cigarettes (compared with no e-cigarettes, no cessation aid, and NRT)^{37,40,48}. There was also evidence in the UK general population of higher odds of quitting successfully when smokers have greater compliance with the recommended guidelines for pharmacotherapy use²⁴. One study in the UK general population found lower odds of quitting successfully for smokers who bought NRT over the counter (compared with no cessation aids)⁴⁹. Other studies in the UK general population found no relationships between quitting success and the use of prescription NRT⁴⁰, NRT bought over the counter⁴⁰, bupropion^{40,45}, or e-cigarette use³⁷. For people with a mental health condition, no relationship with quitting success was found for the use of pharmacotherapy^{32,33,36}, or the number of weeks of NRT, varenicline and bupropion use³³.

Discussion

The review has identified a list of covariates worth considering in plans for the statistical analysis of quitting success following a smoking cessation intervention initiated in a secondary care setting in the UK. The findings support and supplement the previous reviews that have investigated covariates of quitting success, and add to the evaluation framework for hospital based smoking cessation services in the UK⁶ by highlighting the data fields important to consider in “deep dives” into service data to investigate the reasons for variation in quitting outcomes.

Strength and limitations

The strengths of this review lie in the rapid but systematic review approach taken^{11,12} and in the design of the research question and population restrictions to be specific to smoking cessation interventions initiated in a secondary care setting in the UK. The limitations lie in the compromises made as part of the review approach, for example, our focus only on studies published in English, not searching grey literature, limited critical appraisal of the studies found. The review only included studies from the UK and Canada, which was intended to limit the influence of variation in service delivery internationally, while noting our interest was specific to the UK. Whilst this restriction increased relevance, only two studies were identified from a secondary healthcare setting. It is possible that expanding the search worldwide would have identified more covariates specific to understanding the influence of health and the healthcare setting on quitting success. However, healthcare systems differ widely worldwide, and our decisions to limit the scope of this review are in line with recommended best practice for rapid reviews^{11,12}.

Informing real-world data collection: supporting clinical care and public health policy

Improvement of smoking cessation interventions embedded into NHS secondary care services requires the use of real-world data for service monitoring and ongoing evaluation. There will

be incremental improvement in services over time, including attempts to address factors observed to influence the success of quit attempts. This review provides a starting point for understanding what data fields might be important to collect to ensure that sufficient information is available to guide activities aimed at service improvement. The NICE real-world evidence framework⁵⁰ encourages service evaluators to identify the data fields needed through a systematic, transparent and reproducible search. The current review of the covariates of quitting success is part of that systematic approach and could aid the planning of data fields to be collected.

Evidence-based care: trial-based and real-world evidence

When conducting an evaluation of intervention efficacy or comparative effectiveness, be it based on a randomised or non-randomised study design (noting service evaluations are not permitted to randomise patients to treatment assignment), developing a statistical analysis plan is an important step towards reducing potential bias in the evidence base⁵⁰. Service evaluations and associated real-world evidence are often dependent on the real-world data available, hence the importance of considering which covariates to collect data on. For a statistical analysis plan, the interest is usually in adjusting estimates of service outcomes for the influence of confounding variables, but investigations can become more complex by situating covariates within a causal framework for evaluating service outcomes, for example using directed acyclic graphs⁵⁰. The list of covariates identified in the current review could aid the development of a range of plans for statistical analysis to inform the evidence-base, focussed either on association or causality depending on the intention of the analysis and required evidence-base.

Understanding service complexity: informing adaptive logic models

There is increasing recognition in real world implementation and evaluation of healthcare interventions of the complexity of even seemingly “simple” treatments. Healthcare has been described as a complex adaptive system which requires understanding of multiple elements and the way in which they interact, in order to lead to transformation⁵¹. In common with many evaluations, evaluations of tobacco dependence treatment services in the UK draw on a theory of change approach in order to aid understanding of implementation and the effects of the tobacco dependence treatment service on outcomes for smoking and health⁵². The data fields identified during this review help to inform the development of service logic models⁵³, which act as a visual summary of the complexity by which the intervention produces outcomes. These models can help to build our conceptualization and understanding of hypothesized causal links underpinning quitting smoking⁵⁴.

Conclusion

In total, 14 broad categories of covariate were identified as having a statistically significant association with the success in quitting smoking and therefore worth considering in plans for

the statistical analysis of quit success following contact with a smoking cessation intervention initiated within secondary healthcare services in the UK. These covariates also indicate the data fields it might be important to collect as part of the ongoing monitoring and evaluation of such services.

Data availability

Underlying data

All data underlying the results are available as part of the article and no additional source data are required.

Extended data

Open Science Framework: Supplementary information for “Covariates of success in quitting smoking: a systematic review of studies from 2008 to 2021 conducted to inform the statistical analyses of quitting outcomes of a hospital-based tobacco dependence treatment service in the United Kingdom. <https://doi.org/10.17605/OSF.IO/UW8DZ14>.

This project contains the following extended data:

- Supplement 1 – search strategies in full
- Supplement 2 – results tables
 - Supplementary Table S1: Studies excluded at full text screening
 - Supplementary Table S2. Characteristics of included studies.
 - Supplementary Table S3: Participant baseline characteristics of included studies

- Supplementary Table S4: Outcome measurement and analyses in included studies
- Supplementary Table S5. Relationships between covariates and quit success.
- Supplementary Table S6: Covariates of quitting outcomes – full results summary

Reporting guidelines

Open Science Framework: Completed PRISMA checklist for ‘Covariates of success in quitting smoking: a systematic review of studies from 2008 to 2021 conducted to inform the statistical analyses of quitting outcomes of a hospital-based tobacco dependence treatment service in the United Kingdom’. <https://doi.org/10.17605/OSF.IO/UW8DZ14>.

Data are available under the terms of the Creative Commons Attribution 4.0 International license (CC-BY 4.0).

Acknowledgements

The review was initiated as part of a Yorkshire Cancer Research commissioned service evaluation of the QUIT hospital-based tobacco dependence treatment service (<https://sybics-quit.co.uk/>). The authors thank Debbie Robson for support in developing the review and John Holmes for comments to improve the manuscript. An earlier version of this article can be found on medRxiv (doi: <https://doi.org/10.1101/2023.01.10.23284384>).

For the purpose of open access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission.

References

1. Tobacco Advisory Group of the Royal College of Physicians: **Hiding in plain sight: Treating tobacco dependency in the NHS**. 2018. [Reference Source](#)
2. Agrawal S, Mangera Z, Murray RL, et al.: **Successes and Challenges of Implementing Tobacco Dependency Treatment in Health Care Institutions in England**. *Curr Oncol*. 2022; 29(5): 3738–3747. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. UK National Health Service (NHS): **The NHS long term plan**. 2019. [Reference Source](#)
4. Reid RD, Mullen KA, Slovinec D'Angelo ME, et al.: **Smoking cessation for hospitalized smokers: an evaluation of the “Ottawa Model”**. *Nicotine Tob Res*. 2010; 12(1): 11–8. [PubMed Abstract](#) | [Publisher Full Text](#)
5. Evison M, Agrawal S, Conroy M, et al.: **Building the case for comprehensive hospital-based tobacco addiction services: Applying the Ottawa Model to the City of Manchester**. *Lung Cancer*. 2018; 121: 99–100. [PubMed Abstract](#) | [Publisher Full Text](#)
6. Robson D, Richardson S, Howle F, et al.: **Developing and testing a standardised evaluation framework for hospital-initiated tobacco dependence treatment services**. 2020. [Reference Source](#)
7. Vangeli E, Stapleton J, Smit ES, et al.: **Predictors of attempts to stop smoking and their success in adult general population samples: a systematic review**. *Addiction*. 2011; 106(12): 2110–21. [PubMed Abstract](#) | [Publisher Full Text](#)
8. Agrawal S, Mangera Z: **British Thoracic Society smoking cessation audit report: Smoking cessation policy and practice in NHS hospitals: National audit period: 1 April–31 May 2016**. 2016. [Reference Source](#)
9. Devani N, Evison M: **National Smoking Cessation Audit 2021: Management of Tobacco Dependency in Acute Care Trusts: Audit Report. National Audit Period: 1 July - 31 August 2021**. 2021. [Reference Source](#)
10. Mangera Z, Devani N: **National Smoking Cessation Audit 2019. National Audit Period: 1 July - 30 August 2019**. 2020. [Reference Source](#)
11. Tricco AC, Antony J, Zarin W, et al.: **A scoping review of rapid review methods**. *BMC Med*. 2015; 13(1): 224. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Tricco AC, Zarin W, Antony J, et al.: **An international survey and modified Delphi approach revealed numerous rapid review methods**. *J Clin Epidemiol*. 2016; 70: 61–67. [PubMed Abstract](#) | [Publisher Full Text](#)
13. Abrami PC, Borokhovski E, Bernard RM, et al.: **Issues in conducting and disseminating brief reviews of evidence**. *Evidence & Policy: A Journal of Research, Debate and Practice*. 2010; 6(3): 371–389. [Publisher Full Text](#)
14. Hock E, Franklin M, Baxter S, et al.: **Supplementary information for “Covariates of success in quitting smoking: a systematic review of studies from 2008 to 2021 conducted to inform the statistical analyses of quitting**

- outcomes of a hospital-based tobacco dependence treatment service in the United Kingdom". [Dataset], 2023. <http://www.doi.org/10.17605/OSF.IO/UW8DZ>
15. Hefner V: **The SAGE Encyclopedia of Communication Research Methods**. Thousand Oaks, California: SAGE Publications, Inc, 2017.
 16. Szumilas M: **Explaining odds ratios**. *J Can Acad Child Adolesc Psychiatry*. 2010; 19(3): 227–229. [PubMed Abstract](#) | [Free Full Text](#)
 17. Le Grande M, Borland R, Yong HH, *et al.*: **Predictive Power of Dependence Measures for Quitting Smoking. Findings From the 2016 to 2018 ITC Four Country Smoking and Vaping Surveys**. *Nicotine Tob Res*. 2021; 23(2): 276–285. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 18. Lubitz SF, Flitter A, Wileyto EP, *et al.*: **History and Correlates of Smoking Cessation Behaviors Among Smokers With Serious Mental Illness**. *Nicotine Tob Res*. 2020; 22(9): 1492–1499. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 19. Ussher M, Kakar G, Hajek P, *et al.*: **Dependence and motivation to stop smoking as predictors of success of a quit attempt among smokers seeking help to quit**. *Addict Behav*. 2016; 53: 175–80. [PubMed Abstract](#) | [Publisher Full Text](#)
 20. Smit ES, Hoving C, Schelleman-Offermans K, *et al.*: **Predictors of successful and unsuccessful quit attempts among smokers motivated to quit**. *Addict Behav*. 2014; 39(9): 1318–1324. [PubMed Abstract](#) | [Publisher Full Text](#)
 21. Zhou X, Nonnemaker J, Sherrill B, *et al.*: **Attempts to quit smoking and relapse: Factors associated with success or failure from the ATTEMPT cohort study**. *Addict Behav*. 2009; 34(4): 365–73. [PubMed Abstract](#) | [Publisher Full Text](#)
 22. Brose LS, McEwen A, West R: **Does it matter who you see to help you stop smoking? Short-term quit rates across specialist stop smoking practitioners in England**. *Addiction*. 2012; 107(11): 2029–36. [PubMed Abstract](#) | [Publisher Full Text](#)
 23. Brose LS, West R, Stapleton JA: **Comparison of the effectiveness of varenicline and combination nicotine replacement therapy for smoking cessation in clinical practice**. *Mayo Clin Proc*. 2013; 88(3): 226–33. [PubMed Abstract](#) | [Publisher Full Text](#)
 24. Hiscock R, Judge K, Bauld L: **Social inequalities in quitting smoking: what factors mediate the relationship between socioeconomic position and smoking cessation?** *J Public Health (Oxf)*. 2011; 33(1): 39–47. [PubMed Abstract](#) | [Publisher Full Text](#)
 25. McEwen A, West R: **Do implementation issues influence the effectiveness of medications? The case of nicotine replacement therapy and bupropion in UK Stop Smoking Services**. *BMC Public Health*. 2009; 9(1): 28. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 26. Walker N, Gainforth H, Kiparoglou V, *et al.*: **Factors moderating the relative effectiveness of varenicline and nicotine replacement therapy in clients using smoking cessation services**. *Addiction*. 2018; 113(2): 313–324. [PubMed Abstract](#) | [Publisher Full Text](#)
 27. Beenstock J, Lindson-Hawley N, Aveyard P, *et al.*: **Future orientation and smoking cessation: secondary analysis of data from a smoking cessation trial**. *Addiction*. 2014; 109(10): 1732–40. [PubMed Abstract](#) | [Publisher Full Text](#)
 28. Fidler JA, West R: **Enjoyment of smoking and urges to smoke as predictors of attempts and success of attempts to stop smoking: a longitudinal study**. *Drug Alcohol Depend*. 2011; 115(1–2): 30–4. [PubMed Abstract](#) | [Publisher Full Text](#)
 29. Garnett C, Shahab L, Raupach T, *et al.*: **Understanding the Association Between Spontaneous Quit Attempts and Improved Smoking Cessation Success Rates: A Population Survey in England With 6-Month Follow-up**. *Nicotine Tob Res*. 2020; 22(9): 1460–1467. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 30. Kale D, Gilbert HM, Sutton S: **Are predictors of making a quit attempt the same as predictors of 3-month abstinence from smoking? Findings from a sample of smokers recruited for a study of computer-tailored smoking cessation advice in primary care**. *Addiction*. 2015; 110(10): 1653–64. [PubMed Abstract](#) | [Publisher Full Text](#)
 31. Kassim S, Al-Haboubi M, Croucher R: **Short-Term Smoking Cessation in English Resident Adults of Bangladeshi Origin: A Service Review**. *Nicotine Tob Res*. 2016; 18(4): 410–5. [PubMed Abstract](#) | [Publisher Full Text](#)
 32. Masuhara JE, Heah T, Okoli CTC: **Outcomes of a tobacco treatment programme for individuals with severe and persistent mental illness attending a community mental health team**. *J Smok Cessat*. 2014; 9(2): 60–67. [Publisher Full Text](#)
 33. Okoli CTC, Khara M: **Smoking cessation outcomes and predictors among individuals with co-occurring substance use and/or psychiatric disorders**. *J Dual Diagn*. 2014; 10(1): 9–18. [PubMed Abstract](#) | [Publisher Full Text](#)
 34. Khara M, Okoli CTC: **A retrospective review of pilot outcomes from an outpatient tobacco treatment programme within cardiology services**. *J Smok Cessat*. 2015; 10(1): 74–84. [Publisher Full Text](#)
 35. Salman A, Doherty P: **Predictors of Quitting Smoking in Cardiac Rehabilitation**. *J Clin Med*. 2020; 9(8): 2612. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Selby P, Voci SC, Zawertailo LA, *et al.*: **Individualized smoking cessation treatment in an outpatient setting: Predictors of outcome in a sample with psychiatric and addictions co-morbidity**. *Addict Behav*. 2010; 35(9): 811–7. [PubMed Abstract](#) | [Publisher Full Text](#)
 37. Hitchman SC, Brose LS, Brown J, *et al.*: **Associations Between E-Cigarette Type, Frequency of Use, and Quitting Smoking: Findings From a Longitudinal Online Panel Survey in Great Britain**. *Nicotine Tob Res*. 2015; 17(10): 1187–94. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 38. Kotz D, West R: **Explaining the social gradient in smoking cessation: it's not in the trying, but in the succeeding**. *Tob Control*. 2009; 18(1): 43–6. [PubMed Abstract](#) | [Publisher Full Text](#)
 39. Graham H, Hawkins SS, Law C: **Lifecourse influences on women's smoking before, during and after pregnancy**. *Soc Sci Med*. 2010; 70(4): 582–7. [PubMed Abstract](#) | [Publisher Full Text](#)
 40. Jackson SE, Kotz D, West R, *et al.*: **Moderators of real-world effectiveness of smoking cessation aids: a population study**. *Addiction*. 2019; 114(9): 1627–1638. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 41. Zawertailo L, Voci S, Selby P: **Depression status as a predictor of quit success in a real-world effectiveness study of nicotine replacement therapy**. *Psychiatry Res*. 2015; 226(1): 120–7. [PubMed Abstract](#) | [Publisher Full Text](#)
 42. Okoli CTC, Khara M, Torchalla I, *et al.*: **Sex differences in smoking cessation outcomes of a tailored program for individuals with substance use disorders and mental illness**. *Addict Behav*. 2011; 36(5): 523–6. [PubMed Abstract](#) | [Publisher Full Text](#)
 43. Zawertailo LA, Baliunas D, Ivanova A, *et al.*: **Individualized Treatment for Tobacco Dependence in Addictions Treatment Settings: The Role of Current Depressive Symptoms on Outcomes at 3 and 6 Months**. *Nicotine Tob Res*. 2015; 17(8): 937–45. [PubMed Abstract](#) | [Publisher Full Text](#)
 44. Taggar JS, Lewis S, Docherty G, *et al.*: **Do cravings predict smoking cessation in smokers calling a national quit line: secondary analyses from a randomised trial for the utility of 'urges to smoke' measures**. *Subst Abuse Treat Prev Policy*. 2015; 10: 15. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 45. Brose LS, West R, McDermott MS, *et al.*: **What makes for an effective stop-smoking service?** *Thorax*. 2011; 66(10): 924–6. [PubMed Abstract](#) | [Publisher Full Text](#)
 46. Beard E, McNeill A, Aveyard P, *et al.*: **Association between use of nicotine replacement therapy for harm reduction and smoking cessation: a prospective study of English smokers**. *Tob Control*. 2013; 22(2): 118–22. [PubMed Abstract](#) | [Publisher Full Text](#)
 47. Gibson JE, Murray RL, Borland R, *et al.*: **The impact of the United Kingdom's national smoking cessation strategy on quit attempts and use of cessation services: findings from the International Tobacco Control Four Country Survey**. *Nicotine Tob Res*. 2010; 12 Suppl(Suppl 1): S64–71. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 48. Brown J, Beard E, Kotz D, *et al.*: **Real-world effectiveness of e-cigarettes when used to aid smoking cessation: a cross-sectional population study**. *Addiction*. 2014; 109(9): 1531–40. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 49. Kotz D, Brown J, West R: **Prospective cohort study of the effectiveness of smoking cessation treatments used in the "real world"**. *Mayo Clin Proc*. 2014; 89(10): 1360–7. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 50. National Institute for Health and Care Excellence (NICE): **NICE real-world evidence framework. Corporate document [ECD9]**. 2022. [Reference Source](#)
 51. Khan S, Vandermerriis A, Shepherd J, *et al.*: **Embracing uncertainty, managing complexity: applying complexity thinking principles to transformation efforts in healthcare systems**. *BMC Health Serv Res*. 2018; 18(1): 192. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. Mills T, Lawton R, Sheard L: **Advancing complexity science in healthcare research: the logic of logic models**. *BMC Med Res Methodol*. 2019; 19(1): 55. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 53. Rehfuess EA, Booth A, Brereton L, *et al.*: **Towards a taxonomy of logic models in systematic reviews and health technology assessments: a priori, staged, and iterative approaches**. *Res Synth Methods*. 2018; 9(1): 13–24. [PubMed Abstract](#) | [Publisher Full Text](#)
 54. Anderson LM, Petticrew M, Rehfuess E, *et al.*: **Using logic models to capture complexity in systematic reviews**. *Res Synth Methods*. 2011; 2(1): 33–42. [PubMed Abstract](#) | [Publisher Full Text](#)